What Dose of Methamphetamine Do Regular Consumers Use Daily? Estimating Milligrams of Oral Amphetamine Equivalent

Joseph R. Friedman MD, PhD, MPH[1*]; Adam J. Koncsol[2]; Caitlin A. Molina[2]; Ruby Romero[2]; Jasmine Feng[2]; Michelle Poimboeuf[2]; Morgan E. Godvin[2]; Siddarth Puri, MD[3]; Carla Marienfeld MD[1]; Chelsea L. Shover, PhD[2]

[1] *Department of Psychiatry, University of California, San Diego*
[2] *Department of General Internal Medicine and Health Services Research, University of California, Los Angeles*
[3] *Medical Director, County of San Luis Obispo, California*

*Correspondence to:
Joseph Friedman, MD PhD, MPH, University of California, San Diego, 200 W Arbor Dr, San Diego, CA, 92103. jofriedman@health.ucsd.edu

Word Count: 1,071

Abstract

Introduction

The purity, accessibility, and affordability of illicit methamphetamine has increased in recent decades, which has been linked to rising rates of methamphetamine-involved overdoses, psychosis, cardiovascular events, and other health consequences. Nevertheless, information about the quantity of methamphetamine used by regular consumers has been limited, despite the potential clinical utility of exposure quantification.

Methods

From August 2024-April 2025, self-reported daily methamphetamine consumption was assessed among n=68 individuals. Methamphetamine samples (n=112) were analyzed for purity using liquid chromatography-mass spectrometry. Percent bioavailability by route of administration and stimulant equivalency were obtained from literature. A simulation model leveraging bootstrapping was used to estimate MOAE.

Results

The average reported daily methamphetamine consumption was 0.96g (median 0.36g; range 0.1g-4.0g). Average purity was 71.6% (median 75.5%; range 0.1%-95.0%). Given estimated average bioavailability of 52.0% when smoked, 79.3% when insufflated, 67.2% orally or inserted rectally, and a 2:1 amphetamine-methamphetamine equivalency, the average consumer used 1,549.0 MOAE daily (median 516.6; range 1.3-10,112.0).

Discussion

We estimate that consumers of methamphetamine in Los Angeles use an average daily stimulant dose (>1,500 MOAE) that is 25-fold higher than the maximum typical recommended clinical dose of mixed amphetamine salts (60mg). This may help explain the limited efficacy of prescription stimulant treatment for methamphetamine use disorder, which typically employs considerably lower quantities. Given this high dose, these findings shed light on the rising incidence of methamphetamine-related sequalae, such as psychosis, cardiovascular complications, and sudden death. Although exposure quantification is commonplace for alcohol and tobacco use disorders, uncertainties in illicit drug markets has complicated this practice for most illicit drugs. This study supports leveraging emerging information from drug checking programs so that clinicians can approximate exposure quantification.

Introduction

The purity, accessibility, and affordability of illicit methamphetamine has increased in recent decades[1]. This has been linked to rising rates of methamphetamine-involved overdoses, psychosis, psychiatric hospitalizations, cardiovascular events, and other health consequences[2–4]. Nevertheless, information about the actual quantity of methamphetamine used in practice, among people with methamphetamine use disorder, has been limited. This is largely due to the historical difficulty in assessing the purity of illicit substances used by patients. Nevertheless, there is considerable potential clinical utility of exposure quantification, such as understanding how a patient's methamphetamine habit compares to potential replacement doses of prescription stimulants, or assessing the health impacts of dose reductions in illicit stimulant use.

Using recent advancements in drug checking technologies and services, we quantify the purity of methamphetamine samples used by regular consumers in Los Angeles. We combine this with self-reported daily consumption quantities among the same population to estimate the daily Milligrams of Oral Amphetamine Equivalent (MOAE). This quantity is likely to be more easily interpretable by psychiatrists and other clinicians, who are familiar with dosing of mixed amphetamine salts, rather than the quantity of illicit methamphetamine, for which dosing depends on drug concentration, route of administration, and bioavailability.

Methods

Daily MOAE was estimated according to Equation 1, where *Consumption* is the daily milligrams of illicit methamphetamine used, *Purity* is the proportion of active compound in substances sold as illicit "methamphetamine", *Bioavailability* is the route-of-administration-specific proportion of total drug used that is absorbed systemically, $Methamphetamine{:}Amphetamine\ Equivalence$ is the ratio of physiological potency between methamphetamine and mixed amphetamine salts, and $Bioavailability\ Oral\ Amphetamine\ Salts$ is the fraction of mixed amphetamine salts that is absorbed systemically when taken orally.

$$Milligrams\ of\ Oral\ Amphetamine\ Equivalence\ (MOAE) = Consumption \cdot Purity \cdot Bioavailability \cdot$$
$$Methamphetamine{:}Amphetamine\ Equivalence \cdot \frac{1}{Bioavailability\ Oral\ Amphetamine\ Salts}$$

Equation 1. Calculation of estimated mg of oral amphetamine equivalent (MOAE) self-administered among regular consumers in Los Angeles

To estimate the distribution of MOAE combining uncertainty and variation in each of the aforementioned parameter inputs, we employed a bootstrapping approach leveraging 10,000 draws. This allows for the inclusion of uncertainty in each parameter, which is propagated through to the final estimated MOAE.

Methamphetamine purity by weight was assessed using liquid chromatography-mass spectrometry (LC-MS) among n=112 samples collected at a community-based drug checking program in Los Angeles County[5]. Samples were limited to those sold as methamphetamine, and for which clients expected only methamphetamine. Daily methamphetamine consumption

volume and route of administration was assessed by self-report using a survey among n=68

individuals who regularly consume methamphetamine and brought samples for testing.

Bioavailability by route-of-administration and stimulant equivalency factors were drawn from

literature review (see supplemental methods). The UCLA IRB approved this project (IRB-22-

0760) and additionally determined that aspects of this work constituted public health

surveillance and not human subjects research.

Results

The average reported daily methamphetamine consumption was 0.96g (Figure 1). A skewed

distribution was observed with a median of 0.36g. The range was 0.1g-4.0g daily. Estimated

mean and median purity was 71.6% and 75.5%, respectively, with a range from 0.1% to 95.0%.

97   Bioavailability was estimated from literature sources describing experimental data using

human subjects (see supplemental methods for a full description). Average bioavailability of

smoked methamphetamine was estimated at 52.0% (of the total "pipe dose"), reflecting a

uniform distribution with a minimum of 37% and a maximum of 67%. Bioavailability of

insufflated "snorted" methamphetamine was modeled using a normal distribution with a mean

of 79.0% and a standard deviation of 6.7%. Oral bioavailability was estimated using a normal

distribution with a mean of 67.2% and a standard deviation of 1.59%. Rectal bioavailability

was estimated using a normal distribution with a mean of 67.2% and a standard deviation of

20.0%. The amphetamine-methamphetamine equivalency was estimated to be 1:2.

106   Given the above factors, the estimated mean and median MOAE used daily was 1,549.0

107   and 516.6, respectively. The estimated range was 1.3 MOAE to 10,112.0 MOAE daily.

Discussion

We estimate that consumers of methamphetamine in Los Angeles use an average daily stimulant dose (>1,500 MOAE) that is 25-fold higher than the maximum typical recommended clinical dose of mixed amphetamine salts (60mg). This may help explain the limited efficacy of prescription stimulant treatment for methamphetamine use disorder, which typically employs considerably lower quantities[6]. For instance, the 60 to 110mg daily used in trials of amphetamine salts for methamphetamine use disorder would represent 1/14th to 1/25th of the average estimated methamphetamine usage observed in our population[7,8].

Given this high dose, these findings also shed light on the rising incidence of methamphetamine-related sequalae, such as psychosis, cardiovascular complications, and sudden death[1,3], which are much more common with illicit methamphetamine compared to the prescribed use of stimulant therapies[9].

Although exposure quantification is commonplace for alcohol and tobacco use disorders, uncertainties in illicit drug markets has complicated this practice for most illicit drugs. However, this study suggests that by leveraging emerging information from drug checking programs, as well as regularly asking their patients about consumption quantities, clinicians may be able to approximate exposure quantification. This may be useful in understanding the clinical benefits of treatments that result in dose reductions. This can also inform risk stratification for cardiac, psychiatric and other outcomes and personalizing care, including withdrawal management. This approach would also mirror efforts to quantify use and employ usage reduction as a clinical endpoint in addiction-related clinical trials[10].

This study represents an example of how emerging drug checking technologies can

provide consumers with information that can empower them to make safer drug use decisions,

despite the highly uncertain and rapidly-evolving nature of the illicit drug supply.

This work is limited by its use of data from a single city and may not be generalizable to

other contexts. Self-report consumption data is also prone to recall and desirability biases.

Although the LC-MS testing technology used here is generally considered the 'gold standard' for

this type of assessment, it can miss certain contaminants not explicitly tested for.

In sum, we provide the first quantification, to our knowledge, of the daily dose of

methamphetamine used by regular consumers. We defined a novel metric, MOAE, which can

assist psychiatrists and other clinicians to describe methamphetamine consumption in terms of

prescription stimulant equivalent. We find that the average regular consumer uses 25-fold the

recommended maximum dose of amphetamine salts, which may explain the myriad and severe

health harms observed among patients with long-term methamphetamine use. Further study

will be required to assess the degree to which these findings hold true in other populations.

Figure. Distributions, Median, and Mean Grams per Day Consumed, Percent Concentration (Purity), and Estimated mg of Oral Amphetamine Equivalent Self-Administered Among Regular Consumers in LA

For each panel, a dashed vertical line shows the distribution mean, and a dashed solid line shows the distribution mean. Left: The distribution of self-reported grams of methamphetamine ingested per day among regular consumers participating in drug checking services in Los Angeles. Center: Percent concentration (purity) of expected methamphetamine samples provided by clients at drug checking services in Los Angeles. Right: the estimated distribution of milligrams of oral amphetamine equivalent (MOAE) consumed among drug checking clients in Los Angeles.

References

1. Han B, Compton WM, Jones CM, Einstein EB, Volkow ND. Methamphetamine Use, Methamphetamine Use Disorder, and Associated Overdose Deaths Among US Adults. *JAMA Psychiatry*. 2021;78(12):1329. doi:10.1001/jamapsychiatry.2021.2588

2. Rognli EB, Heiberg IH, Jacobsen BK, Høye A, Bramness JG. Transition From Substance-Induced Psychosis to Schizophrenia Spectrum Disorder or Bipolar Disorder. *AJP*. 2023;180(6):437-444. doi:10.1176/appi.ajp.22010076

3. Calcaterra SL, Yamkovoy K, Swathi PA, et al. U.S. trends in methamphetamine-involved psychiatric hospitalizations in the United States, 2015-2019. *Drug Alcohol Depend*. 2024;262:111409. doi:10.1016/j.drugalcdep.2024.111409

4. Friedman J, Shover CL. Charting the fourth wave: Geographic, temporal, race/ethnicity and demographic trends in polysubstance fentanyl overdose deaths in the United States, 2010–2021. *Addiction*. 2023;118(12):2477-2485. doi:10.1111/add.16318

5. Koncsol AJ, Molina CA, Romero R, et al. Purity of consumer-level methamphetamine samples and methamphetamine-adulteration of other drugs: Los Angeles, 2023-2024. Published online March 20, 2025:2025.03.18.25323868. doi:10.1101/2025.03.18.25323868

6. Prescription psychostimulants for the treatment of stimulant use disorder: a systematic review and meta-analysis | Psychopharmacology. Accessed April 26, 2025. https://link.springer.com/article/10.1007/s00213-020-05563-3

7. Galloway GP, Buscemi R, Coyle JR, et al. A Randomized, Placebo-Controlled Trial of Sustained-Release Dextroamphetamine for Treatment of Methamphetamine Addiction. *Clin Pharmacol Ther*. 2011;89(2):276-282. doi:10.1038/clpt.2010.307

8. Longo M, Wickes W, Smout M, Harrison S, Cahill S, White JM. Randomized controlled trial of dexamphetamine maintenance for the treatment of methamphetamine dependence. *Addiction*. 2010;105(1):146-154. doi:10.1111/j.1360-0443.2009.02717.x

9. Moran LV, Skinner JP, Shinn AK, et al. Risk of Incident Psychosis and Mania With Prescription Amphetamines. *AJP*. 2024;181(10):901-909. doi:10.1176/appi.ajp.20230329

10. Amin-Esmaeili M, Farokhnia M, Susukida R, et al. Reduced drug use as an alternative valid outcome in individuals with stimulant use disorders: Findings from 13 multisite randomized clinical trials. *Addiction*. 2024;119(5):833-843. doi:10.1111/add.16409

Acknowledgements

The authors report no conflicts of interest. JRF received funding from the National Institute on Drug Abuse (DA049644) and the National institute of Mental Health (MH101072). AJK received educational support through the NIH/National Center for Advancing Translational Science (NCATS) UCLA CTSI (TL1TR001883). CLS received support from the National Institute on Drug Abuse (K01DA050771). This work was supported by the Centers for Disease Control and Prevention as part of Overdose Data to Action: LOCAL (CDC-RFA-CE-23-0003), and made possible through an equipment grant from the James B. Pendleton Charitable Trust to the UCLA AIDS Institute and UCLA Center for AIDS Research. The funders played no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication. CLS and JF had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Supplemental Methods

Purity

1. Data describing methamphetamine purity and daily quantity ingested among regular consumers were assessed via data generated from anonymous participants accessing a community-based drug checking program, *Drug Checking Los Angeles*. Participants voluntarily provide samples of illicit drug products for testing at four different sites in Los Angeles County, California from August 2024 to April 2025.
2. Samples were analyzed initially in the field with Fourier-transform infrared (FTIR) spectroscopy and immunoassay test strips. Samples were then sent to the National Institute of Standards and Technology (NIST) for secondary laboratory-based qualitative and quantitative testing using direct analysis in real time mass spectrometry (DART-MS) and liquid-chromatography mass spectrometry (LC-MS). Both FTIR and DART-MS assess samples against libraries of over 1,300 substances, including pharmaceutical and illicit drugs, adulterants, cutting and bulking agents, precursor chemicals, and other substances (e.g., adhesives, food products, etc.). The LC-MS quantification panel included twelve substances: fentanyl and fluorofentanyl, fentanyl precursor chemicals (4-ANPP, phenethyl 4-ANPP), heroin, methamphetamine, cocaine, α2-agonists (xylazine and medetomidine), and three common fentanyl adulterants (tetracaine, lidocaine, and Bis(2,2,6,6-tetramethyl-4-piperidyl) sebacate (BTMPS).

3. Of 1,644 total available samples, this study included n=112 that were sold as methamphetamine, expected to contain only methamphetamine, per client self-report, and had quantified results available based on LC-MS.

4. Methamphetamine purity was initially reported as the freebase form which does not account for the additional weight of the hydrochloride salt form. In order to convert this to methamphetamine HCL values (the form that all samples in our dataset were sold as, and the form that medication forms of stimulants typically consist of) concentration values were multiplied by 1.24 to reflect the ratio of the molecular weights between the HCL and freebase forms of methamphetamine.

5. As a result of measurement limitations, concentration at or above 95% was imputed to be 95%. Additionally, samples with methamphetamine concentration below the limit of quantitation were imputed to be 0.1%.

6. The 112 available methamphetamine purity values were resampled to create 10,000 draws representing the distribution of purity using the sample() function in R. A seed was set to ensure the reproducibility of results between code iterations.

Drug Quantity Consumed

1. Information on drug quantity consumed for methamphetamine (and other drugs) were collected through an anonymous, optional survey conducted by trained drug checking staff.

2. Of n=68 participants who regularly consume methamphetamine and brought methamphetamine for testing, n= 46 reported the quantity of methamphetamine they consume in grams and had the option to report the quantity per day, week, or month. Weekly values were converted to days by dividing by 7. Monthly values were converted to daily by dividing by 30. N=22 participants reported quantities in dollars spent on methamphetamine per day or week. Dollar values were converted to grams using a standardized price of $20 per gram, which was known to the drug checking team as a typical 'going rate' for a gram of methamphetamine. This may slightly underestimate the consumption of individuals who spent larger quantities per day and therefore achieve a lower price per gram.

3. The 68 available methamphetamine quantity values (alongside associated route of administration information) were resampled to create 10,000 draws representing the distribution of purity using the sample() function in R.

Bioavailability and Equivalence

1. Literature values were used to estimate credible intervals for all other model parameters, including bioavailability and equivalence factors.

2. Bioavailability of orally consumed methamphetamine was estimated using experimental data from: *Cook, C. E., Jeffcoat, A. R., Sadler, B. M., Hill, J. M., Voyksner, R. D., Pugh, D. E., White, W. R., & Perez-Reyes, M. (1992). Pharmacokinetics of oral methamphetamine and effects of repeated daily dosing in humans. Drug metabolism and disposition: the biological fate of chemicals, 20(6), 856–862* and *Cook CE, Jeffcoat AR, Hill JM, et al. Pharmacokinetics of methamphetamine self-administered to human subjects by smoking S-(+)-methamphetamine hydrochloride. Drug Metab Dispos. 1993;21(4):717-723*. The authors estimated the oral bioavailability of methamphetamine to be 67.2% +/- 3.1%. This quantity was used directly as no other studies were found. Draws were estimated using a normal distribution with a mean of 67.2 and a standard deviation of 1.59.

3. Bioavailability of smoked methamphetamine was estimated from three studies:
    a. *Cook CE, Jeffcoat AR, Hill JM, et al. Pharmacokinetics of methamphetamine self-administered to human subjects by smoking S-(+)-methamphetamine hydrochloride. Drug Metab Dispos. 1993;21(4):717-723*. The quantity of interest is the percent of the "total pipe dose" that is ultimately absorbed. This study estimated that a bioavailability of 66% (90% absorption of 73.1% of drug smoked, 26.9% left in pipe).
    b. *Harris DS, Boxenbaum H, Everhart ET, Sequeira G, Mendelson JE, Jones RT. The bioavailability of intranasal and smoked methamphetamine. Clin Pharmacol Ther. 2003;74(5):475-486.* doi:10.1016/j.clpt.2003.08.002. This study estimated a bioavailability of 37.4% (67% absorption of 55% of drug smoked, 45% left in pipe). Nevertheless, the authors used a method that was not designed to maximize the methamphetamine extracted from the pipe, as each participant took only 2 inhalations of each dose. The authors of this study estimate, in their discussion section, that true absorption of an effective pipe dose is likely to be 37% to 67% based on user smoking skill.
    c. *Perez-Reyes, M., White, R., McDonald, S., Hill, J., Jeffcoat, R., & Cook, C. E. (1990). Pharmacologic effects of methamphetamine vapor inhalation (smoking) in man. NIDA research monograph, 105, 575–577.* This study estimated the bioavailability of "vaporized" (i.e., smoked) methamphetamine to be approximately 50%. This was largely a result of analyzing subjective and

cardiovascular effects under experimental conditions between IV and
smoked methamphetamine.

Given the above, the bioavailability of smoked methamphetamine was
estimated using a uniform distribution with a minimum of 37% and a maximum
of 67%.

4. The bioavailability of insufflated ("snorted") methamphetamine was estimated
using the following study: *Harris DS, Boxenbaum H, Everhart ET, Sequeira G,*
Mendelson JE, Jones RT. The bioavailability of intranasal and smoked
methamphetamine. Clin Pharmacol Ther. 2003;74(5):475-486.
*doi:*10.1016/j.clpt.2003.08.002. This study estimated a bioavailability of 79% +/-
13.1%. Bioavailability was estimated using a normal distribution with a mean of 79.0
and a standard deviation of 6.72%.

5. No literature sources were found describing the bioavailability of "boofed" or
rectally administered methamphetamine. Given that per rectum absorption is often
similar or greater than oral absorption, the distribution was estimated using a
normal distribution with a mean of 67.2% (the same as for PO bioavailability), yet
with a significantly larger standard deviation of 20%, given the lack of data on this
topic.

6. The bioavailability of injected methamphetamine was, by definition, set at 100%.

7. Among the n=68 individuals providing route of administration information alongside
quantity of consumption information (resampled to create 10,000 draws)
bioavailability was calculated for each draw. If an individual reported multiple
routes of administration, then the average of the bioavailability coefficients was
used for that draw. In effect, this assumes that individuals evenly split their
consumption between the different methods they reported. This could
underestimate or overestimate their total absorption depending on their true route
of administration habits and the relative frequencies of each method of
consumption.

8. The methamphetamine to amphetamine potency ratio was assessed using
equivalency tables (https://www.adhdmedcalc.com/), which tend to assign a 2:1
ratio, and based on clinical trials data
(https://www.accessdata.fda.gov/cdrh_docs/reviews/K040133.pdf) . This value of
2.0 was used for calculations.

9. The bioavailability of oral mixed amphetamine salts was assessed using literature
sources. Although no specific experimental data were identified, review sources
generally reported a high-level of absorption, with values ranging from 75%-95%.

Bioavailability was therefore estimated using a uniform distribution ranging from
75% to 95%.

Calculations

323   1. Calculations were performed according to the equation listed in the main text for
324   the 10,000 draws of each parameter, derived as defined above. The distribution of
325   quantity consumed, purity, and MOAE was graphed directly across the draws, and
326   summary statistics were calculated.