IsomiR Utility in ALS Prognostication

Yahel Cohen[1,2]*, Ilan Sinai[1,2]*, Iddo Magen[1,2*], Yehuda Matan Danino[1,2], Joanne Wuu[3], Andrea Malaspina[4], Michael Benatar[3], Eran Hornstein [1,2]

[1]Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel,
[2]Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot, Israel
[3]Department of Neurology and ALS Center, University of Miami Miller School of Medicine, Miami, FL, USA
[4]UCL Queen Square Motor Neuron Disease Center, UCL Queen Square Institute of Neurology, University College London, Queen Square, London, UK

* Authors contributed equally

Correspondence:
Andrea Malaspina (a.malaspina@ucl.ac.uk) and Michael Benatar (mbenatar@med.miami.edu) are corresponding authors for the ABC and PGB cohorts, respectively. Eran Hornstein (eran.hornstein@weizmann.ac.il) is the corresponding author for all other facets of the work.

Abstract

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease characterized by progressive motor neuron loss. IsomiRs are microRNA isoforms that arise from alternative processing or editing events during miRNA biogenesis. While isomiRs may carry distinct biological and clinical relevance, their potential as cell-free biomarkers in neurodegeneration remains largely unexplored. Here, we investigated the prognostic utility of plasma isomiRs in ALS, using next-generation sequencing. We profiled cell-free isomiRs in 154 ALS patients from a British cohort and identified higher levels of one isomiR, let-7g-5p.t, to be associated with longer survival. This finding was independently validated in an international ALS cohort of 200 patients and with two orthogonal approaches. let-7g-5p.t prognostic utility was comparable to that of neurofilament light chain (NfL) or miR-181. These results establish isomiRs as a novel class of blood-based biomarkers in ALS with potential to refine prognostication in clinical trials for neurodegenerative diseases.

Introduction

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease of the human motor neuron system. Heterogeneity of patient progression rate[1] challenges the detection of therapeutic effects in clinical trials[2]. Biofluid-based biomarkers have garnered increasing attention as tools to monitor progression[3,4]. These include urinary p75[5], troponin T2[6] and miRNAs[7]. Additionally, neurofilament light chain protein (NfL)[8] is a strong predictor of the future rate of disease progression, conveying meaningful prognostic information. Accordingly, we have recently demonstrated the utility of NfL in reducing the sample size required for clinical trials in a trial-like population[9].

microRNAs (miRNAs) are small endogenous RNAs that contribute to post-transcriptional silencing of mRNA[10]. miRNAs are critical for motor neuron viability[11] and their biogenesis is known to be dysregulated in ALS[12-16]. Previous studies have shown that cell-free miRNAs and in particular miR-181 might have prognostic value[7,17]. miRNA biogenesis is a multistep process that starts with the transcription of a primary miRNA, which is cleaved by the nuclear microprocessor complex to yield an intermediate double-stranded hairpin precursor (pre-miRNA)[18]. After export of the pre-miRNA from nucleus to cytoplasm, it is cut by Dicer, creating a single strand 22nt guide that is loaded on to the RNA-induced silencing complex (RISC)[19]. IsomiRs display sequence variations that distinguish them from the reference sequence miRNAs (reference miRNA), including 3' or 5' sequence changes, or sequence polymorphism[20-23]. IsomiR production is often the result of aberrant miRNA biogenesis and isomiR levels in tissues or biofluids are often low and heterogeneous[24].

Mutations in the genes encoding for a handful of RNA binding proteins were found to be linked to increased risk of developing ALS, including Fused in Sarcoma (FUS)[25,26], and TAR DNA-binding protein 43 (TDP-43)[27]. Cytoplasmic mislocalization of TDP-43 into inclusions and loss of its nuclear function, are reported in approximately 97% of patients[28]. Accordingly, cryptic exon splicing impacts dozens of mRNAs[29-31], some of which may have value as novel biomarkers relevant to ALS therapy development[32,33]. Interestingly, loss of TDP-43 or FUS also leads to dysregulation in isomiR profiles or miRNA biogenesis[34,35]. Furthermore, miRNA biogenesis proteins, Drosha and Dicer, were suggested to be involved in the disease[11,12,36]. Taken together, changes in isomiRs may be directly or indirectly related to ALS.

The potential of cell-free isomiRs as biomarkers has been suggested in cancer[37-41] and a
recent work by Loher et al. provided profiles and performed analysis of alterations in transfer RNA-
derived fragments (tRFs), ribosomal RNA-derived fragments (rRFs), Y RNA-derived fragments, and
isomiRs in ALS plasma[42].

Here, we test the potential value of isomiRs for ALS prognostication. We profiled cell-free
isomiRs in a British cohort of 154 patients with ALS. Using Cox proportional hazard regression[43-45],
and orthogonally machine learning (ML) survival prediction[43,45-48], we identified 18 isomiR predictors,
which were successfully replicated in an independent international cohort of 200 patients. We further
demonstrate that isomiR-let-7g-5p.t, particularly carries prognostic value with potential to complement
established clinical features and previously described biomarkers, such as miR-181 and NfL. These
findings establish isomiR as a novel family of biomarkers for neurodegeneration and highlight their
potential as reliable predictors of ALS survival.

Methods

Clinical cohorts

Two ALS cohorts were included in this study. The ALS biomarker study (REC Reference: 09/H0703/27 IRAS ID: 47352[49]) cohort which consists of 154 British patients (London, UK) matched in their inclusion criteria to a "trial-like" subset of the Phenotype Genotype Biomarker (PGB) study (registered at clinicaltrials.gov NCT02327845) consisting of 200 patients from 12 U.S. sites and one South African site, associated to the Clinical Research in ALS and Related Disorders for Therapeutic Development (CReATe) Consortium[9,50]. Patients were diagnosed with ALS by experienced ALS neurologists. Inclusion criteria were as follows: time from symptom onset to first blood draw shorter than 36 months, a baseline slow vital capacity (SVC) equal to or above 50%, and at least two study visits with measurement of the revised amyotrophic lateral sclerosis functional rating scale (ALSFRS-R)[51]. Cognitive impairment (ALSci) or behavioral impairment (ALSbi) were determined by applying the revised Strong criteria[52] to Edinburgh Cognitive and Behavioral ALS Screen (ECAS[53]) data. Patients with either neuroinflammatory or neurodegenerative disorders, recent injuries, systemic or organ-specific autoimmune disorders, and recent treatment with steroids, immunosuppressants, or immunoglobulins were excluded from the British cohort. Control cohorts included 99 adult individuals from the ALS biomarker study in Queen Mary Hospital (ethics approval 09/H0703/27) and 56 adult individuals enrolled in the longitudinal FTD cohort studies at UCL (ethics approval 15/0805). All participants provided written consent (or verbal permission for a carer to sign on their behalf) to be enrolled in the ALS biomarker study if they met inclusion criteria. Ethics approval was obtained from East London and the City Research Ethics Committee 1 or the University of Miami central IRB, which serves as the IRB of record for the CReATe Consortium (Internal Protocol ID #20160603). The University of Cape Town Health Sciences Human Research Ethics Committee approved the study in South Africa (REF number 165/2017). Patients were de-identified. Disease severity was assessed with the ALSFRS-R. The progression rate prior to enrollment was estimated using the DeltaFRS, calculated by subtracting the ALSFRS-R score at baseline assessment from a maximum score of 48,

and dividing by the number of months passing from symptom onset to baseline assessment. Any use

of Riluzole was recorded. Index date (follow-up "time zero") was defined as the date of enrollment

and survival time was measured from the index date to events for either mortality, tracheostomy or

permanent assisted ventilation (PAV). The censoring date for survival analysis was documented in a

case where an event was not observed before the study ended. Experimenters were blinded during

the steps of the molecular analysis. Whole blood was collected by venipuncture in EDTA tubes, and

plasma was recovered by centrifugation at room temperature (RT) for 10 min at 1300g- 1750g.

Plasma stored at < −70 °C prior to use.

Small RNA next-generation sequencing

The molecular experimenters were blinded to the identity of the samples. Total RNA was

extracted from 500μl plasma using miRNeasy Micro Kit following the manufacturer's protocol except

adding 2.5ml of Qiazol to each plasma sample (Qiagen, #217084). RNA was quantified with a Qubit

fluorometer using the RNA Broad Range Assay Kit (Thermo Fisher Scientific, #Q10211). Small RNA

next-generation sequencing libraries were prepared using the QIAseq miRNA Library Kit (Qiagen,

#331502) from 7.5 ng of total RNA. If RNA concentration was below the limit of detection, 5μl of

undiluted RNA were used. Precise linear quantification of miRNAs was achieved by 12-nucleotide

long unique molecular identifiers (UMIs)[54]. Samples were randomly assigned to the preparation of

libraries indexed with QIAseq miRNA NGS 48 Index IL (Qiagen, #331595) and sequenced in batches

of 48 or fewer samples. Library concentration was determined with a Qubit fluorometer (dsDNA High

Sensitivity Assay Kit, Thermo Fisher Scientific, #Q32854) and library size with TapeStation D1000

(Agilent). Libraries with different indices were multiplexed and sequenced on a NextSeq 500/550 v2

(Illumina, #20024906) or a NovaSeq 6000 flow cell (Illumina, #20028401), with a 75-bp single read or

paired-end reads with 6-bp index. FASTQ files were de-multiplexed using the user-friendly

transcriptome analysis pipeline[55]. Human miRNAs and isomiRs, as defined by miRBase - V22[56], were

mapped using CLC genomic workbench 22.0 (Qiagen) only to the positive strand. Reads were

trimmed in two consecutive fashions; Phred score[57], based error ($p_{error} = 10^{-\frac{Q}{10}}$) was calculated for

every nucleotide, high values reflect poor read quality. A trimming score defined by the subtraction of

the error from a threshold of 0.05 was set for each nucleotide. Then, a running sum of the nucleotide

errors was calculated per read. The region between the first positive value of the running sum and the

highest value of the running sum was kept with non-positive score reads being removed completely.
Consecutively, Reads' ends were trimmed to keep the maximum length region containing 2 or fewer
ambiguous nucleotides. This was followed by the library's 3'-adaptors
(TGGAATTCTCGGGTGCCAAGG) trimming, with reads that did not contain any adaptors being
excluded from downstream analyses. Reads were filtered based on lengths of 15-55 bps. IsomiRs
were considered as 2bps changes (insertion, deletion, shift, and mismatch) up/downstream from
MirBase reference sequence with a maximum of 2 mismatches allowed in alignment.

IsomiR definitions

142 Reference miRNA refers to the most abundantly expressed sequence variant among all isomiRs
143 derived from a given miRNA gene. The collective set of all isomiRs—often comprising around 50
144 distinct variants—annotated for a specific miRNA gene is defined as the miRNA.

QC and pre-processing of sequencing data

Sequencing data were pre-processed separately for each cohort (British n=154 / PGB n=200)
and yielded ~280K/~228K isomiRs respectively, but 99% of identified isomiRs were extremely rare
and sparsely detected across samples. Therefore, isomiRs were included in the analysis only if their
average expression level across cohort's samples was above 5 unique molecular identifiers (UMIs)
yielding a total of 2046 and 2003 isomiRs in the British and PGB cohorts respectively. As geometric
mean cannot be calculated with missing values, the remaining isomiR missing reads were imputed
using multiple imputation with a denoising autoencoder algorithm (parameters: 40 forward passes,
three layers with 512 nodes each, probability of corruption of 0.95, learning rate of $1e^{-6}$). Missing
values were replaced by the mean isomiR levels across 50 imputed datasets. Clinical features used
for estimation of similarity between patients were age, sex, site of symptom onset, and Riluzole
treatment. Python-V3.9.2 and R-V4.0.4 package *rMIDAS*-V.0.3[58] were used for the imputation. Using
DESeq2[59], libraries were corrected for size by geometric mean, under the assumption that miRNA
counts followed negative binomial distribution[59]. Then, imputed values of original missing values were
transformed back to zero to avoid bias in batch correction. Finally, data were corrected for the library
preparation batch to reduce its potential bias by the R Combat function in *SVA*-V3.38 package[60].
Then, levels of 1480 isomiR identities shared across the two cohorts were standardized and isomiRs
were considered for downstream analyses.
163
Machine learning-based isomiR selection for disease prognostication
165

Biomarker discovery analysis (training) was performed using the British cohort that we have
recently characterized[17] while the PGB cohort was reserved for external validation. The 1,480 isomiR
candidates were randomly sampled with repetitions to generate 3,718 bootstrap subsets, each
containing 20 isomiRs. The sampling process ensured that each isomiR was included in at least 50
distinct subsets. Importantly, any given subset contained one specific isomiR of interest alongside 19
other randomly selected isomiRs, resulting in at least 50 different subset compositions for each
individual isomiR. For each bootstrap set, survival analysis was conducted using three survival
models: Cox regression analysis[43], random survival forest (RSF)[45], and gradient survival boosting
(XGB)[44]. The analyses were performed using the scikit-survival library (version 0.23.0) in Python.
Sequential forward selection (SFS)[61], adds iteratively isomiRs in different combinations to optimize
the concordance index score[62] (C-index, mlxtend library in Python).
The process explored approximately 1,000,000 possible combinations of isomiRs for each

subset $(\sum_{k=0}^{20} \binom{20}{k} = 2^{20} = 1,048,576)$, resulting in roughly 3.9 billion combinations across all 3,718

isomiRs subsets. Each isomiR combination was evaluated using 10-fold cross-validation (CV) on

subsets of the UK (discovery) cohort within each of the three orthogonal survival models. A selection

threshold of ≥90% in the SFS process yielded 84 isomiRs. Subsequently, additional iterations of

feature selection were performed on the 84 isomiRs, generating 222 random subsets for further

analysis. A C-index-based threshold, was determined using the elbow method[63], and applied across

all random subsets. This process identified two final panels, together consisting of 18 unique isomiRs,
which were selected for downstream analyses.
186

isomiRs panel prediction validation

After feature selection, an extreme gradient boosting (XGB) model was trained on the UK
discovery cohort to predict patient survival time from enrollment. Hyperparameter tuning was
conducted using 5-fold cross-validation over three rounds of Optuna optimization[64] (1000 trials per
round, with a random sampler and seed = 42), followed by manual fine-tuning. The optimized model
was trained on the selected isomiR levels from the UK cohort and validated using the corresponding
isomiR levels from the PGB replication cohort. Shapley Additive Explanations (SHAP[65]) were used to
identify feature importance within the model and were calculated on the individual predicted hazard

function. Hazard ratios based on SHAP values were calculated as previously described[66]. Individual
SHAP values were exponentiated across features. Values were averaged separately for all patients
below feature mean level and those above it. The average value of patients above mean levels was
divided by the average value of patients below mean level to yield the hazard ratio. The performance
of the chosen isomiR-based model was compared against alternative models trained on: reference
miRNA levels only, miRNAs levels (defined as the aggregate of all isomiRs within a miRNA
annotation).

NfL assay

NfL levels were quantified using single molecule array (Simoa) technology on the Simoa HD-1
Analyzer (Quanterix) in plasma or serum from 152/200 UK discovery or PGB replication cohorts
respectively. The assay utilized the Simoa Nf-L Advantage Kit (Quanterix, #102258) with standards,
primary and secondary antibodies, and detection ranges specified by the manufacturer, including
defined lower and upper detection limits. Univariate or multivariate Cox regression with or without
clinical features and associations between these and mortality were analyzed as described below.

Feature engineering

For univariate/multivariate Cox regressions isomiR levels and NfL levels were dichotomized
into binary variables based on the median levels in the UK discovery cohort, whereas miR-181 levels
were dichotomized using a previously published threshold[17]. Combinations of isomiR-let-7g-5p.t with
NfL or miR-181 were evaluated by their ratios followed by a dichotomization based on the median
levels in the UK discovery cohort for NfL combination or by an in-house algorithm that evaluated
dichotomization cutpoint by optimizing univariate Cox mortality hazard ratio on the UK discovery
cohort.

miRNA target prediction

mRNA targets of the reference miRNA were retrieved from miRCarta[67], retaining those listed
in at least one of miRTarBase, micro-T-CDS, or TargetScan. Target canonical transcripts sequences
were obtained from Ensembl[68] using *BioPython* V-1.83[69]. miRNA and mRNA sequences were
analyzed using RNA22-v2[70] with sensitivity-optimized parameters (see Supplementary Methods).
Significant hybridization sites were counted and compared between the reference miRNA and the
isomiR. Targets with differing site counts were analyzed for enrichment using Enrichr[71-73] via *GSEApy*
V-1.1.9[74] and tested against an ALS gene list obtained from Eitan et al[75].

Statistical analysis

The student's t-test was applied to assess significant differences in the age at symptom onset or study enrollment between groups. The Mann–Whitney U test was used to evaluate differences in ALSFRS-R, DeltaFRS scores and isomiR-let-7g-5p.t levels between the groups. Differences in the proportions of censored events, sex prevalence or Riluzole treatment, were tested using a chi-squared goodness-of-fit test. One-way ANOVA with Tukey-HSD post-hoc analysis assessed differences in isomiR-let-7g-5p.t levels across ALS sub cohorts and control. The Wilcoxon test signed rank test was used to evaluate enrichment of miRNA targets for ALS genes. Pearson correlation coefficients were calculated to examine the relationships between isomiR levels within/between the two cohorts. Kaplan–Meier survival estimators were analyzed using the log-rank Mantel–Cox test, with survival data censored at a fixed date. Univariate and multivariate Cox proportional hazard models were employed to estimate mortality hazard ratios from the time of enrollment, and model fit was assessed using the C-index. The performance of the XGB model was evaluated using the dynamic time-dependent area under the curve[76] (AUC(t)) across the range of survival periods, the average AUC(t), and the C-index. All statistical analyses were conducted, and visualizations were generated using Python (version 3.9.2), R (version 4.0.4) and BioRender.

Results

Study population and baseline characteristics

IsomiRs are sequence isoforms of miRNAs that arise from differences in processing, editing, or modifications of miRNA precursors (**Figure 1A**). In this study we use the term "reference miRNA" to refer to the most abundantly expressed (predominant) sequence variant among all isomiRs derived from a given miRNA gene. Noteworthy, the literature refers to a "miRNA" as the collective set of all dozens of annotated isomiRs[20].

Our study explores isomiRs as ALS prognostic biomarkers in two independent cohorts of patients with ALS. Clinical data and blood samples were obtained for 154 patients from the ALS biomarker cohort (UK)[49] and for 200 patients from the phenotype-genotype biomarker (PGB) study[50]. Clinical and demographic cohort characteristics are summarized in **Table 1.** Discovery and replication cohorts are clinically comparable. DeltaFRS (Mann-Whitney P-value = 0.18, **Figure S1A**), baseline ALSFRS-R score (Mann-Whitney P-value = 0.24, **Figure S1B**), site of symptom onset (χ^2 goodness of fit P-value = 0.08, **Figure S1C**), Riluzole treatment (χ^2 goodness of fit P-value = 0.34, **Figure S1D**) and C9orf72 repeat expansion status (χ^2 goodness of fit P-value = 0.6). Imbalances in patients' sex distribution (χ^2 goodness of fit P-value = 0.009, **Figure S1E**) and age were noticed (t-test P-value ≤ 0.001, **Figure S1F**). The number of censored events were also statistically different between the cohorts (9% vs 54% in UK and PGB, respectively, χ^2 goodness of fit P-value ≤ 0.001, **Figure S1G**).

Sequencing and data preprocessing were conducted in parallel across 154 and 200 patient samples from the UK and PGB cohorts, respectively. Following noise filtration, 1,480 isomiRs from 287 miRNA genes were co-identified in both the discovery and replication cohorts. The discovery isomiRs study was performed on the British cohort. The PGB was held out as an unseen independent replication cohort.

	UK – discovery (N = 154)	PGB – replication (N = 200)	Overall (N = 354)
Sex:			
Males	91 (59%)	90 (45%)	181 (51%)
Females	63 (41%)	110 (55%)	173 (49%)
Age at symptom onset (years) (Mean ± SD)	62.3 (± 11)	57 (± 12)	59.3 (± 12)
Age at enrollment (Years) (Mean ± SD)	63.9 (± 11)	58.4 (± 12)	60.8 (± 12)
Disease duration at enrollment (months) (Mean ± SD)	19.3 (± 10)	16.5 (± 8)	17.7 (± 9)
Baseline ALSFRS-R (AU) (Median [Q1, Q3])	38 [33, 43]	39 [35, 41]	39 [34, 42]
DeltaFRS (-ALSFRS-R unit / month) (Median [Q1, Q3])	0.6 [0.3, 0.9]	0.6 [0.4, 0.9]	0.6 [0.3, 0.9]
Site of symptom onset			
Bulbar	52 (34%)	55 (28%)	107 (30%)
Non-bulbar	102 (66%)	145 (72%)	247 (70%)
C9orf72 status			
Carrier	12 (8%)	20 (10%)	32 (9%)
Non-carrier	142 (92%)	180 (90%)	322 (91%)
Riluzole treatment			
Yes	109 (71%)	134 (67%)	243 (69%)
No	45 (29%)	66 (33%)	111 (31%)
Survival from symptom onset (months) (Median [Q1, Q3])	33 [26.7, 46.7]	33.4 [22.5, 43.9]	33 [23.7, 44.5]
Survival from enrollment (months) (Median [Q1, Q3])	13.9 [8.9, 23.9]	14.7 [8.5, 24.1]	14.3 [8.7, 24.1]
Censored event			
Yes	9 (6%)	108 (54%)	117 (33%)
No	145 (94%)	92 (46%)	237 (67%)

271
Table 1. Clinical and demographic description of the study cohorts. Abbreviations: Baseline ALSFRS-R -
ALSFRS-R score at baseline assessment. DeltaFRS – change of ALSFRS-R score from symptom onset to
baseline assessment divided by months passed. Age at enrollment – age at baseline assessment. Survival from
symptom onset / enrollment – time from symptom onset or baseline assessment to death/event censored.
Censored event – number and percentage of right-censored cases. SD standard deviation. Q1 - 25th percentile,
Q3 - 75th percentile.
278

isomiR-let-7g-5p.t levels differentiate between short and long survival

To identify potential isomiRs associated with patient survival we dichotomized the 1480

isomiRs that were shared between cohorts, based on their median abundance in the UK discovery

cohort. A univariate Cox regression—used to assess the association between isomiR expression and

time to death—revealed 290 statistically significant isomiRs (P-value ≤ 0.05), 16 of which remained

significant after correction for multiple comparisons (FDR ≤ 0.05, **Figure 1B**). Levels above median of

15 isomiRs were associated with decreased rate of mortality in patients (univariate Cox, HR < 1, 95%

confidence interval (CI) = 0.34–0.77, FDR ≤ 0.05; **Figure 1C**) and levels above median of one isomiR

were associated with increased rate of mortality (univariate Cox, HR = 1.8, 95% (CI) = 1.32–2.6, FDR

≤ 0.05; **Figure 1C**). Next, we tested these associations in the PGB replication cohort by dichotomizing

isomiR levels by the threshold obtained in the UK discovery cohort. Only levels above median (≥246

UMI counts) of isomiR-let-7g-5p.t were found to be associated with decreased rate of mortality in the

PGB replication cohort (univariate Cox, HR = 0.6, 95% CI = 0.39–0.91, P-value = 0.015; **Figure 1D**).

In addition, levels above median of isomiR-let-7g-5p.t were significantly associated with longer

survival in the UK discovery cohort (log rank, χ^2 = 18.7, P-value ≤ 0.001, median survival difference =

8.2 months; **Figure 1E**) and in the PGB replication cohort (log rank, χ^2 = 5.99, P-value ≤ 0.01,

median survival difference = 9.1 months; **Figure 1F**). Mean isomiR-let-7g-5p.t levels were

significantly different in the UK discovery cohort sub cohorts, defined by plasma isomiR-let-7g-5p.t

levels that are either higher or lower than the median isomiR-let-7g-5p.t plasma value (351 vs. 176

UMIs in the above- and below-median, respectively, Mann-Whitney P-value ≤ 0.001, **Figure S2A**).

We found a similar difference in the PGB replication cohort sub cohorts (324 vs. 198 UMIs , Mann-

Whitney P-value ≤ 0.001, **Figure S2A**).

Sub cohorts were clinically comparable within each cohort. Age at symptom onset (Mann-

Whitney P-value ≥ 0.1, **Figure S2B**), baseline ALSFRS-R score (Mann-Whitney P-value ≥ 0.27,

Figure S2C), DeltaFRS (Mann-Whitney P-value ≥ 0.13, **Figure S2D**), sex distribution (χ2 goodness

of fit P-value ≥ 0.1, **Figure S2E**) , site of symptom onset (χ2 goodness of fit P-value ≥ 0.2, **Figure**

S2F), Riluzole treatment (χ2 goodness of fit P-value ≥ 0.15, **Figure S2G**) and C9orf72 repeat

expansion status (7 vs 5 and 8 vs 12 carriers in above/below median groups respectively in each

cohort, χ2 goodness of fit P-value ≥ 0.6). Therefore, isomiR-let-7g-5p.t levels stratify patients

independently of key clinical variables.

We sought to compare the mortality rate associated with isomiRs and the respective miRNAs,

which reflect the aggregate of dozens of expressed isomiRs, assigned to a miRNA gene. While

isomiR-let-7g-5p.t levels were associated with decreased mortality rate, no cognate miRNA was

significantly associated with mortality rate in both cohorts including let-7g-5p (univariate Cox, HR =

0.87, 1.1; 95% CI = 0.63–1.2, 0.65-1.8; P-value > 0.05 in both, respectively **Figure S3**). Accordingly,

plasma levels of cognate miRNA let-7g-5p did not differentiate between short and long survival in the

UK discovery or the PGB replication cohorts (log rank χ^2 = 0.63, 0.09; P-value > 0.05 in both, median

survival difference = 0.3, 3.8 months respectively; **Figure 1G, H**).

Figure 1 – High plasma isomiR-let-7g-5p.t levels predict longer survival in patients with ALS. (A) miRNA
biogenesis yields several forms of miRNA isoforms (isomiRs). Following transcription, the primary miRNA is
cleaved by the nuclear microprocessor, exported as a precursor to the cytoplasm, and further processed by
Dicer into a ~22-nt guide RNA loaded onto the silencing complex (RISC) for target mRNA binding and silencing.

Variability in biogenesis, such as misplacement of processing factors, leads to the generation of isomiRs. **(B)**
Volcano plot of 1480 plasma isomiRs univariate Cox Hazard ratios (HR, x axis) and their significance (y axis).
Red line denotes significance threshold (P-value = 0.05). Only 16 isomiRs remained significant post multiple
comparison correction (black line denotes FDR = 5.2e-4). Forest plots of univariate Cox proportional hazard
ratios of these 16 significant isomiRs in UK discovery **(C)** or PGB replication **(D)** cohorts (HR ≤ 0.6 or HR = 1.8,
P-value ≤ 0.015). Only isomiR-let-7g-5p.t remained significant in both cohorts. The Cumulative survival
(Kaplan–Meier) curves for isomiR-let-7g-5p.t in the **(E)** UK discovery or **(F)** PGB replication cohorts. (UK: 77/77
patients in sub- vs. supra-threshold cohorts. PGB: 104/96 patients in sub- vs. supra-threshold cohorts). Kaplan–
Meier curves for the mature miRNA counterpart, let-7g-5p, in the **(G)** UK discovery or **(H)** PGB replication
cohorts. (UK: 77/77 patients in sub- vs. supra-threshold cohorts. PGB: 40/160 patients in sub- vs. supra-
threshold using the same threshold for dichotomization.)
336

337 **IsomiR-let-7g-5p.t is the best prognostication isomiR in two independent ALS cohorts**
338

Although isomiR-let-7g-5p.t emerged as a promising candidate for ALS prognostication, the high
dimensionality of our dataset motivated us to explore the predictive performance of isomiR
combinations and their potential non-linear interactions, using a machine learning, as employed in our
previous studies[77]. We sampled at random subsets of 20 isomiRs, with repetition, using bootstrapping
so each isomiR within the pool of 1480 would be included in at least 50 distinct subsets. To mitigate
the risk of false positive discoveries inherent in large-scale analyses, the 154 patients of the UK
discovery cohort were divided randomly into 10 bins and a sequential forward selection (SFS) with 10
fold cross-validation technique was performed for each of the isomiR subset[78,79]. 84 isomiRs were
selected within 90% of the subsets in which they were included after SFS (≥45 out of 50, with Cox
regression[43], random survival forest (RSF)[45], or survival gradient boosting (XGB)[44]**, Figure S4A**). Of
these three orthogonal survival models, XGB demonstrated the highest mean concordance index (C-
index) across the 84 isomiRs (**Figure S4B**).
351

Next, we modeled survival of the 84 isomiRs by XGB. IsomiRs were included if their subsets
displayed a C-index higher than a threshold that was defined empirically by the elbow method[63] (>
C-index of 0.74) Reassuringly, isomiR-let-7g-5p.t was detected, along with 17 other isomiRs: isomiR-
let-7e-5p.G9C, isomiR-let-7f-5p.T16C.t, isomiR-let-7i-5p.G5A, isomiR-130b-5p.T, isomiR-142-3p.a,
isomiR-451a.T.A, isomiR-144-3p, isomiR-146a-5p.T22A, isomiR-151a-3p.A9G, isomiR-25-3p.A22T,
isomiR-30d-5p.A20G.CT, isomiR-425-5p.as, isomiR-432-5p, isomiR-486-5p.C, isomiR-486-5p.T9G.T,
isomiR-92a-3p.T1C, isomiR-92a-3p.T22C (**Table S1** for nomenclature; **Figure S4C**).
359

After hyperparameter tuning with 5-fold cross-validation (See supplementary Methods), a
model based on these 18 isomiRs, demonstrated strong prognostic capacity across patient survival
duration of 84 months (time dependent AUC (AUC(t)) values of 0.7 to 0.95; average 0.78; C-index =
0.7; survival duration 2 - 85 months; **Figure 2A**), but survival prediction was less effective in the PGB
replication cohort, (AUC(t) 0.49- 0.9, average, 0.62; C-index = 0.55; survival duration 3–54 months;
Figure 2A). Notably, prediction was better for patients with longer survival times.

We demonstrate higher performance with the model trained on the 18 selected isomiRs,
relative to models trained on the collective set of all annotated isomiRs ("miRNA", average AUC(t) =
0.64/0.53, C-index= 0.59/0.52 in the UK discovery and PGB replication cohorts, respectively) or the
most abundant isomiR ("reference miRNAs" average AUC(t)= 0.68/0.46, and C-index= 0.63/0.47)
Figure 2B, C). Additionally, the summation of all isomiRs annotated to a miRNA gene and ther most
abundant isomiR (miRNA and reference miRNA), displayed poor performance over time (**Figure S5**).

To identify the isomiRs most valuable for survival prediction, we calculated Shapley Additive
Explanations (SHAP) for the model average risk function. High levels of isomiR-let-7g-5p.t were
associated with reduced risk of death and longer survival (**Figure 2D**). In contrast, High levels of
isomiR-92a-3p.T22C, isomiR-92a-3p.T1C, isomiR-let-7f-5p.T16C.t, and isomiR-let-7i-5p.G5A
associate with higher risk and shorter survival. A similar SHAP analysis of the PGB replication cohort
was highly correlated, highlighting the model's robustness (**Figure 2E**). We then derived the monthly
hazard ratios (HRs) from SHAP values as previously described[64]. IsomiR-92a-3p.T22C and isomiR-
let-7g-5p.t levels were associated the most with increased / decreased mortality, respectively (**Figure
2F**).
381

As isomiR expression can vary across cohorts, we assessed abundance and inter-cohort
correlation to evaluate biological reproducibility. Among the 18 selected isomiRs, mean expression
ranged from 4–9 UMIs (UK cohort) and 5–14 UMIs (PGB cohort) and the detection rates were
comparable between cohorts (**Figure S6A, B**). IsomiR-142-3p.a had the highest expression (UK =
974 ± 833 UMIs, PGB = 1984 ± 945 UMIs), followed by isomiR-let-7g-5p.t (UK discovery
cohort: 264 UMIs ± 129; PGB replication cohort: 259 UMIs ± 94). IsomiR abundance was well
correlated between cohorts (r = 0.74, P = 8.1e−28, **Figure S6C, D**) but no correlation was noted

between the abundance of isomiRs and their cognate "miRNA" (collective set of all annotated
isomiRs) nor with the "reference miRNA" (predominant isomiR, **Table S2**).

Lower isomiR-let-7g-5p.t levels in non-neurodegeneration control blood relative to ALS

Having found that isomiR-let-7g-5p.t holds prognostic value for the disease in two orthogonal
approaches, we compared its levels between ALS patients and controls. To do so we reanalyzed
plasma samples from two independent control cohorts from Queen Mary hospital (QMUL - CTL) and
University College of London (UCL - CTL) respectively. isomiR-let-7g-5p.t levels were comparable
across the two ALS cohorts within the same sub cohort (mean difference ≤ 27 UMI, Tukey HSD P-
value ≥ 0.99, **Figure S6**), and although variable across control cohorts (mean difference = 324, Tukey
HSD P-value ≤ 0.001), we observed significantly lower levels in the below-median ALS sub cohorts
compared to both controls and the above-median ALS sub cohorts (mean difference ≥ 369 UMI and
mean difference ≥ 522 UMI respectively, Tukey HSD P-value ≤ 0.001, **Figure S6**). Therefore, isomiR-
let-7g-5p.t is also of potential diagnostic value.

Lastly, after examining isomiR-let-7g-5p.t biomarker potential, we tested if there was a change
in the isomiR's predicted targets compared to its reference miRNA. We retrieved the predicted targets
of miRNA let-7g-5p from miRCarta[67], an online repository, and subjected them to a miRNA-mRNA
hybridization prediction algorithm[70] compared to the reference miRNA let-7g-5p and isomiR-let-7g-
5p.t. Out of potential ~4000 mRNA targets, 3189 had at least one significant hybridization site with
either the reference miRNA let-7g-5p or isomiR-let-7g-5p.t. However, 358 out of 3189 mRNA targets
were predicted to lose at least one hybridization site and 43 were predicted to gain at least one such
isomiR-let-7g-5p.t site (Wilcoxon p-value = 2.4e-47, rank-biserial correlation effect size = 0.67).
Enrichment for targets associated with ALS or was not noticed.

Figure 2 – IsomiR based survival prediction in two independent cohorts. **(A)** Plot of AUC (y axis) in the UK discovery (blue) or PGB replication (purple) cohorts for patient survival, as a function of duration from enrolment (x axis). Bar graphs of **(B)** average AUC across time **(C)** or C-index for a model based on 18 selected isomiRs. SHAP analysis of the contribution of 18 isomiRs to risk prediction in the **(D)** UK discovery or **(E)** PGB replication cohorts. Range of isomiR levels (low–blue to high–red). Positive or negative SHAP values reveal higher or lower effect size of isomiR on the mortality risk (x axis), respectively. **(F)** Plot of HR, on the basis of XGB SHAP values calculated according to Sundrani & Lu[66].

Prognostic Value of Plasma isomiR-let-7g-5p.t in ALS Is Comparable to NfL

As we previously reported[17,80], high (above median) blood NfL levels were significantly associated with shorter survival in the UK discovery cohort (log-rank, χ^2 = 9.4, P-value = 2e-3, median survival difference = 7.1 months; **Figure 3A**) and this was consistent in the PGB replication cohort, when NfL dichotomized according to the median plasma value in the UK cohort (log rank, χ^2 = 17.7, P-value = 2.7e-5, median survival difference = 13.7 months; **Figure 3B**). Above median NfL levels were associated with an increased mortality rate in both cohorts (UK discovery cohort:

univariate Cox, HR = 1.7, 95% CI = 1.2-2.3, P-value = 2.4e-4, C-index = 0.56; PGB replication cohort

Cox HR = 2.4, 95% CI 1.6-3.7, P-value = 4.5e-6, C-index = 0.63; **Figure 3C, D**). In contrast, higher

levels of seven (out of eighteen selected by machine learning, **Figure S3C**) isomiRs were associated

with a decreased mortality rate (univariate Cox, HR < 0.72, 95% CI = 0.37-0.99, P-value ≤ 0.05, C-

index > 0.54; Figure **3C**). However, only isomiR-let-7g-5p.t retained significant HR in univariate COX

regression in the PGB replication cohort (UK discovery cohort: HR = 0.58 95% CI = 0.41-0.81, P-

value = 1.4e-3, C-index = 0.57; PGB replication cohort: univariate Cox, HR = 0.6 95% CI = 0.4-0.9, P-

value = 0.015, C-index = 0.55; **Figure 3C, D**). Accordingly, isomiR-let-7g-5p.t was the only isomiR

that significantly distinguished short and long survival times in both cohorts using the log-rank test

(**Figures S8, S9**).

In multivariate Cox analysis NfL remained significantly associated with increased mortality rate

in both cohorts (UK discovery HR = 1.7, 95% CI = 1.2-2, P-value = 6e-4; PGB replication HR = 3.3,

95% CI = 2.4-5.3, P-value = 2e-7), whereas isomiR-let-7g-5p.t was significant only in the PGB cohort

(multivariate Cox, HR = 0.52, 95% CI = 0.32-0.86, P-value = 0.01; **Figure S10A, B**). Noteworthy,

isomiRs modestly improved the prognostic capacity, compared to a model which only includes NfL, in

both cohorts (C-index = 0.67/0.72 in UK discovery or PGB replication respectively, compared to

0.56/0.63 for NfL alone).

When considering only isomiR-let-7g-5p.t and NfL together, isomiR-let-7g-5p.t showed a

stronger association with risk of mortality than NfL in the UK discovery cohort (HR of NfL in a

multivariate Cox model = 1.64, 95% CI = 1.2-2.3, P-value = 3.9e-4; HR of isomiR-let-7g-5p.t in a

multivariate Cox model = 0.49, 95% CI = 0.35-0.69, P-value = 3.8e-6; **Figure S10C**). Together NfL

and isomiR-let-7g-5p.t yield better prognostication than either marker alone (multivariate Cox C-index

= 0.63 vs. C-index = 0.56 NfL alone or C-index = 0.59 isomiR-let-7g-5p.t alone).

In the PGB cohort, NfL had a stronger association with mortality rate (NfL multivariate Cox,

HR = 3, 95% CI = 1.9-4.6, P-value = 1e-7; isomiR-let-7g-5p.t multivariate Cox, HR = 0.46, 95% CI =

0.29-0.7, P-value = 4.8e-5; **Figure S10D**) but again, both markers together improved survival

prediction compared to either one individually (multivariate Cox C-index = 0.67, NfL alone C-index =

0.63, isomiR-let-7g-5p.t alone C-index = 0.56). Moreover, when combined with relevant clinical

features (baseline ALSFRS-R, site of symptom onset, age at onset), only isomiR-let-7g-5p.t (UK

discovery multivariate Cox, HR = 0.53, 95% CI = 0.38-0.75, P-value = 3e-5; PGB replication

multivariate Cox, HR = 0.52, 95% CI = 0.33-0.83, P-value =6.5e-3) and baseline ALSFRS-R (UK

discovery multivariate Cox, HR = 0.97, 95% CI = 0.94-0.99, P-value = 2.6e-3; PGB replication

multivariate Cox, HR = 0.96, 95% CI = 0.92-0.99, P-value = 0.013) were significantly associated with

mortality risk in both cohorts (**Figure S10E,F**). In contrast, NfL levels association with mortality were

only approaching significance in the UK discovery cohort (multivariate Cox, HR = 1.4, 95% CI = 0.97-

1.94, P-value = 0.08 ;**Figure S10E**) while being significantly associated with mortality in the PGB

cohort with an effect size larger than that of isomiR-let-7g-5p.t (multivariate Cox, HR = 2.5, 95% CI =

1.5-4, P-value ≤ 0.001 ;**Figure S10F**).

Since above-median levels of isomiR-let-7g-5p.t and NfL were associated with opposing

effects on mortality risk, we next assessed their joint prognostic capacity by calculating the ratio

between isomiR-let-7g-5p.t and NfL levels. High values of this ratio slightly improved distinction

between short and long survival compared to NfL alone (UK discovery log-rank, χ^2 = 11.8, P-value =

5.9e-4, median survival difference = 8 months; PGB replication log-rank, χ^2 = 19.8, P-value = 8.8e-6,

median survival difference = 14 months; **Figures 3E, F**). Notably, the ratio showed a stronger inverse

association with mortality risk compared to either marker alone and improved overall prognosis (UK

discovery univariate Cox, HR = 0.56, 95% CI = 0.4-0.78, P-value = 6.2e-4, C-index = 0.58; PGB

replication univariate Cox, HR = 0.4, 95% CI = 0.26-0.6, P-value = 1.8e-5, C-index =0.64; Figure

S11). Taken together, these results suggest that isomiR-let-7g-5p.t performs comparably to NfL and

that combining isomiR-let-7g-5p.t with NfL improved predictive performance, at least as reflected by

the C-index. These findings highlight the value of isomiR-let-7g-5p.t as a potential prognostic

biomarker in ALS.

Figure 3 – isomiR-let-7g-5p.t-based prognostication compared to NfL. **(A)** Cumulative survival (Kaplan–Meier) curves for NfL plasma levels (76/ 68 subthreshold (blue) /suprathreshold (red) patients, respectively, from enrollment in the UK discovery cohorts and similarly, NfL serum levels (60/140 subthreshold (blue) /suprathreshold (red) patients respectively) in **(B)** PGB replication cohort, respectively. Forest plots of univariate Cox proportional hazard ratios of 18 isomiRs, NfL and miR-181 in **(C)** the UK discovery cohort or, **(D)** PGB replication cohort. NfL, miR-181 and isomiR-let-7g-5p.t are highlighted in green. **(E)** Cumulative survival (Kaplan–Meier) for isomiR-let-7g-5p.t/NfL ratio (76/76 subthreshold (blue) / suprathreshold (red) patients respectively) from enrollment in the UK discovery cohorts and **(F)** PGB replication (147/53 subthreshold (blue) / suprathreshold (red) patients respectively)

Prognostic Value of Plasma isomiR-let-7g-5p.t in ALS Is Comparable to miR-181

Our UK cohort was a subset of the cohort where we originally discovered miR-181, a

multiplication of sister miRNAs miR-181a-5p and miR-181b-5p, as a prognostic marker[17]. A univariate

Cox analysis revealed that high miR-181 levels were associated with a >2.1-fold increased risk of

mortality, again demonstrating that isomiR-let-7g-5p.t has an equivalent prognosis capacity to

previously published markers (UK discovery cohort HR = 2.1, 95% CI = 1.4–3.2, P-value ≤ 0.001, C-

index = 0.58; PGB replication cohort HR = 3.0, 95% CI = 1.7–5.4, P-value ≤ 0.001, C-index = 0.55;

Figure 3C, D).

miR-181 levels distinguished between short and long survivors in the UK cohort (log rank on

dichotomized miR-181 levels: χ^2 = 13.7, P-value = 2.1e-4, median survival difference = 7.2 months;

Figure 4A) and also in the PGB cohort (log rank: χ^2 = 15.1, P-value = 1e-4, median survival

difference = 9.7 months; **Figure 4B**). Notably, isomiR-let-7g-5p.t was the only isomiR to remain

statistically significant in both cohorts, in a combined analysis of miR-181 and the 18 isomiR chosen

from the XGB model (UK discovery multivariate Cox, HR = 0.63, 95% CI = 0.4-0.98, P-value = 0.04;

PGB replication multivariate Cox, HR = 0.57, 95% CI = 0.36-0.9, P-value = 0.017; **Figure 4C, D**). To

assess joint prognostic capacity of isomiR-let-7g-5p along with miR-181 we tested isomiR-let-7g-5p /

miR-181 ratio capacity to distinguish between short and long survival. We found a significant

association of the ratio levels to patient survival in both cohorts (UK discovery log rank: χ^2 = 20.6, P-

value = 5.6e-6, median survival difference = 8 months; PGB replication log rank: χ^2 = 7, P-value = 8e-

3 median survival difference = 6 months; **Figures 4E, F**). Moreover, the hazard ratio of isomiR-let-7g-

5p / miR-181 ratio was 0.38 in the discovery and 0.44 in the replication cohort. (UK discovery

univariate Cox, 95% CI = 0.19–0.77, C-index = 0.58, P-value = 8e-6; PGB replication univariate Cox,

95% CI = 0.18-0.8, C-index = 0.53, P-value = 0.01, **Figure S11**). Taken together, these results

highlight the potential of isomiR-let-7g-5p.t as a robust biomarker for ALS survival.

525
Figure 4 – isomiR-let-7g-5p.t-based prognostication compared to miR-181. Cumulative survival (Kaplan–
Meier) for miR-181. (122/32 subthreshold (blue) / suprathreshold (red) patients respectively) in the UK
discovery **(A)** / PGB replication (181 /19 patients subthreshold (blue) / suprathreshold (red) patients
respectively) **(B)** cohorts. Forest plots of multivariate Cox proportional hazard ratios of 18 isomiRs and miR-181
in the UK discovery **(C)** or PGB replication **(D)** cohorts. isomiR-let-7g-5p.t is highlighted in green. **(E)**
Cumulative survival (Kaplan–Meier) for isomiR-let-7g-5p.t/miR-181 ratio (130/24 subthreshold (blue) /
suprathreshold (red) patients respectively) from enrollment in the UK discovery cohorts and **(F)** PGB replication
(89/11 subthreshold (blue) / suprathreshold (red) patients respectively)

Discussion

IsomiRs display small variations of either sequence insertion, deletion or single nucleotide polymorphism that distinguish them from the reference miRNA form. While miRNAs are extensively studied in ALS[12,14-17,81-86], we are first to demonstrate the value of isomiRs as biomarkers for neurodegeneration in two large independent cohorts. Among 1480 studied isomiRs, isomiR-let-7g-5p.t emerges as the most promising prognostic biomarker. High plasma levels of isomiR-let-7g-5p.t are significantly associated with lower mortality rate and longer survival. This effect was not explained by clinical characteristics, suggesting that the information encompassed in isomiR-let-7g-5p.t is not directly associated with a single clinical feature. IsomiR-let-7g-5p.t levels were reduced in two ALS cohorts relative to two control cohorts, suggesting diagnostic potential. Higher plasma levels of isomiR-let-7g-5p.t are comparable but reciprocal to miR-181 and NfL, which we previously associated with ALS prognosis[17,80]. Additionally, the incorporation of isomiR-let-7g-5p.t to either miR-181 or NfL enhances their prognostic value.

Dysregulation of miRNAs in ALS has previously been demonstrated in mice and humans[11,12,15,16]. Numerous studies have proposed miRNA utility as potential biomarkers for prognosis for ALS[7,81,85]. However, isomiRs have been overlooked, in part due to their sequence heterogeneity and their low abundance[24]. However, our work and the work of Loher et al.[42] and notable dysregulation of RNA-binding proteins such as TDP-43, DICER, DROSHA or FUS in ALS[11,12,35,87-89], should prompt additional studies of isomiRs in ALS. In addition, it is possible that isomiRs play a role in other neurodegenerative disorders, with implications ranging from basic research to clinical applications.

Our analysis found that isomiR-let-7g-5p.t accounts for approximately 40% of the reads of let-7g-5p in the plasma and perhaps predominates the let-7g-5p repertoire. In this context, microRNA let-7g-5p has previously been reported to be lowered in abundance in the blood of patients with sporadic ALS (sALS)[82] and in patients with frontotemporal dementia (FTD) due to a *C9orf72* repeat expansion, compared to healthy controls[90]. Loher et al[42] reported hundreds of differentially abundant isomiRs in ALS patients plasma compared to controls, with roughly equal numbers up- and down-regulated. 53 out of 56 Let-7 family isomiRs were elevated in ALS. The association of reduced plasma isomiR-let-7g-5p.t levels to severe disease, observed in our study, suggests a distinct behavior not captured by Loher et al[42]. Let-7g-5p blood levels are also decreased in individuals with Alzheimer's disease (AD)

or mild cognitive impairment (MCI)[91-98], and correlate with AD severity[98]. In addition, let-7g-5p was
found to be downregulated in brain samples of Huntington patients compared to controls[99]. This
could reflect a specific decrease that is mediated, at least in part via isomiR-let-7g-5p.t.

Members of the let-7 miRNA family have been repeatedly linked to neurodegenerative
diseases. In ALS, let-7d and let-7f are reduced in patient plasma, while let-7b is elevated in
leukocytes. In AD, several let-7 members, including let-7b, let-7d, let-7e, and let-7g, are upregulated
in patient biofluids or neuronal vesicles, with let-7b also correlating with disease progression. In
Parkinson's disease, let-7a and let-7f are decreased in plasma, whereas let-7g-3p is increased in
CSF-derived vesicles[84,100-110]. Together, these findings underscore the relevance of the let-7 family in
neurodegeneration and reinforce the significance of isomiR-let-7g-5p.t as a prognostic marker in ALS.

Seventeen other isomiRs displayed potential relevance in our analysis, though to a weaker
extent. Some of the miRNAs associated with these isomiRs (miR-451, miR-144-3p and miR-486-5p)
are linked to oxidative stress[111-114] and inflammation[115-119] which potentially play a role in
neurodegeneration. However, because we analyze the isomiRs in plasma, the tissue expression
origin is unclear.

The mechanisms underlying isomiR levels including isomiR-let-7g-5p.t and their functional
role in disease progression is yet to be fully elucidated. The loss of a single nucleotide at the 3′ end
occurs outside the seed region and has limited impact on target specificity[120-123]. Accordingly, using
RNA22 v2 we found that a subset of predicted targets interacts differently with isomiR-let-7g-5p.t
relative to let-7g-5p. No enrichment was observed for ALS-related pathways.

Despite the promising prognostic potential of isomiRs in ALS, several limitations must be
noted: some uncertainty remains about the reference miRNA, due to high sequence similarity within
the broad family of Let-7 miRNAs. Future advances in sequencing technologies and alignment
algorithms may improve isomiR resolution and assignment. Larger and more diverse population
cohorts would strengthen the generalizability of the findings and help assess isomiRs robustness
across different ALS subtypes and disease stages. Last, given the availability and strong prognostic
performance of NfL, the practical utility of isomiRs as standalone biomarkers may be limited until an
effective way to measure them is developed. These limitations highlight the need for continued
research on isomiRs in ALS.

Together, our results highlight the potential of isomiR-let-7g-5p.t as a prognostic biomarker in
ALS, potentially offering comparable value to clinical features, miR-181, and NfL. Moreover, isomiR-
let-7g-5p.t enhances effective prognostication in conjunction with these markers. These findings
warrant further exploration of the utility of isomiR-let-7g-5p.t and other isomiRs in neurodegenerative
diseases.

Supplementary tables and figures

Mutation	Nomenclature	Reference sequence	Mutated sequence	Comment
Reference	Let-7g-5p	TGAGGTAGTAGTTTGTACAGTT	TGAGGTAGTAGTTTGTACAGTT	No change
3' deletion	Let-7g-5p.tt	TGAGGTAGTAGTTTGTACAG**TT**	TGAGGTAGTAGTTTGTACAG	Lowercase letter (n)
3' insertion	Let-7g-5p.CC	TGAGGTAGTAGTTTGTACAG	TGAGGTAGTAGTTTGTACAG**CC**	Uppercase letter (N)
5' deletion	Let-7g-5p.stg	**TG**AGGTAGTAGTTTGTACAGTT	AGGTAGTAGTTTGTACAGTT	Addition of "s"
5' insertion	Let-7g-5p.sGG	TGAGGTAGTAGTTTGTACAGTT	**GG**TGAGGTAGTAGTTTGTACAGTT	Addition of "s"
Base exchange	Let-7g-5p.G2C.T16G	TGAGGTAGTAGTTTGTACAGTT	T**C**AGGTAGTAGTTTG**G**ACAGTT	Position relative to reference sequence

Table S1 - IsomiR Nomenclature
Mutations are separated by a dot. Examples of mutations are highlighted in bold and red.
Base exchanges are represented by the original nucleotide, followed by its position in the reference
sequence, and then the new nucleotide (e.g., A12G).
Reference miRNAs do not have any suffixes. Deletions are indicated by lowercase letter suffixes.
Insertions are indicated by uppercase letter suffixes.
5′ indels are prefixed with the letter "s"; otherwise, indels are assumed to be at the 3′ end. The
position of base exchange in indels remains relative to the reference sequence.
616

	miRNA UK discovery	miRNA PGB replication	reference miRNA UK discovery	reference miRNA PGB replication
hsa-let-7e-5p	545.76 ± 350.51	561.41 ± 244.80	290.93 ± 225.13	303.17 ± 137.28
hsa-let-7f-5p	9434.27 ± 4849.83	11554.54 ± 4143.22	5908.90 ± 4210.59	7920.46 ± 3218.58
hsa-let-7g-5p	992.89 ± 372.96	1173.12 ± 307.08	555.03 ± 385.22	713.87 ± 264.32
hsa-let-7i-5p	12677.13 ± 5929.79	14073.11 ± 4257.76	9116.78 ± 5659.07	9864.22 ± 3331.09
hsa-miR-130b-5p	8.26 ± 5.39	9.40 ± 4.97	nan	nan
hsa-miR-142-3p	1180.20 ± 776.49	2430.42 ± 1049.35	61.75 ± 44.68	123.52 ± 63.52
hsa-miR-144-3p	24.58 ± 27.28	16.31 ± 18.94	8.63 ± 9.66	5.56 ± 6.45
hsa-miR-146a-5p	2270.01 ± 1118.95	6507.35 ± 1719.87	1200.03 ± 835.11	3592.34 ± 1241.20
hsa-miR-151a-3p	1480.06 ± 677.72	4782.99 ± 1337.09	802.72 ± 467.30	2963.98 ± 869.57
hsa-miR-25-3p	1865.72 ± 1072.00	2025.82 ± 669.98	1363.26 ± 744.40	1630.85 ± 599.57
hsa-miR-30d-5p	2920.10 ± 861.54	4553.77 ± 999.95	275.84 ± 146.96	399.31 ± 133.34
hsa-miR-425-5p	865.75 ± 427.35	914.48 ± 244.07	129.94 ± 90.52	213.61 ± 80.55
hsa-miR-432-5p	653.12 ± 706.14	640.87 ± 407.83	241.62 ± 214.92	266.80 ± 176.47
hsa-miR-451a	1416.05 ± 1993.90	550.35 ± 745.91	257.16 ± 461.63	133.97 ± 225.36
hsa-miR-486-5p	25375.59 ± 29731.65	7693.64 ± 7295.35	9280.17 ± 8542.15	2768.21 ± 2999.47
hsa-miR-92a-3p	7033.59 ± 6283.45	4454.01 ± 2302.21	4381.68 ± 3579.75	3375.72 ± 1824.18

Table S2 – plasma levels of reference miRNA associated to the 18 isomiRs studied. Levels of reference miRNAs and miRNA (integrated of all isomiRs associated to a defined miRNA gene), across patients in the UK discovery and the PGB replication cohorts. Units are UMIs, unique molecular identifiers. Mean plasma levels and standard deviation.

Figure S1 – Comparison of the UK discovery and PGB replication cohorts metadata. Boxplot of **(A)** DeltaFRS, measured as the reduction in the ALSFRS-R score per month or **(B)** baseline ALSFRS-R score at study enrollment. Pie charts depicting **(C)** SSO - site of symptom onset **(D)** Riluzole treatment, or **(E)** sex for the cohorts. **(F)** Boxplots of age at study enrollment and age at symptom onset. *** P-value ≤ 0.001, two-sided

Student's t-test. **(G)** Pie charts showing the number of censored and deceased events for the UK discovery

(left) and the PGB replication (right) cohorts.

Figure S2 – Above/below median levels of isomiR-let-7g-5p.t are not correlated to clinical features.

Boxplot of **(A)** isomiR-let-7g-5p.t levels **(B)** Age at disease symptom onset, **(C)** baseline ALSFRS-R score or **(D)**

DeltaFRS, measured as the reduction in the ALSFRS-R score per month in above/below median groups across

cohorts. Pie charts of **(E)** Sex, **(F)**SSO - site of symptom onset or **(G)** Riluzole treatment above/below median

groups across cohorts. *** P-value ≤ 0.001, two-sided Mann-Whitney U test.

634
635

636 **Figure S3 – miRNAs, a term reflecting on the integration of all isomiRs associated to a miRNA gene**
637 **display poor prognostication capacity.** Forest plots of univariate Cox proportional hazard ratios of 14 miRNA
638 form of the 16 FDR significant isomiRs (FDR ≤ 0.05) in UK discovery **(A)** or PGB replication **(B)** cohorts.

Figure S4 - Feature selection process yielded 18 candidate isomiRs as optimal predictors for survival.
(A) The workflow outlines the multi-step feature selection strategy applied to 1,480 isomiRs profiled in a UK discovery cohort of 154 ALS patients. Initially, three different survival models—Cox proportional hazards (COX), Random Survival Forest (RSF), and survival-boosting (XGB) —were each applied across repeated data splits, composed of 20 isomiR. For each model, sequential feature selection (SFS) was performed to identify isomiRs contributing most to predictive performance. IsomiRs selected in at least 90% of the SFS runs were retained, yielding a set of 84 highly stable candidates. This refined set underwent a second round of XGB modeling. The top-scoring model from this phase, based on concordance index (C-index), identified a panel of 18 isomiRs that supports survival outcomes. **(B)** Scatter plot of the mean concordance index (C-index) across 3,718 isomiR subsets (y axis) and percentage of times isomiRs were selected in sequential forward selection (SFS) in three survival analyses: Cox (green), RSF (blue), or XGB (purple). **(C)** A plot of C-index (y axis) for sampling of 20 isomiRS from a total of 84, with repetition, using XGB. 18 isomiRs within two subsets, above the red dashed line, displayed best C-index.

654
655 **Figure S5 - Cognate miRNAs and their reference miRNA display ML-based poor prognostic capacity.**
656 Time-dependent AUC (y axis) of **(A)** reference miRNA (predominant isomiR), **(B)** miRNA (integration of all
657 isomiRs associated to a miRNA gene) or isomiRs (dashed line) in the UK discovery (blue) or PGB replication
658 (purple) across cohort patient survival duration (x axis).

Figure S6 – isomiR characteristics across the discovery and replication cohorts. (A) Boxplots of corrected isomiR levels (y-axis) in the UK discovery cohort (blue) and the PGB replication cohort (purple). The 1st and 99th percentiles of isomiR levels are denoted by whiskers. Extreme values are marked by circles **(B)** Bar plot of missing isomiR values in patients from the UK discovery (blue) and PGB replication (purple) cohorts. Heatmaps of the correlation between plasma isomiR levels in **(C)** the UK discovery cohort and **(D)** the PGB replication cohort clustered by hierarchical clustering according to Euclidean distance and average linkage by UK discovery data. Pearson's correlation r from -1.0 (complete negative correlation, clue) to 1.0 (complete correlation, red).

Figure S7 – isomiR-let-7g-5p.t plasma levels are reduced in ALS patients compared to controls. Boxplot

of isomiR-let-7g-5p.t levels in plasma of controls from two independent cohorts (green & yellow) and ALS sub

groups above (light blue / light purple) or below (dark blue / purple) in the UK discovery or PGB - replication

cohorts respectively. * P-value ≤ 0.05, ** P-value ≤ 0.01, *** P-value ≤ 0.001, Tukey's HSD post-hoc test.

Figure S8 - Cumulative survival (Kaplan–Meier) curves for selected isomiRs in the UK discovery cohort. Data dichotomized by UK discovery isomiR median values. Subthreshold or suprathreshold groups are denoted in blue / red, respectively. Group median survival and Log rank statistics are presented.

Figure S9 - Cumulative survival (Kaplan–Meier) curves for selected isomiRs in the PGB replication cohort. Data dichotomized by UK discovery isomiR median values. Subthreshold or suprathreshold groups are denoted in blue / red, respectively. Group median survival and Log rank statistics are presented.

682
683 **Figure S10 - Prognostic performance of joint isomiR-let-7g-5p.t and NfL.** Forest plots of multivariate Cox
684 proportional hazard ratios of 18 isomiRs and NfL in (**A**) the UK discovery or **(B)** PGB replication cohorts.
685 Multivariate and univariate Cox hazard ratios of NfL and isomiR-let-7g-5p.t in the UK discovery **(C)** or PGB
686 replication **(D)** cohorts Multivariate Cox hazard ratios of NfL, isomiR-let-7g-5p.t and clinical features in the UK
687 discovery **(E)** or PGB replication **(F)**.

688
689 **Figure S11 - Ratios of isomiR-let-7g-5p.t to NfL or miR-181 improve mortality association over single**
690 **markers** Forest plot of univariate Cox proportional hazard ratios of isomiR-let-7g-5p.t, NfL, miR-181 and the
691 ratios of isomiR-let-7g-5p.t,/NfL or isomiR-let-7g-5p.t/miR-181, in both cohorts.

692   <u>Supplementary methods</u>
693
RNA-V2 parameters set to increased sensitivity were:

Number seed nucleotides mismatched tolerated (seedmismatchsallowed) = 2

Seed size (seedsize) = 8, minimum paired bases between miRNA and mRNA outside of seed region

(minpairmatchs) = 10, maximum binding energy to be considered as for match (minenergy) = -10, no

restriction for G:U wobbles (maxguwobbles) = -1, sensitivity-specificity threshold set to 87% and 25%

respectively (threshold=3000000)

700
701
702   Models for feature selection were fitted with the following parameters:
703   -   <u>Cox</u>: without regularization, ties = 'breslow' and a tolerance of $1e^{-9}$.
704   -   <u>Random Survival Forest</u>: n_estimators = 2, max_depth = 4, max_samples = 0.25,
705   n_estimators = 20, min_samples_split = 2, min_samples_leaf = 2, max_features = 0.5,
706   max_samples = 0.5, min_weight_fraction_leaf = 0.0, max_leaf_nodes = None, bootstrap =
707   True, random_state = 42).
708   -   <u>Survival Gradient Boosting</u>: n_estimators = 20, learning_rate = 0.5, max_depth = 2,
709   subsample = 0.3, loss = 'coxph', criterion = 'friedman_mse', min_samples_split = 2,
710   min_samples_leaf = 1, min_weight_fraction_leaf = 0.0, min_impurity_decrease = 0.0,
711   max_features = None, max_leaf_nodes = None, validation_fraction = 0.1, n_iter_no_change =
712   None, tol = 0.0001, dropout_rate = 0.3, ccp_alpha = 0.0,random_state = 0)
713
Survival Gradient Boosting hyperparameters post optuna and manuel tuning: n_estimators = 95,
learning_rate = 0.1, max_depth = 35 subsample = 0.4545454545454545, loss = 'coxph', criterion =
'friedman_mse', min_samples_split = 60, min_samples_leaf = 20, min_weight_fraction_leaf =
0.060461799614030046, min_impurity_decrease = 0.6, max_features = 0.1111111111111111,
max_leaf_nodes = 100, validation_fraction = 0.1, n_iter_no_change = None, tol = 0.0001,
dropout_rate = 0.016, verbose = 1, ccp_alpha = 0.1, random_state = 42.

Acknowledgments

Eran Hornstein is the Mondry Family Professorial Chair and Head of the Nella and Leon Benoziyo Centre for Neurological Diseases and of the Andi and Larry Wolfe Centre for Neuroimmunology and Neuromodulation. We acknowledge patients for their contributions. We thank ALS biomarker and the Phenotype-Genotype biomarker studies co-workers and their contributions to the biobanking project, the Crown Genomics Team of Nancy and Stephen Grand Israel National Center for Personalized Medicine for small RNA sequencing. The authors thank Dr. Isidore Rigoutsos and Phillipe Loher from Thomas Jefferson University for their assistance with the revised manuscript. The authors thank participants in the CReATe Consortium's Phenotype-Genotype-Biomarker (PGB1) study; research staff at each clinical site; and the CReATe Consortium project management and data management teams, genomics sub-core, and biorepository

Funding

Research at the Hornstein lab is funded by Andi and Larry Wolfe Centre for Neuroimmunology and Neuromodulation; the Binational Science Foundation (BSF); Association Française Contre les Myopathies (AFM) grants 24882, 28680; Muscular Dystrophy Association (MDA) grant 1280000; Target ALS; Israel Science Foundation (ISF 3497/21, 424/22); ALS Canada; Minna-James-Heineman Stiftung through Minerva, Minerva Foundation, with funding from the Federal German Ministry for Education and Research; Robert Packard Center for ALS Research at Johns Hopkins; McGill University; EU - ERA-Net; Radala Foundation for ALS Research; Additional support generously provided by the Kekst Family Institute for Medical Genetics. Weizmann SABRA - Yeda-Sela - WRC Program, the Estate of Emile Mimran, and The Maurice and Vivienne Wohl Biology Endowment. Nella and Leon Benoziyo Center for Neurological Diseases, Goldhirsh-Yellin Foundation. Dr. Sydney Brenner and friends, Weizmann - Center for Research on Neurodegeneration,. Redhill Foundation – Sam and Jean Rothberg Charitable Trust Dr. Dvora and Haim Teitelbaum Endowment Fund and Weizmann – Institute for Artificial Intelligence - Acceleration Grant. The CReATe Consortium (U54NS092091) is part of the NIH Rare Diseases Clinical Research Network (RDCRN), an initiative of the Office of Rare Diseases Research (ORDR), NCATS. This Consortium is funded through a collaboration between NCATS and the NINDS. This work was also supported by a Clinical Trial Readiness grant (U01NS107027) from NINDS and by a grant from the ALS Association to support the CReATe Biorepository (grant ID 16-TACL-242).

References

1. Pfohl, S.R., Kim, R.B., Coan, G.S., and Mitchell, C.S. (2018). Unraveling the complexity of amyotrophic lateral sclerosis survival prediction. Front. Neuroinform. *12*. 10.3389/fninf.2018.00036.

2. Beghi, E., Chiò, A., Couratier, P., Esteban, J., Hardiman, O., Logroscino, G., Millul, A., Mitchell, D., Preux, P.-M., Pupillo, E., et al. (2011). The epidemiology and treatment of ALS: focus on the heterogeneity of the disease and critical appraisal of therapeutic trials. Amyotroph. Lateral Scler. *12*, 1-10. 10.3109/17482968.2010.502940.

3. Benatar, M., Boylan, K., Jeromin, A., Rutkove, S.B., Berry, J., Atassi, N., and Bruijn, L. (2016). ALS biomarkers for therapy development: State of the field and future directions. Muscle Nerve *53*, 169-182. 10.1002/mus.24979.

4. Turner, M.R., Bowser, R., Bruijn, L., Dupuis, L., Ludolph, A., McGrath, M., Manfredi, G., Maragakis, N., Miller, R.G., Pullman, S.L., et al. (2013). Mechanisms, models and biomarkers in amyotrophic lateral sclerosis. Amyotroph. Lateral Scler. Frontotemporal Degener. *14 Suppl 1*, 19-32. 10.3109/21678421.2013.778554.

5. Shepheard, S.R., Wuu, J., Cardoso, M., Wiklendt, L., Dinning, P.G., Chataway, T., Schultz, D., Benatar, M., and Rogers, M.-L. (2017). Urinary p75ECD: A prognostic, disease progression, and pharmacodynamic biomarker in ALS. Neurology *88*, 1137-1143. 10.1212/WNL.0000000000003741.

6. Dergai, O., Wuu, J., Koziczak-Holbro, M., Malaspina, A., Granit, V., Hernandez, J.P., Cooley, A., Sachdev, R., Yu, L., Bidinosti, M., et al. (2025). Skeletal muscle biomarkers of amyotrophic lateral sclerosis: a large-scale, multi-cohort proteomic study. medRxiv. 10.1101/2025.04.23.25326161.

7. Dobrowolny, G., Martone, J., Lepore, E., Casola, I., Petrucci, A., Inghilleri, M., Morlando, M., Colantoni, A., Scicchitano, B.M., Calvo, A., et al. (2021). A longitudinal study defined circulating microRNAs as reliable biomarkers for disease prognosis and progression in ALS human patients. Cell Death Discov *7*, 4. 10.1038/s41420-020-00397-6.

8. Benatar, M., Wuu, J., and Turner, M.R. (2023). Neurofilament light chain in drug development for amyotrophic lateral sclerosis: a critical appraisal. Brain *146*, 2711-2716. 10.1093/brain/awac394.

9. Benatar, M., Macklin, E.A., Malaspina, A., Rogers, M.-L., Hornstein, E., Lombardi, V., Renfrey, D., Shepheard, S., Magen, I., Cohen, Y., et al. (2024). Prognostic clinical and biological markers for amyotrophic lateral sclerosis disease progression: validation and implications for clinical trial design and analysis. EBioMedicine *108*, 105323. 10.1016/j.ebiom.2024.105323.

10. Lee, R.C., Feinbaum, R.L., and Ambros, V. (1993). The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. Cell *75*, 843-854. 10.1016/0092-8674(93)90529-y.

11. Haramati, S., Chapnik, E., Sztainberg, Y., Eilam, R., Zwang, R., Gershoni, N., McGlinn, E., Heiser, P.W., Wills, A.-M., Wirguin, I., et al. (2010). miRNA malfunction causes spinal motor neuron disease. Proc. Natl. Acad. Sci. U. S. A. *107*, 13111-13116. 10.1073/pnas.1006151107.

12. Emde, A., Eitan, C., Liou, L.-L., Libby, R.T., Rivkin, N., Magen, I., Reichenstein, I., Oppenheim, H., Eilam, R., Silvestroni, A., et al. (2015). Dysregulated miRNA biogenesis downstream of cellular stress and ALS-causing mutations: a new mechanism for ALS. EMBO J. *34*, 2633-2651. 10.15252/embj.201490493.

13. Katsu, M., Hama, Y., Utsumi, J., Takashina, K., Yasumatsu, H., Mori, F., Wakabayashi, K., Shoji, M., and Sasaki, H. (2019). MicroRNA expression profiles of neuron-derived extracellular

811    vesicles in plasma from patients with amyotrophic lateral sclerosis. Neurosci. Lett. *708*,
812    134176. 10.1016/j.neulet.2019.03.048.
813 14. Reichenstein, I., Eitan, C., Diaz-Garcia, S., Haim, G., Magen, I., Siany, A., Hoye, M.L., Rivkin,
814    N., Olender, T., Toth, B., et al. (2019). Human genetics and neuropathology suggest a link
815    between miR-218 and amyotrophic lateral sclerosis pathophysiology. Sci Transl Med *11*.
816    10.1126/scitranslmed.aav5264.
817 15. Toivonen, J.M., Manzano, R., Oliván, S., Zaragoza, P., García-Redondo, A., and Osta, R.
818    (2014). MicroRNA-206: a potential circulating biomarker candidate for amyotrophic lateral
819    sclerosis. PLoS One *9*, e89065. 10.1371/journal.pone.0089065.
820 16. Wakabayashi, K., Mori, F., Kakita, A., Takahashi, H., Utsumi, J., and Sasaki, H. (2014).
821    Analysis of microRNA from archived formalin-fixed paraffin-embedded specimens of
822    amyotrophic lateral sclerosis. Acta Neuropathol Commun *2*, 173. 10.1186/s40478-014-0173-z.
823 17. Magen, I., Yacovzada, N.S., Yanowski, E., Coenen-Stass, A., Grosskreutz, J., Lu, C.-H.,
824    Greensmith, L., Malaspina, A., Fratta, P., and Hornstein, E. (2021). Circulating miR-181 is a
825    prognostic biomarker for amyotrophic lateral sclerosis. Nat. Neurosci. *24*, 1534-1541.
826    10.1038/s41593-021-00936-z.
827 18. Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Rådmark, O., Kim, S.,
828    and Kim, V.N. (2003). The nuclear RNase III Drosha initiates microRNA processing. Nature
829    *425*, 415-419. 10.1038/nature01957.
830 19. Bernstein, E., Caudy, A.A., Hammond, S.M., and Hannon, G.J. (2001). Role for a bidentate
831    ribonuclease in the initiation step of RNA interference. Nature *409*, 363-366.
832    10.1038/35053110.
833 20. Guo, L., and Chen, F. (2014). A challenge for miRNA: multiple isomiRs in miRNAomics. Gene
834    *544*, 1-7. 10.1016/j.gene.2014.04.039.
835 21. Morin, R.D., O'Connor, M.D., Griffith, M., Kuchenbauer, F., Delaney, A., Prabhu, A.-L., Zhao,
836    Y., McDonald, H., Zeng, T., Hirst, M., et al. (2008). Application of massively parallel
837    sequencing to microRNA profiling and discovery in human embryonic stem cells. Genome
838    Res. *18*, 610-621. 10.1101/gr.7179508.
839 22. Newman, M.A., Mani, V., and Hammond, S.M. (2011). Deep sequencing of microRNA
840    precursors reveals extensive 3' end modification. RNA *17*, 1795-1803. 10.1261/rna.2713611.
841 23. Ma, H., Wu, Y., Niu, Q., Zhang, J., Jia, G., Manjunath, N., and Wu, H. (2016). A sliding-bulge
842    structure at the Dicer processing site of pre-miRNAs regulates alternative Dicer processing to
843    generate 5'-isomiRs. Heliyon *2*, e00148. 10.1016/j.heliyon.2016.e00148.
844 24. Neilsen, C.T., Goodall, G.J., and Bracken, C.P. (2012). IsomiRs--the overlooked repertoire in
845    the dynamic microRNAome. Trends Genet. *28*, 544-549. 10.1016/j.tig.2012.07.005.
846 25. Vance, C., Rogelj, B., Hortobágyi, T., De Vos, K.J., Nishimura, A.L., Sreedharan, J., Hu, X.,
847    Smith, B., Ruddy, D., Wright, P., et al. (2009). Mutations in FUS, an RNA processing protein,
848    cause familial amyotrophic lateral sclerosis type 6. Science *323*, 1208-1211.
849    10.1126/science.1165942.
850 26. Kwiatkowski, T.J., Jr., Bosco, D.A., Leclerc, A.L., Tamrazian, E., Vanderburg, C.R., Russ, C.,
851    Davis, A., Gilchrist, J., Kasarskis, E.J., Munsat, T., et al. (2009). Mutations in the FUS/TLS
852    gene on chromosome 16 cause familial amyotrophic lateral sclerosis. Science *323*, 1205-
853    1208. 10.1126/science.1166066.
854 27. Sreedharan, J., Blair, I.P., Tripathi, V.B., Hu, X., Vance, C., Rogelj, B., Ackerley, S., Durnall,
855    J.C., Williams, K.L., Buratti, E., et al. (2008). TDP-43 mutations in familial and sporadic
856    amyotrophic lateral sclerosis. Science *319*, 1668-1672. 10.1126/science.1154584.
857 28. Hardiman, O., Al-Chalabi, A., Chio, A., Corr, E.M., Logroscino, G., Robberecht, W., Shaw,
858    P.J., Simmons, Z., and van den Berg, L.H. (2017). Amyotrophic lateral sclerosis. Nature
859    Reviews Disease Primers *3*, 1-19. 10.1038/nrdp.2017.71.

860   29.   Ma, X.R., Prudencio, M., Koike, Y., Vatsavayai, S.C., Kim, G., Harbinski, F., Briner, A.,
861   Rodriguez, C.M., Guo, C., Akiyama, T., et al. (2022). TDP-43 represses cryptic exon inclusion
862   in the FTD-ALS gene UNC13A. Nature *603*, 124-130. 10.1038/s41586-022-04424-7.
863   30.   Keuss, M.J., Harley, P., Ryadnov, E., Jackson, R.E., Zanovello, M., Wilkins, O.G., Barattucci,
864   S., Mehta, P.R., Oliveira, M.G., Parkes, J.E., et al. (2024). Loss of TDP-43 induces synaptic
865   dysfunction that is rescued by UNC13A splice-switching ASOs. bioRxivorg.
866   10.1101/2024.06.20.599684.
867   31.   Melamed, Z.e., López-Erauskin, J., Baughn, M.W., Zhang, O., Drenner, K., Sun, Y.,
868   Freyermuth, F., McMahon, M.A., Beccari, M.S., Artates, J.W., et al. (2019). Premature
869   polyadenylation-mediated loss of stathmin-2 is a hallmark of TDP-43-dependent
870   neurodegeneration. Nature Neuroscience *22*, 180-190. 10.1038/s41593-018-0293-z.
871   32.   Calliari, A., Daughrity, L.M., Albagli, E.A., Castellanos Otero, P., Yue, M., Jansen-West, K.,
872   Islam, N.N., Caulfield, T., Rawlinson, B., DeTure, M., et al. (2024). HDGFL2 cryptic proteins
873   report presence of TDP-43 pathology in neurodegenerative diseases. Molecular
874   Neurodegeneration *19*, 1-5. 10.1186/s13024-024-00718-8.
875   33.   Irwin, K.E., Jasin, P., Braunstein, K.E., Sinha, I.R., Garret, M.A., Bowden, K.D., Chang, K.,
876   Troncoso, J.C., Moghekar, A., Oh, E.S., et al. (2024). A fluid biomarker reveals loss of TDP-43
877   splicing repression in presymptomatic ALS–FTD. Nature Medicine *30*, 382-393.
878   10.1038/s41591-023-02788-5.
879   34.   Chen, X., Fan, Z., McGee, W., Chen, M., Kong, R., Wen, P., Xiao, T., Chen, X., Liu, J., Zhu,
880   L., et al. (2018). TDP-43 regulates cancer-associated microRNAs. Protein Cell *9*, 848-866.
881   10.1007/s13238-017-0480-9.
882   35.   Morlando, M., Dini Modigliani, S., Torrelli, G., Rosa, A., Di Carlo, V., Caffarelli, E., and
883   Bozzoni, I. (2012). FUS stimulates microRNA biogenesis by facilitating co-transcriptional
884   Drosha recruitment: FUS participates in microRNA biogenesis. EMBO J. *31*, 4502-4510.
885   10.1038/emboj.2012.319.
886   36.   Porta, S., Kwong, L.K., Trojanowski, J.Q., and Lee, V.M.Y. (2015). Drosha inclusions are new
887   components of dipeptide-repeat protein aggregates in FTLD-TDP and ALS C9orf72 expansion
888   cases. J. Neuropathol. Exp. Neurol. *74*, 380-387. 10.1097/NEN.0000000000000182.
889   37.   Ibuki, Y., Nishiyama, Y., Tsutani, Y., Emi, M., Hamai, Y., Okada, M., and Tahara, H. (2020).
890   Circulating microRNA/isomiRs as novel biomarkers of esophageal squamous cell carcinoma.
891   PLoS One *15*, e0231116. 10.1371/journal.pone.0231116.
892   38.   Koppers-Lalic, D., Hackenberg, M., de Menezes, R., Misovic, B., Wachalska, M., Geldof, A.,
893   Zini, N., de Reijke, T., Wurdinger, T., Vis, A., et al. (2016). Non‐invasive prostate cancer
894   detection by measuring miRNA variants (isomiRs) in urine extracellular vesicles. Oncotarget
895   *7*, 22566-22578. 10.18632/oncotarget.8124.
896   39.   Aiso, T., and Ueda, M. (2023). 5'‐isomiR is the most abundant sequence of miR‐1246, a
897   candidate biomarker of lung cancer, in serum. Mol. Med. Rep. *27*. 10.3892/mmr.2023.12979.
898   40.   Wang, S., Zheng, Z., Chen, P., and Wu, M. (2019). Tumor classification and biomarker
899   discovery based on the 5'isomiR expression level. BMC Cancer *19*, 127. 10.1186/s12885-
900   019-5340-y.
901   41.   Lan, C., Peng, H., McGowan, E.M., Hutvagner, G., and Li, J. (2018). An isomiR expression
902   panel based novel breast cancer classification approach using improved mutual information.
903   BMC Med. Genomics *11*, 118. 10.1186/s12920-018-0434-y.
904   42.   Loher, P., Londin, E., Ilieva, H., Pasinelli, P., and Rigoutsos, I. (2025). Re-Analyses of
905   Samples From Amyotrophic Lateral Sclerosis Patients and Controls Identify Many Novel Small
906   RNAs With Diagnostic And Prognostic Potential. Molecular Neurobiology *62*, 8135-8149.
907   10.1007/s12035-025-04747-2.
908   43.   Cox, D.R. (1972). Regression Models and Life-Tables. J. R. Stat. Soc. Series B Stat.
909   Methodol. *34*, 187-202. 10.1111/j.2517-6161.1972.tb00899.x.

44. Hothorn, T., Bühlmann, P., Dudoit, S., Molinaro, A., and Van Der Laan, M.J. (2005). Survival
ensembles. Biostatistics *7*, 355-373. 10.1093/biostatistics/kxj011.
45. Ishwaran, H., Gerds, T.A., Kogalur, U.B., Moore, R.D., Gange, S.J., and Lau, B.M. (2014).
Random survival forests for competing risks. Biostatistics *15*, 757-773.
10.1093/biostatistics/kxu010.
46. Breiman, L. (1996). Bagging predictors. Mach. Learn. *24*, 123-140. 10.1007/BF00058655.
47. Džeroski, S., and Ženko, B. (2004). Is Combining Classifiers with Stacking Better than
Selecting the Best One? Mach. Learn. *54*, 255-273. 10.1023/B:MACH.0000015881.36452.6e.
48. Malviya, R., Ghinea, G., Dhanaraj, R.K., Balusamy, B., and Sundram, S. (2022). Deep
Learning for Targeted Treatments: Transformation in Healthcare (John Wiley & Sons).
49. Faure, H., and Hrynaszkiewicz, I. (2011). The ISRCTN Register: achievements and
challenges 8 years on: The ISRCTN Register. J. Evid. Based Med. *4*, 188-192.
10.1111/j.1756-5391.2011.01138.x.
50. ClinicalTrials.gov/study/NCT02327845. https://clinicaltrials.gov/study/NCT02327845.
51. Cedarbaum, J.M., Stambler, N., Malta, E., Fuller, C., Hilt, D., Thurmond, B., and Nakanishi, A.
(1999). The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments
of respiratory function. BDNF ALS Study Group (Phase III). J. Neurol. Sci. *169*, 13-21.
10.1016/s0022-510x(99)00210-5.
52. Strong, M.J., Abrahams, S., Goldstein, L.H., Woolley, S., McLaughlin, P., Snowden, J.,
Mioshi, E., Roberts-South, A., Benatar, M., HortobáGyi, T., et al. (2017). Amyotrophic lateral
sclerosis - frontotemporal spectrum disorder (ALS-FTSD): Revised diagnostic criteria.
Amyotroph Lateral Scler Frontotemporal Degener *18*, 153-174.
10.1080/21678421.2016.1267768.
53. Abrahams, S., Newton, J., Niven, E., Foley, J., and Bak, T.H. (2014). Screening for cognition
and behaviour changes in ALS. Amyotroph. Lateral Scler. Frontotemporal Degener. *15*, 9-14.
10.3109/21678421.2013.805784.
54. Coenen-Stass, A.M.L., Magen, I., Brooks, T., Ben-Dov, I.Z., Greensmith, L., Hornstein, E.,
and Fratta, P. (2018). Evaluation of methodologies for microRNA biomarker detection by next
generation sequencing. RNA Biol. *15*, 1133-1145. 10.1080/15476286.2018.1514236.
55. Kohen, R., Barlev, J., Hornung, G., Stelzer, G., Feldmesser, E., Kogan, K., Safran, M., and
Leshkowitz, D. (2019). UTAP: User-friendly Transcriptome Analysis Pipeline. BMC
Bioinformatics *20*, 154. 10.1186/s12859-019-2728-2.
56. Kozomara, A., and Griffiths-Jones, S. (2014). miRBase: annotating high confidence
microRNAs using deep sequencing data. Nucleic Acids Res. *42*, D68-73.
10.1093/nar/gkt1181.
57. Ewing, B., and Green, P. (1998). Base-calling of automated sequencer traces using phred. II.
Error probabilities. Genome Res. *8*, 186-194. 10.1101/gr.8.3.186.
58. Lall, R., and Robinson, T. Efficient Multiple Imputation for Diverse Data in Python and R:
MIDASpy and rMIDAS.
59. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. 10.1186/s13059-014-0550-
8.
60. Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E., and Storey, J.D. (2012). The sva package
for removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics *28*, 882-883. 10.1093/bioinformatics/bts034.
61. Pudil, P., Novovičová, J., and Kittler, J. (1994). Floating search methods in feature selection.
Pattern Recognit. Lett. *15*, 1119-1125. 10.1016/0167-8655(94)90127-9.
62. Raschka, S. (2018). MLxtend: Providing machine learning and data science utilities and
extensions to Python's scientific computing stack. Journal of Open Source Software *3*, 638.
10.21105/joss.00638.

63. Thorndike, R.L. (1953). Who belongs in the family? Psychometrika *18*, 267-276.
10.1007/BF02289263.
64. Akiba, T., Sano, S., Yanase, T., Ohta, T., and Koyama, M. (2019). Optuna: A next-generation
hyperparameter optimization framework. Proceedings of the 25th ACM SIGKDD International
Conference on Knowledge Discovery & Data Mining. ACM.
65. Lundberg, S., and Lee, S.-I. (2017). A unified approach to interpreting model predictions. arXiv
[cs.AI].
66. Sundrani, S., and Lu, J. (2021). Computing the hazard ratios associated with explanatory
variables using machine learning models of survival data. JCO Clin. Cancer Inform. *5*, 364-
378. 10.1200/CCI.20.00172.
67. Backes, C., Fehlmann, T., Kern, F., Kehl, T., Lenhof, H.P., Meese, E., and Keller, A. (2018).
miRCarta: a central repository for collecting miRNA candidates. Nucleic Acids Res *46*, D160-
D167. 10.1093/nar/gkx851.
68. Yates, A.D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R.,
Armean, I.M., Azov, A.G., Bennett, R., et al. (2020). Ensembl 2020. Nucleic Acids Res *48*,
D682-D688. 10.1093/nar/gkz966.
69. Cock, P.J., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I.,
Hamelryck, T., Kauff, F., Wilczynski, B., and de Hoon, M.J. (2009). Biopython: freely available
Python tools for computational molecular biology and bioinformatics. Bioinformatics *25*, 1422-
1423. 10.1093/bioinformatics/btp163.
70. Miranda, K.C., Huynh, T., Tay, Y., Ang, Y.S., Tam, W.L., Thomson, A.M., Lim, B., and
Rigoutsos, I. (2006). A pattern-based method for the identification of MicroRNA binding sites
and their corresponding heteroduplexes. Cell *126*, 1203-1217. 10.1016/j.cell.2006.07.031.
71. Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and
Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. BMC Bioinformatics *14*, 128. 10.1186/1471-2105-14-128.
72. Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev,
S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a comprehensive gene
set enrichment analysis web server 2016 update. Nucleic Acids Res *44*, W90-97.
10.1093/nar/gkw377.
73. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann,
A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene Set Knowledge
Discovery with Enrichr. Curr Protoc *1*, e90. 10.1002/cpz1.90.
74. Fang, Z., Liu, X., and Peltz, G. (2023). GSEApy: a comprehensive package for performing
gene set enrichment analysis in Python. Bioinformatics *39*. 10.1093/bioinformatics/btac757.
75. Eitan, C., Siany, A., Barkan, E., Olender, T., van Eijk, K.R., Moisse, M., Farhan, S.M.K.,
Danino, Y.M., Yanowski, E., Marmor-Kollet, H., et al. (2022). Whole-genome sequencing
reveals that variants in the Interleukin 18 Receptor Accessory Protein 3'UTR protect against
ALS. Nat Neurosci *25*, 433-445. 10.1038/s41593-022-01040-6.
76. Lambert, J., and Chevret, S. (2016). Summary measure of discrimination in survival models
based on cumulative/dynamic time-dependent ROC curves. Stat Methods Med Res *25*, 2088-
2102. 10.1177/0962280213515571.
77. Magen, I., Yacovzada, N.-S., Warren, J.D., Heller, C., Swift, I., Bobeva, Y., Malaspina, A.,
Rohrer, J.D., Fratta, P., and Hornstein, E. (2023). microRNA-based predictor for diagnosis of
frontotemporal dementia. Neuropathol. Appl. Neurobiol., e12916. 10.1111/nan.12916.
78. Charilaou, P., and Battat, R. (2022). Machine learning models and over-fitting considerations.
World J Gastroenterol *28*, 605-607. 10.3748/wjg.v28.i5.605.
79. Taylor, J.M.G., Ankerst, D.P., and Andridge, R.R. (2008). Validation of biomarker-based risk
prediction models. Clin Cancer Res *14*, 5977-5983. 10.1158/1078-0432.CCR-07-4534.

80. Lu, C.-H., Macdonald-Wallis, C., Gray, E., Pearce, N., Petzold, A., Norgren, N., Giovannoni,
G., Fratta, P., Sidle, K., Fish, M., et al. (2015). Neurofilament light chain: A prognostic
biomarker in amyotrophic lateral sclerosis. Neurology *84*, 2247-2257.
10.1212/WNL.0000000000001642.
81. Koval, E.D., Shaner, C., Zhang, P., du Maine, X., Fischer, K., Tay, J., Chau, B.N., Wu, G.F.,
and Miller, T.M. (2013). Method for widespread microRNA-155 inhibition prolongs survival in
ALS-model mice. Hum Mol Genet *22*, 4127-4135. 10.1093/hmg/ddt261.
82. Liguori, M., Nuzziello, N., Introna, A., Consiglio, A., Licciulli, F., D'Errico, E., Scarafino, A.,
Distaso, E., and Simone, I.L. (2018). Dysregulation of MicroRNAs and Target Genes
Networks in Peripheral Blood of Patients With Sporadic Amyotrophic Lateral Sclerosis. Front.
Mol. Neurosci. *11*, 288. 10.3389/fnmol.2018.00288.
83. Maimon, R., Ionescu, A., Bonnie, A., Sweetat, S., Wald-Altman, S., Inbar, S., Gradus, T.,
Trotti, D., Weil, M., Behar, O., and Perlson, E. (2018). miR126-5p Downregulation Facilitates
Axon Degeneration and NMJ Disruption via a Non-Cell-Autonomous Mechanism in ALS. J
Neurosci *38*, 5478-5494. 10.1523/JNEUROSCI.3037-17.2018.
84. Raheja, R., Regev, K., Healy, B.C., Mazzola, M.A., Beynon, V., Von Glehn, F., Paul, A., Diaz-
Cruz, C., Gholipour, T., Glanz, B.I., et al. (2018). Correlating serum micrornas and clinical
parameters in amyotrophic lateral sclerosis. Muscle Nerve *58*, 261-269. 10.1002/mus.26106.
85. Saucier, D., Wajnberg, G., Roy, J., Beauregard, A.-P., Chacko, S., Crapoulet, N., Fournier, S.,
Ghosh, A., Lewis, S.M., Marrero, A., et al. (2019). Identification of a circulating miRNA
signature in extracellular vesicles collected from amyotrophic lateral sclerosis patients. Brain
Res. *1708*, 100-108. 10.1016/j.brainres.2018.12.016.
86. de Andrade, H.M.T., de Albuquerque, M., Avansini, S.H., de S Rocha, C., Dogini, D.B., Nucci,
A., Carvalho, B., Lopes-Cendes, I., and França, M.C., Jr. (2016). MicroRNAs-424 and 206 are
potential prognostic markers in spinal onset amyotrophic lateral sclerosis. J. Neurol. Sci. *368*,
19-24. 10.1016/j.jns.2016.06.046.
87. Kawahara, Y., and Mieda-Sato, A. (2012). TDP-43 promotes microRNA biogenesis as a
component of the Drosha and Dicer complexes. Proc. Natl. Acad. Sci. U. S. A. *109*, 3347-
3352. 10.1073/pnas.1112427109.
88. Kapeli, K., Martinez, F.J., and Yeo, G.W. (2017). Genetic mutations in RNA-binding proteins
and their roles in ALS. Hum. Genet. *136*, 1193-1214. 10.1007/s00439-017-1830-7.
89. Kour, S., Fortuna, T., Anderson, E.N., Mawrie, D., Bilstein, J., Sivasubramanian, R., Ward, C.,
Roy, R., Rajasundaram, D., Sterneckert, J., and Pandey, U.B. (2023). Drosha-dependent
microRNAs modulate FUS-mediated neurodegeneration in vivo. Nucleic Acids Res *51*, 11258-
11276. 10.1093/nar/gkad774.
90. Kmetzsch, V., Latouche, M., Saracino, D., Rinaldi, D., Camuzat, A., Gareau, T., French
Research Network on, F.A., Le Ber, I., Colliot, O., and Becker, E. (2022). MicroRNA
signatures in genetic frontotemporal dementia and amyotrophic lateral sclerosis. Ann. Clin.
Transl. Neurol. *9*, 1778-1791. 10.1002/acn3.51674.
91. Kafshdooz, T., Farajnia, S., Sharifi, R., and Najmi, S. (2023). Hsa-let-7g-5p, a circulating
microRNA, as a biomarker for Alzheimer's disease. Inform. Med. Unlocked *38*, 101203.
10.1016/j.imu.2023.101203.
92. Tan, L., Yu, J.-T., Tan, M.-S., Liu, Q.-Y., Wang, H.-F., Zhang, W., Jiang, T., and Tan, L.
(2014). Genome-wide serum microRNA expression profiling identifies serum biomarkers for
Alzheimer's disease. J. Alzheimers. Dis. *40*, 1017-1027. 10.3233/JAD-132144.
93. Zhang, X., Wang, Y., Liu, W., Wang, T., Wang, L., Hao, L., Ju, M., and Xiao, R. (2021). Diet
quality, gut microbiota, and microRNAs associated with mild cognitive impairment in middle-
aged and elderly Chinese population. Am. J. Clin. Nutr. *114*, 429-440. 10.1093/ajcn/nqab078.
94. Zhang, X., Xi, Y., Yu, H., An, Y., Wang, Y., Tao, L., Wang, Y., Liu, W., Wang, T., and Xiao, R.
(2019). 27-hydroxycholesterol promotes Aβ accumulation via altering Aβ metabolism in mild

cognitive impairment patients and APP/PS1 mice: 27-hydroxycholesterol leads to cognitive dysfunction via Aβ. Brain Pathol. *29*, 558-573. 10.1111/bpa.12698.

95. Mendes-Silva, A.P., Pereira, K.S., Tolentino-Araujo, G.T., Nicolau, E.d.S., Silva-Ferreira, C.M., Teixeira, A.L., and Diniz, B.S. (2016). Shared biologic pathways between Alzheimer disease and major depression: A systematic review of MicroRNA expression studies. Am. J. Geriatr. Psychiatry *24*, 903-912. 10.1016/j.jagp.2016.07.017.

96. Kumar, P., Dezso, Z., MacKenzie, C., Oestreicher, J., Agoulnik, S., Byrne, M., Bernier, F., Yanagimachi, M., Aoshima, K., and Oda, Y. (2013). Circulating miRNA biomarkers for Alzheimer's disease. PLoS One *8*, e69807. 10.1371/journal.pone.0069807.

97. Satoh, J.-I., Kino, Y., and Niida, S. (2015). MicroRNA-Seq data analysis pipeline to identify blood biomarkers for Alzheimer's disease from public data. Biomark. Insights *10*, 21-31. 10.4137/BMI.S25132.

98. Aharon, A., Spector, P., Ahmad, R.S., Horrany, N., Sabbach, A., Brenner, B., and Aharon-Peretz, J. (2020). Extracellular vesicles of Alzheimer's disease patients as a biomarker for disease progression. Mol. Neurobiol. *57*, 4156-4169. 10.1007/s12035-020-02013-1.

99. Martí, E., Pantano, L., Bañez-Coronel, M., Llorens, F., Miñones-Moyano, E., Porta, S., Sumoy, L., Ferrer, I., and Estivill, X. (2010). A myriad of miRNA variants in control and Huntington's disease brain regions detected by massively parallel sequencing. Nucleic Acids Res. *38*, 7219-7235. 10.1093/nar/gkq575.

100. Daneshafrooz, N., Joghataei, M.T., Mehdizadeh, M., Alavi, A., Barati, M., Panahi, B., Teimourian, S., and Zamani, B. (2022). Identification of let-7f and miR-338 as plasma-based biomarkers for sporadic amyotrophic lateral sclerosis using meta-analysis and empirical validation. Sci Rep *12*, 1373. 10.1038/s41598-022-05067-4.

101. Vrabec, K., Bostjancic, E., Koritnik, B., Leonardis, L., Dolenc Groselj, L., Zidar, J., Rogelj, B., Glavac, D., and Ravnik-Glavac, M. (2018). Differential Expression of Several miRNAs and the Host Genes AATK and DNM2 in Leukocytes of Sporadic ALS Patients. Front Mol Neurosci *11*, 106. 10.3389/fnmol.2018.00106.

102. Lehmann, S.M., Kruger, C., Park, B., Derkow, K., Rosenberger, K., Baumgart, J., Trimbuch, T., Eom, G., Hinz, M., Kaul, D., et al. (2012). An unconventional role for miRNA: let-7 activates Toll-like receptor 7 and causes neurodegeneration. Nat Neurosci *15*, 827-835. 10.1038/nn.3113.

103. Poursaei, E., Abolghasemi, M., Bornehdeli, S., Shanehbandi, D., Asadi, M., Sadeghzadeh, M., Rahmanpour, D., and Sadeh, R.N. (2022). Evaluation of hsa-let-7d-5p, hsa-let-7g-5p and hsa-miR-15b-5p plasma levels in patients with Alzheimer's disease. Psychiatr Genet *32*, 25-29. 10.1097/YPG.0000000000000303.

104. Durur, D.Y., Tastan, B., Ugur Tufekci, K., Olcum, M., Uzuner, H., Karakulah, G., Yener, G., and Genc, S. (2022). Alteration of miRNAs in Small Neuron-Derived Extracellular Vesicles of Alzheimer's Disease Patients and the Effect of Extracellular Vesicles on Microglial Immune Responses. J Mol Neurosci *72*, 1182-1194. 10.1007/s12031-022-02012-y.

105. Liu, Y., He, X., Li, Y., and Wang, T. (2018). Cerebrospinal fluid CD4+ T lymphocyte-derived miRNA-let-7b can enhances the diagnostic performance of Alzheimer's disease biomarkers. Biochem Biophys Res Commun *495*, 1144-1150. 10.1016/j.bbrc.2017.11.122.

106. van Harten, A.C., Mulders, J., Scheltens, P., van der Flier, W.M., and Oudejans, C.B. (2015). Differential Expression of microRNA in Cerebrospinal Fluid as a Potential Novel Biomarker for Alzheimer's Disease. J Alzheimers Dis *47*, 243-252. 10.3233/JAD-140075.

107. Gamez-Valero, A., Campdelacreu, J., Vilas, D., Ispierto, L., Rene, R., Alvarez, R., Armengol, M.P., Borras, F.E., and Beyer, K. (2019). Exploratory study on microRNA profiles from plasma-derived extracellular vesicles in Alzheimer's disease and dementia with Lewy bodies. Transl Neurodegener *8*, 31. 10.1186/s40035-019-0169-5.

108. Samareh, A., Nematollahi, M.H., Pourghadamyari, H., Meimand, H.A.E., Shabani, M., and Asadikaram, G. (2024). Diagnostic Potential of NEAT1, hsa-let-7a-5p, and miR-506-3p in Early-stage Parkinson's Disease. Curr Med Chem. 10.2174/0109298673336756241016063552.

109. Awuson-David, B., Williams, A.C., Wright, B., Hill, L.J., and Di Pietro, V. (2023). Common microRNA regulated pathways in Alzheimer's and Parkinson's disease. Front Neurosci *17*, 1228927. 10.3389/fnins.2023.1228927.

110. Gui, Y., Liu, H., Zhang, L., Lv, W., and Hu, X. (2015). Altered microRNA profiles in cerebrospinal fluid exosome in Parkinson disease and Alzheimer disease. Oncotarget *6*, 37043-37053. 10.18632/oncotarget.6158.

111. Gan, X., Zhao, H., Wei, Y., Jiang, Q., Wen, C., and Ying, Y. (2020). Role of miR-92a-3p, oxidative stress, and p38MAPK/NF-κB pathway in rats with central venous catheter related thrombosis. BMC Cardiovascular Disorders *20*, 1-8. 10.1186/s12872-020-01436-x.

112. Ma, N., Li, S., Lin, C., Cheng, X., and Meng, Z. (2021). Mesenchymal stem cell conditioned medium attenuates oxidative stress injury in hepatocytes partly by regulating the miR-486-5p/PIM1 axis and the TGF-β/Smad pathway. Bioengineered *12*, 6434-6447. 10.1080/21655979.2021.1972196.

113. Liu, Q., Hu, Y., Zhang, M., Yan, Y., Yu, H., and Ge, L. (2018). microRNA-451 protects neurons against ischemia/reperfusion injury-induced cell death by targeting CELF2. Neuropsychiatr. Dis. Treat. *14*, 2773-2782. 10.2147/NDT.S173632.

114. Strycharz, J., Wróblewski, A., Zieleniak, A., Świderska, E., Matyjas, T., Rucińska, M., Pomorski, L., Czarny, P., Szemraj, J., Drzewoski, J., and Śliwińska, A. (2021). Visceral adipose tissue of prediabetic and diabetic females shares a set of similarly upregulated microRNAs functionally annotated to inflammation, oxidative stress and insulin signaling. Antioxidants (Basel) *10*, 101. 10.3390/antiox10010101.

115. Lin, Z., Ge, J., Wang, Z., Ren, J., Wang, X., Xiong, H., Gao, J., Zhang, Y., and Zhang, Q. (2017). Let-7e modulates the inflammatory response in vascular endothelial cells through ceRNA crosstalk. Scientific Reports *7*, 1-12. 10.1038/srep42498.

116. Tan, W., Gu, Z., Leng, J., Zou, X., Chen, H., Min, F., Zhou, W., Zhang, L., and Li, G. (2019). Let-7f-5p ameliorates inflammation by targeting NLRP3 in bone marrow-derived mesenchymal stem cells in patients with systemic lupus erythematosus. Biomed Pharmacother *118*, 109313. 10.1016/j.biopha.2019.109313.

117. Luo, Q., Zhu, J., Zhang, Q., Xie, J., Yi, C., and Li, T. (2020). MicroRNA-486-5p Promotes Acute Lung Injury via Inducing Inflammation and Apoptosis by Targeting OTUD7B. Inflammation *43*, 975-984. 10.1007/s10753-020-01183-3.

118. Mandolesi, G., De Vito, F., Musella, A., Gentile, A., Bullitta, S., Fresegna, D., Sepman, H., Di Sanza, C., Haji, N., Mori, F., et al. (2017). MiR-142-3p is a key regulator of IL-1β-dependent synaptopathy in neuroinflammation. J. Neurosci. *37*, 546-561. 10.1523/jneurosci.0851-16.2017.

119. Thomas, J.J., Harp, K.O., Bashi, A., Hood, J.L., Botchway, F., Wilson, M.D., Thompson, W.E., Stiles, J.K., and Driss, A. (2022). MiR-451a and let-7i-5p loaded extracellular vesicles attenuate heme-induced inflammation in hiPSC-derived endothelial cells. Front. Immunol. *13*, 1082414. 10.3389/fimmu.2022.1082414.

120. Dika, E., Broseghini, E., Porcellini, E., Lambertini, M., Riefolo, M., Durante, G., Loher, P., Roncarati, R., Bassi, C., Misciali, C., et al. (2021). Unraveling the role of microRNA/isomiR network in multiple primary melanoma pathogenesis. Cell Death Dis *12*, 473. 10.1038/s41419-021-03764-y.

121. Lewis, B.P., Shih, I.H., Jones-Rhoades, M.W., Bartel, D.P., and Burge, C.B. (2003). Prediction of mammalian microRNA targets. Cell *115*, 787-798. 10.1016/s0092-8674(03)01018-3.

1157   122.   Wong, L.L., Fadzil, A.B., Chen, Q., Rademaker, M.T., Charles, C.J., Richards, A.M., and
1158   Wang, P. (2023). Interrogating the Role of miR-125b and Its 3'isomiRs in Protection against
1159   Hypoxia. Int J Mol Sci *24*. 10.3390/ijms242116015.
1160   123.   Yu, F., Pillman, K.A., Neilsen, C.T., Toubia, J., Lawrence, D.M., Tsykin, A., Gantier, M.P.,
1161   Callen, D.F., Goodall, G.J., and Bracken, C.P. (2017). Naturally existing isoforms of miR-222
1162   have distinct functions. Nucleic Acids Res *45*, 11371-11385. 10.1093/nar/gkx788.
1163