Graphical Abstract

Enhancing Liver Fibrosis Measurement: Deep Learning and Uncertainty Analysis Across Multi-Centre Cohorts

Marta Wojciechowska, Stefano Malacrino, Dylan Windell, Emma L. Culver, Jessica K. Dyson on behalf of the UKAIH consortium, Jens Rittscher

Standardized Quantification of Liver Fibrosis Across Multi-Centre Cohorts

Cohort Collection

N = **686** biopsies

4 individual studies

20+ healthcare centres

Methodology

PSR staining variability assessment

WSI → Collagen → Uncertainty

Collagen quantification & uncertainty estimation
through deep learning ensembles

Findings

Variability in staining affects CPA quantification

Need for **standardised** protocols

Improved **reliability** of CPA measurements

Framework for future digital pathology studies

Highlights

Enhancing Liver Fibrosis Measurement: Deep Learning and Uncertainty Analysis Across Multi-Centre Cohorts

Marta Wojciechowska, Stefano Malacrino, Dylan Windell, Emma L. Culver, Jessica K. Dyson on behalf of the UKAIH consortium, Jens Rittscher

- A retrospective cohort of liver biopsies collected from over 20 healthcare centres has been assembled.

- The cohort is characterized on the basis of collagen staining used for liver fibrosis assessment.

- A computational pipeline for the quantification of collagen from liver histology slides has been developed and applied to the described cohorts.

- Uncertainty estimation is evaluated as a method to build trust in deep-learning based collagen predictions.

Enhancing Liver Fibrosis Measurement: Deep Learning and Uncertainty Analysis Across Multi-Centre Cohorts

Marta Wojciechowska[a,b], Stefano Malacrino[c], Dylan Windell[d], Emma L. Culver[a,e], Jessica K. Dyson on behalf of the UKAIH consortium[f], Jens Rittscher[a,b,g]

[a]*Nuffield Department of Medicine, University of Oxford, Oxford, UK*
[b]*Big Data Institute, University of Oxford, Oxford, UK*
[c]*Nuffield Department of Surgical Sciences, University of Oxford, Oxford, UK*
[d]*Perspectum Ltd., Oxford, UK*
[e]*Translational Gastroenterology and Liver Unit, John Radcliffe Hospital, Oxford, UK*
[f]*Translational and Clinical Research Institute, Newcastle University and Liver Unit, Freeman Hospital, Newcastle upon Tyne, UK*
[g]*Department of Engineering Science, University of Oxford, Oxford, UK*

Abstract

The introduction of digital pathology has revolutionised the way in which histology-based measurements can support large, multi-centre studies. However, pooling data from various centres often reveals significant differences in specimen quality, particularly regarding histological staining protocols. These variations present challenges in reliably quantifying features from stained tissue sections using image analysis. In this study, we investigate the statistical variation of measuring fibrosis across a liver cohort composed of four individual studies from 20 clinical sites across Europe and North America.

In a first step, we apply colour consistency measurements to analyse staining variability across this diverse cohort. Subsequently, a learnt segmentation model is used to quantify the collagen proportionate area (CPA) and employed uncertainty mapping to evaluate the quality of the segmentations. Our analysis highlights a lack of standardisation in PicroSirius Red (PSR) staining practices, revealing significant variability in staining protocols across institutions. The deconvolution of the staining of the digitised slides identified the different numbers and types of counterstains used, leading to potentially incomparable results.

Email address: `marta.wojciechowska@ndm.ox.ac.uk` (Marta Wojciechowska)

Our analysis highlights the need for standardised staining protocols to ensure reliable collagen quantification in liver biopsies. The tools and methodologies presented here can be applied to perform slide colour quality control in digital pathology studies, thus enhancing the comparability and reproducibility of fibrosis assessment in the liver and other tissues.

Keywords: digital pathology, liver, collagen staining, collagen quantification, quality assurance, uncertainty

---

1. Introduction

Fibrosis is increasingly recognised as a core functional component of the tissue microenvironment. Collagen proportionate area (CPA) is an important image-based measure of fibrosis in whole slide images (WSIs) of liver tissue samples [1]. CPA has already been shown to be an independent predictor of long-term clinical outcomes in patients with metabolic-associated fatty liver disease (MAFLD) [2] and alcoholic liver disease (ALD) [3]. That is, the Liver Forum Case Definitions Working Group [4] has proposed changing CPA as a treatment trial endpoint. Despite its potential, the clinical application of CPA as an imaging biomarker faces substantial challenges. Published findings on CPA often lack consistency, mainly due to variations in collagen detection and quantification methodologies [5]. These discrepancies hinder the comparability of the results between studies and undermine the reliability of CPA as a robust biomarker for liver fibrosis.

In engineering and scientific disciplines, any measurement process is typically characterised by the sources of variability of each step of the process.

However, in the field of digital pathology, the practice of reporting uncertainty values in the computation of stain-positive tissue areas is not commonly observed. This omission largely stems from the complexity of the measurement task, which integrates elements from biochemistry, microscopy, metrology, and computer vision.

To address these challenges, we present an in-depth study of the factors influencing the measurement of collagen percentage in liver tissue samples. We introduce a novel approach in estimating the uncertainty of CPA measurements using an ensemble of deep-learning networks. By combining advanced computational methods with robust statistical analysis, our study aims to enhance the accuracy and reliability of collagen quantification in liver biopsies.

Our work is based on a comprehensive liver cohort comprising four separate studies conducted at more than 20 clinical sites in Europe and North America. This multicenter cohort provides a diverse dataset that reflects the real-world variability in histological staining practices. We also introduce a publicly available tool to assess stain color consistency at the cohort level, which can be applied to perform quality control in digital pathology studies.

In summary, this paper addresses the critical issue of staining variability in collagen quantification and proposes a methodological framework to improve the standardisation and reliability of CPA as an imaging biomarker for liver fibrosis. By doing so, we aim to facilitate the broader adoption of CPA in clinical practice and enhance the comparability of findings across different studies.

1.1. Causes of uncertainty in CPA measurement

First, it must be acknowledged that any biopsy sample may not be representative of the examined organ as a whole. This limitation, caused by the small size of the acquired specimen, affects all histology-derived metrics and predictions, and has been addressed elsewhere [6]. However, from the moment of acquisition of the biopsy, there are still a number of confounding factors that need to be considered to obtain a reliable measure of the collagen area in the analyzed sample.

The process of CPA measurement can be broken down into three separate detection and quantification tasks: 1) collagen in-situ localisation (i.e. chemical staining), 2) the acquisition of the digital image and 3) the quantification of the stained area relative to the area of the tissue section. Each of these processes is subject to its own sources of error, which collectively contribute to the overall uncertainty of the measurement.

1.2. In-situ localisation of collagen

Liver fibrosis is a condition characterised by the pathological deposition of collagen fibres within the liver lobule, and is therefore assessed using collagen-binding stains. Other connective tissue fibres, such as elastin fibres are also present in fibrotic structures. In comparison, their quantity is typically small[7]. Several stainings for identifying connective tissue in general and collagen specifically have been independently developed and traditionally used in pathology studies and in the clinic [5]. These include: PicroSirius Red (PSR), Masson's Trichrome staining for connective tissue, van Gieson's stain, Orcein and others. Each of these stainings is used in pathology to

make the structure of connective fibres visible and can be used to visually assess the grade of fibrosis. However, the only histochemical stain which is known to be collagen-specific and so can be used to quantify collagen is Sirius Red.

PicroSirius Red (PSR) is a staining developed in the 1960's with the purpose of selectively staining collagen in histology samples [8]. The primary agents in this staining are Sirius red F3B (CI 35780, Direct red 80) and picric acid (2,4,6-trinitrophenol, TNP). In the presence of picric acid, Sirius red F3B selectively binds with type I collagen (thicker collagen fibres), and type III collagen (thinner fibrillar collagen/reticulin fibres) [9]. Because of the selective staining for type I and III collagen fibres and its sensitivity to fine collagen strands, the PSR protocol is preferred for the quantification of collagen in histology slides. In this paper, only Sirius red-stained slides were used in the development of digital pathology methods.

In spite of its wide-spread use as a collagen-specific stain, there isn't a single standard protocol for PSR staining, which is used globally across institutions. The classical form of PSR staining includes only the mixture of Sirius Red and picric acid for collagen detection [8]. However, laboratories often add additional stains to the same tissue slice, to simultaneously examine other forms of tissue pathology. Commonly, two other stains may be added: a nuclear stain (a haematoxylin) and a cellular plasma stain (typically a green dye). This lack of PSR staining standardisation between healthcare institutions leads to very significant differences in the colour of the stained slides. This phenomenon can be well observed within the UK-AIH cohort where 20 NHS UK hospitals contributed liver FFPE slides prepared within their pathology laboratories (Fig. 1). The figure illustrates how deep the visual differences between PSR-stained slides can be when comparing samples collected at different sites.

While PSR staining is known to create differential contrast between collagen strands and other tissue components, the exact sensitivity and specificity of PSR stain binding to collagen in a liver tissue section is not known [9]. Additionally, parameters such as the freshness of the specimen, stain concentration, stain exposure time, and ambient temperature can all affect the intensity of staining. It is not clear how the interactions of different chemicals and even the order in which different stains are applied affects the sensitivity and specificity of stain-collagen binding. Finally, histological stains are known to fade over the course of years and therefore retrospectively collected slides may not be comparable to freshly obtained ones in terms of colour

Figure 1: **A visual representation of a selection of Sirius red-stained liver slides from the UK-AIH subcohort.** This image illustrates how the use of different protocols for PSR collagen staining results in dramatic differences in slide appearance within the cohort collected retrospectively from 20 NHS hospitals. The images are not to scale. In each digital slide the background was removed and colours adjusted through histogram equalisation for visual clarity.

intensity.

1.3. Digital sampling

The thickness of an individual collagen fibre is well below the x,y imaging resolution of a typical WSI scanner [10, 11]. Collagen fibres which can be seen in WSIs are aggregates of many collagen fibrils and are characterised by highly complex fractal-like shapes. Because of this fractal-like appearance of the analysed shape the measure of collagen area is intrinsically tied to the x,y resolution of the microscope objective lens and the size of the pixel of the imaging sensor [12]. The dependency of the quantified collagen area on the imaging magnification is significant. Which is why fractal metrics of dimensionality have been successfully applied to predict the overall stage of fibrosis [13].

Beyond the imaging resolution in the xy-plane, parameters such as slice thickness, lighting conditions and sensor exposure time directly affect the intensity of the colour of the digitised slide [14]. Clinical WSIs are typically acquired at either 20x or 40x magnification, which corresponds to x,y pixel sizes of approximately 0.5 x 0.5 µm, and 0.2 x 0.2 µm, respectively (see

Tab. 2). In contrast, typical clinical biopsies are sliced at the thickness range between 4 and 10 µm. Hence a WSI pixel can contain a wide range of collagen thicknesses within the corresponding z volume of the tissue [15].

The final colour intensity of an imaged pixel is therefore a function of 1) the thickness of the tissue slice, 2) the thickness of the imaged collagen fibre in the tissue volume and 3) the relative intensity of collagen staining.

1.4. Segmentation errors

Finally, at any resolution the results of CPA quantification depend on the quality of the segmentation of collagen fibres. Fig. 2 illustrates the difficulty of annotating collagen on an example tile from a SR stained liver biopsy specimen. For now we focus on the data of the PREV study and only consider Sirius Red staining. The stain is present in the entire tissue section and can shown some spatial heterogeneity. Typically, collagen fibres have an intensely red colour. But it can be seen that other regions within the tissue parenchyma (cell cytoplasm and even more so cell nuclei) are also coloured pink. Unspecific staining is a factor that needs to be taken into account. Compression artifacts are another source of noise that needs to be considered. This illustrates why annotating collagen fibres in liver histology slides is burdened with high inter- and intra-observer variability [16].

Figure 2: **Limitations of creating human annotations of collagen fibres in microscopy images.** The task is highly subjective and characterised by high inter- and intra-observer variability, particularly for thin collagen strands and in tissue slides with poor staining quality. While regions with thick, intensely stained collagen are captured, thinner fibres were missed by the annotator (MW).

Performing manual or even semi-automatic annotations of collagen area

is a tedious task which can easily be automated by the employment of image analysis methods. Simple annotation tools often rely on the manual selection of a pixel intensity threshold by the annotating pathologist, such pipelines are known to be vulnerable to stain intensity gradients which are often found in histology slides. Deep learning segmentation models can be trained against pathologist segmentations to be robust to minor and even major differences in stain colour.

2. Materials and Methods

2.1. Cohorts and Data Acquisition

Four separate datasets of liver digital slides are used in this paper, henceforth referred to by the names of their respective original studies. Table 1 contains a brief summary of the cohort characteristics.

Table 1: **An overview of the study cohorts.**

Cohort	N Slides	N Sites	Disease	Country
CALM [17]	153	2	Mix	UK
HepaT1ca [18]	44	1	Liver Cancer, MAFLD	UK
PREV [19]	279	1	MAFLD	USA
UK-AIH [20]	210	20	AIH	UK

In total, 686 digitised biopsies of mixed etiologies were gathered and analysed from a combined total of over 20 clinical sites. In each of the studies, liver biopsies or surgical resections were collected as part of standard care, with the exception of the PREV cohort which consists entirely of volunteers. Slides stained with H&E and PSR stainings were included in the analysis.

The samples within each of the three prospective cohorts in this paper: CALM, HepaT1ca and PREV have been processed according to internally agreed processing and staining protocols. The specimens from all of these studies have been stained with automatic stainers. In contrast to the above cohorts, the UK-AIH study is a retrospective study and slides dating back to 1998 from 20 hospital sites in the UK were selected. The FFPE slides in this cohort were processed according to internal standards within each of the participating hospitals at the time they were collected.

The CALM (ISCRTN39463479), HepaT1ca (NCT03213314) and PREV (NCT03142867) cohorts were made available by Perspectum Ltd., Oxford,

UK. The UK-AIH cohort (IRAS ID: 144806) was made available by the UK-AIH consortium funded by the NIHR Rare Diseases Translational Research Collaboration (RD-TRC) and coordinated by Newcastle University and Newcastle upon Tyne Hospitals NHS Foundation Trust. The UK-AIH study cohort was supported by LiverNorth, a UK-based patient support group and registered charity (No. 1087226). Further information about study design and data acquisition is available in the respective original publications [17–20].

2.2. Slide digitisation

The glass slides were digitised using whole slide imaging (WSI) scanners. Each cohort was scanned internally using parameters chosen by the individuals conducting the respective studies. Key parameters of the image acquisition are summarised in Table 2.

Table 2: **Overview of digital slide parameters for each of the cohorts.**

Cohort	Scanner	Pixel Size at 20x [μm]	File Format	Compression
CALM	Hamamatsu C12000-02	0.4533, 0.4533	*.ndpi	JPEG *level unknown*
HepaT1ca	Hamamatsu C12000-02	0.4533, 0.4533	*.ndpi	JPEG *level unknown*
PREV	Aperio *model unknown*	0.4954, 0.4954	*.svs	Aperio JP2000 RGB
UK-AIH	Leica SCN400	0.5000, 0.5000	*.scn	JPEG *level unknown*

Scanners from several WSI manufacturers were used for data acquisition. WSI systems from different vendors are equipped with different optical instruments and digital image sensors. These differences in hardware mean that the whole-slide images were acquired at slightly different resolutions, given the same nominal magnification (e.g. 20x). Similarly, scanning the same histology slide with two WSI scanners from different vendors produces digital images with slightly different colours [21]. This can be a result of both hardware differences and the colour calibration of a particular scanner. Each vendor also typically uses a proprietary image format.

All slides were processed using the vendor-neutral libraries: OpenSlide [22] and wsi-reader [1]. Wsi-reader is a Python library developed by SM within the Rittscher group at the University of Oxford, designed to offer a unified interface for the most common WSI libraries, such as openslide, tifffile[2] and Philips Pathology SDK.

2.3. Sirius red staining for collagen

To identify the stains used in each of the respective studies in the absence of staining protocols, stain deconvolution was performed on selected digitised slides from each cohort [23]. Stain deconvolution is a transformation of the image from the RGB colourspace into a colourspace in which each of the three image channels encodes the proportion of a colour of a single stain. This processing was performed using QuPath software [24].

The stain deconvolution of the Sirius Red staining, shown in Figure 3, is used as an example to demonstrate the staining variability. Although a red collagen dye was used in each of the centres, it is obvious that different counter stains were used. This is in stark contrast to the results of stain deconvolution applied to the corresponding H&E slices from the same samples (Figure 4).

In the H&E-stained slides across all study cohorts is more standardised. In the PSR-labelled slides, the number of dyes used was ranging from one (just Sirius red) to three (i.e. Sirius red and two additional stains). In the case of the non-homogeneous UK-AIH cohort, different numbers and colours of stains were used for PSR staining.

2.4. Colour characterisation of digital slides

In order to visualise the spectrum of Sirius red-stained digitised FFPE section appearances, we here propose the concept of a primary colour of a digital slide (Fig. 5).

One of the most commonly used approaches to quantisation of colours in digital images is through the employment of median cut algorithm [25]. Median cut algorithm is a method of sorting data of an arbitrary number of dimensions into series of sets by recursively cutting each set of data at the median point along the longest dimension. When applied to an image

---

[1]https://github.com/stefano-malacrino/wsi-reader
[2]https://github.com/cgohlke/tifffile

Figure 3: **The results of stain deconvolution on example digital PSR slides from the analysed cohorts.** a) CALM: the black colour of some collagen fibres is a result of non-specific binding of an unknown green stain to fibres stained with Sirius red. The slides contain a pale yellow residuum. b) HepaT1ca: Sirius red and two unknown stains, olive and blue. Upon close inspection one can see that cell nuclei are non-specifically stained red along with the collagen. c) PREV: only Sirius red stain was detected. d) UK-AIH: slide stained with Sirius red and a purple haematoxylin with an empty residual channel.

Figure 4: **The results of stain deconvolution on example digital H&E slides from the four study cohorts.** In each of the cohorts a haematoxylin and an eosin were used to produce H&E slides. The residual channel is empty or nearly empty in all of the slides. Slight differences in stain hue of both haematoxylin and eosin, as well as scan background colour can be seen between samples from different cohorts.

Input WSI

$y_0 = 10176$ px

$x_0 = 16256$ px

Downsampled image

$y_1 = 636$ px

$x_1 = 1016$ px

Downsample level identification

Background removal
(Otsu thresholding)

Colour palette
(RGB coordinates)

(193, 205, 187)   (161, 137, 156)   (180, 166, 178)

Primary colour of the digitised slide

Median cut algorithm

Pre-processed image

Figure 5: **Identification of the primary colour of a digital slide.** First, an appropriate downsample level is identified in the WSI pyramid. The slide image is then read from the downsampled level of the WSI file. Tissue/background intensity-based threshold is identified with the Otsu thresholding algorithm (applied to the greyscale-converted image). Background pixels are replaced with white values. Finally, the median cut algorithm is applied to thus pre-processed image to identify the primary RGB value in the digital slide.

histogram, it outputs the most commonly occurring pixel values. Here, the WSI is first pre-processed by extraction of a downsampled version of the image and background removal. A pixel intensity threshold between tissue and background is established using the Otsu method applied to a grey-level version of the image [26]. All pixels of value above the established intensity threshold are replaced with white values. Median cut algorithm is applied to the pre-processed image to extract a three-colour colour palette, and the RGB values of the most commonly occurring colour encode the primary colour of the digital slide.

The primary colour has been computed for all of the SR and H&E-stained digitised slides available to this project. For purposes of comparison the RGB values of the primary slide colours were converted to the more perceptually

Figure 6: **Primary colour distribution in digital slides of PSR-labelled sections.**
Samples from the individual cohorts form separate colour clusters.

Combined Cohort H&E

Figure 7: **Primary colour distribution in digital slides of H&E-stained sections.**

uniform CIELAB colourspace [27]. The CIELAB colourspace expresses each colour as three values: L^* for lightness, a^* where negative values indicate green and positive values indicate red, and b^* where negative values indicate blue and positive values indicate yellow. The results of primary colour estimation for the WSIs from all four cohorts are shown in Figures 6 and 7.

3. Computational Methods

3.1. Experimental Data

Table 3 summarises information about the training and validation data in this paper. Cases from the three training cohorts (CALM, PREV, UK-AIH) were manually selected using the obtained PSR stain distribution (Fig. 6) as a guide, to maximise the variation in colour in the experimental dataset. In total 38 whole slide images of liver biopsy sections were densely annotated by MW using QuPath software [24]. The pipeline for semi-automatic annotation of the slides with human in the loop is shown in Figure 8. Tiles containing less than 5% of annotated collagen area were discarded from the training and validation set to improve the stability of model training. All tiles from the HepaT1ca study were retained as an independent (unseen) validation set. A total of 2367 annotated 512x512 px tiles were used in the analysis. Of these 1970 were used for model training and 397 for model validation.

Table 3: **A summary of training and validation data for each of the study cohorts.** Images from CALM, PREV and UK-AIH cohorts were used for method development. HepaT1ca was retained as an independent validation dataset.

Cohort	CALM	PREV	UK-AIH	HepaT1ca
N Annotated Cases	12	13	13	5
N Train Tiles	442	1172	356	-
N Validation Tiles	53	141	52	151
Mean Collagen %	21%	15%	24%	-

3.2. Segmentation Models

The U-Net is a popular deep learning model used for medical image segmentation [28]. We trained three variants of the U-Net model for the col-

Figure 8: **Collagen annotation with human in the loop.** Firstly, a representative region is cropped from the full resolution level of the WSI file. The annotator trains a built-in random trees (RT) pixel classifier to categorise individual pixels within the image as either: "collagen" or "ignore". The results of the RT classification are carefully examined visually. If needed, the human annotations are refined, and the process is repeated until the classifier produces a satisfactory pixel-level classification for the entire image. The annotation is then exported as a binary mask.

lagen segmentation task using the implementation available at [3]. The first two models (here called U-Net Tiny and U-Net Mini) are each down-sized versions of the original U-Net. An Attention U-Net [29] was trained for the same task. The parameters of the individual networks are summarised in Table 4.

Each of the trained segmentation networks has an input which matches the size of 512x512 px RGB image tiles. The ReLU activation function was used in each segmentation model. U-Net Tiny and U-Net Mini used max pooling as the pooling strategy and a sigmoid activation function at the network output. Each of the models used bilinear unpooling. The models were trained using binary crossentropy loss and Adam optimiser. Each of

---

[3]`https://github.com/yingkaisha/keras-unet-collection`

Table 4: **A summary of the parameters of the trained segmentation models.**

Parameter	U-Net Tiny	U-Net Mini	Att U-Net
Depth	3	3	3
Number of Filters	[8, 16, 32]	[32, 64, 128]	[32, 64, 128]
Stacked Layers (Down)	2	2	2
Stacked Layers (Up)	2	2	2
Trainable Parameters	32 665	517 729	612 691

the networks was trained from scratch with a randomised initialisation seed and a batch size of 8. A Quadro RTX 6000 GPU core was used for all of the experiments conducted.

3.3. Uncertainty Estimation

In order to estimate the real uncertainty of PSR measurement, one would need to perform an uncertainty estimation for each of the steps in sample processing and image acquisition separately. Unfortunately, as only digitised slides were made available to this project, such an analysis goes beyond the scope of this paper. Here we propose a way of estimating the uncertainty inherent to the digital image itself, which can be inferred by the image segmentation deep network.

Uncertainty estimation in the context of deep learning can improve the quality of predictions [30, 31]. Bayesian inference is a statistical framework which can be used to quantify the uncertainty present in deep-learning models. The concept is based on Bayes' theorem, which links the posterior distribution of our observations to the prior probability of the hypothesis and the likelihood of the data given the hypothesis:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} \tag{1}$$

where: P(A—B) is the posterior probability of hypothesis A given the observed data B, P(B—A) is the likelihood of the observed data B given hypothesis A, P(A) is the prior probability of hypothesis A and P(B) is the marginal likelihood of the observed data B.

In the Bayesian framework can be used to quantify uncertainty. However, exact Bayesian inference with deep neural networks is computationally

intractable. It can be approximated by Bayesian Neural Networks (BNNs), which are neural networks with probabilistic weights and biases. In the BNN framework, the weights and biases of a network are treated as random variables with a prior distribution over their values. Bayesian inference is then used in the process of training the network to update the prior distribution with new data. The posterior distribution over weights and biases obtained during network training can be used to compute the predictive distribution of the model. This predictive distribution of the model is a representation of the range of possible model outputs given an input. The uncertainty of the model can be inferred from the model's predictive distribution and can be a measure of the model's confidence in the predictions. An alternative approach to Bayesian neural networks is Bayesian model averaging, in which predictions of multiple models are averaged. In practice, several methods of applying Bayesian inference to uncertainty estimation in deep learning have been proposed. Those include Markov chain Monte Carlo [32], variational inference [33], Monte Carlo dropout (MC dropout) [30] and deep ensembles [34].

Predictive uncertainty can be decomposed into two components: aleatoric, which captures inherent randomness in the data (such as noise in the labels), and epistemic, which represents the uncertainty of the model (e.g. when making predictions on out-of-distribution samples) [35]. Kendall et al. [31] proposed an uncertainty decomposition method using a neural network that outputs two values representing the mean and variance of the prediction and using Monte Carlo dropout to approximate Bayesian inference. A similar approach was used by Kwon et al. [36] who proposed a new method for estimating aleatoric and epistemic uncertainties using Monte Carlo dropout. Fort et al. [37] recently demonstrated that deep learning ensembles with random initialisations allow for a more robust approach to uncertainty estimation. This is attributed to the fact that deep ensembles explore entirely different modes, while methods which are based on exploring the subspace of a single model (such as MC dropout) converge to a single mode.

Following [34], we used deep ensembles as an alternative to Bayesian inference. In our experiments, all models were trained for the same duration of 30 epochs. We trained separate ensembles of M = 10 models for each of the studied cohorts and an additional one for the combined cohort (CALM + PREV + UK-AIH). Overall 12 ensembles of M = 10 models each were trained. Slides from the HepaT1ca cohort were retained as a separate validation set.

Given an ensemble Ω of M trained models $\Omega = \{\omega_m\}_{m=1}^M$, we computed the pixel-wise collagen likelihood as the average $\hat{p}$ of the ensemble predictions p_m for an input pixel i.

$$p_m = p(y|i, \omega_m) \tag{2}$$

$$\hat{p} = \frac{1}{M} \sum_{m=1}^M p_m \tag{3}$$

We then used the method proposed by [36] to estimate the pixel-wise predictive uncertainty for an input pixel i and the output prediction y as the variance of y given i over the predictive distribution $p(y|i, \Omega)$:

$$\mathrm{Var}_{p(y|i,\Omega)}(y) = \underbrace{\frac{1}{M} \sum_{m=1}^M [p_m(1-p_m)]}_{\text{aleatoric}} + \underbrace{\frac{1}{M} \sum_{m=1}^M (p_m - \hat{p})^2}_{\text{epistemic}} \tag{4}$$

Two approaches for computing collagen percentage area in a tile were employed; the first made use of the mean ensemble prediction P and the other of the voted ensemble segmentation B. The mean collagen prediction P for image I with dimensions W, H is given by the equation:

$$P = \frac{1}{WH} \sum_{x=1}^W \sum_{y=1}^H \hat{p}(x, y) \tag{5}$$

Collagen segmentation value $\hat{b}$ was obtained for each pixel via thresholding of the corresponding ensemble prediction at collagen likelihood equal or higher than 0.5. Let $\hat{p}(x, y)$ be the ensemble prediction value for pixel i at location (x, y) in the image I, and let $\hat{b}(x, y)$ be the corresponding binary value after thresholding. Hence $\hat{b}(x, y)$ can be defined as:

$$\hat{b}(x, y) = \begin{cases} 1, & \text{if } \hat{p}(x, y) \geq 0.5 \\ 0, & \text{otherwise .} \end{cases} \tag{6}$$

The segmented collagen area proportion B is given by:

$$B = \frac{1}{WH} \sum_{x=1}^{W} \sum_{y=1}^{H} \hat{b}(x,y) \ . \tag{7}$$

The uncertainty in the mean tile collagen prediction is reported as the interquartile range (IQR) of the ensemble collagen predictions P. Similarly, the uncertainty of the segmented collagen area in a tile is reported as the IQR of the collagen segmentation maps produced by thresholding of the predictions by each of the models in the ensemble.

4. Results

The validation metrics for the trained segmentation models are summarised in Table 5. We provide the scores separately for ensembles trained and validated on the individual study cohorts and for the ensemble models trained on the pooled cohort (CALM + PREV + UK-AIH).

Table 5: **Segmentation network Dice scores on validation tiles** (M = 10 ensemble average ± standard deviation).

Training Set	CALM	PREV	UK-AIH	HepaT1ca
Individual Cohorts				
U-Net Tiny	0.847 ± 0.022	0.877 ± 0.012	0.793 ± 0.036	-
U-Net Mini	$\mathbf{0.905 \pm 0.026}$	$\mathbf{0.882 \pm 0.005}$	$\mathbf{0.852 \pm 0.019}$	-
Att U-Net	0.820 ± 0.022	0.825 ± 0.059	0.759 ± 0.070	-
Pooled Cohort				
U-Net Tiny	0.823 ± 0.021	0.861 ± 0.013	0.767 ± 0.037	$\mathbf{0.594 \pm 0.075}$
U-Net Mini	$\mathbf{0.834 \pm 0.029}$	$\mathbf{0.864 \pm 0.014}$	$\mathbf{0.795 \pm 0.035}$	0.555 ± 0.104
Att U-Net	0.785 ± 0.036	0.834 ± 0.017	0.623 ± 0.067	0.539 ± 0.123

The U-Net Mini has achieved the highest validation metric scores in all of the assigned tasks. It can be seen that for all model architectures there is a drop in Dice scores between the ensembles trained on individual cohorts and those trained on the pooled cohort. The higher Dice scores for individual cohorts can be explained by the models over-fitting to the data within each of the cohorts. It can be seen that while very high Dice scores are achieved

for the CALM and PREV cohorts, performance of each of the network architectures is slightly worse for the UK-AIH cohort. This can be explained by the fact that the UK-AIH samples were collected retrospectively from a large number of hospitals and are characterised by frequent staining artefacts and large intra-cohort differences in slide colour. As can be expected, the performance of the segmentation models is significantly lower for the unseen cohort (HepaT1ca).

Figure 9 shows model predictions for an example tile from the CALM cohort. It can be seen that the human annotation used as ground truth for model validation contains noisy false positive pixels around the annotated fibres. These noisy labels can be attributed to falsely positive labelling by the annotator (MW) of the JPEG compression-induced noise in the original WSI. All prediction maps generated by the tested U-Net models provide a realistic representation of the collagen pattern in the original image. It can be seen that neither of the trained segmentation models has replicated the image compression noise in the final prediction, which is a desirable outcome. It is difficult to notice any visual differences in the quality of the prediction maps output by each of the models. This shows that each of the tested networks is appropriate for the task of collagen segmentation.

4.1. Uncertainty computation

Fig. 10 shows the results of model predictions and uncertainty maps for the U-Net Mini ensemble trained on the pooled cohort. It can be seen that almost all of the aleatoric uncertainty is concentrated on the boundaries of collagen fibres. This is expected, as boundary pixels are commonly affected by partial volume effects. Because of this phenomenon, the aleatoric uncertainty is directly tied to the length of the collagen outline in the tile. We observe that tiles with little to no collagen are characterised by uncertainty values near to 0 and tiles with much collagen have high inherent collagen area uncertainty.

Table 6 presents the mean ensemble uncertainty scores obtained from model predictions on the validation set. It can be seen that the values of epistemic uncertainty are very low compared to the values of aleatoric uncertainty for all of the image ensembles. We conclude that the uncertainty of the collagen area measurement primarily depends on the characteristics of a given image. The variability of the measured collagen percentage due to differences in predictions by individual segmentation models within each ensemble is relatively very small.

Image

Figure 9: **Collagen prediction maps output by the tested segmentation U-Net models.** The human annotation used as ground truth for model validation contains noisy false positive pixels around the annotated fibres. Each prediction map shows an output prediction from a single model trained on the pooled cohort. It is difficult to see any differences in the quality of the collagen maps predicted by different model types.

Figure 10: **The results of U-Net Mini ensemble model prediction with uncertainty estimations for examples of unseen tiles from each of the cohorts.** Tiles with low contrast of stained collagen and unseen colours are characterised by high ensemble model uncertainty.

Table 6: **Ensemble uncertainty scores on the validation set for each of the trained model types** (M = 10 ensemble average ± standard deviation).

Model ensemble	U Aleatoric	U Epistemic	U Combined
U-Net Tiny	0.020 ± 0.025	0.002 ± 0.003	0.022 ± 0.027
U-Net Mini	0.017 ± 0.022	0.002 ± 0.003	0.019 ± 0.025
Att U-Net	0.024 ± 0.036	0.003 ± 0.006	0.027 ± 0.041

Figure 11: **An example image prediction tile with high epistemic uncertainty.** The air bubble visible in the original tile is highlighted in the epistemic uncertainty map, which visualises out of distribution data.

Figure 11 shows ensemble model prediction and uncertainties for a tile with a common imaging artefact, an air bubble. The bubble can be clearly seen in the epistemic uncertainty map, meaning that some of the models in the ensemble have classified its pixels as regions of collagen, and some did not. The visualisation of epistemic uncertainty can provide the pathologist with a map of regions in the slide which were not represented in the model training set, and therefore are likely to contain prediction errors.

4.2. Interpreting uncertainty

The results of uncertainty scores computed for each of the cohorts are provided in Table 7.

Table 7: **Ensemble prediction uncertainty scores for validation samples from each of the cohorts, U-Net Mini** (M = 10 ensemble average ± standard deviation).

Training Set	CALM	PREV	UK-AIH	HepaT1ca
Individual Cohorts				
U Aleatoric	0.016 ± 0.016	0.006 ± 0.008	0.024 ± 0.020	-
U Epistemic	0.002 ± 0.003	0.001 ± 0.003	0.003 ± 0.005	-
Pooled Cohort				
U Aleatoric	0.026 ± 0.028	0.008 ± 0.011	0.044 ± 0.033	0.018 ± 0.022
U Epistemic	0.002 ± 0.002	0.001 ± 0.001	0.001 ± 0.005	0.002 ± 0.004

The PREV cohort has the lowest aleatoric uncertainty values of the cohorts studied. The aleatoric uncertainty is highly correlated to the area of collagen in an image (see Fig. A.13) and therefore the lower uncertainty in the PREV cohort can be partly explained by the lower content of collagen in this cohort overall (see Table 3).

Interestingly, after cohort pooling the mean aleatoric uncertainty scores increase for all the models, especially in the cohorts where more than one stain colour was used (i.e. CALM and UK-AIH). This indicates that the segmentation models introduced to different variants of collagen staining have lower confidence at the collagen boundaries where more than one stain is present. Such interpretation would correspond to the intuition that placing additional stains on the tissue specimen introduces additional sources of variability in the data. The highest aleatoric uncertainty values are reported for the UK-AIH cohort in which the purple haematoxylin staining was used in the PSR slides. Collagen tiles in this cohort have overall a lower visual contrast than those from other cohorts.

It can be seen that the correlation between the average prediction P and the segmented collagen B is equal to 1, meaning that these measures can be considered equivalent to each other (Fig. A.13). This means that it may be acceptable to present pathologists with the collagen prediction maps rather than the binarised segmentation maps. The prediction maps have a closer resemblance to the original input image as they capture the colour intensity gradients characteristic of collagen fibres of different thicknesses. Collagen prediction maps are also visually more similar to the images acquired using SHG/TPEF microscopy, where similarly, an inferred collagen density map rather than a binary segmentation map is obtained [38]. Such continu-

ous collagen prediction maps could be interpreted as a kind of "ideal" stain deconvolution, where only pixels containing any collagen can have positive values, and regions of tissue which have no collagen (i.e. cell cytoplasm, cell nuclei) have a collagen probability of 0.

5. Discussion

The problem of accurately performing histological quantification and inference from differently stained slides is a known issue, to which several approaches can be found within the digital pathology literature. The first of them is the concept of staining normalisation.

The first attempts to normalise stain intensity between histology slides relied on the process of stain deconvolution. Stain deconvolution is the transformation of a tissue image from the RGB colourspace into a representation in which each image channel encodes the proportion of a particular histological stain [23]. Stain deconvolution can be used directly for stain content quantification in each pixel, in stains which conform to the Beer-Lambert law, i.e. those in which the relation between the stain concentration and light absorbance is linear.

To perform stain deconvolution (SD) on a digital slide, one needs to know the colour fingerprints of the particular pure stains, represented as stain vectors in the RGB colourspace. A stain vector can be measured directly for a given combination of stain and tissue scanner or alternatively, it can be estimated from a stained digital slide. Such an estimated stain vector can then be applied to deconvolve the stain channels of a histology image and to perform a colour normalisation between slides [39]. Deconvolution-based colour normalisation has been successfully used to correct for minor stain representation differences between WSI scanners from different vendors [21, 40].

There are several known limitations of applying stain deconvolution to whole-slide images. Firstly, WSI scanners typically make use of poly-chromatic illumination, as opposed to white light. The assumption of SD is that the light is white, which can lead to significant deconvolution errors [41]. Furthermore, automated stain vector estimation is known to fail in slides which contain only very small areas stained with a particular stain.

From the SD results presented in our study, it can be seen that stain-deconvolved slides (Figure 3) still need to be further processed before they can be used for collagen quantification. Especially regions of non-specific

stain binding in the tissue (e.g. cell nuclei) need to be identified and excluded. This means that the computed stain distribution map needs to be either binarised using a stain intensity threshold or segmented using a deep-learning approach. Performing colour deconvolution adds an additional step into the analysis, which needs to be justified given the high performance of the state-of-the-art deep-learning models. Our results demonstrate that even a relatively simple deep learning network, such as the U-Net, can compensate for the use of different scanners and even major differences in sample colour, as long as sufficiently varied data is used for model training.

Independently of stain deconvolution, the concept of colour-transfer has been developed. The first methods of automatic colour transfer between images were developed in the early 21st century using simple statistical approaches, such as image histogram transfer [42]. Subsequently, deep-learning based methods were developed to achieve the same goal. Sparse autoencoders have been successfully applied to normalise colours between H&E slides without the need for estimating a stain matrix [43]. Finally, the introduction of generative autoencoders and the CycleGAN model [44] in particular has opened a new direction for digital pathology research.

Autoencoders are a type of artificial neural network which can learn a compressed representation of an input such as an image in an unsupervised manner. An autoencoder is built of two main parts; an encoder which encodes the compressed data representation and a decoder which reconstructs the original input. The whole autoencoder is trained using a process called backpropagation in which the network aims to minimise the difference between the input and the output.

Generative autoencoders are a special type of autoencoders which have the ability to generate new data similar to the received input data. In addition to an encoder and a decoder, these networks additionally contain a module called a generator, which accepts random noise as an input and is trained to generate new data which resembles the original input.

Generative autoencoders have been successfully applied to normalising between different variants of H&E staining and different scanners [45]. Similar models have enabled the generation of realistically looking coloured images from unstained tissue slides [46]. The concept has been further extended to predict the distribution of specialist stains from the basic H&E-stained slide [47].

Most recent deep-learning models are capable of creating very realistic looking colour-transformations between different stainings [48]. However, it

doesn't mean that such simulated images carry biologically meaningful and clinically usable information.

The use of special stainings in pathology workflows enables the pathologist to examine the distribution of substances (such as glycogen, tissue iron or viral antigens) which cannot be seen in the basic H&E staining. While such histological generated stain-like images may appear realistic to an untrained eye, they shouldn't be used to infer information which isn't embedded in the source image.

6. Conclusions

In this study, we introduced a novel cohort of PSR-stained liver biopsies obtained from four distinct studies across multiple healthcare institutions. Our analysis revealed significant and previously unseen differences in histological staining protocols among these institutions, specifically highlighting the variability in PSR staining compared to more standardised methods like H&E staining. This variability poses a challenge for the standardisation and application of digital pathology.

Our findings indicate that differences in CPA quantification results among pathologists are influenced by the variability in collagen staining and the high subjectivity of manual assessments. However, we suggest that the primary limitations in CPA quantification stem from the fractal-like shape of collagen fibres, lengthy object boundaries, partial-volume effects in the z-axis, and the imaging resolution of digitised slides.

We successfully applied a U-Net based semantic segmentation model to identify tissue collagen in a large multi-centre cohort. This study is unique in its scope, encompassing four independent groups of participants with samples processed using different staining protocols, resulting in diverse tissue specimen colours. While this study treated the slides as equivalent in staining, it is essential to recognise that this assumption may not hold true in practice.

We employed a Bayesian inference framework to estimate the uncertainty in the predicted collagen maps and quantified collagen areas. This method provides uncertainty estimates at the pixel and tile levels, though these estimates likely underestimate the true uncertainty of the entire collagen percentage measurement process, which is also influenced by errors in preceding tissue processing steps.

Despite its limitations, our approach demonstrates significant utility. The uncertainty estimates align broadly with previously published literature, and the separation of segmentation uncertainty into aleatoric and epistemic components provides valuable insights. These uncertainty maps offer pathologists a direct interpretation of model performance on different sample regions, particularly highlighting out-of-distribution data such as artefacts and novel staining colours. This approach can also indicate when additional training may be necessary for reliable collagen segmentation in new cohorts.

Previous advancements in stain transfer allow to generate realistic histology images with various staining patterns. However, these innovations have not substantially improved the quantification of stain-positive regions in clinical data, as simulated stains may not accurately reflect the biological state of the tissue. Our study demonstrates that a standard U-Net model is sufficient for this task, and we emphasise the importance of providing measurement error estimates.

Without such error bounds, quantitative results may lack a meaningful interpretation. Careful application of these estimates is crucial for diagnostic purposes. When CPA is measured from human annotations, the error is typically estimated via annotator disagreement. We propose that an ensemble of deep-learning segmentation models can generate likely annotations, modelling the distribution of possible quantification results. Similar uncertainty estimations should be applied to all stain quantification tasks in modern digital pathology, including immunohistochemistry (IHC) stain quantification.

In conclusion, our study highlights the need for standardized staining protocols and the integration of uncertainty estimation methods to enhance the reliability and comparability of digital pathology measurements. These advancements are essential for the broader adoption and clinical application of CPA and other histological metrics in liver fibrosis and beyond.

7. Acknowledgements

The computational aspects of this research were supported by the Wellcome Trust Core Award Grant Number 203141/Z/16/Z and the NIHR Oxford BRC. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

We are indebted to all the members of the clinical and research teams in centres who have recruited to UK-AIH and are key to the ongoing success of the study. The principal investigators and recruiting centres are:

Dr Shirley English (Aberdeen Royal Infirmary), Dr Graeme Alexander/Dr George Mells (Addenbrooke's Hospital), Dr Debabrata Majumdar (St Peters Hospital), Dr Vinay Sathyanarayana (Barnsley Hospital), Professor John Ramage (Basingstoke North Hampshire Hospital), Dr Christopher Shorrock (Blackpool Victoria Hospital), Dr James Maggs (Buckinghamshire Hospital), Dr David Elphick (Chesterfield Royal Hospital), Dr Chris Macdonald (Cumberland Infirmary), Professor Matthew Cramp (Derriford Hospital), Dr Joanne Sayer (Doncaster Royal Infirmary), Dr James Jupp (Dorset County Hospital), Dr Jessica Dyson (Freeman Hospital), Dr Coral Hollywood (Gloucestershire Royal Hospital), Dr Alexandra Daley (Heartlands Hospital), Dr Lynsey Corless (Hull Royal Infirmary), Dr Darren Craig (James Cook University Hospital), Dr Emma Culver (John Radcliffe Hospital, Oxford), Professor Michael Heneghan (King's College Hospital), Dr Sharat Misra (King's Mill Hospital), Dr Chris Corbett (New Cross Hospital), Professor John Dillon (Ninewells Hospital), Dr Simon Rushbrook (Norfolk and Norwich University Hospital), Dr Thomas Lee (North Tyneside General Hospital), Dr Nicholas M Sharaer (Poole Hospital), Dr Kara Rye (Princess Royal Hospital), Dr Andrew Fowell (Queen Alexandra Hospital, Portsmouth), Dr Andrea Broad/Dr Dina Mansour (Queen Elizabeth Hospital, Gateshead), Dr Andy Douds (Queen Elizabeth Hospital, King's Lynn), Dr Stephen Ryder (Queen's Medical Centre), Dr Richard Keld (Royal Albert Edward Infirmary), Dr Earl Williams (Royal Bournemouth Hospital), Dr William Stableforth (Royal Cornwall Hospital), Dr Andrew Austin (Royal Derby Hospital), Professor Dermot Gleeson (Royal Hallamshire Hospital), Dr Kenneth Simpson (Royal Infirmary of Edinburgh), Dr Imran Patanwala (Royal Liverpool University Hospital), Dr Alison Brind (Royal Stoke University Hospital), Dr Shanika de Silva (Russells Hall Hospital), Dr Aqueel Jamil (Salisbury District Hospital), Dr Saket Singhal (Sandwell General Hospital), Dr Chin Lye Ch'ng (Singleton Hospital), Dr Joanne Topping (South Tyneside District Hospital), Dr Mark Wright (Southampton General Hospital), Dr Talal Valliani (Southmead Hospital), Dr Rebecca Jones (St. James's University Hospital, Leeds), Dr Harriet Mitchison (Sunderland Royal Hospital), Dr Douglas Thorburn (The Royal Free Hospital), Professor Aftab Ala (The Royal Surrey County Hospital), Dr Ye Htun Oo (University Hospital Birmingham NHS Foundation Trust), Dr Sushma Sakena/Dr Francisco Porras-Perez (University Hospital of North Durham), Prof Jane Metcalf/Dr Stephen Mitchell (University Hospital of North Tees), Dr Esther Unitt/Dr Victoria Gordon (University Hospitals Coventry Warwick), and Dr Jeremy Shearman (Warwick Hospital, South

Warwickshire NHS Foundation Trust).

JKD is supported by the NIHR Newcastle Biomedical Research Centre.

Declaration of funding interests: The UK-AIH study is funded by Liver-North (a national patient charity) and previously by the NIHR Rare Diseases Translational Research Collaboration. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

8. Code Availability

The source code for the slide colour analysis tool is available under: github.com/mkatw/slide_colour_palette.

Appendix A. Appendix

Appendix A.1. Processing time

Table A.8 shows a comparison of the training and inference times for each of the tested models. The times are reported for the processing of the training set (training over 30 epochs) and of the validation set, respectively. It can be seen that tile processing time is strongly related to the size of the network used, with U-Net Tiny outperforming other models in our study. Processing time is an important factor to consider when choosing deep learning models for use in clinical practice.

Table A.8: **Training and inference time comparison for the trained models** (M = 10 ensemble average ± standard deviation).

Model	Training [s]	Inference [s]
U-Net Tiny	417.7 ± 16.0	1.3 ± 0.2
U-Net Mini	1358.8 ± 24.5	2.5 ± 0.6
Att U-Net	1590.4 ± 37.0	3.3 ± 0.8

Appendix A.2. Correlation statistics

Figures A.12 and A.13 show the correlation statistics for computed collagen areas (the average prediction P and the segmented collagen area B), as well as uncertainties for the models trained on the individual cohorts and the model trained on the pooled cohort, respectively.

Figure A.12: **Correlation coefficients and corresponding statistical significance values for the prediction parameters of the model trained on the individual cohorts.** The values are reported for predictions obtained using the U-Net Mini ensemble.

Figure A.13: **Correlation coefficients and corresponding statistical significance values for the prediction parameters of the model trained on the pooled cohort.** The values are reported for predictions obtained using the U-Net Mini ensemble.

References

[1] Roberta Forlano et al. "NAFLD: Time to apply quantitation in liver biopsies as endpoints in clinical trials". In: *Journal of Hepatology* 74.1 (2021), pp. 241–242.

[2] Elena Buzzetti et al. "Collagen proportionate area is an independent predictor of long-term outcome in patients with non-alcoholic fatty liver disease". In: *Alimentary Pharmacology and Therapeutics* 49.9 (2019), pp. 1214–1222.

[3] Mads Israelsen et al. "Collagen proportionate area predicts clinical outcomes in patients with alcohol-related liver disease". In: *Alimentary Pharmacology and Therapeutics* 52.11-12 (2020), pp. 1728–1739.

[4] Amanda Cheung et al. "Defining Improvement in Nonalcoholic Steatohepatitis for Treatment Trial Endpoints: Recommendations From the Liver Forum". In: *Hepatology* 70.5 (2019), pp. 1841–1855.

[5] Alexandros Arjmand et al. "Quantification of liver fibrosis-a comparative study". In: *Applied Sciences (Switzerland)* 10.2 (2020), pp. 1–34.

[6] Qiang Li et al. "Virtual liver needle biopsy from reconstructed three-dimensional histopathological images: Quantification of sampling error". In: *Computers in Biology and Medicine* 147 (Aug. 2022).

[7] Timothy J. Kendall et al. "Hepatic elastin content is predictive of adverse outcome in advanced fibrotic liver disease". In: *Histopathology* 73.1 (2018), pp. 90–100.

[8] Holde Puchtler and Faye Sweat. "Histochemical specifity of staining methods for connective tissue fibers: Resorcin-fuchsin and van Gieson's picro-fuchsin". In: *Histochemie* 4 (1964), pp. 24–34.

[9] R. W. Dapson et al. "Certification procedures for sirius red F3B (CI 35780, Direct red 80)". In: *Biotechnic and Histochemistry* 86.3 (2011), pp. 133–139.

[10] Stéphane Bancelin et al. "Determination of collagen fibril size via absolute measurements of second-harmonic generation signals". In: *Nature Communications* 5 (2014), pp. 1–8.

[11] Seyed Mohammad Siadat et al. "Measuring collagen fibril diameter with differential interference contrast microscopy". In: *Journal of Structural Biology* 213.1 (2021), p. 107697.

[12] Benoit B Mandelbrot. *The fractal geometry of nature*. eng. [Rev. ed.] New York: W.H. Freeman, 1983.

[13] Nicola Dioguardi et al. "Liver fibrosis and tissue architectural change measurement using fractal-rectified metrics and Hurst's exponent". In: *World Journal of Gastroenterology* 12.14 (2006), pp. 2187–2194.

[14] Ender Gunes Yegin, Korkut Yegin, and Osman Cavit Ozdogan. "Digital image analysis in liver fibrosis : basic requirements and clinical implementation". In: *Biotechnology & Biotechnological Equipment* 30.4 (2016), pp. 653–660.

[15] Cyrill Matenaers et al. "Practicable methods for histological section thickness measurement in quantitative stereological analyses". In: *PLoS ONE* 13.2 (2018), pp. 1–21.

[16] Stuart Astbury et al. "Reliable computational quantification of liver fibrosis is compromised by inherent staining variation". In: *Journal of Pathology: Clinical Research* 7.5 (2021), pp. 471–481.

[17] Natasha Mcdonald et al. "Multiparametric magnetic resonance imaging for quantitation of liver disease : a two-centre cross- sectional observational study". In: *Scientific Reports* February (2018), pp. 1–10.

[18] Damian J. Mole et al. "Study protocol: HepaT1ca - An observational clinical cohort study to quantify liver health in surgical candidates for liver malignancies". In: *BMC Cancer* 18.1 (2018), pp. 1–7.

[19] Stephen A. Harrison et al. "Prospective evaluation of the prevalence of non-alcoholic fatty liver disease and steatohepatitis in a large middle-aged US cohort". In: *Journal of Hepatology* 75.2 (2021), pp. 284–291.

[20] Jessica K. Dyson et al. "Inequity of care provision and outcome disparity in autoimmune hepatitis in the United Kingdom". In: *Alimentary Pharmacology and Therapeutics* 48.9 (2018), pp. 951–960.

[21] Abhishek Vahadane et al. "Structure-preserved color normalization for histological images". In: *Proceedings - International Symposium on Biomedical Imaging* 2015-July (2015), pp. 1012–1015.

[22] Adam Goode et al. "OpenSlide: A vendor-neutral software foundation for digital pathology". In: *Journal of Pathology Informatics* 4.1 (2013), p. 27.

[23] A. C. Ruifrok and D. A. Johnston. "Quantification of histochemical staining by color deconvolution". In: *Analytical and Quantitative Cytology and Histology* 23.4 (2001), pp. 291–299.

[24] Peter Bankhead et al. "QuPath: Open source software for digital pathology image analysis". In: *Scientific Reports* 7.1 (2017), p. 16878.

[25] Paul Heckbert. "Color Image Quantization for Frame Buffer Display". In: *Computer Graphics* 16.3 (1982), pp. 157–166.

[26] Nobuyuki Otsu. "A Threshold Selection Method from Gray-Level Histograms". In: *IEEE Transactions on Systems, Man, and Cybernetics* C.1 (1979), pp. 62–66.

[27] Stéfan Van Der Walt et al. "Scikit-image: Image processing in python". In: *PeerJ* 2014.1 (2014), pp. 1–18.

[28] Olaf Ronneberger, Philipp Fischer, and Thomas Brox. *U-Net: Convolutional Networks for Biomedical Image Segmentation*. Ed. by Nassir Navab et al. Vol. 9351. Lecture Notes in Computer Science. Cham: Springer International Publishing, 2015, pp. 234–241.

[29] Ozan Oktay et al. "Attention U-Net: Learning Where to Look for the Pancreas". In: Midl (2018). arXiv: 1804.03999.

[30] Yarin Gal and Zoubin Ghahramani. "Dropout as a Bayesian Approximation: Representing Model Uncertainty in Deep Learning". In: *Proceedings of The 33rd International Conference on Machine Learning*. Ed. by Maria Florina Balcan and Kilian Q. Weinberger. Vol. 48. Proceedings of Machine Learning Research. New York, New York, USA: PMLR, 2016, pp. 1050–1059.

[31] Alex Kendall and Yarin Gal. "What Uncertainties Do We Need in Bayesian Deep Learning for Computer Vision?" In: *Advances in Neural Information Processing Systems*. Ed. by I. Guyon et al. Vol. 30. Curran Associates, Inc., 2017.

[32] Radford M Neal. *Bayesian Learning for Neural Networks*. 1996.

[33] Alex Graves. "Practical Variational Inference for Neural Networks". In: *Advances in Neural Information Processing Systems*. Ed. by J. Shawe-Taylor et al. Vol. 24. Curran Associates, Inc., 2011.

[34] Balaji Lakshminarayanan, Alexander Pritzel, and Charles Blundell. "Simple and Scalable Predictive Uncertainty Estimation Using Deep Ensembles". In: *Proceedings of the 31st International Conference on Neural Information Processing Systems*. NIPS'17. Long Beach, California, USA: Curran Associates Inc., 2017, pp. 6405–6416.

[35] Armen Der Kiureghian and Ove Ditlevsen. "Aleatory or epistemic? Does it matter?" In: *Structural Safety* 31.2 (2009). Risk Acceptance and Risk Communication, pp. 105–112.

[36] Yongchan Kwon et al. "Uncertainty quantification using Bayesian neural networks in classification: Application to biomedical image segmentation". In: *Computational Statistics and Data Analysis* 142 (2020), p. 106816.

[37] Stanislav Fort, Huiyi Hu, and Balaji Lakshminarayanan. "Deep Ensembles: A Loss Landscape Perspective". In: (2019), pp. 1–15. arXiv: `1912.02757`.

[38] Shuoyu Xu et al. "QFibrosis: A fully-quantitative innovative method incorporating histological features to facilitate accurate fibrosis scoring in animal model and chronic hepatitis B patients". In: *Journal of Hepatology* 61.2 (2014), pp. 260–269.

[39] Marc Macenko et al. "A method for normalizing histology slides for quantitative analysis". In: *Proceedings - 2009 IEEE International Symposium on Biomedical Imaging: From Nano to Macro, ISBI 2009* (2009), pp. 1107–1110.

[40] Adnan Mujahid Khan et al. "A nonlinear mapping approach to stain normalization in digital histopathology images using image-specific color deconvolution". In: *IEEE Transactions on Biomedical Engineering* 61.6 (2014), pp. 1729–1738.

[41] Peter Haub and Tobias Meckel. "A model based survey of colour deconvolution in diagnostic brightfield microscopy: Error estimation and spectral consideration". In: *Scientific Reports* 5.February (2015), pp. 1–15.

[42] Erik Reinhard et al. "Color transfer between images". In: *IEEE Computer Graphics and Applications* 21.5 (2001), pp. 34–41.

[43] Andrew Janowczyk, Ajay Basavanhally, and Anant Madabhushi. "Stain Normalization using Sparse AutoEncoders (StaNoSA): Application to digital pathology". In: *Computerized Medical Imaging and Graphics* 57 (2017), pp. 50–61.

[44] Jun-yan Zhu et al. "Unpaired Image-to-Image Translation using Cycle-Consistent Adversarial Networks". In: (2017). arXiv: `1703.10593`.

[45] M. Tarek Shaban et al. "Staingan: Stain style transfer for digital histological images". In: *Proceedings - International Symposium on Biomedical Imaging* 2019-April (2019), pp. 953–956.

[46] Yair Rivenson et al. "PhaseStain: the digital staining of label-free quantitative phase microscopy images using deep learning". In: *Light: Science and Applications* 8.1 (2019). arXiv: `1807.07701`.

[47] Kevin de Haan et al. "Deep learning-based transformation of H&E stained tissues into special stains". In: *Nature Communications* 12.1 (2021), pp. 1–13.

[48] Nicolas Wagner et al. *Federated Stain Normalization for Computational Pathology*. Vol. 1. Springer Nature Switzerland, 2022, pp. 14–23.