Title: Geospatial Accessibility to Primary and Community Health Centres in India

Short title: PHC CHC access India

Authors: Shreyas Patil[1,2], Aamir Miyajiwala[1], Anoushka Arora[1,3], Ankit Raj[1], Harsh

Thakkar[1,], Mukta S. Takalikar[4], Siddhesh Zadey[1,5]

Affiliations:

1 Association of Socially Applicable Research (ASAR), Pune, Maharashtra, India

2 Department of Community Medicine and Family Medicine, All India Institute of Medical

Sciences, Bibinagar, Telangana, India

3 Smt. Nathiba Hargovandas Lakhmichand Municipal Medical College, Ahmedabad, Gujarat,

India

4 Department of Computer Engineering, SCTR's Pune Institute of Computer Technology,

Pune, Maharashtra, India

5 Dr. D. Y. Patil Dental College, Hospital, and Research Centre, Dr. D. Y. Patil Vidyapeeth,

Pune, Maharashtra, India

Corresponding Author: Siddhesh Zadey BSMS, MScGH;

ASAR Office Address: D2 Sai Heritage, New DP Road, Aundh, Pune, Maharashtra, India

411007.

Email: sidzadey@asarforindia.org

Article type: Original Research Article

Abstract

Background: Accessibility to public healthcare in India is poor. Data on access to public health facilities in terms of travel time and geographical scope are limited.

Methods: Primary health centre (PHC) and community health centre (CHC) locations were taken from the Geographic Information System dataset of the Pradhan Mantri Gram Sadak Yojana 2021. Motorized and walking travel-time friction surface rasters were taken from the Malaria Atlas Project 2020. We examined the densities, travel times, and Access Population Coverage (APC) for PHCs and CHCs in India up to the district level. For walking APCs, we used a 30-minute time threshold for PHCs and a 60-minute threshold for CHCs. For motorized APCs, we used a 60-minute threshold for PHCs and a 120-minute threshold for CHCs.

Results: In 2021, India had 23.22 PHCs and 3.07 CHCs per million people. Median PHC travel time was 79.53 (walking) and 16.77 minutes (motorised transport). CHC travel time was 227.53 (walking) and 36.23 minutes (motorised transport). PHC APCs for walking and motorised transport were 18.69% and 97.97%, respectively. CHC APCs for walking and motorised transport were 10.89% and 98.88%, respectively. Wide variations were seen in rural-urban analysis.

Conclusion: Our study found marked heterogeneity in timely geographic access, with access by walking being better in rural India and by motorised transport being better in urban India.

Keywords: Primary Healthcare, Secondary Healthcare, Geographic Access, Population Coverage, Motorized Transport, Travel Time

1 Background

India is currently the most populous country in the world, being home to over 1.4 billion individuals.(1) Safeguarding the health of this huge population has several challenges and requires immense resources with their commensurate distribution. Geographical factors like the size of the country, the 7th largest in the world, affect the distribution of these resources. Variations in terrain exacerbate these difficulties, particularly in hilly or mountainous regions.(2) The Alma-Ata declaration propounded the goal of "Health for All" recognising the importance of primary health care in attaining this objective.(3) In India, primary health care is rendered mainly by the primary health centres (PHCs), the first point of contact of the community with a trained medical officer. Above the PHCs in the continuum of care are the community health centres (CHCs), which are the first level of referral and provide specialist care. The Indian Public Health Standards (IPHS) 2022 recommend the availability of one PHC for every 50,000 people in urban areas, every 30,000 people in rural areas, and every 20,000 people in tribal/hilly areas. The recommendations for CHCs are that one facility must be available for every 80,000 people in tribal/hilly areas and for every 120,000 people in rural areas. For urban areas, the recommended population norms are one CHC for every 500,000 people in metro areas and every 250,000 people in non-metro regions.(4)

The number of health facilities in terms of a defined population size is referred to as the density of health facilities and is a measure of availability. On the other hand, accessibility is a spatial measure of 'closeness' or geographical proximity with an element of time.(5,6) A mere availability of health facilities will not fulfil the health needs of a population if geographic access is difficult due to reasons such as distance, terrain, or costs. Geographical distance and travel times are significant barriers to accessibility, universalization of health care, and thus utilisation of health services, particularly in rural settings.(6–13) Moreover,

timely geographic access to healthcare is essential to ensure optimal health outcomes.(14–16)

Longer travel times have a harmful influence on child mortality rates.(17,18) They are also

associated with an increased risk of stillbirths, maternal morbidity, and mortality.(19–22)

Increased distances to healthcare facilities can also negatively impact neonatal and maternal

outcomes by decreasing the probability of women undergoing institutional delivery.(23,24)

Distances do not necessarily indicate geographic access better due to various factors such as

terrain. Time taken to travel distances, therefore, serves as a more accurate measure.

Furthermore, longer travelling times impose additional burdens like higher travel costs and

longer loss of wages.(25–27) Additionally, access, geography, and timeliness have been

identified as forms of moral capital when healthcare systems are viewed as moral economies.

Taking these into consideration can improve healthcare systems by making them more

equitable, effective, and efficient.(28)

It is estimated that 44.2% or 3.16 billion people around the globe would not be able to access

healthcare facilities within one hour if they travel on foot. This figure drops to 8.9% or 646

million people with access to motorised transport.(29) Other studies in low- and middle-

income countries (LMIC) settings have also shown that a significant share of individuals are

not able to access healthcare in a timely manner.(30,31) Travel times are particularly higher

in rural areas, with motorised transport availability and ownership improving access.(32–35)

Similarly, in India, it has been observed that public healthcare accessibility is low in rural

areas, particularly for those with no access to motorised transport. (36) However, studies on

geographic access to public healthcare facilities in terms of travel time in the Indian context

are few and limited in geographical breadth. The current study aims to estimate the travel

times and the proportion of the population with timely geographic access to PHCs and CHCs

in India.

2 Methods

2.1 Data sources

To comprehensively map the densities of PHCs and CHCs and their travel times, the study

necessitated collecting, extracting, and collating data from multiple sources. The geolocation

data for the PHCs and CHCs were extracted from the Geographic Information System (GIS)

dataset of the Pradhan Mantri Gram Sadak Yojana (PMGSY).(37) The PMGSY is a wholly

centrally sponsored scheme by the Government of India (GoI) to provide all-weather

connectivity to unconnected habitations as a poverty alleviation strategy. We utilized the 2021

PMGSY GIS dataset and extracted geo-coordinates and other related geographical

information for the health facilities under study, i.e., the PHCs and CHCs. PMGSY GIS data

on PHCs and CHCs were not available for 7 states/UTs, i.e., Andaman & Nicobar Islands,

Chandigarh, Dadra & Nagar Haveli and Daman & Diu (DNH and DD), Delhi, Lakshadweep,

Puducherry, and Goa. At the district level, data on PHCs and CHCs were missing for 60 and

134 districts, respectively. The Malaria Atlas Project (MAP) data for the year 2019 were used

to obtain the motorized and walking friction rasters for every square kilometre (1 km2).(38)

GADM v3.6 data were used to delineate the administrative boundaries of India, while high-

resolution (1 sq. km.) United Nations (UN) adjusted population counts from the WorldPop

dataset for 2020 for India served as our source of population data.(39,40) The use of global

data sources such as the MAP for travel time friction surfaces ensures that our findings are

comparable to studies using these data sources from other countries and multi-country

assessments. Validation of MAP-derived travel time rasters for certain geographies, including

India, has been presented in original sources.(29,41)

2.2 Outcomes

The study had three outcomes. The first outcome was to calculate the density of health

facilities, i.e., the number of PHCs and CHCs for every million population in India. The

second outcome was to determine the median travel time to the closest PHC and CHC, from

each 1 km2 raster in India. The third outcome was to estimate the proportion of the population

with timely geographic access, also called Access Population Coverage (APC), to a PHC and

a CHC. By walk, timely geographic access was defined as access within 30 minutes for PHCs

and within 60 minutes for CHCs. For motorised transport, timely geographic access meant

access within 60 minutes for PHCs, and 120 minutes for CHCs. These outcomes were

calculated for India, 29 states and union territories (UTs), and 675 districts for PHCs and 601

districts for CHCs, and their urban and rural areas as well.

2.3. Data Analysis

For geocoding of the primary and secondary healthcare facilities (PHCs and CHCs),

addresses were cleaned manually for improved machine readability. The Google Maps'

application programming interface (API) and the 'Awesome Table' add-on for Google Sheets

for machine or API-based geocoding were used. For locations that returned multiple sets of

coordinates, the ones with the most relevant address string were chosen. Geo-coordinates

were used to identify and remove duplicates and points extending beyond the latitude and

longitude limits of India.

An implementation of the Dijkstra algorithm was utilised to compute the minimum time

required to traverse the friction surface from every 1 km^2 pixel (grid cell) on the map to the

geo-coordinates of every PHC and CHC. The algorithm was implemented for two modes of

transport, i.e., walking and motorised transport. The proportion of the population with access

to a PHC within 30 minutes by walking and 60 minutes by motorised transport, and the

proportion with access to a CHC within 60 minutes by walking and within 120 minutes by

motorised transport were the proxy variables for timely geographic access.

A binary accessibility raster was created with '1's for pixels that fulfilled the timeliness

criterion of each proxy variable and '0's for cases otherwise. This raster was then overlaid on

the population raster (extent matched). The population figures at each pixel were multiplied

with the weights (i.e., 1s and 0s) to get the population number with timely geographic access.

Proportions or percentages of the population with timely geographic access were obtained by

dividing these numbers by the overall population.

Although PMGSY GIS data on PHCs and CHCs were missing for seven states/UTs and for a

subset of 134 districts for CHC and 60 districts for PHC, these areas were retained during the

computation of travel time and APC to preserve spatial continuity and consistency across

geographies. However, to avoid reporting estimates based on incomplete facility data, all

districts and states/UTs with missing PHC/CHC locations were explicitly masked in the

outputs. Specifically, their values were set to NA, they were removed from all tabular

summaries, and they were greyed out in all maps. As a result, no district- or state-level

estimates are reported or interpreted for these areas, and they do not contribute to any

reported summary statistics.

The workflow has been described in Figure 1. The same workflow has been discussed for

investigating healthcare accessibility under the project, IndoHealMap.(42–47) Analysis was

performed in R (Version 4.3.3).

3 Results

3.1 Density of healthcare facilities

3.1.1 Primary Health Centres (PHCs)

According to the PMGSY GIS 2021 dataset, there were 31,958 PHCs in India with a national

average density of 23.22 centres per million population (cpm). Among states/UTs,

Maharashtra (3445) had the highest number of PHCs (3445) while the UT of Ladakh (31) had

the lowest number of PHCs. The density of PHCs exceeded the national average in 25/29

(86.21%) states/UTs. The density of PHCs among states/UTs ranged from 8.46 cpm in West

Bengal to 124.88 cpm in the state of Arunachal Pradesh. In rural areas, the density of PHCs

ranged from 10.62 in West Bengal to 130.18 cpm in Arunachal Pradesh. In urban areas, the

density ranged from 0 in 3 states (Arunachal Pradesh, Mizoram, and Ladakh) to 38.77 in

Manipur. At the district level, the number of PHCs varied from 1 in Khagaria, Bihar, to 386

in Pune, Maharashtra. PHC density at the district level surpassed the national average in

391/675 (57.93%), and ranged from 0.49 cpm in Khagaria, Bihar, to 638.23 cpm in Anjaw,

Arunachal Pradesh (**Figure 2**).

3.1.2 Community Health Centres (CHCs)

The total number of CHCs at the national level was 4237 centres. Among states/UTs, the

number varied from 2 centres in Sikkim to 760 centres in Uttar Pradesh. The national average

density of CHCs was estimated to be 3.07 cpm. The density of CHCs was found to be better

than the national average in 16/29 (55.17%) states/UTs. The density of CHCs ranged from

0.16 cpm in West Bengal to 36.25 cpm in Arunachal Pradesh. In rural areas, the density

ranged from 0.13 cpm in West Bengal to 37.2 in Arunachal Pradesh. In urban areas, the

density ranged from 0 in Ladakh to 14.3 in Arunachal Pradesh. At the district level, the

number of CHCs varied from 1 in 71/601 (11.83%) districts to 95 in Ahmednagar,

Maharashtra. The density at the district level exceeded the national average in 353/601

(58.74%). The lowest and the highest density of PHCs were 0.08 cpm in South Twenty Four

Parganas and 127.64 cpm in Anjaw, Arunachal Pradesh **(Figure 2)**.

3.2 Travel time to the nearest PHC

3.2.1 By walking

At the national level, the median time taken to access a PHC by walking was found to be

79.53 (IQR: 45.88-134.00) minutes. The travel time was shorter than the national median in

15/29 (51.72%) states/UTs. The travel times ranged from 47.95 (IQR: 28.86- 73.05) minutes

in Punjab to 852.32 (IQR: 477.10-1338.35) minutes in the UT of Ladakh. At the district level,

409/675 (60.59%) districts had a shorter travel time by walking compared to the national

value. The walking time to a PHC was the shortest in Nalanda, Bihar [25.06 (IQR: 14.97-

41.02) minutes] and the longest in Leh, Ladakh [858.47 (IQR: 501.86-1359.10) minutes]. The

travel times for rural and urban regions are summarized in **Table 1.**

3.2.2 By motorised transport

At the national level, the median time taken to access a PHC by motorized transport was

found to be 16.77 (IQR: 7.79-35.10) minutes. Among states/UTs, 13/29 (44.82%) had a

shorter travel time compared to the national median. The state of Kerala had the shortest

median travel time, 6.39 (IQR: 3.56-13.64) minutes, and the UT of Ladakh had the longest

time, 311.80 (IQR: 126.65- 601.51) minutes. At the district level, the travel time was shorter

than the national median in 413/675 (61.18%) districts. The travel time ranged from 2.68

(IQR: 1.48- 28.98) minutes in Pulwama, Jammu & Kashmir, to 604.11 (IQR: 318.51- 650.93)

minutes in Anjaw, Arunachal Pradesh. The travel times for rural and urban regions are

summarized in **Table 1.**

3.3 Median travel time to the nearest CHC

3.3.1 By walking

At the national level, the median time to reach the nearest CHC by walk was found to be

227.53 (IQR: 130.80-384.07) minutes. At the state/UT level, 14/29 (48.27%) had a shorter

travel time compared to the national median. The median travel time varied from 107.40

(IQR: 64.04-183.95) minutes in Kerala to 1774 (IQR: 1028.23-2513.59) minutes in the UT of

Ladakh. The median travel time was shorter than the national value in 399/601 (66.39%)

districts. The median time on foot was found to be the shortest in the Bongaigaon district of

Assam at 61.63 (IQR: 37.22-92.41) minutes and the longest in the Leh district [2059.95 (IQR:

1488.33-2707.40) minutes] of Ladakh. The travel times for rural and urban regions are

summarized in **Table 1.**

3.3.2 By motorised transport

At the national level, the median time to access a CHC by motorised transport was estimated

to be 36.23 (IQR: 20.22-63.92) minutes. The travel time was shorter than the national median

in 14/29 (48.27%) states/UTs. The median time ranged from 12.66 (IQR: 7.44-25.28) minutes

in Kerala to 396.83 (IQR: 218.94- 682.48) minutes in Ladakh. The median travel time was

shorter than the national value in 395/601 (65.72%) districts. The median time in districts

varied from 7.58 (IQR: 5.03-10.49) minutes in Kottayam, Kerala, to 1008.83 (IQR: 437.32-

1560.25) minutes in Kurung Kumey of Arunachal Pradesh. The travel times for rural and

urban regions are summarized in **Table 1.**

.4 Access Population Coverage (APC) for PHCs

3.4.1 By walking

By walking, timely geographic access to a PHC meant being able to access one within 30

minutes. At the national level, the APC was estimated to be 18.69%. At the state/UT level,

14/29 (48.28%) had an APC greater than the national median. APC ranged from 5.2% in

Mizoram to 43.71% in Manipur. Among districts, 297/675 (44%) had an APC greater than

the national average, and greater than 50% in 18/675 (2.66%) districts. APC varied from

0.78% in Upper Siang, Arunachal Pradesh, to 69.68% in Bishnupur, Manipur **(Figure 3)**.

APC was less than 50% in all states/UTs. The APCs for urban and rural regions have been

summarized in **Table 2**. APC was less than 50% in all rural and urban areas except urban

Manipur (69.3%). APC was 0% in urban Mizoram.

3.4.2 By motorised transport

By motorised transport, timely geographic access to a PHC meant being able to access one

within 60 minutes. At the national level, the APC was estimated to be 97.97%. At the

state/UT level, 13/29 (44.83%) states/UTs had an APC greater than the national average. APC

was greater than 50% in 28/29 (96.55%) states/UTs, and ranged from 44.70% in Ladakh to

99.98% in Haryana. APC was greater than the national average in 382/675 (56.59%) districts

and greater than 50% in 664/675 (98.37%) districts. APC ranged from 13.39% in Anjaw,

Arunachal Pradesh, to 100% in 64/675 (9.48%) districts **(Figure 3)**. The APCs for urban and

rural regions have been summarized in **Table 2.** APC was greater than 50% in urban areas of

all states/UTs. In rural areas, the APC was less than 50% in Ladakh (44.6%).

3.5. Access Population Coverage (APC) for CHCs

3.5.1 By walking

By walking, timely geographic access to a CHC meant being able to access one within 60

minutes. At the national level, the APC was estimated to be 10.89%. At the state/UT level,

13/29 (44.83%) had an APC greater than the national average. APC ranged from 1.5% in

West Bengal to 29.39% for the state of Kerala. Among districts, 277/601 (46.09%) had an

APC greater than the national average, while APC was found to be greater than 50% in 2/601

(0.33%) districts. APC varied from 0% in 2/601 (0.33%) districts to 52.20% in Dhanbad,

Jharkhand **(Figure 3)**. APC was estimated to be 57/731 (7.8%) districts. The APCs for urban

and rural regions have been summarized in **Table 2**. APC was less than 1% in urban

Mizoram.

3.5.2 By motorised transport

By motorised transport, timely geographic access to a CHC meant being able to access one

within 120 minutes. At the national level, the APC was estimated to be 98.88%. Among

states/UTs, APC was greater than the national average in 13/29 (44.83%) states/UTs and

greater than 50% in 28/29 (96.55%) states/UTs, respectively, and ranged from 49.65% in

Ladakh to 99.99 in Haryana. Among districts, 416/601 (69.22%) had an APC greater than the

national average, while APC was found to be greater than 50% in 593/601 (98.67%) districts.

The APCs for urban and rural regions have been summarized in **Table 2.**

4 Discussion

4.1 Summary and interpretation of findings

The main aim of this study was to map timely geographic access to primary and secondary

public healthcare facilities across geographic scales from the national to the district levels in

India. Arunachal Pradesh, Himachal Pradesh, Nagaland, and Ladakh were observed to have

the highest densities of PHCs and CHCs. The lowest densities for PHCs were seen in West

Bengal, Bihar, and Uttar Pradesh, and the lowest densities for CHCs were observed in West

Bengal, Tamil Nadu, and Maharashtra. Travel times by walking were the longest in

states/UTs with difficult terrain, including Arunachal Pradesh, Ladakh, Mizoram, and Sikkim.

Given that rural areas have greater PHCs and CHCs per million than urban areas, urban areas

fared better than rural areas for median travel time, possibly due to a dearth of pedestrian-

friendly infrastructure in rural areas. The average road density for India in 2018-19 was

5296.3 per 1000 km^2 for urban areas in comparison to 1458.1 for rural areas.(48) This could

explain the relatively shorter travel times observed for urban areas by motorised transport.

However, even in the case of motorised transport, travel times were the longest in the

states/UTs of Arunachal Pradesh, Ladakh, Mizoram, and Sikkim. Therefore, it can be seen

that the nature of the terrain in these regions tends to slow down movement in general,

whether on foot or by motorised transport.

Travel time, a summary statistic, cannot showcase a holistic picture of geographic access.

Hence, we looked at access population coverage (APC) as a measure of geographic access.

Geographic access by walking was found to be poor. Overall, approximately 1118 million or

81.31% Indians, 798 million or 79.49% rural Indians, and 320 million or 86.19% urban

Indians were beyond the 30-minute threshold. Similarly, approximately 1225 million or

89.11% Indians, 888 million or 88.42% rural Indians, and 337 million or 90.95% urban

Indians were beyond the 60-minute threshold for CHCs. Geographic access by walking was

low in Arunachal Pradesh, Ladakh, and Mizoram. This could have resulted from the long

travel times in Arunachal Pradesh, Ladakh, and Mizoram. Geographic access by walking was

heterogeneous across states/UTs and districts, but was generally better in rural areas

compared to urban areas, despite the longer travel times. This could be credited to

improvements in the rural health infrastructure brought about by sustained investments over

the last two decades through the National Rural Health Mission (NRHM).(49) On the whole,

these figures depict an alarming situation, as they indicate that hundreds of millions of Indians

lack geographic access to primary and secondary healthcare, resulting in significant

repercussions for their health.

Approximately 28 million Indians, 27 million rural Indians, and 1 million urban Indians were

still beyond the 60-minute threshold by motorised transport for PHCs. For CHCs,

approximately 15 million, 14 million rural Indians, and 1 million urban Indians were beyond

the 120-minute threshold. Although these percentages appear small, they translate into a

sizable number of Indians lacking geographic access and merit a deeper look for action.

Geographic access was on the lower end for Ladakh, Arunachal Pradesh, and Mizoram,

possibly due to the long travel times in these regions. The small administrative areas of these

UTs and well-developed road networks, due to their urban nature, could have facilitated quick

access to primary and secondary healthcare facilities beyond their borders. There was marked

heterogeneity in geographic access by motorised transport across states/UTs and districts, but

it was generally better in urban areas.

4.2 Strengths and limitations

Our study has the following strengths. First, using the PMGSY GIS dataset allowed us to

analyse precisely geocoded locations of primary and secondary healthcare facilities. This

enabled us to comprehensively map travel times and geographic access using APC. Second,

travel times and APCs were estimated for two modes of travel: walking and motorised

transport, and two regions: rural and urban. Third, estimates were calculated up to the district

level, lending further strength to our study.

There were certain limitations as well. First, data related to primary and secondary health

facilities were not available for several states/UTs and districts. Second, our study is also

limited by the lack of information on the functional status of the health facilities. Third, the

estimates for walking do not account for groups like the elderly or disabled, who would lack

access despite geographical and temporal proximity. Fourth, the APCs for motorised transport

assume that everybody has access to it, the unavailability of which would preclude an

individual from accessing a health facility on time. Fifth, the travel time raster data from

MAP is from 2019. India has observed remarkable growth in roadway infrastructure, which

can potentially reduce travel times and improve APC. Hence, our estimates for travel time

estimates are likely higher, and those for APC values are lower than India's current reality. In

other words, we present a conservative picture of timely geographic access. However, we

used the most recent available travel-time raster data and updated facility data. Future studies

can use similar methods to update the estimates, once travel time rasters are available for

2025. Sixth, since we did not have access to resource metrics required to inform model

weights, such as the number of beds, healthcare workers, or surgical volumes, we used a

parsimonious approach instead of advanced models like Floating Catchment Area (FCA) or

Rational Agent Modeling (RAM). Future studies should employ FCA or RAM to assess

measures of geographic access that include resource metrics. Seventh, due to limited data on

time thresholds, we adopted arbitrary thresholds. Most literature in the past has focused on a

distance-based threshold. The Lancet Commission on Global Surgery suggested a threshold

of 120 minutes was associated with reduced maternal mortality.(50) This guided our selection

of the motorized travel time threshold for CHCs, where institutional deliveries are a key

service. The rationale for the remaining thresholds was two-fold: first, they must be less than

120 minutes; and second, the access to PHCs should be higher than that to CHCs, as the

former are primary-level care facilities. Lastly, our approach used either walking or motorized

transport. We acknowledge that travel to a healthcare facility often necessitates more complex

of different transportation modes. However, travel time raster data to consider such complex

travel scenarios remains unavailable. Hence, our analyses may not reflect the realistic travel

patterns in India. Even so, the walking and motorized modes of transport provide the upper

and lower bounds of travel times. Travel time could be as short as that of motorized transport,

assuming optimal conditions, or as long as the time required to walk to the nearest healthcare

facility. The true travel time would lie between the two bounds. Contingent on data

availability, future studies should explore hybrid travel models for a more accurate

assessment of timely geographic access.

4.3 Implications and Takeaways

Increased travel times and low geographic access can result in reduced utilization of health

services and worsen health outcomes. Issues with healthcare access also impede the pursuit of

Universal Health Coverage (UHC), and deepen inequities.(9,13–15,18–22,24,30) We

identified three states/UTs that must be the focus of future policy and efforts like APC-guided

facility placement and improvements in pedestrian and road infrastructure, as they were

consistently found to have long travel times and low geographic access: Arunachal Pradesh,

Mizoram, and Ladakh. Our findings indicate that the current spatial distribution of primary

and secondary healthcare facilities in India does not favour geographic access for a major

portion of the Indian population by walking. A spatial coverage modelling study done in one

district of Jammu & Kashmir, India, reported that access to ambulatory and immunization

care within one hour of walking was 17%.(36) A similar study from Western Province,

Rwanda, found that 26.6% of the population could access the primary health care network by

walking within one hour.(34) This divergence in findings across studies can be explained by

the differences in study settings and the methodologies used to determine geographic access.

Adding new healthcare facilities or strategically relocating existing ones, using APC as a

guidance tool, can improve walkability.

Geographic access by motorised transport to primary and secondary healthcare facilities in

the current study was high. This phenomenon of high geographic access by motorised

transport has been noted in other studies as well.(29,33,36) However, geographic access was

lower in rural areas compared to urban areas. This can be addressed by improving public road

transport and strengthening ambulance networks in these areas. These measures are faster to

implement and are a pragmatic stopgap until a more equitable APC-guided establishment of

future healthcare facilities can be achieved over a long-term horizon. While improving timely

geographic access through enhancements to public road transport and ambulance networks is

a crucial, faster-to-implement measure, it primarily addresses the resistance in transport, not

the fundamental lack of service availability. For timely geographic access, the proximity and

availability of facilities remain the most decisive factors in increasing utilization and

achieving true timely geographic access. Therefore, the long-term, structural solution remains

the establishment of new facilities or optimization of locations guided by APC. Furthermore,

our study also lends support to the use of APC as a measure of geographic access over travel

times. We noticed that the travel times for walking in urban areas were shorter compared to

rural areas, but APC values were lower for urban areas. This is because APC takes into

account population size, density, and spatial distribution of health facilities, providing a more

holistic view of geographic access. Finally, it is necessary to highlight that future research

must incorporate the functional status of the health facilities as well as population attributes

that facilitate or impede geographic access. This will result in more precise estimates of

geographic access to health facilities and better inform policy making and programmes.

5 Conclusion

This study finds several gaps in timely geographic access to primary and secondary healthcare

facilities in India, particularly for walking. Arunachal Pradesh, Mizoram, and Ladakh, with

long travel times and low geographic access, should be the priority areas for future

interventions. Focusing on APC-based facility establishment and pedestrian infrastructure can

help improve timely geographic access by walking. Additionally, rural areas must be

prioritized for improvements in road transport and ambulance networks. Addressing these

disparities in timely geographic access is essential for improving health outcomes and

achieving UHC.

6 List of Abbreviations

1. PHC: Primary Health Centre

2. CHC: Community Health Centre

3. IPHS: Indian Public Health Standards

4. LMIC: Low-and Middle-Income Countries

5. GIS: Geographic Information System

6. PMGSY: Pradhan Mantri Gram Sadak Yojana

7. GoI: Government of India

8. MAP: Malaria Atlas Project

9. UN: United Nations

10. APC: Access Population Coverage

11. UT: Union Territory

12. API: Application Programming Interface

13. RHS: Rural Health Statistics

7 Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: The datasets used and/or analysed during the

current

study is available from the corresponding author on reasonable request.

Competing interests: Siddhesh Zadey is the co-founding director of the Association for

Socially Applicable Research (ASAR). He also represents ASAR at the G4 Alliance

Permanent Council, and serves as the Chair of the SOTA Care in Asia Working Group, The

G4 Alliance, a Fellow of the Lancet Citizens' Commission on Reimagining India's Health

System, and a Drafting Committee Member for Maharashtra State Mental Health Policy.

Other authors declare no competing interests.

Funding: None

Authors' contributions:

Study concept and design: Siddhesh Zadey

Acquisition, analysis, or interpretation of data: Shreyas Patil, Harsh Thakkar, Aamir

Miyajiwala, Anoushka Arora, Harsh Thakkar, Ankit Raj, Siddhesh Zadey

Drafting of the manuscript: Shreyas Patil, Anoushka Arora

Critical revision of the manuscript for important intellectual content: All authors

Statistical analysis: Aamir Miyajiwala, Harsh Thakkar, Ankit Raj, Siddhesh Zadey

Administrative, technical, or material support: Siddhesh Zadey

Study supervision: Siddhesh Zadey

Acknowledgements: None

8 Bibliography

1. UN DESA Policy Brief No. 153: India overtakes China as the world's most populous country | Department of Economic and Social Affairs [Internet]. [cited 2024 Mar 18]. Available from: https://www.un.org/development/desa/dpad/publication/un-desa-policy-brief-no-153-india-overtakes-china-as-the-worlds-most-populous-country/

2. The hilly terrain: Discussing health in Manipur [Internet]. [cited 2024 Jun 29]. Available from: https://www.orfonline.org/expert-speak/health-manipur-hilly-terrain

3. International Conference on Primary Health Care. Declaration of Alma-Ata. WHO Chron. 1978 Nov;32(11):428–30.

4. IPHS 2022 Guidelines | National Health Systems Resource Centre [Internet]. [cited 2024 Jun 29]. Available from: https://nhsrcindia.org/IPHS2022/iphs-2022-guidelines

5. McLaughlin CG, Wyszewianski L. Access to care: remembering old lessons. Health Serv Res. 2002 Dec;37(6):1441–3.

6. Forsyth A. Geographic Healthcare Access and Place. Health and Places Initiative; 2014 Sep.

7. Garchitorena A, Ihantamalala FA, Révillion C, Cordier LF, Randriamihaja M, Razafinjato B, et al. Geographic barriers to achieving universal health coverage: evidence from rural Madagascar. Health Policy Plan. 2021 Nov 11;36(10):1659–70.

8. Varela C, Young S, Mkandawire N, Groen RS, Banza L, Viste A. TRANSPORTATION BARRIERS TO ACCESS HEALTH CARE FOR SURGICAL CONDITIONS IN MALAWI a cross sectional nationwide household survey. BMC Public Health. 2019 Mar 5;19(1):264.

9. Ipa M, Laksono AD, Wulandari RD. The Role of Travel Time on Hospital Utilization in the Islands Area: A Cross-Sectional Study in the Maluku Region, Indonesia, in 2018. Indian J Community Med. 2023 Apr 7;48(2):269–73.

10. Al-Taiar A, Clark A, Longenecker JC, Whitty CJM. Physical accessibility and utilization of health services in Yemen. Int J Health Geogr. 2010 Jul 21;9:38.

11. Müller I, Smith T, Mellor S, Rare L, Genton B. The effect of distance from home on attendance at a small rural health centre in Papua New Guinea. Int J Epidemiol. 1998 Oct;27(5):878–84.

12. Stock R. Distance and the utilization of health facilities in rural Nigeria. Soc Sci Med. 1983;17(9):563–70.

13. Panciera R, Khan A, Rizvi SJR, Ahmed S, Ahmed T, Islam R, et al. The influence of travel time on emergency obstetric care seeking behavior in the urban poor of Bangladesh: a GIS study. BMC Pregnancy Childbirth. 2016 Aug 22;16(1):240.

14. Barry R, Rehm J, de Oliveira C, Gozdyra P, Chen S, Kurdyak P. The relationship between rurality, travel time to care and death by suicide. BMC Psychiatry. 2023 May 17;23(1):345.

15.   Konerding U, Bowen T, Elkhuizen SG, Faubel R, Forte P, Karampli E, et al. The
impact of travel distance, travel time and waiting time on health-related quality of life
of diabetes patients: An investigation in six European countries. Diabetes Res Clin
Pract. 2017 Apr;126:16–24.

16.   Kelly C, Hulme C, Farragher T, Clarke G. Are differences in travel time or distance to
healthcare for adults in global north countries associated with an impact on health
outcomes? A systematic review. BMJ Open. 2016 Nov 24;6(11):e013059.

17.   Okwaraji YB, Cousens S, Berhane Y, Mulholland K, Edmond K. Effect of
geographical access to health facilities on child mortality in rural Ethiopia: a
community based cross sectional study. PLoS ONE. 2012 Mar 12;7(3):e33564.

18.   Manongi R, Mtei F, Mtove G, Nadjm B, Muro F, Alegana V, et al. Inpatient child
mortality by travel time to hospital in a rural area of Tanzania. Trop Med Int Health.
2014 May;19(5):555–62.

19.   Örtqvist AK, Haas J, Ahlberg M, Norman M, Stephansson O. Association between
travel time to delivery unit and unplanned out-of-hospital birth, infant morbidity and
mortality: A population-based cohort study. Acta Obstet Gynecol Scand. 2021
Aug;100(8):1478–89.

20.   Luke S, Hobbs A, Mak S, Der K, Pederson A, Schummers L. Travel Time to Delivery,
Antenatal Care, and Birth Outcomes: A Population-Based Cohort of Uncomplicated
Pregnancies in British Columbia, 2012-2019. J Obstet Gynaecol Can. 2022
Aug;44(8):886–94.

21.   Ravelli ACJ, Jager KJ, de Groot MH, Erwich JJHM, Rijninks-van Driel GC, Tromp M,
et al. Travel time from home to hospital and adverse perinatal outcomes in women at
term in the Netherlands. BJOG. 2011 Mar;118(4):457–65.

22.   Hanson C, Cox J, Mbaruku G, Manzi F, Gabrysch S, Schellenberg D, et al. Maternal
mortality and distance to facility-based obstetric care in rural southern Tanzania: a
secondary analysis of cross-sectional census data in 226 000 households. Lancet Glob
Health. 2015 Jul;3(7):e387-95.

23.   Kumar S, Dansereau EA, Murray CJL. Does distance matter for institutional delivery
in rural India? Appl Econ. 2014 Nov 22;46(33):4091–103.

24.   McGuire F, Kreif N, Smith PC. The effect of distance on maternal institutional
delivery choice: Evidence from Malawi. Health Econ. 2021 Sep;30(9):2144–67.

25.   Mahajan A, Tirakotai W, Masayaanon P. The hidden costs of universal health
coverage: solutions from the fight against catastrophic healthcare expenditure in
Thailand. BMJ Glob Health. 2023 Jun;8(6).

26.   Rocque GB, Williams CP, Miller HD, Azuero A, Wheeler SB, Pisu M, et al. Impact of
travel time on health care costs and resource use by phase of care for older patients
with cancer. J Clin Oncol. 2019 Aug 1;37(22):1935–45.

27.   Kornelsen J, Khowaja AR, Av-Gay G, Sullivan E, Parajulee A, Dunnebacke M, et al.

The rural tax: comprehensive out-of-pocket costs associated with patient travel in British Columbia. BMC Health Serv Res. 2021 Aug 21;21(1):854.

28. Bartosch P, Jaye C, Crampton P. Moral economy and moral capital: A new approach to understanding health systems. Soc Sci Med. 2024 Jul;352:117016.

29. Weiss DJ, Nelson A, Vargas-Ruiz CA, Gligorić K, Bavadekar S, Gabrilovich E, et al. Global maps of travel time to healthcare facilities. Nat Med. 2020 Dec;26(12):1835–8.

30. Amoah-Nuamah J, Agyemang-Duah W, Prosper Ninorb G, Gladstone Ekeme B. Analysis of spatial distribution of health care facilities and its effects on access to primary healthcare in rural communities in kpandai district, ghana. Cogent Public Health. 2023 Dec 31;10(1).

31. Palk L, Okano JT, Dullie L, Blower S. Travel time to health-care facilities, mode of transportation, and HIV elimination in Malawi: a geospatial modelling analysis. Lancet Glob Health. 2020 Dec;8(12):e1555–64.

32. Carrasco-Escobar G, Manrique E, Tello-Lizarraga K, Miranda JJ. Travel time to health facilities as a marker of geographical accessibility across heterogeneous land coverage in peru. Front Public Health. 2020 Sep 16;8:498.

33. Ouko JJO, Gachari MK, Sichangi AW, Alegana V. Geographic information system☐based evaluation of spatial accessibility to maternal health facilities in Siaya County, Kenya. Geographical Research. 2019 Aug;57(3):286–98.

34. Huerta Munoz U, Källestål C. Geographical accessibility and spatial coverage modeling of the primary health care network in the Western Province of Rwanda. Int J Health Geogr. 2012 Sep 17;11:40.

35. Comber AJ, Brunsdon C, Radburn R. A spatial analysis of variations in health access: linking geography, socio-economic status and access perceptions. Int J Health Geogr. 2011 Jul 25;10:44.

36. Verma VR, Dash U. Geographical accessibility and spatial coverage modelling of public health care network in rural and remote India. PLoS ONE. 2020 Oct 21;15(10):e0239326.

37. Home | National Rural Infrastructure Development Agency (NRIDA), Government of India [Internet]. [cited 2024 Jun 30]. Available from: https://pmgsy.nic.in/

38. MAP [Internet]. Maria Atlas. [cited 2024 Jun 30]. Available from: https://malariaatlas.org/our-work/

39. GADM. GADM maps and data [Internet]. GADM. [cited 2021 Mar 16]. Available from: https://gadm.org/index.html

40. Global 1km Population total adjusted to match the corresponding UNPD estimate. University of Southampton. 2020;

41. Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber A, et al. A global map of travel time to cities to assess inequalities in accessibility in 2015. Nature. 2018

Jan 18;553(7688):333–6.

42.   Jadhav T, Vissoci JRN, Zadey S. Measuring timely geographical access to surgical
care in India: a geospatial modelling study. Lancet Glob Health. 2022 Mar;10 Suppl
1:S29.

43.   Zadey S. Measuring Access to Surgical Care in Rural India: Synthesis of Data and
Novel Index. 2021;

44.   Thakkar PSH, Patil A, Chauhan P, Chembon P, Joseph SA, Rana S, et al. Access to
palliative care in India: situational analysis and modeling of access from public
healthcare centers. Ecancermedicalscience. 2025 Nov 19;19.

45.   Poddar S, Gorkar A, Passi VR, Thakkar H, Andhale A, Puntambekar V, et al.
Geographical access to COVID-19 testing centers in India. BMC Health Serv Res.
2025 Nov 28;25(1):1551.

46.   Urs G, Miyajiwala A, Raj A, Zadey S. Timely Geographic Access to Blood Banking
Facilities: A Pan-India Modeling Study. medRxiv. 2025 May 19;

47.   Thakkar H, Reddy C, Passi VR, Miyajiwala A, Kale S, Raj A, et al. Assessing
Population-level Accessibility to Medical College Hospitals in India: A Geospatial
Modeling Study. Cold Spring Harbor Laboratory. 2024;

48.   Ministry of Road Transport & Highways, Government of India. Basic Road Statistics
of India 2018-19. Ministry of Road Transport & Highways, Government of India;
2019.

49.   National Rural Health Mission ☐:: National Health Mission [Internet]. [cited 2025 Feb
16]. Available from:
https://nhm.gov.in/index1.php?lang=1&level=1&lid=49&sublinkid=969

50.   Meara JG, Leather AJM, Hagander L, Alkire BC, Alonso N, Ameh EA, et al. Global
Surgery 2030: evidence and solutions for achieving health, welfare, and economic
development. Lancet. 2015 Aug 8;386(9993):569–624.

Figure Captions

Figure 1: Raster-based estimation pipeline used for creating criteria-specific regional

population counts.

Figure 2: Map depicting the variation in density of a) PHCs and b) CHCs across districts in

India.

Figure 3: Maps depicting the APC in the districts of India for a) PHCs by walking, b)

PHCs by motorised transport, c) CHCs by walking, and d) CHCs by motorised transport.

Tables for "Geospatial Accessibility to Primary and Community Health Centres in India

Table 1. Summary of travel times for rural and urban regions.

National level				
Median (IQR) travel time (in minutes)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	79.8 (46.12-134.28)	60.77 (34.66-94.58)	228.36 (131.55-384.50)	173.65 (96.94-275.93)
Motorised transport	17.16 (8.04-35.61)	6.99 (3.82-11.22)	36.80 (20.68-64.36)	18.26 (10.55-29.29)
State/UT level				
States/UTs with travel time shorter than national median, n (%)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	14/29 (48.28%)	18/29 (62.07%)	15/29 (51.72%)	15/29 (51.72%)
Motorised transport	12/29 (41.38%)	15/29 (51.72%)	14/29 (48.28%)	15/29 (51.72%)
Shortest median (IQR) travel time (in minutes)				

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Punjab: 47.92 (28.76 - 72.84)	Manipur: 25.11 (11.08 - 35.24)	Kerala: 112.31 (64.29-199.52)	Kerala: 88.92 (57.80-128.72)
Motorised transport	Kerala: 6.97 (3.79-17.91)	Manipur: 2.50 (1.58 - 4.14)	Kerala: 13.69 (7.77-30.88)	Kerala: 9.39 (6.15-13.48)
Longest median (IQR) travel time (in minutes)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Ladakh: 852.06 (477.00 -1337.78)	Mizoram: 154.12 (133.73-170.86)	Ladakh: 1773.88 (1027.79-2513.11)	Mizoram: 863.19 (450.79-897.74)
Motorised transport	Ladakh: 311.74 (126.63-601.20)	Mizoram: 11.70 (9.83-13.55)	Ladakh: 396.73 (218.88-681.95)	Mizoram: 65.78 (40.40-27.88)
District level				
Districts with travel time shorter than the national median, n (%)				

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	410/674 (60.83%)	261/504 (51.78%)	390/565 (69.02%)	250/387 (64.60%)
Motorised transport	406/674 (60.24%)	281/504 (55.75%)	376/565 (66.55%)	257/387 (66.41%)
Shortest median (IQR) travel time (in minutes)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Nalanda, Bihar: 25.06 (14.97-40.30)	Bishnupur, Manipur: 11.07 (10.12-20.23)	Alappuzha, Kerala: 57.74 (37.36-84.46)	Changlang, ARP[2]: 13.60 (9.88-21.03)
Motorised transport	Pulwama, J&K[1]: 2.87 (1.56-32.54)	Bishnupur, Manipur: 1.63 (0.79-2.41)	Alappuzha, Kerala: 6.7 (4.11-10.01)	Tuensang, Nagaland: 1.68 (0.86-1.24)
Longest median (IQR) travel time (in minutes)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Leh, Ladakh: 858.06 (501.72-1358.29)	Singrauli, MP[3]: 176.79 (130.21-196.24)	Leh, Ladakh: 2059.70 (1487.61-2706.89)	Dhule, Maharashtra: 862.40 (422.29-888.83)

Motorised transport	Anjaw, ARP[2]: 600.27 (316.82-965.44)	Punch, J&K[1]: 22.12 (2.62-68.03)	KK[4], ARP[2]: 1007.826 (437.17-1558.69)	Dhule, Maharashtra 67.61 (45.15-70.43)

[1]J&K: Jammu & Kashmir
[2]ARP: Arunachal Pradesh
[3]MP: Madhya Pradesh
[4]KK: Kurung Kumey

Table 2: Summary of access population coverage (APC) for rural and urban regions.

National level				
Access Population Coverage (APC), %				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	20.51%	13.81%	11.58%	9.05%
Motorised transport	97.36%	99.73%	98.62%	99.66%
State/UT level				
States/UTs with APC greater than the national average, n (%)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	17/29 (58.62%)	8/29 (27.59%)	15/29 (51.72%)	12/29 (41.38%)
Motorised transport	12/29	29/29 (100%)	13/29 (44.83%)	26/29 (89.65%)

	(41.38%)			
States/UTs with APC = 0, n (%)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	0/29 (0.0%)	1/29 (3.49%)	0/29 (0.0%)	1/29 (3.49%)
Motorised transport	0/29 (0.0%)	0/29 (0.0%)	0/29 (0.0%)	0/29 (0.0%)
States/UTs with APC ≥ 50% n (%)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	0/29 (0.0%)	1/29 (3.49%)	0/29 (0.0%)	0/29 (0.0%)
Motorised transport	28/29 (96.55%)	29/29 (100%)	28/29 (96.55%)	29/29 (100%)
States/UTs with APC = 100% n (%)				
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)

Walking	0/29 (0.0%)	0/29 (0.0%)	0/29 (0.0%)	0/29 (0.0%)
Motorised transport	0/29 (0.0%)	6/29 (20.69%)	0/29 (0.0%)	11/29 (37.93%)

State with the lowest access population coverage (APC), %

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Mizoram: 5.93%	Mizoram: 0.0%	West Bengal: 1.15%	Mizoram: 0.71%
Motorised transport	Arunachal Pradesh: 64.06%	Assam: 98.38%	Ladakh: 49.54%	Tripura: 97.98%

State with the highest access population coverage (APC), %

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Kerala:	Manipur:	Kerala:	Jharkhand:

	34.75%	69.35%	34.31%	35.35%
Motorised transport	Karnataka: 99.58%	6 states/UTs: 100%	Haryana: 99.99	11 states/UTs: 100%

District level

Districts with APC greater than the national average, n (%)

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking Motorised transport	318/675 (47.11%) 364/675 (53.93%)	256/644 (39.75%) 620/644 (96.27%)	284/601 (47.25%) 405/601 (67.5%)	257/577 (44.54%) 557/577 (96.53%)

Districts with APC = 0, n (%)

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	0/675 (0.0%)	48/644 (7.45%)	2/601 (0.33%)	74/577 (12.83%)

Motorised transport	0/675 (0.0%)	0/644 (0.0%)	0/601 (0.0%)	0/577 (0.0%)

Districts with APC ≥ 50% n (%)			

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking Motorised transport	18/675 (2.66%) 664/675 (98.37%)	40/644 (6.21%) 644/644 (100.0%)	3/601 (0.49%) 593/601 (98.66%)	42/577 (7.28%) 577/577 (100%)

Districts with APC = 100% n (%)			

	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking Motorised transport	0/675 (0%) 64/675 (9.48%)	0/644 (0.0%) 520/644 (80.75%)	0/601 (0.0%) 165/601 (27.45%)	1/577 (0.17%) 506/577 (87.69%)

		District with the lowest access population coverage (APC), %		
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking Motorised transport	Upper Siang, Arunachal Pradesh: 0.78% Anjaw, Arunachal Pradesh: 16.43%	48 districts: 0% Karimganj, Assam: 66.72%	2 districts: 0.0% Anjaw, Arunachal Pradesh: 17.36%	74 districts: 0.0% Karimganj, Assam: 66.73%
		District with the highest access population coverage (APC), %		
	PHC (Rural)	PHC (Urban)	CHC (Rural)	CHC (Urban)
Walking	Bishnupur, Manipur:	Bishnupur, Manipur: 95.08%	Bongaigaon, Assam: 52.67%	Changlang, Arunachal Pradesh:

| Motorised transport | 65.9% 64 districts: 100% | 520 districts: 100% | 165 districts: 100% | 100% 506 districts: 100% |

a

PHC-centers per million population

- 0.5 to 14.1
- 14.1 to 22.3
- 22.3 to 31.3
- 31.3 to 50.8
- 50.8 to 638.2
- Missing

b

CHC-centers per million population

- 0.1 to 1.6
- 1.6 to 3.0
- 3.0 to 4.6
- 4.6 to 8.0
- 8.0 to 127.6
- Missing