Cohort Profile: Baseline characteristics and design of the McMaster Monitoring My Mobility (MacM3) Study, a prospective digital mobility cohort of community-dwelling older Canadians from Southern Ontario

Marla K Beauchamp[1,2,3], Renata Kirkwood[1], Cody Cooper[1], William McIlroy[4], Karen Van Ooteghem[4], Kit Beyer[4,], Julie Richardson[1,2], Ayse Kuspinar[1,2], Paul D. McNicholas[5,2], K. Bruce Newbold[6,2], Darren M. Scott[6,2], Parminder Raina[7,2], Qiyin Fang[8,2], Paula Gardner[9,2], Manaf Zargoush[10,2], Jinhui Ma[7,2], Sachi O'Hoski[1,3], Talha Rafiq[7,2], on behalf of the MacM3 investigators

ABSTRACT

Purpose: The McMaster Monitoring My Mobility (MacM3) study aims to understand trajectories of mobility decline in later life using multi-sensor wearable technology. To our knowledge, MacM3 is the first major cohort to combine accelerometry and GPS to track real-world mobility in community-dwelling older adults.

Participants: Between May 2022 and May 2024, MacM3 recruited 1,555 community-dwelling older adults (mean age 73.9 years, SD = 5.5) from Hamilton and Toronto, Ontario. Of the cohort, 66.3% were female, 62.4% married/partnered, 75.3% had post-secondary education, and 63.0% had ≥3 comorbidities. Most were Canadian born (69.4%) and White/Caucasian (88.2%), with greater ethnocultural diversity observed at the Toronto site.

Findings to date: At baseline, 56.7% of participants reported no mobility limitations, 15.9% had preclinical limitations, and 27.4% had minor mobility limitations. Mean gait speed for the total sample was 1.23 m/s, with a mean Timed Up and Go time of 9.4 seconds and a 5x sit-to-stand time of 13.0 seconds. A total of 1,301 participants had valid wrist-worn device data, and 1,008 participants who agreed to wear the thigh-worn device had valid data (≥7 days with ≥10 hours of wear per day). Step count data (n = 1008) revealed a mean of 8,437 steps per day (SD = 2,943), with 5,073 steps in the lowest quartile and 12,303 steps in the highest.

Future plans: Ongoing work aims to develop predictive models of mobility decline by integrating wearable, clinical, and environmental data. Pipeline enhancements will enable GPS/inertial measurement unit fusion to explore mobility-environment interactions and support aging-in-place tools.

INTRODUCTION

In the 2015 World Report on Ageing and Health, the World Health Organization defines healthy aging as the 'process of developing and maintaining the functional ability that enables well-being in older age'.[1] Central to this concept is mobility — the ability to move, walk, and use transportation within our homes and communities.[2] Mobility is a cornerstone of healthy aging, reported to be of utmost importance to older people as they age, and has been labeled by many aging experts as the '6th vital sign'.[3–5] With population aging, the number of older adults living with mobility limitations will rise dramatically over the next four decades. In Canada, one in two older adults report reduced mobility compared to the previous decade, with 24-33% facing challenges in conducting daily activities that require physical capabilities, such as climbing stairs and walking.[6–8] These limitations can signal early neurological, musculoskeletal, or cardiorespiratory health changes and predict adverse outcomes like falls, hospitalization, institutionalization, and mortality.[9–12]

Despite ongoing efforts to understand predictors and trajectories of mobility decline in older adulthood, knowledge gaps remain, particularly in understanding the varied trajectories of mobility changes across different age groups and functional levels.[13–18] Previous studies have predominantly focused on older populations (age 70+) already experiencing mobility limitations or relied on self-reported measures of function,[19,20] which may not accurately capture day-to-day mobility challenges.[13,14,16–18] As such, our understanding of the actual level and types of mobility undertaken by individuals in their daily lives is limited.[21] By expanding our research to include a broader age range and incorporating device-based measures of real-world mobility, we can effectively identify early signs of decline and inform interventions to delay or prevent the onset of disability.

Advancements in wearable technology have revolutionized our ability to continuously monitor unsupervised mobility in individuals' daily lives, both at home and in the community ('free-living').[22–24] This innovation has generated substantial interest and great potential to utilize wearables for prevention, health management, and personalized care, especially among older adults who increasingly recognize the value of self-monitoring their mobility as part of their health regimen.[25,26] Our consultations with Canadian stakeholders (i.e., older adults, caregivers,

and industry and community partners) confirmed that older adults have a strong interest in using such technology to monitor their mobility and enhance their health. Given that mobility data obtained using wearable technology (i.e., digital mobility measures) reflect real-world functioning, these measures hold exceptional potential for monitoring and predicting health outcomes and promoting aging-in-place.[27,28]

In 2022, we established the McMaster Monitoring My Mobility (MacM3) research platform, with the primary objective of characterizing later-life trajectories of early mobility decline through longitudinal, multi-sensor-based data collected from a large cohort of older adults exhibiting differing levels of mobility at baseline. This pioneering initiative represents the first major longitudinal study to evaluate real-world mobility with frequent repeated measures in community dwelling older adults using a wearable system that integrates GPS (Global Positioning Systems) and accelerometry.[29] Our methodology centers on a wrist-worn device that synchronizes time-based accelerometry with GPS data, complemented by a thigh-worn accelerometer. The unique combination of accelerometry and geolocation data provides a holistic view of mobility, capturing interactions between individuals and their environments. Such a comprehensive approach enhances our capacity to detect early signs of mobility decline and develop strategies to mitigate mobility-related adverse health impacts in older adulthood.

COHORT DESCRIPTION

Study design and recruitment

MacM3 is a prospective cohort study with 24 months of follow-up initiated in May 2022 at the Hamilton, Ontario site and expanded to the Dixon Hall social services site in Toronto in January 2023. The study targeted individuals aged 65 years and older residing within a 50 km radius of McMaster University in Hamilton or Dixon Hall in Toronto. All participants provided verbal informed consent. Eligibility criteria included proficient in English or either English or Mandarin (at Dixon Hall) and classification into one of three mobility groups based on the concept of preclinical mobility limitation[30] measured by the Preclinical Mobility Limitation classification (PCML)[20]: 1) no mobility limitation, 2) preclinical mobility limitation, or 3) minor mobility limitation. The PCML classification is based on self-reported difficulty with three mobility tasks:

walking 0.5 km, walking 2 km, and climbing one flight of stairs. For each task, participants are asked whether they are: (1) able to manage without difficulty, (2) able to manage with some difficulty, (3) able to manage with a great deal of difficulty, (4) able to manage only with the help of another person, or (5) unable to manage even with help. Participants who report being able to manage any task without difficulty (response 1) are further asked about task modification using a set of follow-up questions. These include: "I am able to walk 0.5 km the same way I always have," "I need to rest in the middle of the walk," "I walk 0.5 km less often than I used to," "It takes me longer to walk 0.5 km than it used to," and "I am more tired after walking 0.5 km than I used to be." Participants who report no task modification—i.e., they perform all tasks the same way as before—are classified as having **no mobility limitation**. Those who report modification of any task (e.g., reduced frequency, slower performance, fatigue, or need for rest) are classified as having a **preclinical mobility limitation**. Individuals who report being able to manage a task with some difficulty (response 2) are classified as having a **minor mobility limitation**, while those who report great difficulty, needing help, or being unable to complete a task (responses 3, 4, or 5) are classified as having a **major mobility limitation**. While individuals with major mobility limitation were excluded at baseline, they could be reclassified into this category during follow-up assessments. Additional exclusion criteria included inability to communicate in required languages, significant visual or hearing impairments that could affect safety during assessments and plans to relocate outside the study area within two years.

We engaged in a multi-pronged recruitment strategy aimed at enhancing the diversity of participants in the study (in ethnicity, race, income level, culture, ability, and region), and ensuring the data reflects diverse participant experiences. Our recruitment strategy employed a combination of community-based approaches, including word of mouth, advertisements in local newspapers, and online platforms. Targeted mail campaigns were conducted using Canada Post Snap Admail, focusing on postal routes and codes that matched the study's demographic in lower-income areas, and advertisements were placed on relevant public transportation routes. Our recruitment strategy was designed to reach adults aged 65 and older—including males, females, individuals of any gender identity, and people living in both higher- and lower-income areas—to ensure a diverse and inclusive sample.

Our second recruitment site, Dixon Hall, serves a large, diverse population, including many individuals who are underrepresented in research, particularly those facing social challenges such as poverty, isolation, and health inequities. Our partnership with Dixon Hall, facilitated by the McMaster Institute for Research on Aging (MIRA) | Dixon Hall Centre, allows us to engage with a population that is often excluded from research studies, ensuring that the data collected are more generalizable to the broader community of older adults. In order to build trust and access, we embedded researchers at the Dixon Hall site, recruiting participants active in social and recreation groups via a snowball method, where participants invited peers to participate. While the Hamilton participants tend to be more homogenous (Caucasian and middle-income), Dixon Hall participants had more diversity in race, income level, ethnicity, and primary spoken language (see Table 1). Additionally, a small subset of participants ($n = 148$) was recruited from the MIRA Intergenerational Study (MIRA-iGeN) platform, which includes 5,000 individuals pre-recruited by Canadian Viewpoint Inc. through phone, online, and mail-out methods.

Our sample size was designed to support MacM3's main objectives: to compare mobility patterns among older adults with varying levels of early mobility limitation; identify distinct mobility trajectories and their characteristics; assess the predictive value of digital mobility measures for adverse outcomes; and model risk factors for the onset and progression of mobility decline in older Canadians. To allow for trajectory analyses and to account for potential seasonal changes in activity patterns, we incorporated frequent repeated follow-ups (every 4 months).

Based on our original objectives, our target sample size was 1,500 participants, grounded in both feasibility and predictive validity considerations. Prior studies examining the accuracy of mobility measures for predicting health outcomes have reported area under the curve (AUC) values ranging from 0.6 to 0.8.[31,32] To ensure adequate power (80%) to detect a significant deviation from the null hypothesis (AUC = 0.5; two-sided test at $p < 0.05$), we based our calculations on the most conservative expected AUC of 0.6. Based on conservative expected incidence rates for key outcomes: at least 1 fall (20% over 1 year), visit to the emergency room (12% over 1 year), overnight hospital stays (8% over 2 years), and progression to major mobility

disability (10% over 2 years), a sample of approximately 740 participants would be needed to detect progression to major mobility limitation over two years, and around 634 participants to detect emergency room visits over the same period. Accounting for an anticipated dropout rate of 20% and the potential for missing device-based data of up to 20%, the required sample increases to approximately 1,156 for progression to major mobility disability and 991 for ER visits. This sample size is also sufficient for detecting small and clinically relevant differences between the three mobility level groups based on the PCML for step count (400-600 steps) and for trips per week (4-6 trips). Our flowchart (Figure 3) updated on September 23, 2025, highlights that we completed our 12-month follow-up with 1,378 active participants.

Study Procedures

Participants in MacM3 are asked to wear a consumer grade smartwatch with Global Navigation Satellite System (GNSS) and activity monitoring capabilities, the TicWatch Pro 3 Ultra GPS[29] on the wrist for 10 days at baseline and at 4-month intervals for a total of seven time points of data collection over 24 months. At the time of consent, participants were also asked if they would be willing to wear a second (optional) activity monitoring device, the ActiGraph wGT3X-BT[29,33] or the Axivity X3,[34] on the thigh concurrently with the wrist-worn device. At baseline, 12 months, and 24 months, participants complete a more in-depth series of questionnaires about their health and mobility through a combination of telephone interviews and online surveys as well as an in-person physical assessment. In between the yearly assessments, devices are mailed to participants' homes every 4 months and then returned to the study site via pre-paid shipping. To monitor health outcomes (e.g., new diagnoses, healthcare utilization, injuries, falls), we use quarterly phone calls and monthly mail-in fall diaries.[35] Figure 1 provides an overview of the MacM3 study design.

Study measures
Our data collection encompasses all three "tenses" of mobility/function in line with the model first proposed by Glass[36] and outlined more recently by members of our team based on the World Report on Ageing and Health[37]: enacted (i.e., actual mobility in the community measured via wearables), hypothetical (i.e., perceived ability measured via self-reported questionnaires),

Figure 1. MacM3 study overview.

and experimental (i.e., locomotor capacity for mobility measured via in-person physical assessments). Self-reported measures of mobility provide information on what patients "can do"; performance-based testing provides information on what patients "could do"; and importantly, the sensor data on enacted/actual mobility provide information on what patients actually "do" in the real-world.[37]

Mobility Measures

<u>Enacted mobility</u>

The MacM3 study employs a multi-sensor, multi-nodal approach for studying enacted and free-living mobility, integrating time-synchronized accelerometry with GPS in a wrist-worn device alongside a second thigh-worn accelerometer.[29] Device selections and wear locations for the current cohort were made after extensive lab-based testing, consultation with our older adult Stakeholder Advisory Committee, and pilot testing to determine their reliability and technical validity.[29]

Participants are asked to wear a TicWatch Pro 3 Ultra GPS on their non-dominant wrist and either the ActiGraph GT3X-BT or an Axivity AX3 on the thigh for 10 consecutive days during waking hours at baseline and every 4 months throughout the study. The **wrist-worn device** (TicWatch by Mobvoi) is a smartwatch that uses the Wear OS operating system by Google[TM]

and has a built-in tri-axial accelerometer, gyroscope, heart rate sensor, and multi-GNSS (GPS, GLONASS, Galileo, QZSS, and Beidou) for logging position coordinates and time. As the watch was intended to be used in our study as a stand-alone data logging device without a cellphone, GNSS data are collected at different intervals depending on the predicted activity state of the device using Google's Activity Recognition API. To improve battery life, we configured the device to collect location coordinates every 5 seconds when the watch is in an active motion state (e.g., walking, in a vehicle) with good satellite reception, and less frequently when the device is still or there is poor satellite reception.[29] We validated the TicWatch against a gold-standard stand-alone GPS device (Qstarz GPS) and validated its accelerometer output against the thigh-worn ActiGraph device. In this validation study, we found that the TicWatch models were valid devices for capturing both GPS and raw accelerometer data, supporting their use as reliable tools for assessing real-life mobility in older adults.[29]

The second device is **a thigh-worn accelerometer** (either the ActiGraph wGT3X-BT or Axivity AX3) secured to the upper thigh using an adhesive patch (Hypafix). The ActiGraph wGT3X-BT[38,39] (ActiGraph, Pensacola, FL) and Axivity AX3[34] (Axivity Ltd, United Kingdom) are commonly used in epidemiological research for measuring physical activity[40–42] and body posture, with thigh placement considered the gold standard for accurately distinguishing between sitting, standing, and walking.[29,43–46] Both devices have built-in triaxial accelerometers for recording high-resolution raw acceleration data and long-lasting batteries that eliminate the need for recharging during the data collection period. In our validation study, the ActiGraph demonstrated high accuracy in detecting body posture. Full details of this validation study have been published elsewhere.[29]

A custom data collection application, Ivy, was designed by our team exclusively for the TicWatch to optimize data quality and battery life by collecting data from onboard sensors, including the accelerometer, gyroscope, geolocation, and heart rate.[29] A companion app, Clover, supports Ivy on Windows by automating the setup of both the TicWatch and the Axivity device, streamlining data downloads and device preparation. The ActiGraph device, on the other hand, is initialized using the ActiLife proprietary software. Additionally, Clover integrates

information on the date, time, and completion status of data downloads with our REDCap data collection system for all devices, including the ActiGraph.

The primary digitally-derived mobility measures of interest for MacM3 can be classified into five core domains: *activity* (amount of time spent in sedentary, light, moderate and vigorous physical activity),[47,48] *volume-related features of walking* (steps/day, walking bout lengths and duration), *spatiotemporal features of walking* (gait cadence, variability, velocity, and stride width during long bouts),[49] *postural* (time spent lying down, sitting, standing, and frequency of transitions),[44,50] and *geospatial features* (trip frequency, trip duration, trip mode, maximum distance from home, and minimum convex hull/life space area).[51,52] The derivation of these digital mobility measures has been described in previous technical validation papers by our team and others.[37,53–55] Notably, with integrated GNSS and accelerometry data, we are able to uniquely distinguish and characterize indoor and outdoor mobility (bouts, trips) and patterns of mobility outside of the home.

All measures are processed using the NiMBaLWear[53] pipeline which is a modular, open-source wearable sensor analytic framework designed to accommodate a wide range of sensors, signal types, and body wear locations. NiMBalWear was developed to address the limitations of device-specific processing by providing a standardized, flexible approach for analyzing raw accelerometry data across different platforms. This framework unifies the measurement and analysis of multiple health-related domains—such as mobility, posture, and physical activity—by optimizing data extraction from individual sensors and enabling data fusion across multiple sensor types and locations when applicable. By focusing on raw signal processing, NiMBaLWear ensures device-agnostic harmonization during the feature extraction stage, allowing for consistent, comparable metrics regardless of brand or configuration. This is particularly important for longitudinal studies or multi-site research using different devices, as it supports reproducible and scalable analysis without reliance on proprietary algorithms or cut-points. Currently, the NiMBaLWear platform also includes a Mobility module focused on IMU-based outcomes, which is being expanded to incorporate GNSS+ analytics.

NiMBaLWear-based processing of thigh- and wrist-worn accelerometry data

Step counts and walking bouts are derived from the thigh-worn accelerometer. NiMBaLWear applies wavelet-based time-frequency analysis to the triaxial acceleration signal to detect unilateral steps.[56,57] Step-related events are identified based on the frequency content of the acceleration profile, capturing both the swing and stance phases of walking. Foot contact is used to define the timing of each step following a detected swing phase. Consecutive steps are then grouped into walking bouts according to predefined timing criteria (e.g., a minimum number of steps within a maximum interval). This method offers greater precision than traditional threshold-based techniques and has demonstrated improved accuracy in detecting steps and walking patterns, particularly in older adults. The step count data reported in Table 5 align with findings from other large cohort studies[58,59] that applied machine learning methods to estimate step count from wrist-worn devices and observed similar trends—namely, a decline in step counts with advancing age and increased mobility limitations.

Postural behaviour is derived from the same thigh-worn triaxial accelerometer using algorithms based on the orientation of the device relative to gravity. Specifically, posture was classified as either standing or sitting/lying based on the acceleration due to gravity recorded along each axis of the accelerometer, assuming consistent device placement on the thigh. Postural transitions between standing and sitting/lying were detected by identifying periods of sustained change in thigh orientation that met predefined criteria. Once a transition was detected, all subsequent time points were classified according to the last identified posture. This method allows us to determine body posture continuously throughout the wear period using a single device, with classification based solely on orientation and transition patterns.

Activity intensity, including sedentary behavior (SED), light physical activity (LPA), and moderate-to-vigorous physical activity (MVPA), is derived from the wrist device. We applied epoch-independent cut-points validated by Fraysse et al.,[60] which classify activity based on raw acceleration (in mg). Specifically, thresholds of 42.5 mg (non-dominant wrist) and 62.5 mg (dominant wrist) were used for light activity, while 98.0 mg (non-dominant) and 92.5 mg (dominant) defined moderate activity. These cut-points have been applied in other aging

cohorts and support comparability across studies examining accelerometry-derived activity patterns.

Hypothetical mobility

Perceived mobility is assessed through various self-report instruments developed to capture different mobility domains among older adults.[61,62] The Late-Life Function and Disability Instrument (LLFDI)[63,64] evaluates both physical function mobility and participation in valued life roles, encompassing tasks from basic physical actions to community activities. The Mobility Assessment Tool – short form (MAT-sf)[65] is a 10-item computer-administered measure of perceived mobility using video technology. The Life-Space Assessment (LSA)[66] measures life-space mobility – the distance, frequency and independence perceived by individuals as they move through their communities. Transportation mobility is evaluated through questions from the Canadian Longitudinal Study on Aging (CLSA)[67] Transportation, Mobility, Migration (TRA)[68] module, while driving behavior is assessed using the Driving Questionnaire, Driving Comfort Scale (DCS)[69,70] as well as selected questions from the Perceived Driving Abilities (PDA) Scale.[69]

Experimental mobility

Participants undergo in-person physical assessments at either the Hamilton or Toronto location at baseline, 12-months, and 24-months follow-up. These assessments include commonly used and well-validated performance-based tests of mobility and physical function, namely the Timed Up and Go (TUG)[71] usual and dual-task tests, the repeated sit-to-stand test (5 x STS), tests of gait speed[72] (both usual and fast pace) and single-leg balance[73]. The 400-meter walk test[74,75] evaluates aerobic capacity, while muscle power is estimated with the stair-climbing power test.[76,77] Other physical measures collected at the in-person assessment include the Clinical Frailty Scale (CFS),[78] Visual Acuity (VA) test,[79] spirometry[80] for pulmonary function evaluation, and handgrip strength[81] assessment. Participants are permitted to use their mobility aids during the TUG, gait speed, stair-climbing, and 400-meter walk tests. Use of mobility aids was documented and recorded in the study database for future analyses.

Demographic and health measures

As part of the yearly in-person assessments, participants also complete a series of

questionnaires related to their basic health and the possible physical, psychosocial, cognitive, financial, and environmental contributors to mobility. This encompasses demographics, such as socioeconomic status and lifestyle factors, as well as health-related behaviors. We assess physical activity through the Physical Activity Scale for the Elderly (PASE),[82] nutrition risk using the Seniors in the Community: Risk Evaluation for Eating and Nutrition (SCREEN-8),[83] pain using the Brief Pain Inventory-Short Form[84], and health status with the EuroQol-5D-5L (EQ-5D-5L).[85] Additionally, cognitive[86] and hearing[87] function, social support,[88–90] and mental health[91,92] are assessed through validated tools. These measures are captured using a combination of telephone interviews and online surveys at baseline, 12-month, and 24-month follow-ups.

To monitor health and mobility outcomes longitudinally, participants are contacted at 4-month intervals from baseline until the conclusion of the study. During these calls, the PCML[19,20] is re-administered to track changes over time and to identify the development of mobility disability. Participants also rate their perceived changes in mobility using the Global Rating of Change (GRC)[93] scale and answer questions about healthcare utilization, including hospitalizations, emergency room visits, primary care visits, rehabilitation service use, motor vehicle collisions, new diagnoses, living situation updates, and general and mental health status.

Falls are measured prospectively using monthly fall diary calendars, which are provided to participants along with pre-addressed and pre-stamped postcards. This method of data collection for falls is regarded as the gold standard by international expert consensus.[94] As part of the data quality checks, the research team calls participants to rectify missing data and confirm reported falls, ensuring they fit the definition of a fall[95] and documenting circumstances. Participants who report falls are requested to complete a standardized case report form via telephone to collect detailed information, with fall-related injuries categorized as serious, moderate, or minor based on symptoms and healthcare use. Participants are encouraged to report injurious falls promptly for accurate documentation. Table 1 provides an overview of the measures and corresponding assessment time points.

Table 1. Outcome measures and time points of the MacM3 cohort study.

Model of Assessment	Outcomes		Follow-ups						
			Base-line	4M	8M	12M	16M	20M	24M
	Measures								
Free-living Mobility Protocol	Wearables	Accelerometry and GPS data	✓	✓	✓	✓	✓	✓	✓
Phone Call Assessments	Mobility/Function	Late Life Function and Disability Instrument (LLFDI)	✓			✓			✓
		Life-space assessment (LSA)	✓			✓			✓
		Mobility Screening questions	✓	✓	✓	✓	✓	✓	✓
		Global Rating Change scale (GRC scale)	✓	✓	✓	✓	✓	✓	✓
	Driving Mobility	Transportation mobility questions (TRA)	✓			✓			✓
		Driving Comfort Scale (DCS)	✓			✓			✓
		Perceiving Driving Abilities (PDA)	✓			✓			✓
In-Person Assessments	Performance-based tests and other measures	Mobility Assessment Tool (MAT-sf); TUG (Regular/Dual-task) Repeated Sit-to-Stand Test (5 x STS); Gait Speed 3 meters (usual /fast pace); Single Leg Stance; Stair climbing power test; 400-meter walk; Visual Acuity test; Lung Function; Clinical Frailty Scale (CFS) and Grip Strength	✓			✓			✓
	Comprehensive Health and Well-being Measures								
Phone Call Assessments	Demographics, socioeconomic and health status	Age, country of birth, race and ethnicity, education	✓						
		Height, weight, sex and gender, marital status, employment situation, income, housing and living situation, self-reported chronic conditions, smoking, alcohol use, care giving and care receiving, medication, nutritional supplements and cannabis use, pain, balance, fall history and fear of falling, neighborhood environment, technology	✓			✓			✓
		Health care utilization, new diagnoses, general health, mental health, car crashes, mobility aids, falls and near falls information, and injuries	✓	✓	✓	✓	✓	✓	✓
	COVID-19	COVID-19 and vaccination status	✓	✓	✓	✓	✓	✓	✓
	Cognition	Montreal Cognitive Assessment (MoCA) (telephone version)	✓			✓			✓
	Physical Activity	Physical Activity Scale for the Elderly (PASE)	✓			✓			✓

			1	2	3	4	5	6	7
	Social Support	*Medical Outcome Study Social Support Survey (MOS-SSS)	✓			✓			✓
	Depression	*The Centre for Epidemiological Studies Depression Scale (CES-D-10)	✓			✓			✓
	Hearing	*Hearing Handicap Inventory for the Elderly Screening Version (HHIE-s)	✓			✓			✓
	Nutrition	*Risk Evaluation for Eating and Nutrition (SCREEN-8)	✓			✓			✓
	Quality of Life	*EuroQol-5D-5L (EQ-5D-5L)	✓			✓			✓
	Anxiety	*General Anxiety Disorder (GAD-7)	✓			✓			✓
	Loneliness	*20-item Loneliness scale (UCLA)	✓			✓			✓
	Social Network	*Lubben Social Network Scale (LSNS 6)	✓			✓			✓
	Pain	*Brief Pain Inventory				✓			✓
Monthly Diaries	Falls	Occurrence of falls; injurious falls	Monthly						

M = month; *Option to complete during the in-person physical assessment or at home in a survey format.

Patient and Public Involvement

Older adult stakeholders (n=7) were actively engaged throughout the development of the MacM3 cohort study. They provided valuable feedback on various aspects of the project, including the study concept (interest and perceived value of mobility monitoring) and data collection methods (i.e., selection of the wearable device and wear location). While older adult stakeholders were not directly involved in defining the research questions or selecting outcome measures, their feedback informed the perceived value, feasibility, and burden of study procedures.

We plan to involve both stakeholder and community advisory groups in the interpretation and dissemination of study findings, including selecting relevant information and appropriate formats for sharing results with participants and wider communities. We will continue to consult with our adult partners regarding the interpretation of study findings, and strategies for translating research outcomes into practical applications. This ongoing engagement helps ensure the research remains relevant to the needs and concerns of the community it seeks to benefit. To maintain this strong connection, the project plans to meet with stakeholders at least once a year, offering a forum for continued dialogue and input.

As part of our partnership with the Dixon Hall Centre in Toronto, we also consulted extensively with older adults and social service providers in the community to ensure our data collection methods were as inclusive as possible. A community advisory group was assembled (11 members) who continue to provide input on all stages of our research at the Centre which has resulted in key modifications to our protocol to facilitate participant recruitment and retention including: options for retuning the device in person, Mandarin speaking assessors, drop-in hours for trouble shooting, gift cards for returning devices on time, option to complete online survey in-person, and distributing flyers that outlined our commitment to addressing the underrepresentation of diverse older adults in research—while also emphasizing safety and security. This comprehensive approach to stakeholder involvement ensures that the MacM3 cohort reflects a wide range of experiences and perspectives, strengthening the overall impact

and generalizability of the research.

Results

Participants characteristics

We aimed to enroll a total of 1500 participants, and as of May 2024, recruitment was completed with 1209 participants enrolled at the Hamilton site and 346 at the Dixon Hall site, for a total of 1555 participants. Figure 2 displays the geographic distribution of participant addresses, demonstrating a concentration of recruitment in Hamilton and Toronto, Ontario, aligning with the study's targeted catchment area. The inset highlights the dense and well-distributed recruitment pattern within Hamilton, indicating strong geographic coverage across the city. Based on Statistics Canada's Population Centre and Rural Area Classification, most participants (n = 1,416) reside in large population centers. A smaller number live in medium (n = 5) and small (n = 57) population centers, and 76 participants were recruited from rural areas. One participant's address was missing and could not be classified.

Figure 2. Geographic distribution of MacM3 participants.

Figure 3 illustrates the flow of study participants from recruitment through each follow-up period, highlighting participant retention and attrition trends over time. A total of 2,135 individuals expressed interest in the MacM3 study. Of these, 140 were deemed ineligible—primarily due to major mobility limitations (n=117), based on our PCML screening survey. Additionally, 440 individuals declined to participate. As a result, 1,555 participants consented to the MacM3 study. Baseline assessments were completed for all enrolled participants, along with 4-, 8-, and 12-month follow-ups, with 1,378 participants completing the 12-month assessment. The study continues to demonstrate good retention, with an overall attrition rate of just 16.0%. The highest dropout rate occurred at the beginning of the study, between baseline and 4 months, where many participants (n=57, 3.7%) dropped out after completing all components of the baseline assessment. The most common reasons for dropout included participants losing interest, becoming unreachable, device discomfort or experiencing health-related challenges. Mortality has been reported in 12 participants (0.7%) during follow-up to date. As of the latest follow-up, 859 participants have successfully completed their 24-month follow-up; data collection is still ongoing. The study's overall high retention rate underscores the feasibility of long-term data collection using wearable devices and highlights the effectiveness of our engagement strategies in sustaining participant involvement (Figure 3).

Table 2 presents a comprehensive overview of the sociodemographic, lifestyle, and health characteristics of the MacM3 participants at baseline. Among the total cohort (n = 1555), 66.3% were female, and the mean age at baseline was 73.9 years (SD = 5.5). A majority of participants (62.4%) were married or living with a partner, and 75.3% had attained a post-secondary diploma. Additionally, 59.8% reported never having smoked, and 83.3% were regular drinkers. In terms of health-related measures, 63.0% reported having three or more comorbidities, with cataracts, osteoarthritis and hypertension being the most reported conditions. Furthermore, 81.2% were taking prescribed medication(s), and 44.3% and 41.7% rated their general health and mental health, respectively, as 'very good.'

More than two-thirds of the participants were born in Canada (69.4%). Most identified as White/Caucasian (88.2%), with smaller proportions identifying as Chinese (7.0%) or another ethnic background such as Black, Japanese, South Asian, Latin American, North American

Figure 3. Flow of study participants and timeline of data collection in the MacM3 study, from recruitment through September 2025. Grey areas indicate completed follow-up periods.

Indigenous, Métis, Arab, or Korean (5.7%). Ethnic diversity was more pronounced at the Dixon Hall site, where less than half of the participants were born in Canada. In terms of cultural background, participants frequently reported English (40.0%), Scottish (25.1%), Irish (24.1%), and Canadian (14.5%) heritage, with others identifying German, Chinese, French, Italian, Polish, Dutch, or other cultural backgrounds. Cultural diversity was also more evident at the Dixon Hall site, where 39.0% of male participants and 27.0% of females reported Chinese cultural heritage, and a wide range of other cultural identities were selected, including Ukrainian, Métis, and North American Indigenous backgrounds.

Table 2. Selected baseline characteristics of the MacM3 participants.

			Hamilton site		Dixon Hall site	
		Total	Males *n* (%)	Females *n* (%)	Males *n* (%)	Females *n* (%)
Sample size, *n* (%)		1555	408 (33.7)	801 (66.3)	83 (24.0)	263 (76.0)
Age (years), mean (SD)		73.9 (5.5)	74.4 (5.7)	73.5 (5.4)	74.8 (5.8)	73.7 (5.5)
BMI (kg/m^2), mean (SD)		26.3 (4.8)	27.0 (4.2)	26.4 (4.9)	25.8 (5.9)	25.2 (5.0)
Country of Birth, *n* (%)	Born in Canada	1,079 (69.4)	296 (72.6)	620 (77.4)	35 (42.2)	128 (48.7)
Ethnicity *(participants could select multiple ethnicities)*	White/Caucasian	1369 (88.2)	388 (95.1)	769 (96.2)	48 (58.5)	164 (62.4)
	Chinese	108 (7.0)	1 (0.2)	4 (0.5)	32 (39.0)	71 (27.0)
	Other (Black, Japanese, South Asian, Latin American, North American Indian, Metis, Arab, Korean, other)	88 (5.7)	25 (6.1)	31 (3.9)	2 (2.4)	30 (11.4)
Cultural backgrounds, *n* (%) *(participants could select multiple cultural identities)*	English	622 (40.0)	165 (40.4)	372 (46.4)	17 (20.5)	68 (25.9)
	Scottish	391 (25.1)	99 (24.3)	237 (29.6)	14 (16.9)	41 (15.6)
	Irish	375 (24.1)	109 (26.7)	211 (26.3)	5 (6.0)	50 (19.0)
	Canadian	226 (14.5)	70 (17.2)	142 (17.7)	4 (4.8)	10 (3.8)
	German	157 (10.1)	48 (11.8)	80 (10.0)	6 (7.2)	23 (8.8)
	Chinese	109 (7.0)	1 (0.2)	5 (0.6)	32 (38.6)	71 (27.0)
	French	104 (6.7)	29 (7.1)	54 (6.7)	1 (1.2)	20 (7.6)
	Italian	83 (5.3)	23 (5.6)	52 (6.5)	1 (1.2)	7 (2.7)
	Polish	83 (5.3)	22 (5.4)	38 (4.7)	8 (9.6)	15 (5.7)
	Dutch	67 (4.3)	19 (4.7)	38 (4.7)	2 (2.4)	8 (3.0)
	Other (Ukrainian, Jewish, Welsh, South Asian, Norwegian, Portuguese, Metis, North American Indian, other)	470 (30.2)	130 (31.9)	220 (27.5)	21 (25.3)	99 (37.6)
Marital Status, *n* (%)	Married/living with a partner	970 (62.4)	337 (82.6)	456 (57.0)	67 (80.7)	110 (41.8)
	Widowed	229 (14.7)	22 (5.4)	158 (19.8)	4 (4.8)	45 (17.1)
	Divorced	174 (11.2)	15 (3.7)	110 (13.8)	8 (9.6)	41 (15.6)
	Separated	36 (2.3)	9 (2.2)	16 (2.0)	0 (0.0)	11 (4.2)
	Single	145 (9.3)	25 (6.1)	60 (7.5)	4 (4.8)	56 (21.3)

Education, n (%)	Secondary school incomplete	54 (3.5)	12 (2.9)	26 (3.3)	2 (2.4)	14 (5.3)
	Secondary school completed	158 (10.2)	53 (13.0)	74 (9.2)	6 (7.2)	25 (9.5)
	Some post-secondary education	172 (11.1)	44 (10.8)	103 (12.9)	7 (8.4)	18 (6.8)
	Post-secondary diploma	1,171 (75.3)	299 (73.3)	598 (74.7)	68 (81.9)	206 (78.3)
*Work status, n (%)		1321	338	641	84	262
(participants could select multiple work status)	Retired	1084 (82.0)	285 (84.3)	541 (84.4)	60 (71.4)	198 (75.6)
	Working (part time or full time)	125 (9.5)	37 (10.9)	57 (8.9)	11 (13.1)	20 (7.6)
	Volunteer	75 (5.7)	7 (2.1)	27 (4.2)	11 (13.1)	30 (11.5)
	Other	37 (2.8)	9 (2.7)	16 (2.5)	2 (2.4)	14 (5.3)
Household income, n (%)	< $20,000	86 (5.5)	3 (0.7)	5 (0.6)	21 (25.3)	57 (21.7)
	$20,000-49,999	331 (21.3)	57 (14.0)	189 (23.6)	15 (18.1)	70 (26.6)
	$50,000-$99,999	654 (42.1)	184 (45.1)	369 (46.1)	20 (24.1)	81 (30.8)
	≥ $100,000	399 (25.7)	147 (36.0)	178 (22.2)	26 (31.3)	48 (18.3)
	Don't know/Prefer not to answer	85 (5.5)	17 (4.2)	60 (7.5)	1 (1.2)	7 (2.7)
Smoking status, n (%)	Current smoker	31 (2.0)	11 (2.7)	16 (2.0)	1 (1.2)	3 (1.1)
	Former smoker	594 (38.2)	195 (47.8)	298 (37.2)	36 (43.4)	65 (24.7)
	Never smoked	930 (59.8)	202 (49.5)	487 (60.8)	46 (55.4)	195 (74.1)
Alcohol consumption past year, n (%)	Regular drinker (> once a month)	1,041 (83.3)	315 (88.0)	552 (81.7)	42 (87.5)	132 (78.6)
	Occasional drinker (< once a month)	206 (16.5)	42 (11.7)	122 (18.1)	6 (12.5)	36 (21.4)
	Non-drinker	3 (0.2)	1 (0.3)	2 (0.3)	0 (0.0)	0 (0.0)
Comorbidities, n (%)	None	54 (3.5)	21 (5.15)	21 (2.6)	2 (2.4)	10 (3.8)
	< 3	522 (33.6)	126 (30.9)	269 (33.6)	30 (36.1)	97 (36.9)
	≥ 3	979 (63.0)	261 (64.0)	511 (63.8)	51 (61.5)	156 (59.3)
Common conditions, n (%)	Cataracts	684 (44.0)	158 (38.7)	358 (44.7)	40 (48.2)	128 (48.7)
	Osteoarthritis	652 (41.9)	145 (35.5)	381 (47.6)	24 (28.9)	102 (38.8)
	Hypertension	568 (36.5)	188 (46.1)	262 (32.7)	36 (43.4)	82 (31.2)
	Back Pain	268 (17.2)	68 (16.7)	141 (17.6)	15 (18.1)	44 (16.7)

	Osteoporosis	247 (15.9)	11 (2.7)	151 (18.9)	11 (13.3)	74 (28.1)
	Cancer	246 (15.8)	79 (19.4)	98 (12.2)	17 (20.5)	52 (19.8)
	Diabetes	174 (11.2)	73 (17.9)	68 (8.5)	15 (18.1)	18 (6.8)
	Depression	170 (10.9)	30 (7.4)	94 (11.7)	5 (6.0)	41 (15.6)
Prescribed medication, *n* (%)	Yes	1,261 (81.2)	355 (87.5)	652 (81.5)	71 (85.5)	183 (69.6)
	No	291 (18.8)	51 (12.5)	148 (18.5)	12 (14.5)	80 (30.4)
Number of prescribed medications, *n* (%)	0	291 (18.8)	51 (12.5)	148 (18.5)	12 (14.5)	80 (30.4)
	1	312 (20.1)	56 (13.8)	169 (21.1)	13 (15.7)	74 (28.1)
	2	265 (17.1)	62 (15.3)	143 (17.9)	15 (18.1)	45 (17.1)
	3	243 (15.7)	66 (16.3)	132 (16.5)	18 (21.7)	27 (10.3)
	4	169 (10.9)	53 (13.1)	88 (11.0)	8 (9.6)	20 (7.6)
	5 or more	272 (17.5)	118 (29.1)	120 (15.0)	17 (20.5)	17 (6.5)
Self-rated general health, *n* (%)	Excellent	258 (16.6)	58 (14.2)	134 (16.8)	7 (8.4)	59 (22.6)
	Very good	687 (44.3)	184 (45.1)	368 (46.0)	27 (32.5)	108 (41.4)
	Good	474 (30.5)	129 (31.6)	248 (31.0)	36 (43.4)	61 (23.4)
	Fair	123 (7.9)	34 (8.3)	49 (6.1)	11 (13.3)	29 (11.1)
	Poor	10 (0.6)	3 (0.7)	1 (0.1)	2 (2.4)	4 (1.5)
Self-rated mental health, *n* (%)	Excellent	286 (18.4)	77 (18.9)	129 (16.1)	18 (21.7)	62 (23.9)
	Very good	647 (41.7)	177 (43.4)	332 (41.5)	26 (31.3)	112 (43.1)
	Good	515 (33.2)	130 (31.9)	295 (36.8)	28 (33.7)	62 (23.9)
	Fair	89 (5.7)	21 (5.1)	41 (5.1)	7 (8.4)	20 (7.7)
	Poor	15 (1.0)	3 (0.7)	4 (0.5)	4 (4.8)	4 (1.5)

Findings to date

Table 3 describes the self-reported mobility and performance-based test results at baseline. Most participants (93.8%) did not use mobility aids, and 37.6% reported experiencing a fall in the previous year. According to the PCML classification, 56.7% of participants had no mobility limitations, while 15.9% had preclinical mobility limitations, and 27.4% had minor mobility limitations at baseline.

For the performance-based tests, the mean of the best right- or left-hand grip strength in the overall cohort was 27.3 kg (SD = 9.4). The mean gait speed was 1.23 m/s (SD = 0.02), the mean Timed Up and Go test time was 9.4 seconds (SD = 2.5), and the mean 5x sit-to-stand test time was 13.0 seconds (SD = 3.8). Results are also stratified by sex. Males generally demonstrated higher grip strength and stair climbing power, while other functional measures were relatively consistent across groups. Overall, these results reflect a relatively high-functioning sample of older adults at baseline, which is consistent with our recruitment strategy.[96]

Preliminary results - wearables data

At baseline, data from 1412 participants were processed. Wear time in our study was defined as at least 10 hours per day on a minimum of 7 days. Based on this criterion, 62 participants had no valid wrist wear data, and 49 wore the wrist device for fewer than 7 valid days. A total of 1,301 participants had valid wrist-worn device data, with an average wear time of 13.6 hours per day (range: 7.2 to 18.3 hours) over 10.2 days. For the thigh-worn device, 309 participants declined to wear it at baseline, and 95 did not meet the wear-time criteria. As a result, data from 1,008 participants are presented. On average, participants wore the thigh device for 13.6 hours per day (range: 6.1 to 22.5 hours) over approximately 10.6 days.

Table 3. Baseline self-reported mobility, falls and performance-based tests results.

		Total Sample n (%)	Males n (%)	Females n (%)
Self-reported mobility, *n*		1555	491	1064
Preclinical Mobility Disability Scale	No mobility limitation	881 (56.7)	298 (60.7)	583 (54.8)
	Preclinical mobility limitation	248 (15.9)	67 (13.6)	181 (17.0)
	Minor mobility limitation	426 (27.4)	126 (25.7)	300 (28.2)
Mobility aids, *n*		1553	490	1063
	Yes	96 (6.2)	30 (6.1)	66 (6.2)
	No	1457 (93.8)	460 (93.9)	997 (93.8)
Falls in the past year, *n*		1555	491	1064
	Yes	584 (37.6)	169 (34.4)	415 (39.0)
	No	971 (62.4)	322 (65.6)	649 (61.0)
Performance-based tests				
	[a]Grip Strength (Kg), *n*	1504	479	1025
	mean ± SD	27.3 (9.4)	37.2 (8.3)	22.7 (5.5)
	Gait speed (m/s), *n*	1506	480	1026
	mean ± SD	1.23 ± 0.2	1.24 (0.2)	1.22 (0.2)
	Timed up and go (s), *n*	1505	479	1026 9.4
	mean ± SD	9.4 ± 2.5	9.5 (2.5)	(2.6)
	5 × sit-to-stand (s), *n*	1472	466	1006
	mean ± SD	13.0 ± 3.8	12.9 (3.5)	13.0 (4.0)
	[a]Single leg stance (s), *n*	1467	463	1004
	mean ± SD	29.6 (22.7)	26.4 (22.1)	31.1 (22.8)
	[b]400 meters walk test (min), *n*	1109	381	728
	mean ± SD	4.9 ± 0.9	4.8 (1.0)	4.9 (0.9)
	[b]Stair climbing Power (W), *n*	1140	389	751
	mean ± SD	258.1 (84.0)	313.3 (89.9)	229.5 (64.2)

[a]best of left and right test; [b]The 400 meters walk test and the stair climbing power test are not conducted at DH site.; W = watts

Table 4 presents the mean and median daily step counts by quartile. Step counts ranged from 5,073 steps in the lowest quartile to 12,303 steps in the highest. Participants with higher activity levels (Q3: 72.9 years; Q4: 72.7 years) were slightly younger than those with lower activity (Q1: 74.6 years; Q2: 74.7 years). Mobility status also varied with activity level: 73.8% of those in the highest quartile reported no mobility limitations, versus only 36.9% in the lowest quartile. Wear-time compliance was consistent across all quartiles, with a median of 10 compliant days and average daily wear time ranging from 13.8 to 14.6 hours. These data are consistent with findings from other large cohort studies showing a decline in step counts with advancing age and increasing mobility limitations.[97,98]

Trip data from 1,422 participants were processed using the algorithms proposed by Fillekes et al.[29,51] , as described in Table 5. Applying the study's compliance criteria of ≥7 days of wrist wear with ≥10 hours per day, data from 1,323 participants are presented. On average participants wore the wrist device for 12.1 (SD = 1.3) hours. In terms of mobility outcomes, participants recorded an average of 2.0 active trips per day (SD = 1.3) and 1.8 passive trips per day (SD = 1.2). Active and passive trips ranged from fewer than one trip in the lowest quartile to over three trips in the highest quartiles. Participants in the highest quartile were, on average, younger and had fewer mobility limitations.

Table 4. Baseline characteristics of participants by quartiles of mean steps per day derived from the thigh devices (n=1008).

Characteristics	Total Sample	Quartiles of mean steps per day, mean (SD)			
		Quartile 1 (Lowest)	Quartile 2	Quartile 3	Quartile 4 (Highest)
Total participants, n	1008	252	252	252	252
Age (years), mean (SD)	73.7 (5.4)	74.6 (5.5)	74.7 (5.5)	72.9 (5.2)	72.7 (4.9)
Mobility score, n (%)					
No mob limitation	582 (57.7)	93 (36.9)	149 (59.1)	154 (61.1)	186 (73.8)
Preclinical	166 (16.5)	49 (19.4)	41 (16.3)	47 (18.7)	29 (11.5)
Minor mob limitation	260 (25.8)	110 (43.7)	62 (24.6)	51 (20.2)	37 (14.7)
No. of Days with wear (≥7 days)					
Mean (SD)	9.9 (1.2)	9.7 (1.2)	9.8 (1.0)	10.0 (1.5)	10.0 (1.0)
median	10	10	10	10	10
Hours of wear on compliant days (≥ 10h)					
Mean (SD)	14.1 (1.4)	13.8 (1.4)	14.1 (1.3)	14.1 (1.3)	14.6 (1.2)
Median	14.1	13.7	14.0	14.1	14.6
Steps/day					
Mean (SD)	8437 (2943)	5073 (962)	7306 (490)	9065 (560)	12303 (2283)
Median	8175	5278	7318	8971	11694

Table 5. Baseline characteristics of trip mode derived from the wrist device (n=1323).

Characteristics	Total Sample	Quartiles of active trip mode, mean (SD)			
		Quartile 1 (Lowest)	Quartile 2	Quartile 3	Quartile 4 (Highest)
Total participants, n	1323	349	341	318	315
Age (years), mean (SD)	73.8 (5.3)	74.9 (5.6)	73.0 (5.3)	73.7 (4.9)	73.5 (5.4)
Mobility score, n (%)					
No mob limitation	772 (58.4)	171 (49.0)	211 (61.9)	190 (59.7)	200 (63.5)
Preclinical	209 (15.8)	52 (14.9)	49 (14.4)	60 (18.9)	48 (15.2)
Minor mob limitation	342 (25.8)	126 (36.1)	81 (23.7)	68 (21.4)	67 (21.3)
No. of Days with wear (≥7 days)					
Mean (SD)	9.8 (4.5)	9.9 (1.6)	10.0 (1.7)	9.8 (1.3)	9.7 (1.1)
median	10	10	10	10	10
H/d of wear on compliant days (≥ 10h)					
Mean (SD)	12.1 (1.3)	11.9 (1.3)	12.0 (1.4)	12.1 (1.3)	12.3 (1.4)
Median	12.0	11.8	11.9	12.1	12.3
Active trips/day					
Mean (SD)	2.0 (1.3)	0.5 (0.3)	1.5 (0.3)	2.4 (0.3)	3.7 (0.8)
Median	1.9	0.6	1.5	2.4	3.6
		Quartiles of passive trip mode, mean (SD)			
		Quartile 1 (Lowest)	Quartile 2	Quartile 3	Quartile 4 (Highest)
Total participants, n	1323	336	359	313	315
Age (years), mean (SD)	73.8 (5.3)	74.7 (5.7)	74.4 (5.6)	73.2 (4.6)	72.7 (5.1)
Mobility score, n (%)					
No mob limitation	772 (58.4)	185 (55.0)	193 (53.8)	191 (61.0)	203 (64.4)
Preclinical	209 (15.8)	54 (16.1)	56 (15.6)	53 (17.0)	46 (14.6)
Minor mob limitation	342 (25.8)	97 (28.9)	110 (30.6)	69 (22.0)	66 (21.0)
No. of Days with wear (≥7 days)					
Mean (SD)	9.8 (4.5)	9.9 (1.4)	9.8 (1.5)	9.8 (1.2)	9.9 (1.7)

median	10	10	10	10	10
H/d of wear on compliant days (≥ 10h)					
Mean (SD)	12.1 (1.3)	11.8 (1.4)	12.1 (1.3)	12.2 (1.3)	12.2 (1.3)
Median	12.0	11.5	12.0	12.1	12.2
Passive trips/day					
Mean (SD)	1.8 (1.2)	0.3 (0.3)	1.4 (0.3)	2.2 (0.2)	3.5 (0.7)
Median	1.8	0.3	1.4	2.2	3.3

Strengths and limitations

To our knowledge, MacM3 is the **first longitudinal study designed to understand trajectories of mobility decline in a generalizable sample of community-dwelling older adults using a body-worn monitoring system that integrates GNSS and accelerometry**. The main strengths of the MacM3 study include its repeated measures and extended follow-up period, its innovative approach to mobility monitoring with advanced wearable technologies, the richness of the health and mobility data collected, and the cohort's high participation and retention rate to date.

Although the use of wearable technology to measure real-world mobility holds great promise for improving our understanding of healthy aging, there are several challenges in this field. There is wide variability in the protocols used and parameters reported from wearable technology, with few studies evaluating their validity.[18, 32, 33] Wear location of a sensor, for example, is critical to generating valid and sensitive outcomes.[34] Additionally, much of the literature on wearable technology to date is from small cross-sectional studies, specific disease groups, or single sensor designs that can limit accuracy, generalizability and precision in the study of mobility in older adults.[35] In recognition of some of these issues, a large European consortium—Mobilise-D— was formed to validate digital mobility measures for health and regulatory agencies. Mobilise-D is conducting a series of clinical validation studies in four specific disease cohorts (each n = 600) focusing on accelerometry-derived mobility measures such as gait speed.[18] Our work complements that of Mobilise-D, as MacM3 will be the first cohort dedicated to evaluating digital mobility measures in large sample of community-dwelling older adults. Additionally, based on our extensive pilot testing,[37] we have innovated a novel multi-sensor, multi-nodal approach that combines GNSS and accelerometry to optimize the potential for identifying early changes in mobility that could inform prevention and early intervention strategies from health, environmental, and social perspectives. This is a significant step forward in the study of real-world mobility in community-dwelling older adults, as GPS technology can provide an understanding of spatial properties of mobility patterns and environmental influences but lacks the motion and temporal fidelity to provide details of

movement or walking. In contrast, accelerometry data provides details of movement and walking kinematics with continuous millisecond resolution but lacks spatial context. By temporally aligning accelerometry and GPS data, we leverage their complementary strengths to enhance the collection and interpretation of comprehensive mobility data, particularly for understanding environment-movement interactions. Furthermore, our cohort's size, diversity (race, ethnicity, income, and ability), and range of health conditions and mobility levels will allow us to validate digital mobility measures with broad applicability for early risk assessment and intervention in community-dwelling older adults.

The main limitations of MacM3 include our convenience sample with recruitment from only 2 sites in Ontario, Canada, with a large proportion of individuals with post-secondary education and higher income levels. Our sample also consists of more females than males. The gender imbalance may similarly affect the generalizability of the findings, as males and females may have different patterns of mobility, health outcomes and health utilization. Another limitation is that wearing the thigh device was optional in this study, leading to the absence of key gait and posture-related behavior data for participants who chose not to wear it. Nevertheless, this limitation is mitigated by the high percentage of participants who agreed to wear the device, maintaining the overall usefulness of the dataset for studying gait and posture.

Future Plans

Our future research will focus on leveraging the rich, longitudinal, multi-sensor data from MacM3 to develop predictive models of mobility decline. These models will enable early identification of individuals at risk of developing mobility limitations, guiding targeted interventions to prevent or mitigate disability. A key goal is to explore the integration of wearable-derived mobility metrics with clinical and environmental factors to personalize strategies for promoting functional independence.

Opportunities for extending the follow-up in MacM3 will also be sought to enhance the usefulness of the data platform and allow for rigorous validation of mobility measures obtained through multi-sensor wearable technology for monitoring and predicting long-term health outcomes in older adults.

Another critical avenue involves advancing the analytic capabilities of our open-source data processing pipeline, enhancing the fusion of GPS and IMU data to investigate the interaction between mobility behaviors and environmental contexts. By incorporating machine learning techniques and stakeholder input, we aim to translate these innovations into actionable tools for clinicians, caregivers, and older adults themselves. Ultimately, these efforts will contribute to developing scalable solutions that support aging-in-place, improve health outcomes, and enhance quality of life for aging populations.

Collaboration

We encourage collaboration on projects that align with the goals of our study, particularly those focused on mobility and healthy aging. The dataset includes wearable sensor data (accelerometry and GPS), questionnaire responses, demographic and health information, as well as data from physical assessments. Researchers interested in accessing the data will be required to submit a proposal outlining their intended use, which will be reviewed by the study team. Data will be shared in a de-identified format whenever possible. However, certain restrictions may apply—particularly for GPS data—due to the sensitive nature of location information. Access to such data will require additional justification and safeguards to protect participant confidentiality.

Authors affiliation

[1]School of Rehabilitation Science, McMaster University, Hamilton, Ontario, Canada
[2]McMaster Institute for Research on Aging (MIRA), McMaster University, Hamilton, Ontario, Canada
[3]MIRA|Dixon Hall Centre, McMaster University, Hamilton, Ontario, Canada
[4]Department of Kinesiology and Health Sciences, University of Waterloo, Waterloo, Ontario, Canada
[5] Department of Mathematics and Statistics, McMaster University, Hamilton, Ontario, Canada
[6]School of Earth, Environment & Society, McMaster University, Hamilton, Ontario, Canada
[7]Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada
[8]Department of Engineering Physics, McMaster University, Hamilton, Ontario, Canada
[9]Faculty of Humanities, McMaster University
[10]Faculty of Business, Health Policy & Management, McMaster University, Hamilton, Ontario Canada

MacM3 Investigators (Collaborators)

Jamal Deen, Electrical and Computer Engineering Department, McMaster University; Vanina Dal Bello-Haas, McMaster University; Rebecca Ganann, School of Nursing, McMaster University; Jack Guralnik, Department of Epidemiology and Public Health, University of Maryland School of Medicine; Lori Letts, School of Rehabilitation Science, McMaster University; Sarrah Lal, Faculty of Health Sciences, McMaster University; Stuart Phillips, Department of Kinesiology, McMaster University, McMaster University; Nigar Sekercioglu, Health Research Methods, Evidence, and Impact, McMaster University; Kathryn Sibley, Max Rady College of Medicine, University of Manitoba.

Authors Contribution: MKB, JR, AK, KBN, DMS, PR, QF, PG, MZ, PDM, and MZ conceived the study. MKB, JR, AK, PDM, KBN, DMS, PR, QF, PM, MZ, JD, VDBH, RG, JG, LL, SL, SP, NS and KS contributed to the study design. MKB, RK, CC and SO were responsible for the study planning and data collection. MKB, RK, CC, WM, KVO and KB were responsible for developing the protocol for analyzing the wearables data. MKB, RK, JM, and TR were responsible for the analysis. MKB, RK and TH were responsible for writing the initial draft of the manuscript, and subsequent drafts were reviewed by all authors listed. All authors had input on the interpretation and reporting of study findings. All authors provided approval for the published version of this manuscript. MKB is the guarantor of the work and accepts full responsibility for the integrity of the data and the accuracy of the analysis.

Acknowledgements: The authors would like to thank the study coordinators, Tara Noble and Lisette Machado, and the research assistants—Kathleen Romanoski, Jasmyn Stoffers, Rhea Sinha, Mae Vidal, Shalini Jaisankar Sumathi and Catherine Deng — for their support throughout data collection and participant coordination.

Funding Statement: This work was supported by major research program funding from the Labarge Centre for Mobility in Aging within the McMaster Institute for Research on Aging at McMaster University, and by AGE-WELL NCE Inc.—a member of Canada's Networks of Centres of Excellence program (award numbers AWCAT-2019-139 and AWCRP-2020-13). Marla Beauchamp holds a Tier 2 Canada Research Chair in Mobility, Aging, and Chronic Disease (CRC-2020-00043) and an Early Researcher Award from the Government of Ontario.

Competing interests: All authors have no competing interest to declare.

Ethics approval: This study was approved by the Hamilton Integrated Research Ethics Board (HiREB), McMaster University (Project number: 13905).

Data availability statement: Access to MacM3 data is currently unavailable to other researchers. We are working on developing processes to make de-identified data available to qualified researchers for approved analyses, following the review and approval of a study proposal, data use agreement, and ethical approval. For more information, please contact the corresponding author at beaucm1@mcmaster.ca.

ORCID ids

Marla K Beauchamp: https://orcid.org/0000-0003-2843-388X
Renata Kirkwood: https://orcid.org/0000-0002-8320-3342
William McIlroy: https://orcid.org/0000-0003-2319-1746
Julie Richardson: https://orcid.org/0000-0001-6356-0492
Ayse Kuspinar: https://orcid.org/0000-0002-3886-0407
Paul D. McNicholas: https://orcid.org/0000-0002-2482-523X
K Bruce Newbold: https://orcid.org/0000-0002-6441-7863
Darren M. Scott: https://orcid.org/0000-0002-1545-3737
Qiyin Fang: https://orcid.org/0000-0003-2786-9884
Paula Gardner: https://orcid.org/0000-0002-2190-8021

References

1.   World Report on Ageing and Health Geneva, Switzerland: Geneva, Switzerland: World Health Organization, 2015. http://apps.who.int/iris/bitstream/10665/186463/1/9789240694811%5Feng.pdf. (2015).

2.   Striuk, M. I., Striuk, M. I., Striuk, A. M. & Semerikov, S. O. Mobility in the information society: a holistic model. *Educational Technology Quarterly* **2023**, 277–301 (2023).

3.   Brabrand, M., Kellett, J., Opio, M., Cooksley, T. & Nickel, C. H. Should impaired mobility on presentation be a vital sign? *Acta Anaesthesiol Scand* **62**, 945–952 (2018).

4.   Middleton, A., Fritz, S. L. & Lusardi, M. Walking speed: the functional vital sign. *J Aging Phys Act* **23**, 314–22 (2015).

5.   Richardson, J. *et al.* Monitoring physical functioning as the sixth vital sign: evaluating patient and practice engagement in chronic illness care in a primary care setting--a quasi-experimental design. *BMC Fam Pract* **13**, 29 (2012).

6.   Now and Tomorrow Excellence in Everything We Do Disability Facts about Seniors. https://publications.gc.ca/collections/collection_2011/rhdcc-hrsdc/HS64-11-1-2010-eng.pdf.

7.   Public Health Agency of Canada. The Chief Public Health Officer's Report on The State of Public Health in Canada 2010 – Growing older: Adding life to years 2010. https://www.canada.ca/en/public-health/corporate/publications/chief-public-health-officer-reports-state-public-health-canada/annual-report-on-state-public-health-canada-2010.html.

8.   Statistics Canada. Canadians with a mobility disability 2020. https://www.canada.ca/en/public-health/corporate/publications/chief-public-health-officer-reports-state-public-health-canada/annual-report-on-state-public-health-canada-2010.html.

9.   Population Projections for Canada, Provinces and Territories, 2005 to 2031 . https://www150.statcan.gc.ca/n1/pub/91-520-x/91-520-x2005001-eng.pdf.

10.   Hirvensalo, M., Rantanen, T. & Heikkinen, E. Mobility difficulties and physical activity as predictors of mortality and loss of independence in the community-living older population. *J Am Geriatr Soc* **48**, 493–8 (2000).

11.   Penninx, B. W. J. H. *et al.* Lower extremity performance in nondisabled older persons as a predictor of subsequent hospitalization. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* **55**, M691–M697 (2000).

12.   Von Bonsdorff, M., Rantanen, T., Laukkanen, P., Suutama, T. & Heikkinen, E. Mobility limitations and cognitive deficits as predictors of institutionalization among community-dwelling older people. *Gerontology* **52**, 359–365 (2006).

13.   Jonkman, N. H. *et al.* Predicting Trajectories of Functional Decline in 60- to 70-Year-Old People. *Gerontology* **64**, 212 (2018).

14. Han, L. *et al.* Dynamics of Functional Aging Based on Latent-class Trajectories of Activities of Daily Living. (2012) doi:10.1016/j.annepidem.2012.11.010.

15. Ferrucci, L. *et al.* Age-related change in mobility: Perspectives from life course epidemiology and geroscience. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* **71**, 1184–1194 (2016).

16. Martin, L. G., Zimmer, Z. & Lee, J. Foundations of Activity of Daily Living Trajectories of Older Americans. (2015) doi:10.1093/geronb/gbv074.

17. Gill, T. M., Gahbauer, E. A., Han, L. & Allore, H. G. Trajectories of Disability in the Last Year of Life. *N Engl J Med* **362**, 1173 (2010).

18. Buurman, B. M. *et al.* Trajectories of disability among older persons before and after a hospitalization leading to a skilled nursing facility admission HHS Public Access. *J Am Med Dir Assoc* **17**, 225–231 (2016).

19. Chaves, P. H. M., Garrett, E. S. & Fried, L. P. Predicting the risk of mobility difficulty in older women with screening nomograms: The Women's Health and Aging Study II. *Arch Intern Med* **160**, 2525–2533 (2000).

20. Mänty, M. *et al.* Construct and Predictive Validity of a Self-Reported Measure of Preclinical Mobility Limitation. *Arch Phys Med Rehabil* **88**, 1108–1113 (2007).

21. Giannouli, E., Bock, O., Mellone, S. & Zijlstra, W. Mobility in old age: Capacity is not performance. *Biomed Res Int* **2016**, (2016).

22. Breasail, M. *et al.* Wearable gps and accelerometer technologies for monitoring mobility and physical activity in neurodegenerative disorders: A systematic review. *Sensors* **21**, (2021).

23. Saner, H. Wearable Sensors for Assisted Living in Elderly People. *Front. ICT* **5**, 1 (2018).

24. Kheirkhahan, M. *et al.* Actigraphy features for predicting mobility disability in older adults HHS Public Access. *Physiol Meas* **37**, 1813–1833 (2016).

25. Van Ooteghem, K. *et al.* User-centered design of feedback regarding health-related behaviors derived from wearables: An approach targeting older adults and persons living with neurodegenerative disease. *Digit Health* **9**, (2023).

26. Mikolaizak, A. S. *et al.* Connecting real-world digital mobility assessment to clinical outcomes for regulatory and clinical endorsement–the Mobilise-D study protocol. *PLoS One* **17**, (2022).

27. Stamatakis, E. *et al.* Association of wearable device-measured vigorous intermittent lifestyle physical activity with mortality. *Nature Medicine 2022 28:12* **28**, 2521–2529 (2022).

28. Paluch, A. E. *et al.* Daily steps and all-cause mortality: a meta-analysis of 15 international cohorts. *Lancet Public Health* **7**, e219 (2022).

29. Beauchamp, M. *et al.* Monitoring mobility in older adults using a Global Positioning System (GPS) smartwatch and accelerometer: A validation study. *PLoS One* **18**, 0296159 (2023).

30. Richardson, J. *et al.* Defining and Measuring Preclinical Mobility Limitation: An Expert Consensus Exercise Informed by a Scoping Review. *J Gerontol A Biol Sci Med Sci* **2023**, 1641–1650 (2023).

31. Viccaro, L. J., Perera, S. & Studenski, S. A. Is timed up and go better than gait speed in predicting health, function, and falls in older adults? *J Am Geriatr Soc* **59**, 887–892 (2011).

32. Beauchamp, M. K. *et al.* Mobility screening for fall prediction in the Canadian Longitudinal Study on Aging (CLSA): Implications for fall prevention in the decade of healthy ageing. *Age Ageing* **51**, (2022).

33. Lee, L. F. R. & Dall, P. M. Concurrent agreement between ActiGraph⌷ and activPAL⌷ in measuring moderate to vigorous intensity physical activity for adults. *Med Eng Phys* **74**, 82–88 (2019).

34. De Craemer, M. & Verbestel, V. Comparison of Outcomes Derived from the ActiGraph GT3X+ and the Axivity AX3 Accelerometer to Objectively Measure 24-Hour Movement Behaviors in Adults: A Cross-Sectional Study. *Public Health* **19**, 271 (2022).

35. Lamb, S. E., Jørstad-Stein, E. C., Hauer, K. & Becker, C. Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* **53**, 1618–22 (2005).

36. Glass, T. A. Conjugating the 'Tenses' of Function: Discordance Among Hypothetical, Experimental, and Enacted Function in Older Adults. *Gerontologist* **38**, 101–112 (1998).

37. Beauchamp, M. K. *et al.* A unified framework for the measurement of mobility in older persons. *Age and Ageing* vol. 52 IV82–IV85 Preprint at https://doi.org/10.1093/ageing/afad125 (2023).

38. Curtis S. Florence, PhD1, Gwen Bergen, PhD1, Adam Atherly, PhD2, Elizabeth Burns, MPH1, Judy Stevens, PhD1, and Cynthia Drake, M. Actigraphy features for predicting mobility disability in older adults. *Physiol Behav* **66**, 693–698 (2018).

39. Giurgiu, M. Assessment of 24-hour physical behaviour in adults via wearables: a systematic review of validation studies under laboratory conditions. *International Journal of Behavioral Nutrition and Physical Activity* **20**, 68 (2023).

40. Wu, W. J., Yu, H. Bin, Tai, W. H., Zhang, R. & Hao, W. Y. Validity of Actigraph for Measuring Energy Expenditure in Healthy Adults: A Systematic Review and Meta-Analysis. *Sensors (Basel)* **23**, 8545 (2023).

41. Matthews, C. E. *et al.* Measurement of Active and Sedentary Behavior in Context of Large Epidemiologic Studies. *Med Sci Sports Exerc* **50**, 266 (2018).

42. Bammann, K., Thomson, N. K., Albrecht, B. M., Buchan, D. S. & Easton, C. Generation and validation of ActiGraph GT3X+ accelerometer cut-points for assessing physical activity intensity in older adults. The OUTDOOR ACTIVE validation study. *PLoS One* **16**, e0252615 (2021).

43. Lyden, K., John, D., Dall, P. & Granat, M. H. Differentiating sitting and lying using a thigh-worn accelerometer. *Med Sci Sports Exerc* **48**, 742–747 (2016).

44. Steeves, J. A. *et al.* Ability of thigh-worn actigraph and activpal monitors to classify posture and motion. *Med Sci Sports Exerc* **47**, 952–959 (2015).

45. Kalisch, T. *et al.* Measuring sedentary behavior using waist- and thigh-worn accelerometers and inclinometers – are the results comparable? *Ther Adv Musculoskelet Dis* **14**, (2022).

46. Stevens, M. L. *et al.* Thigh-worn accelerometry for measuring movement and posture across the 24-hour cycle: a scoping review and expert statement. *BMJ Open Sport — Exercise Medicine* **6**, 874 (2020).

47. Kwan, R. Y. C., Liu, J. Y. W., Lee, D., Tse, C. Y. A. & Lee, P. H. A validation study of the use of smartphones and wrist-worn ActiGraphs to measure physical activity at different levels of intensity and step rates in older people. *Gait Posture* **82**, 306–312 (2020).

48. Sasaki, J. E. *et al.* Performance of Activity Classification Algorithms in Free-living Older Adults. *Med Sci Sports Exerc* **48**, 941 (2016).

49. Urbanek, J. K. *et al.* Validation of Gait Characteristics Extracted From Raw Accelerometry During Walking Against Measures of Physical Function, Mobility, Fatigability, and Fitness. *J Gerontol A Biol Sci Med Sci* **73**, 676–681 (2018).

50. Clark, B., Winker, E., Ahmadi, M. & Trost, S. Comparison of Three Algorithms Using Thigh-Worn Accelerometers for Classifying Sitting, Standing, and Stepping in Free-Living Office Workers. *J Meas Phys Behav* **4**, 89–95 (2021).

51. Fillekes, M. P., Giannouli, E., Kim, E.-K., Zijlstra, W. & Weibel, R. Towards a comprehensive set of GPS-based indicators reflecting the multidimensional nature of daily mobility for applications in health and aging research. *Int J Health Geogr* **18**, 17 (2019).

52. Brusilovskiy, E. *et al.* Examining the relationship between community mobility and participation using GPS and self-report data. *Soc Sci Med* **265**, 113539 (2020).

53. Beyer, K. B. *et al.* NiMBaLWear analytics pipeline for wearable sensors: a modular, open-source platform for evaluating multiple domains of health and behaviour. *BMC Digital Health* **2**, (2024).

54. Weber, K. S., Godkin, F. E., Cornish, B. F., McIlroy, W. E. & Van Ooteghem, K. Wrist Accelerometer Estimates of Physical Activity Intensity During Walking in Older Adults and People Living With Complex Health Conditions: Retrospective Observational Data Analysis Study. *JMIR Form Res* **7**, e41685 (2023).

55. Vert, A. *et al.* Detecting accelerometer non-wear periods using change in acceleration combined with rate-of-change in temperature. *BMC Med Res Methodol* **22**, 147 (2022).

56. Gourrame, K. *et al.* Parkinson's disease classification with CWNN: Using wavelet transformations and IMU data fusion for improved accuracy. *Technology and Health Care* **31**, 2447–2455 (2023).

57. Straczkiewicz, M., Huang, E. J. & Onnela, J. P. A "one-size-fits-most" walking recognition method for smartphones, smartwatches, and wearable accelerometers. *NPJ Digit Med* **6**, (2023).

58. Osuka, Y., Chan, L. L. Y., Brodie, M. A., Okubo, Y. & Lord, S. R. A Wrist-Worn Wearable Device Can Identify Frailty in Middle-Aged and Older Adults: The UK Biobank Study. *J Am Med Dir Assoc* **25**, (2024).

59.   Del Pozo Cruz, B., Ahmadi, M. N., Lee, I. M. & Stamatakis, E. Prospective Associations of Daily Step Counts and Intensity with Cancer and Cardiovascular Disease Incidence and Mortality and All-Cause Mortality. *JAMA Intern Med* **182**, 1139–1148 (2022).

60.   Fraysse, F. *et al.* Physical Activity Intensity Cut-Points for Wrist-Worn GENEActiv in Older Adults. *Front Sports Act Living* **2**, (2021).

61.   Kuspinar, A. *et al.* Assessing the measurement properties of life-space mobility measures in community-dwelling older adults: a systematic review. *Age Ageing* **52**, IV86–IV99 (2023).

62.   Beauchamp, M. *et al.* Measures of perceived mobility ability in community-dwelling older adults: a systematic review of psychometric properties. *Age Ageing* **52**, IV100–IV111 (2023).

63.   Jette, A. M. *et al.* Late Life Function and Disability Instrument: I. Development and Evaluation of the Disability Component. *The Journals of Gerontology: Series A* **57**, M209–M216 (2002).

64.   Haley, S. M. *et al.* Late life function and disability instrument: II. Development and evaluation of the function component. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* **57**, 209–216 (2002).

65.   Rejeski, W. J. *et al.* The MAT-sf: identifying risk for major mobility disability. *J Gerontol A Biol Sci Med Sci* **70**, 641–646 (2015).

66.   Stalvey, B. T., Owsley, C., Sloane, M. E. & Ball, K. The life space questionnaire: A measure of the extent of mobility of older adults. *Journal of Applied Gerontology* **18**, 460–478 (1999).

67.   Raina, P. *et al.* Cohort Profile: The Canadian Longitudinal Study on Aging (CLSA). *International Epidemiological Association International Journal of Epidemiology* 1752–1753 (2019) doi:10.1093/ije/dyz173.

68.   Marshall, S. C. *et al.* Protocol for Candrive II/Ozcandrive, a multicentre prospective older driver cohort study. *Accid Anal Prev* **61**, 245–252 (2013).

69.   Myers, A. M., Paradis, J. A. & Blanchard, R. A. Conceptualizing and Measuring Confidence in Older Drivers: Development of the Day and Night Driving Comfort Scales. (2008) doi:10.1016/j.apmr.2007.09.037.

70.   MacDonald L, Myers AM & Blanchard RA. Correspondence Among Older Drivers' Perceptions, Abilities, and Behaviors | Article | NursingCenter. *Top Geriatr Rehabil* **24**, 239 (2008).

71.   Podsiadlo, D. & Richardson, S. The Timed "Up & Go": A Test of Basic Functional Mobility for Frail Elderly Persons. *J Am Geriatr Soc* **39**, 142–148 (1991).

72.   Bohannon, R. W. Comfortable and maximum walking speed of adults aged 20-79 years: Reference values and determinants. *Age Ageing* **26**, 15–19 (1997).

73.   Springer, B. A., Marin, R., Cyhan, T., Roberts, H. & Gill, N. W. Normative values for the unipedal stance test with eyes open and closed. *J Geriatr Phys Ther* **30**, 8–15 (2007).

74. Vasunilashorn, S. *et al.* Use of the short physical performance battery score to predict loss of ability to walk 400 meters: Analysis from the InCHIANTI study. *Journals of Gerontology - Series A Biological Sciences and Medical Sciences* **64**, 223–229 (2009).

75. Sayers, S. P. *et al. Concordance and Discordance between Two Measures of Lower Extremity Function: 400 Meter Self-Paced Walk and SPPB. Aging Clin Exp Res* vol. 18 (2006).

76. Ni, M., Brown LG, Lawler, D. & Bean, J. Reliability, Validity, and Minimal Detectable Change of Four-Step Stair Climb Power Test in Community-Dwelling Older Adults. *Phys Ther* **97**, 767–773 (2017).

77. Bean, J. F., Kiely, D. K., LaRose, S., Alian, J. & Frontera, W. R. Is Stair Climb Power a Clinically Relevant Measure of Leg Power Impairments in At-Risk Older Adults? (2007) doi:10.1016/j.apmr.2007.02.004.

78. Rockwood, K. *et al.* A global clinical measure of fitness and frailty in elderly people. *Can Med Assoc J* **173**, 489 (2005).

79. Lord, S. R. & Dayhew, J. Visual Risk Factors for Falls in Older People. *J Am Geriatr Soc* **49**, 508–515 (2001).

80. Rabe, K. F. *et al.* Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med* **176**, 532–555 (2007).

81. Dudzińska-Griszek, J., Szuster, K. & Szewieczek, J. Grip strength as a frailty diagnostic component in geriatric inpatients. *Clin Interv Aging* **12**, 1151–1157 (2017).

82. Washburn, R. A., Smith, K. W., Jette, A. M. & Janney, C. A. The physical activity scale for the elderly (PASE): Development and evaluation. *J Clin Epidemiol* **46**, 153–162 (1993).

83. Keller, H. H., Goy, R. & Kane, S. L. Validity and reliability of SCREEN II (Seniors in the Community: Risk evaluation for eating and nutrition, Version II). *Eur J Clin Nutr* **59**, 1149–1157 (2005).

84. Im, D. D., Jambaulikar, G. D., Kikut, A., Gale, J. & Weiner, S. G. Brief Pain Inventory–Short Form: A New Method for Assessing Pain in the Emergency Department. *Pain Medicine* **21**, 3263–3269 (2020).

85. Xie, F. *et al.* A Time Trade-off-derived Value Set of the EQ-5D-5L for Canada. *Med Care* **54**, 98–105 (2016).

86. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *J Am Geriatr Soc* **53**, 695–699 (2005).

87. Lichtenstein, M. J., Bess, F. H. & Logan, S. A. Diagnostic Performance of the Hearing Handicap Inventory for the Elderly (Screening Version) Against Differing Definitions of Hearing Loss. *Ear Hear* **9**, (1988).

88. Priede, A. *et al.* The factor structure of the Medical Outcomes Study–Social Support Survey: A comparison of different models in a sample of recently diagnosed cancer patients. *J Psychosom Res* **108**, 32–38 (2018).

89.   Russell, D. W. UCLA Loneliness Scale (Version 3): Reliability, Validity, and Factor Structure. *J Pers Assess* **66**, 20–40 (1996).

90.   Lubben, J. *et al.* Performance of an Abbreviated Version of the Lubben Social Network Scale Among Three European Community-Dwelling Older Adult Populations. *Gerontologist* **46**, 503–513 (2006).

91.   Williams, N. The GAD-7 questionnaire. *Occup Med (Chic Ill)* **64**, 224 (2014).

92.   Andresen, E. M., Malmgren, J. A., Carter, W. B. & Patrick, D. L. Screening for depression in well older adults: evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *Am J Prev Med* **10**, 77–84 (1994).

93.   Kamper, S. J., Maher, C. G. & Mackay, G. Global Rating of Change Scales: A Review of Strengths and Weaknesses and Considerations for Design. *Journal of Manual & Manipulative Therapy* **17**, 163–170 (2009).

94.   Montero-Odasso, M. *et al.* World guidelines for falls prevention and management for older adults: a global initiative the Task Force on Global Guidelines for Falls in Older Adults. *Age Ageing* **51**, 1–36 (2022).

95.   Masud, T; Morris, R. Epidemiology of Falls (world report).pdf. **30**, 3–7 (2001).

96.   Mayhew, A. J. *et al.* Normative values for grip strength, gait speed, timed up and go, single leg balance, and chair rise derived from the Canadian longitudinal study on ageing. *Age Ageing* **52**, 1–11 (2023).

97.   Lee, I.-M. *et al.* Association of Step Volume and Intensity With All-Cause Mortality in Older Women. *JAMA Intern Med* **02215**, (2019).

98.   Chan, L. L. Y., Lord, S. R. & Brodie, M. A. Daily-Life Walking Speed, Quality and Quantity Derived from a Wrist Motion Sensor: Large-Scale Normative Data for Middle-Aged and Older Adults. (2024) doi:10.3390/s24165159.