Childhood Maltreatment and Mental-Physical Multimorbidity: A Quasi-Experimental Analysis of Canadian Survey Data Using Entropy Balancing

Andrew J. Fullerton[1], Alpamys Issanov[1], Rafael Meza[1] [2]
1. School of Population and Public Health, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada.
2. Department of Population Health Sciences, BC Cancer Research Institute, Vancouver, BC, Canada.

ABSTRACT
Objectives: To explore childhood maltreatment as a risk factor for mental-physical multimorbidity across disease areas and examine gender as a potential effect modifier.

Methods: We analyzed data from the 2022 Mental Health and Access to Care Survey. We described sample characteristics with unweighted counts, survey-weighted percentages, and weighted chi-square tests. Missing data were addressed via multiple imputation. Entropy balancing adjusted for age, gender, LGBTQ2+ identity, visible minority group, and immigration status and multinomial logistic regression was used to estimate associations between the number of childhood maltreatment subtypes reported and physical (≥ 2 physical conditions but no mental), mental (≥ 2 mental conditions but no physical), and mental-physical (≥ 1 mental and physical condition) multimorbidity. Survey weights were applied during both entropy balancing and regression modeling. Effect modification by gender was examined and sub-analyses of mental-cardiometabolic, mental-inflammatory, and mental-somatic multimorbidity were conducted.

Results: 8,967 respondents were included. Mental-physical multimorbidity increased with maltreatment exposure: 3.4% (none, n=4647), 6.3% (1 type, n=2804), 10.1% (2 types, n=1208), and 18.2% (3 types, n=308). Adjusted odds ratios for mental-physical multimorbidity ranged from 2.15 (95% CI:1.90-2.44) for 1 type to 8.72 (95% CI:7.01-10.85) for 3 types compared to physical (aOR=1.31-2.00) and mental (aOR=1.90-3.63) multimorbidity. Men showed higher odds of mental-physical multimorbidity at high exposure levels (aOR=6.14, 95% CI:4.90-7.70 in women; aOR=13.96, 95% CI:9.58-20.34 in men) with varying effect sizes across disease areas.

Conclusion: Childhood maltreatment shows a strong dose-response association with mental-physical multimorbidity. Further research is needed to clarify biopsychosocial mechanisms and gender-specific pathways.

KEYWORDS: multimorbidity, comorbidity, adverse childhood experiences, childhood trauma, childhood abuse

INTRODUCTION

Multimorbidity has been shown to worsen the prognosis of all present diseases while making treatment more complicated, more costly, and less effective. These challenges are even greater for individuals with co-occurring mental and physical health conditions due to the mental-physical dichotomy of many diagnostic frameworks and limited clinical overlap between non-psychiatric and psychiatric specialists (Halstead et al., 2024). With an estimated prevalence of 8.4% in Canada as of 2014, individuals with mental-physical multimorbidity – the co-occurrence of at least one mental and one physical chronic health condition – report higher rates of suicidal ideation, more adverse health outcomes, greater healthcare utilization and more unmet healthcare needs across the life course compared to those with only physical or mental multimorbidity (Dai et al., 2020). Given its substantial burden, identifying modifiable early life risk factors for mental-physical multimorbidity is an important public health priority. While childhood maltreatment is an established early life risk factor for various mental and physical health conditions, its role in mental-physical multimorbidity remains underexplored in Canada.

Childhood maltreatment is thought to affect a variety of health outcomes through increased risk of health-harming behaviours and chronic dysregulation of neuroregulatory pathways such as the hypothalamic-pituitary-adrenal (HPA) axis (Hughes et al., 2017; Metzler et al., 2016). Recent studies using data from the Canadian Community Health Survey (CCHS) have validated the association between childhood maltreatment and increased risk of diabetes (Shields et al., 2016a), chronic obstructive pulmonary disease (Shields et al., 2016b), arthritis (Badley et al., 2019), cancer (Hovdestad et al., 2020), and mental health conditions including anxiety, depression, and substance use disorder (Afifi et al., 2014). Other research in Canada has identified childhood maltreatment as a key risk factor for psychiatric comorbidity in individuals with multiple sclerosis, arthritis, and inflammatory bowel disease (Wan et al., 2022; O'Mahony et al., 2024). To our knowledge, however, only one study (England-Mason et al., 2018) has assessed childhood maltreatment as a risk factor for multimorbidity across health domains in a representative sample of the Canadian population. Consistent with other evidence, their findings suggest a dose-response relationship between the number of maltreatment types experienced and multimorbidity in adulthood.

While England-Mason et al. (2018) provided initial evidence of the association, the role of childhood maltreatment in mental-physical multimorbidity has not been explored in-depth. Despite recent evidence indicating climbing multimorbidity rates among Canadian men (Kone et al., 2021; Ferris et al., 2025), no work has been published examining potential gender differences. Our study addresses this gap by examining the association between childhood maltreatment and mental-physical multimorbidity using weighted, nationally representative data from the 2022 Mental Health and Access to Care Survey (MHACS). This work aims to provide a foundation for future research into the early life mechanisms underlying mental-physical multimorbidity and inform more targeted prevention efforts. Building on England-Mason et al. (2018), we aim to explore childhood maltreatment as a risk factor for mental-physical multimorbidity across disease areas and examine gender as a potential effect modifier.

METHODS
Data Source

MHACS is a multi-stage stratified survey designed to assess the mental health, service use, and functioning of Canadians including individuals aged 15 and older as of March 1, 2022, residing in the ten Canadian provinces (Statistics Canada, 2023). The sampling frame is representative of the Canadian population excluding full-time Canadian Forces members, individuals living on reserves or Indigenous settlements, and those residing in collective dwellings. Out of 39,485 units selected from the 2021 census, 9,861 individuals responded to the survey, yielding an overall response rate of around 25%. Data were collected between March 2022 and July 2022 using computer-assisted telephone interviews. Further details are available from Statistics Canada. All data used in this analysis were obtained through the Abacus Data Network (Abacus Data Network, n.d.).

Analytic Sample
The sample included respondents who were administered the childhood experiences section of MHACS. Invalid responses and non-responses were imputed using multiple imputation, resulting in an analytic sample of 8,967 individuals aged 20 and older.

Study Variables
Childhood Maltreatment
For consistency with prior research and the Canadian Child Maltreatment Surveillance Indicator framework (Campeau et al., 2020), we derived an exposure variable measuring the number of childhood maltreatment subtypes reported ranging from none to three (physical abuse, sexual abuse, and exposure to domestic violence). Respondents reported childhood maltreatment experiences before age 16 using items from the Childhood Experiences of Violence Questionnaire (Walsh et al., 2008) with additional items for sexual abuse (Statistics Canada, 2023). Physical abuse was classified as present if respondents reported at least one occurrence of being physically attacked. Sexual abuse was classified as present if respondents reported at least one instance of forced unwanted sexual activity or touching. Exposure to domestic violence was classified as present if respondents reported witnessing a parent or guardian physically harm another adult in the home.

Mental-Physical Multimorbidity Status
We derived the outcome variable, mental-physical multimorbidity status, based on self-reported reported chronic conditions across two domains: mental and physical health. To enable comparisons between mental-physical and single-domain multimorbidity, the variable was constructed as a multilevel nominal variable with four states: no multimorbidity (reference group), physical multimorbidity (≥ 2 physical conditions but no mental), mental multimorbidity (≥ 2 mental conditions but no physical), and mental-physical multimorbidity (≥ 1 mental and physical condition).

Mental health conditions were assessed using the WHO 12-Month Composite International Diagnostic Interview criteria (Statistics Canada, 2023) and included major depression, generalized anxiety disorder, social phobia, bipolar I/II, hypomania, and substance use disorders (alcohol, cannabis, and other drugs). Physical health conditions were self-reported and included hypertension, diabetes, heart disease, asthma, arthritis, inflammatory bowel disease (IBD), chronic obstructive pulmonary disease (COPD), chronic fatigue syndrome, fibromyalgia, back pain, cancer, and chronic migraines.

Covariates

We used a directed acyclic graph (Fig 1) to represent hypothesized relationships between variables and identify a minimally sufficient adjustment set: age, gender, LGBTQ2+ identity, visible minority group, and immigration status. Health-related behaviours and adulthood socioeconomic factors were identified as mediators and omitted from the analytic model.

Fig 1. Directed acyclic graph presenting relationships between exposure, outcome, and measured covariates based on a priori understanding. Blue nodes indicate mediator variables while green edges indicate causal paths. White nodes indicate adjusted variables and black edges indicate blocked paths.

Statistical Analysis

We described respondent characteristics and multimorbidity prevalence using unweighted counts and survey-weighted percentages stratified by the number of childhood maltreatment subtypes reported. We assessed associations between characteristics and exposure groups using survey-weighted chi-square tests.

To adjust for confounding, we used entropy balancing to create a reweighted pseudopopulation balanced on age, gender, LGBTQ2+ identity, visible minority group, and immigration status. Entropy balancing is a reweighting method that achieves covariate balance by directly matching covariate distributions across treatment groups, ensuring balance on specified moments (e.g., mean, variance, and skewness) while minimizing divergence from uniform base weights via an optimization function (Hainmueller et al., 2012). Since all included covariates were categorical, balance was achieved on proportions with a constraint tolerance of 1e-10 and 1000 maximum iterations.

We estimated associations between childhood maltreatment and mental-physical multimorbidity status using multinomial logistic regression. To examine effect modification, we added a gender-by-maltreatment interaction after confirming covariate balance across gender (standardized mean difference < 0.1). Sampling weights were applied during both entropy balancing and regression modeling to maintain population representativeness. Variance was estimated using a bootstrap estimator with replicate weights provided by Statistics Canada to account for survey design. All analyses were conducted in R 4.4.1.

Missing Data

After determining that data were not missing completely at random, data were assumed to be missing at random in approximately 2.41% of the unweighted sample. We addressed missing values using multiple imputation by chained equations with 10 imputations and 5 iterations. The imputation model included all non-derived study variables and auxiliary variables (self-rated mental health, self-rated physical health, family member mental health, household income, occupational status, marital status, and educational attainment). Following imputation, derived variables were created and entropy balancing and regression modeling were performed on each imputed dataset. Final estimates were pooled using Rubin's rules (Rubin, 1987).

Cluster-Specific Analyses

To examine variations across previously identified clusters of co-occurring mental and physical disease, we conducted sub-analyses of mental-cardiometabolic (co-occurring mental illness and diabetes, heart disease, or hypertension), mental-inflammatory (co-occurring mental illness and asthma, arthritis, COPD, or IBD), and mental-somatic multimorbidity (co-occurring mental illness and chronic fatigue syndrome, back pain, chronic migraines, or fibromyalgia). Clusters were constructed based on shared risk factors, pathophysiology, and previously identified links to childhood maltreatment (Baltramonaityte et al., 2023; Souama et al., 2023; Wan et al., 2022; O'Mahony et al., 2024; England-Mason et al., 2018).

Ethics Approval

This secondary data analysis did not require institutional ethics review (Canadian Institutes of Health Research, 2022).

RESULTS

Descriptive Statistics

Out of 9,861 MHACS respondents, 8,967 individuals (90.9%) were included in the analysis. Age, visible minority group, and immigration status distributions were consistent across all levels of exposure to childhood maltreatment (Table 1). Notably, the proportion of women increased with exposure to childhood maltreatment (1 type: 44.4%, 2 types: 50.0%, 3 types: 73.6%) as did the proportion identifying as LGBTQ2+ (1 type: 5.7%, 2 types: 7.5%, 3 types: 17.3%).

Table 1. *Characteristics of Canadian adults (> 20 years of age) stratified by the number of types of childhood maltreatment experiences reported before 16 years of age based on the 2022 Mental Health and Access to Care Survey.*

Variable	None[1] (n = 4647)	1 maltreatment type[1] (n = 2804)	2 maltreatment types[1] (n = 1208)	3 maltreatment types[1] (n = 308)	p
Age (%)					<0.001[2]
20 to 24 years	643 (9.8)	307 (7.1)	169 (7.8)	34 (7.0)	
25 to 29 years	291 (7.7)	175 (6.0)	69 (5.5)	21 (7.3)	
30 to 34 years	359 (9.0)	181 (7.4)	84 (8.0)	23 (7.9)	
35 to 44 years	678 (16.8)	465 (19.1)	190 (17.6)	46 (15.5)	

45 to 54 years	600 (13.6)	366 (14.3)	176 (15.3)	61 (18.9)	
55 to 64 years	647 (17.0)	437 (19.6)	196 (21.3)	54 (23.5)	
65 years or older	1,429 (26.0)	873 (26.4)	324 (24.5)	69 (19.9)	
Gender (%)³ ⁴					<0.001²
Women+	2,439 (54.3)	1,193 (44.4)	575 (50.0)	219 (73.6)	
Men+	2,200 (45.7)	1,607 (55.6)	631 (50.0)	89 (26.4)	
LGBTQ2+					<0.001²
Cisgender/Heterosexual	4,242 (95.8)	2,624 (94.3)	1,112 (92.5)	257 (82.7)	
LGBTQ2+	185 (4.2)	152 (5.7)	84 (7.5)	47 (17.3)	
Visible minority group (%)³					<0.001²
Not a visible minority	2,382 (72.6)	1,576 (76.5)	607 (70.8)	166 (72.4)	
Black	377 (3.1)	305 (3.7)	163 (5.3)	34 (3.9)	
Chinese	446 (5.6)	281 (5.3)	107 (4.1)	28 (5.2)	
Filipino	381 (2.8)	177 (2.1)	87 (2.9)	19 (2.5)	
South Asian	500 (8.5)	225 (5.7)	136 (8.9)	27 (6.5)	
Other visible minority	308 (7.4)	183 (6.7)	91 (8.1)	28 (9.5)	
Immigration status (%)³					0.2²
Immigrant	2,023 (32.1)	1,187 (29.6)	546 (32.4)	124 (31.4)	
Canadian-born	2,419 (67.9)	1,600 (70.4)	658 (67.6)	183 (68.6)	

Unweighted counts and survey-weighted percentages are reported; column totals reflect unweighted population including missingness.
[1] Maltreatment types included are physical abuse, sexual abuse, and exposure to domestic violence.
[2] Weighted chi-square test used to calculate p-values.
[3] Missing: Gender (0.15%); Visible minority group (3.75%); Immigration status (2.36%); LGBTQ2+ (2.99%).
[4] Mental Health and Access to Care Survey (MHACS) used random imputation to distribute non-binary persons into either Men+ or Women+.

Table 2 reports the estimated prevalence of mental-physical multimorbidity (5.7%), mental multimorbidity (10.7%), and physical multimorbidity (24.8%) in the Canadian population. The prevalence of mental-physical multimorbidity increased from 3.4% among those who reported no maltreatment (n=4647) to 18.2% among those who reported 3 maltreatment types (n=308), showing a clear gradient. Physical multimorbidity was elevated in those who reported any maltreatment (1 type: 26.8%; 2 types: 27.5%; 3 types: 25.8%) compared to none (22.7%). Mental multimorbidity prevalence increased steadily with exposure to childhood maltreatment from 8.2% (no maltreatment) to 18.6% (3 types).

Table 2. *Prevalence of multimorbidity by number of childhood maltreatment types experienced in Canadian individuals 20 years of age and older based on the 2022 Mental Health and Access to Care Survey.*

Childhood maltreatment[2]	None n (%)[1]	Physical n (%)[1]	Mental n (%)[1]	Mental-physical n (%)[1]	p
None (n = 4647)	3083 (65.6)	1075 (22.7)	357 (8.2)	132 (3.4)	<0.001[3]
1 maltreatment type (n = 2804)	1562 (54.7)	745 (26.8)	343 (12.3)	154 (6.3)	
2 maltreatment types (n = 1208)	610 (48.5)	314 (27.5)	176 (13.9)	108 (10.1)	
3 maltreatment types (n = 308)	127 (37.4)	77 (25.8)	56 (18.6)	48 (18.2)	
Overall	5382 (58.8)	2211 (24.8)	932 (10.7)	442 (5.7)	

Unweighted counts and survey-weighted percentages are reported; row totals reflect unweighted population.
[1] Counts and percentages are pooled across imputations.
[2] Maltreatment types included are physical abuse, sexual abuse, and exposure to domestic violence.
[3] Weighted chi-square test used to calculate p-values.

Childhood Maltreatment and Mental-Physical Multimorbidity

As shown in Table 3, we found higher odds of all multimorbidity states with each additional childhood maltreatment type reported compared to those reporting no maltreatment in both unadjusted and adjusted analyses. The unadjusted odds ratios (OR) ranged from OR=2.19 (95% CI:1.93-2.48) with 1 reported maltreatment type to OR=9.25 (95% CI:7.6-11.26) with 3 reported maltreatment types for mental-physical multimorbidity. Estimates ranged from OR=1.41 (95% CI:1.35-1.48) to OR=1.99 (95% CI:1.75-2.26) for physical multimorbidity and from OR=1.79 (95% CI:1.58-2.02) to OR=3.94 (95% CI:3.22-4.83) for mental multimorbidity.

Table 3. *Odds of physical multimorbidity, mental multimorbidity, and mental-physical multimorbidity among Canadian individuals over 20 years of age who reported exposure to 1 or more types of childhood maltreatment compared to those who did not report exposure to any types of childhood maltreatment based on the 2022 Mental Health and Access to Care Survey[1].*

Childhood maltreatment[3]	Unadjusted								Adjusted[2]					
	Physical		Mental		Mental-physical				Physical		Mental		Mental-physical	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p			OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
1 type[4]	1.41 (1.35, 1.48)	<0.001	1.79 (1.58, 2.02)	<0.001	2.19 (1.93, 2.48)	<0.001			1.31 (1.26, 1.37)	<0.001	1.9 (1.68, 2.15)	<0.001	2.15 (1.9, 2.44)	<0.001
2 types[4]	1.64 (1.53, 1.76)	<0.001	2.28 (2.02, 2.58)	<0.001	3.93 (3.34, 4.62)	<0.001			1.59 (1.48, 1.7)	<0.001	2.48 (2.18, 2.81)	<0.001	3.88 (3.33, 4.52)	<0.001
3 types[4]	1.99 (1.75, 2.26)	<0.001	3.94 (3.22, 4.83)	<0.001	9.25 (7.6, 11.26)	<0.001			2.00 (1.74, 2.3)	<0.001	3.63 (2.87, 4.59)	<0.001	8.72 (7.01, 10.85)	<0.001

OR, odds ratio; CI, confidence interval; p, p-value.

[1] Two separate survey-weighted multinomial logistic regressions with bootstrap variance estimators are summarized. The adjusted model was first balanced on age, gender, LGBTQ2+ status, visible minority group and immigration status across treatment groups via entropy balancing.

[2] Survey weights were used in the estimation of balancing weights and in the outcome regression to maintain population representativeness and account for non-response.

[3] Childhood maltreatment types included: physical abuse, sexual abuse, and exposure to domestic violence.

[4] Reference group: no childhood maltreatment reported.

Associations remained robust after balancing the sample on age, gender, LGBTQ2+ identity, visible minority group and immigration status. Mental-physical multimorbidity exhibited the strongest graded association with adjusted odds ratios (aOR) ranging from aOR=2.15 (95% CI: 1.9-2.44) with 1 maltreatment type to aOR=8.72 (95% CI: 7.01-10.85) with 3 maltreatment types. Odds ratios ranged from aOR=1.31 (95% CI: 1.26-1.37) to aOR=2.00 (95% CI: 1.74-2.3) for physical multimorbidity and from aOR=1.9 (95% CI: 1.68-2.15) to aOR=3.63 (95% CI: 2.87, 4.59) for mental multimorbidity. All associations were statistically significant (p<0.001) with no confidence interval overlap between exposure levels, demonstrating a progressive effect increase across all multimorbidity states with increasing maltreatment exposure (Fig 2).

Fig 2. *Forest plot of the adjusted relationship (odds ratios) between number of types (physical abuse, sexual abuse, exposure to domestic violence) of childhood maltreatment reported before 16 years of age and multimorbidity among Canadian adults (> 20 years of age) by multimorbidity state based on the 2022 Mental Health and Access to Care Survey.*

Effect Modification by Gender

Table 4 presents the gender-specific effect estimates of childhood maltreatment on multimorbidity from a single model with a gender-by-maltreatment interaction. While interactions were not significant at the lowest level of exposure to childhood maltreatment, they were significant for both mental-physical multimorbidity and physical multimorbidity at higher exposure levels. For mental-physical multimorbidity, the effect in women ranged from aOR=2.13 (95% CI:1.81-2.51) with 1 maltreatment type to aOR=6.14 (95% CI:4.90-7.70) with 3 types. In men, associations were comparable at low exposure (aOR=2.21, 95% CI:1.76-2.78) but stronger at high exposure (aOR=13.96, 95% CI:9.58-20.34), indicating magnified gender differences with increasing maltreatment exposure (Fig 3).

Table 4. *Gender-specific odds of physical multimorbidity, mental multimorbidity, and mental-physical multimorbidity among Canadian individuals over 20 years of age who reported exposure to 1 or more types of childhood maltreatment compared to those who did not report exposure to any types of childhood maltreatment based on the 2022 Mental Health and Access to Care Survey[1].*

Childhood maltreatment[3]	Men							Women					
	Physical		Mental		Mental-physical			Physical		Mental		Mental-physical	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
1 type[4]	1.37 (1.28, 1.46)	<0.001	1.96 (1.7, 2.25)	<0.001	2.21 (1.76, 2.78)	<0.001		1.27 (1.19, 1.36)	<0.001	1.87 (1.58, 2.2)	<0.001	2.13 (1.81, 2.51)	<0.001
2 types[4]	1.74 (1.58, 1.92)**	<0.001	2.27 (1.96, 2.62)	<0.001	4.7 (3.57, 6.18)	<0.001		1.45 (1.32, 1.6)**	<0.001	2.64 (2.2, 3.18)	<0.001	3.49 (2.84, 4.29)	<0.001
3 types[4]	2.45 (1.96, 3.05)**	<0.001	3.94 (2.65, 5.85)	<0.001	13.96 (9.58, 20.34)***	<0.001		1.64 (1.38, 1.94)**	<0.001	3.36 (2.68, 4.21)	<0.001	6.14 (4.9, 7.7)***	<0.001

OR, odds ratio; CI, confidence interval; p, p-value; *, significant interaction at 0.05 level; **, significant interaction at 0.01 level; ***, significant interaction at 0.001 level.

[1] Results from a single survey-weighted multinomial logistic regression with a bootstrap variance estimator are summarized. The model was first balanced on age, gender, LGBTQ2+ status, visible minority group and immigration status across treatment groups via entropy balancing; the outcome regression included an interaction between gender and childhood maltreatment.

[2] Survey weights were used in the estimation of balancing weights and in the outcome regression to maintain population representativeness and account for non-response.

[3] Childhood maltreatment types included: physical abuse, sexual abuse, and exposure to domestic violence.

[4] Reference group: no childhood maltreatment reported.

Fig 3. *Forest plot of the adjusted relationship (odds ratios) between number of types (physical abuse, sexual abuse, exposure to domestic violence) of childhood maltreatment reported before 16 years of age and multimorbidity among Canadian adults (> 20 years of age) by multimorbidity state and gender based on the 2022 Mental Health and Access to Care Survey.*

Associations were more modest for physical multimorbidity. Effect estimates in women ranged from aOR=1.27 (95% CI:1.19-1.36) with 1 reported maltreatment type to aOR=1.64 (95% CI:1.38-1.94) with 3 types. In men, effect estimates ranged from aOR=1.37 (95% CI:1.28-1.46) with 1 maltreatment type to aOR=2.45 (95% CI:1.96-3.05) with 3. Effect modification was weaker and non-significant for mental multimorbidity across all exposure levels ranging from aOR=1.87 (95% CI:1.58-2.2) to aOR=3.36 (95% CI:2.68-4.21) in women and from aOR=1.96 (95% CI:1.7-2.25) to aOR=3.94 (95% CI:2.65-5.85) in men.

Cluster-Specific Analyses
In cluster-specific models, findings were consistent with the main analysis (Table 5 and Fig 4). However, patterns varied significantly when effect modification was considered (Table 6). In the mental-cardiometabolic cluster, odds of mental-physical multimorbidity were much higher in men (aOR=7.28, 95% CI:2.81-18.87) than women (aOR=0.62, 95% CI:0.29-1.33) but converged at higher levels of exposure (Fig 5). In the mental-inflammatory cluster, effect modification was not significant except in physical multimorbidity where women exhibited higher odds of physical multimorbidity with 3 maltreatment types (aOR=2.44, 95% CI:1.87-3.18 in women; aOR=1.39, 95% CI:0.91-2.13 in men). In the mental-somatic cluster, stronger associations were observed in women with 1 maltreatment type (aOR =2.82, 95% CI:2.14-3.71 in women; aOR=0.99, 95% CI:0.65-1.5 in men). However, associations were stronger in men with 3 maltreatment types (aOR=6.45, 95% CI:4.57-9.1 in women; aOR=14.61, 95% CI:9.92-21.52 in men).

Table 5. *Odds of cluster-specific physical multimorbidity, mental multimorbidity, and mental-physical multimorbidity among Canadian individuals over 20 years of age who reported exposure to 1 or more types of childhood maltreatment compared to those who did not report exposure to any types of childhood maltreatment based on the 2022 Mental Health and Access to Care Survey[1].*

	Unadjusted							Adjusted[2]					
Mental-cardiometabolic multimorbidity[2]													
Childhood maltreatment[3]	**Physical**		**Mental**		**Mental-physical**			**Physical**		**Mental**		**Mental-physical**	
	OR (95% CI)	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**		**OR (95% CI)**	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**
1 type[4]	1.05 (0.98, 1.13)	0.17	1.71 (1.56, 1.88)	<0.001	2.2 (1.48, 3.26)	<0.001		0.96 (0.89, 1.03)	0.27	1.82 (1.66, 2.00)	<0.001	1.81 (1.23, 2.67)	0.004
2 types[4]	1.43 (1.29, 1.6)	<0.001	2.4 (2.12, 2.72)	<0.001	3.85 (2.66, 5.58)	<0.001		1.32 (1.19, 1.46)	<0.001	2.53 (2.23, 2.88)	<0.001	3.66 (2.54, 5.27)	<0.001
3 types[4]	1.26 (1.01, 1.56)	0.004	4.21 (3.63, 4.89)	<0.001	11.63 (7.83, 17.29)	<0.001		1.5 (1.2, 1.87)	0.002	4.05 (3.4, 4.84)	<0.001	9.25 (6.08, 14.06)	<0.001
Mental-inflammatory multimorbidity[2]													
Childhood maltreatment[3]	**Physical**		**Mental**		**Mental-physical**			**Physical**		**Mental**		**Mental-physical**	
	OR (95% CI)	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**		**OR (95% CI)**	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**
1 type[4]	1.39 (1.24, 1.56)	<0.001	1.75 (1.59, 1.93)	<0.001	1.76 (1.32, 2.34)	<0.001		1.34 (1.2, 1.5)	<0.001	1.87 (1.69, 2.07)	<0.001	1.72 (1.31, 2.24)	<0.001
2 types[4]	1.67 (1.45, 1.93)	<0.001	2.41 (2.14, 2.71)	<0.001	2.93 (2.1, 4.09)	<0.001		1.77 (1.53, 2.05)	<0.001	2.58 (2.28, 2.92)	<0.001	2.88 (2.1, 3.95)	<0.001
3 types[4]	2.29 (1.87, 2.79)	<0.001	3.85 (3.29, 4.5)	<0.001	13.86 (10.61, 18.09)	<0.001		2.04 (1.65, 2.52)	<0.001	3.63 (3, 4.4)	<0.001	11.49 (8.49, 15.55)	<0.001
Mental-somatic multimorbidity[2]													
Childhood maltreatment[3]	**Physical**		**Mental**		**Mental-physical**			**Physical**		**Mental**		**Mental-physical**	
	OR (95% CI)	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**		**OR (95% CI)**	**p**	**OR (95% CI)**	**p**	**OR (95% CI)**	**p**
1 type[4]	1.33 (1.19, 1.49)	<0.001	1.74 (1.58, 1.91)	<0.001	1.8 (1.47, 2.2)	<0.001		1.32 (1.19, 1.47)	<0.001	1.82 (1.65, 2.0)	<0.001	2.1 (1.71, 2.57)	<0.001
2 types[4]	1.55 (1.31, 1.85)	<0.001	2.3 (2.03, 2.61)	<0.001	3.42 (2.7, 4.32)	<0.001		1.56 (1.32, 1.83)	<0.001	2.43 (2.14, 2.77)	<0.001	3.64 (2.92, 4.54)	<0.001

3 types[4]	2.95 (2.32, 3.74)	<0.0001	4.17 (3.49, 4.98)	<0.0001	8.95 (7.02, 11.42)	<0.0001		2.76 (2.17, 3.51)	0.002	3.74 (3.03, 4.62)	<0.0001	9.48 (7.39, 12.16)	<0.0001

OR, odds ratio; CI, confidence interval; p, p-value.

[1] Separate survey-weighted multinomial logistic regressions using a bootstrap variance estimator are summarized. The adjusted models were first balanced on age, gender, LGBTQ2+ status, visible minority group and immigration status across treatment groups via entropy balancing. Survey weights were used in the estimation of balancing weights and in the outcome regression to maintain population representativeness and account for non-response.

[2] Diseases included in each cluster are mental-inflammatory: asthma, chronic obstructive pulmonary disease, arthritis, inflammatory bowel disease; mental-somatic: chronic fatigue syndrome, back pain, chronic migraines, fibromyalgia; mental-cardiometabolic: diabetes, heart disease, and hypertension. All mental health conditions were included in each cluster.

[3] Childhood maltreatment types included: physical abuse, sexual abuse, and exposure to domestic violence.

[4] Reference group: no childhood maltreatment reported.

Fig 4. *Forest plot of the adjusted relationship (odds ratios) between number of types (physical abuse, sexual abuse, exposure to domestic violence) of childhood maltreatment reported before 16 years of age and multimorbidity among Canadian adults (> 20 years of age) by multimorbidity state and mental-physical disease cluster based on the 2022 Mental Health and Access to Care Survey.*

Table 6. *Gender-specific odds of cluster-specific physical multimorbidity, mental multimorbidity, and mental-physical multimorbidity among Canadian individuals over 20 years of age who reported exposure to 1 or more types of childhood maltreatment compared to those who did not*

report exposure to any types of childhood maltreatment based on the 2022 Mental Health and Access to Care Survey[1].

Childhood maltreatment[3]	Men Physical		Men Mental		Men Mental-physical			Women Physical		Women Mental		Women Mental-physical	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p		OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Mental-cardiometabolic multimorbidity[2]													
1 type[4]	0.96 (0.87, 1.06)	0.43	1.76 (1.57, 1.98)	<0.001	7.28 (2.81, 18.87)**	<0.001		0.95 (0.84, 1.08)	0.43	1.87 (1.65, 2.13)	<0.001	0.62 (0.29, 1.33)**	0.2
2 types[4]	1.5 (1.32, 1.72)**	<0.001	2.38 (2.03, 2.78)	<0.001	12.98 (4.47, 37.71)**	<0.001		1.06 (0.86, 1.29)**	0.6	2.65 (2.21, 3.19)	<0.001	1.64 (0.82, 3.31)**	0.15
3 types[4]	1.4 (0.95, 2.05)	0.09	4.73 (3.37, 6.64)	<0.001	18.36 (6.47, 52.06)	<0.001		1.61 (1.16, 2.24)	0.005	3.59 (2.99, 4.32)	<0.001	7.41 (4.38, 12.53)	<0.001
Mental-inflammatory multimorbidity[2]													
1 type[4]	1.32 (1.1, 1.59)	0.003	1.88 (1.65, 2.14)	<0.001	1.81 (1.14, 2.88)	0.01		1.36 (1.21, 1.53)	<0.001	1.87 (1.65, 2.13)	<0.001	1.69 (1.24, 2.3)	0.003
2 types[4]	1.51 (1.18, 1.93)	0.001	2.45 (2.12, 2.84)	<0.001	2.79 (1.69, 4.63)	<0.001		1.98 (1.65, 2.37)	<0.001	2.72 (2.26, 3.27)	<0.001	3.00 (2.13, 4.22)	0.001
3 types[4]	1.39 (0.91, 2.13)*	0.12	4.22 (2.93, 6.08)	<0.001	13.29 (7.9, 22.35)	<0.001		2.44 (1.87, 3.18)*	<0.001	3.25 (2.72, 3.88)	<0.001	10.78 (7.87, 14.77)	<0.001
Mental-somatic multimorbidity[2]													
1 type[4]	1.08 (0.86, 1.36)*	0.49	1.95 (1.74, 2.19)	<0.001	0.99 (0.65, 1.5)***	0.95		1.47 (1.28, 1.68)*	<0.001	1.74 (1.52, 1.98)	<0.001	2.82 (2.14, 3.71)***	<0.001
2 types[4]	1.74 (1.32, 2.29)	0.008	2.44 (2.08, 2.86)	<0.001	2.86 (2.01, 4.07)	<0.001		1.48 (1.22, 1.81)	<0.001	2.46 (2.06, 2.95)	<0.001	4.2 (3.2, 5.53)	<0.001
3 types[4]	4.28 (2.82, 6.5)*	0.04	4.12 (2.79, 6.09)	<0.001	14.61 (9.92, 21.52)**	<0.001		1.98 (1.43, 2.74)*	<0.001	3.47 (2.83, 4.26)	<0.001	6.45 (4.57, 9.1)**	<0.001

OR, odds ratio; CI, confidence interval; p, p-value; *, significant interaction at 0.05 level; **, significant interaction at 0.01 level; ***, significant interaction at 0.001 level.

[1] Results from single survey-weighted multinomial logistic regressions with a bootstrap variance estimator are summarized. The models were first balanced on age, gender, LGBTQ2+ status, visible minority group and

immigration status across treatment groups via entropy balancing; an interaction between gender and childhood maltreatment was included in the outcome regression. Survey weights were used in the estimation of balancing weights and in the outcome regression to maintain population representativeness and account for non-response.

[2] Diseases included in each cluster are mental-inflammatory: asthma, chronic obstructive pulmonary disease, arthritis, inflammatory bowel disease; mental-somatic: chronic fatigue syndrome, back pain, chronic migraines, fibromyalgia; mental-cardiometabolic: diabetes, heart disease, and hypertension. All mental health conditions were included in each cluster.

[3] Childhood maltreatment types included: physical abuse, sexual abuse, and exposure to domestic violence.

[4] Reference group: no childhood maltreatment reported.

Fig 5. Forest plot of the adjusted relationship (odds ratios) between number of types (physical abuse, sexual abuse, exposure to domestic violence) of childhood maltreatment reported before 16 years of age and multimorbidity among Canadian adults (> 20 years of age) by multimorbidity state, mental-physical disease cluster, and gender based on the 2022 Mental Health and Access to Care Survey (MHACS).

DISCUSSION

Our findings demonstrate a robust dose-response association between childhood maltreatment and adulthood multimorbidity with particularly strong effects for mental-physical multimorbidity. Although effect sizes varied, this pattern persisted in gender-specific analyses and held across mental-cardiometabolic, mental-inflammatory, and mental-somatic clusters.

These findings are consistent with prior Canadian research. Using 2012 CCHS data and an identically structured exposure variable, England-Mason et al. (2018) reported a similar dose-response pattern. For co-occurring mental and physical conditions (excluding chronic pain/fatigue), their adjusted odds ratios ranged from 1.98 (95% CI:1.38-2.85) with 1 maltreatment type to 6.51 (95% CI:3.09-13.70) with all types. For co-occurring mental and chronic pain/fatigue conditions, they ranged from 1.44 (95% CI: 1.21-1.73) to 3.38 (95% CI:1.94-5.87), and for the co-occurrence of all three conditions they ranged from 2.63 (95% CI: 1.93-3.58) to 10.96 (95% CI:6.12-19.64). Comparatively, our adjusted estimates range from 2.15 (95% CI:1.90-2.44) with 1 type to 8.72 (95% CI:7.01-10.85) with all types for mental-physical multimorbidity, and from 2.1 (95% CI:1.71-2.57) to 9.48 (95% CI:7.39-12.16) for mental-somatic multimorbidity. Although not directly comparable due to England-Mason et al.'s distinct outcome specification and additional behavioural and socioeconomic adjustments, the overall dose-response pattern is consistent. More broadly, our results align with related research in Canada and the UK demonstrating strong associations between childhood maltreatment and both mental-cardiometabolic (Souama et al., 2023) and mental-inflammatory multimorbidity (Wan et al., 2022; O'Mahony et al., 2024) as well as existing biopsychosocial evidence linking childhood maltreatment to adulthood morbidity (Hughes et al., 2017; Metzler et al., 2016; Baltramonaityte et al., 2023).

We also identified novel gender-specific patterns with men showing particularly strong associations at higher maltreatment exposure levels (127.4% stronger in men reporting all maltreatment types). Importantly, these gender differences varied across disease clusters: effect modification was strongest in the mental-cardiometabolic cluster (147.8% stronger in men reporting all maltreatment types), weaker in mental-somatic (126.5% stronger in men reporting all maltreatment types), and minimal in the mental-inflammatory cluster (23.2% stronger in men reporting all maltreatment types). Although there is a paucity of comparable research in Canada, our findings contrast with a recent meta-analysis in the UK which did not find any evidence of sex-based differences in the association between childhood maltreatment and co-occurring depression and cardiometabolic disease (Souama et al., 2023). While the biological mechanisms involved in mental-physical multimorbidity may differ from those driving the development of individual conditions, it is noteworthy that these findings also contrast with prior Canadian research which has reported stronger associations between maltreatment and individual mental (Gallo et al., 2018) and physical (Wan et al., 2022) morbidities in women. Our results are, however, generally consistent with recent evidence from Ontario and British Columbia indicating that the incidence and prevalence of multimorbidity are increasing at greater rates among Canadian men (Kone et al., 2021; Ferris et al., 2025).

While gender-specific pathways linking childhood maltreatment to mental-physical multimorbidity remain underexplored in Canada, several potential mechanisms may explain the gender differences observed in our analysis. Recent Canadian estimates based on MHACS indicate higher prevalence of mood and anxiety disorders among women and higher prevalence of substance use disorders among men (Stephenson, 2023). Meanwhile, established evidence in the US has linked increased incidence of internalizing disorders (e.g. anxiety and mood disorders) in women to inflammatory and chronic pain conditions, and increased incidence of externalizing disorders (e.g. substance use disorders) in men to cardiometabolic diseases (Needham & Hill, 2010). Alongside evidence indicating that men are more likely to exhibit

impaired cardiometabolic functioning while women are more likely to exhibit neuroendocrine and immune dysregulation (Longpré-Poirer et al., 2022), these associations provide some plausible explanations for our findings. Other research suggests that men may be more likely to engage in health-harming behaviours associated with multimorbidity (Punjani et al., 2018) and less likely to access healthcare services compared to women (Thompson et al., 2016). Childhood maltreatment may exacerbate these existing biological and behavioural vulnerabilities in men, resulting in an accumulation of risk factors for multimorbidity across the life course. These mechanisms alone, however, are likely insufficient in explaining the increased effect sizes seen among men in our analysis. Further research is needed to understand the gender-specific effects of childhood maltreatment on mental-physical multimorbidity.

Strengths and Limitations
To our knowledge, this study is the first in Canada to report gender-specific associations between childhood maltreatment and mental-physical multimorbidity using a nationally representative sample. However, several limitations must be considered. First, relevant early life covariates such as childhood household conditions were not measured and no proxy measures could be identified. Additionally, the broad analytic focus of our study prohibited disease-specific adjustments. Although post-hoc E-value estimation indicates that unmeasured confounders would need to both be more prevalent among those exposed to maltreatment and increase the risk of mental-physical multimorbidity by a factor of 2.29 for those reporting 1 type of maltreatment and 16.92 for those reporting all types to fully explain the observed associations, some residual confounding is anticipated. Second, the cross-sectional design prohibited investigation of multimorbidity onset and precludes causal interpretation. Third, it is important to note that the gender variable in the MHACS dataset includes non-binary individuals who were randomly imputed into the "Men+" and "Women+" categories (Statistics Canada, 2023), and that men were underrepresented at the highest exposure level. Although we adjusted for LGBTQ2+ identity and the latter is reflected in the uncertainty associated with our estimates, these factors should be considered when interpreting our gender-specific findings. Finally, while our exposure variable aligns with existing maltreatment surveillance frameworks in Canada and implicitly captures aspects of both cumulative and subtype-specific exposure, it does not explicitly disentangle their contributions.

Conclusion
While our findings indicate that childhood maltreatment is associated with increased odds of mental-physical multimorbidity, they do not suggest that early adversity leads deterministically to multimorbidity. To support intervention development, further research is needed. Longitudinal studies should aim to clarify the gender-specific biopsychosocial pathways linking childhood maltreatment to mental-physical multimorbidity and identify opportunities for prevention and mitigation. To prevent the progression of morbidity into multimorbidity and multimorbidity into mental-physical multimorbidity, more work is needed to understand the related onset patterns. By preventing further harm and creating additional supports for those affected by childhood maltreatment, it may be possible to reduce the public health burden of mental-physical multimorbidity.

References

Abacus Data Network. (n.d.) *Abacus Data Library.* https://abacus.library.ubc.ca/

Afifi, T. O., MacMillan, H. L., Boyle, M., Taillieu, T., Cheung, K., & Sareen, J. (2014). Child abuse and mental disorders in Canada. *CMAJ: Canadian Medical Association journal = journal de l'Association medicale canadienne*, *186*(9), E324–E332. https://doi.org/10.1503/cmaj.131792

Badley, E. M., Shields, M., O'Donnell, S., Hovdestad, W. E., & Tonmyr, L. (2019). Childhood Maltreatment as a Risk Factor for Arthritis: Findings From a Population-Based Survey of Canadian Adults. *Arthritis care & research*, *71*(10), 1366–1371. https://doi.org/10.1002/acr.23776

Baltramonaityte, V., Pingault, J., Cecil, C. a. M., Choudhary, P., Järvelin, M., Penninx, B. W. J. H., Felix, J., Sebert, S., Milaneschi, Y., & Walton, E. (2023). A multivariate genome-wide association study of psycho-cardiometabolic multimorbidity. *PLoS Genetics*, *19*(6), e1010508. https://doi.org/10.1371/journal.pgen.1010508

Campeau, A., Qadri, S., Barakat, F., Williams, G., Hovdestad, W., Shahid, M., & Lary, T. (2020). At-a-glance - the Child Maltreatment Surveillance Indicator Framework. *Health Promotion and Chronic Disease Prevention in Canada*, *40*(2), 58–61. https://doi.org/10.24095/hpcdp.40.2.04

Canadian Institutes of Health Research. (2022). Ethical Conduct for Research Involving Humans. https://ethics.gc.ca/eng/documents/tcps2-2022-en.pdf

Dai, H., Mei, Z., An, A., & Wu, J. (2020). Epidemiology of physical and mental comorbidity in Canada and implications for health-related quality of life, suicidal ideation, and healthcare utilization: A nationwide cross-sectional study. *Journal of Affective Disorders*, *263*, 209–215. https://doi.org/10.1016/j.jad.2019.11.146

England-Mason, G., Casey, R., Ferro, M., MacMillan, H. L., Tonmyr, L., & Gonzalez, A. (2018). Child maltreatment and adult multimorbidity: results from the Canadian Community Health Survey. *Canadian journal of public health = Revue canadienne de sante publique*, *109*(4), 561–572. https://doi.org/10.17269/s41997-018-0069-y

Ferris, J. K., Prangnell, A., Wagar, B., Choi, A., Simkin, J., Woods, R. R., Sbihi, H., Harder, K., & Smolina, K. (2025). Trends and inequalities in multimorbidity from 2001/2002 to 2019/2020: A population-based study in British Columbia. *PubMed*, *36*(3), 3–16. https://doi.org/10.25318/82-003-x202500300001-eng

Gallo, E. a. G., Munhoz, T. N., De Mola, C. L., & Murray, J. (2018). Gender differences in the effects of childhood maltreatment on adult depression and anxiety: A systematic review and meta-analysis. *Child Abuse & Neglect*, *79*, 107–114. https://doi.org/10.1016/j.chiabu.2018.01.003

Hainmueller, J. (2012). Entropy Balancing for Causal Effects: A Multivariate Reweighting Method to Produce Balanced Samples in Observational Studies. *Political Analysis*, *20*(1), 25–46. doi:10.1093/pan/mpr025

Halstead, S., Sartorius, N., Every-Palmer, S., Siddiqi, N., De Girolamo, G., Siskind, D., & Warren, N. (2024). Physical multimorbidity and mental illness: A global challenge. *Australian & New Zealand Journal of Psychiatry*, *58*(4), 293–296. https://doi.org/10.1177/00048674241235587

Hovdestad, W. E., Shields, M., Shaw, A., & Tonmyr, L. (2020). Childhood maltreatment as a risk factor for cancer: findings from a population-based survey of Canadian adults. *BMC Cancer*, *20*(1). https://doi.org/10.1186/s12885-019-6481-8

Hughes, K., Bellis, M. A., Hardcastle, K. A., Sethi, D., Butchart, A., Mikton, C., Jones, L., & Dunne, M. P. (2017). The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet. Public health*, *2*(8), e356–e366. https://doi.org/10.1016/S2468-2667(17)30118-4

Kone, A. P., Mondor, L., Maxwell, C., Kabir, U. S., Rosella, L. C., & Wodchis, W. P. (2021). Rising burden of multimorbidity and related socio-demographic factors: a repeated cross-sectional study of Ontarians. *Canadian journal of public health = Revue canadienne de sante publique*, *112*(4), 737–747. https://doi.org/10.17269/s41997-021-00474-y

Longpré-Poirier, C., Dougoud, J., Jacmin-Park, S., Moussaoui, F., Vilme, J., Desjardins, G., Cartier, L., Cipriani, E., Kerr, P., Le Page, C., & Juster, R. P. (2022). Sex and Gender and Allostatic Mechanisms of Cardiovascular Risk and Disease. *The Canadian journal of cardiology*, *38*(12), 1812–1827. https://doi.org/10.1016/j.cjca.2022.09.011

Metzler, M., Merrick, M. T., Klevens, J., Ports, K. A., & Ford, D. C. (2016). Adverse childhood experiences and life opportunities: Shifting the narrative. *Children and Youth Services Review*, *72*, 141–149. https://doi.org/10.1016/j.childyouth.2016.10.021

Needham, B., & Hill, T. D. (2010). Do gender differences in mental health contribute to gender differences in physical health? *Social Science & Medicine*, *71*(8), 1472–1479. https://doi.org/10.1016/j.socscimed.2010.07.016

O'Mahony, J., Bernstein, C. N., & Marrie, R. A. (2024). Adverse childhood experiences and psychiatric comorbidity in multiple sclerosis, inflammatory bowel disease, and rheumatoid arthritis in the Canadian longitudinal study on aging. *Journal of psychosomatic research*, *187*, 111893. https://doi.org/10.1016/j.jpsychores.2024.111893

Punjani, N., Flannigan, R., Oliffe, J. L., McCreary, D. R., Black, N., & Goldenberg, S. L. (2018). Unhealthy Behaviors Among Canadian Men Are Predictors of Comorbidities: Implications for Clinical Practice. *American journal of men's health*, *12*(6), 2183–2193. https://doi.org/10.1177/1557988318799022

Rubin, D. B. (1987). Multiple imputation for nonresponse in surveys. In *Wiley series in probability and statistics*. https://doi.org/10.1002/9780470316696

Shields, M. E., Hovdestad, W. E., Pelletier, C., Dykxhoorn, J. L., O'Donnell, S. C., & Tonmyr, L. (2016a). Childhood maltreatment as a risk factor for diabetes: findings from a population-based survey of Canadian adults. *BMC public health*, *16*(1), 879. https://doi.org/10.1186/s12889-016-3491-1

Shields, M. E., Hovdestad, W. E., Gilbert, C. P., & Tonmyr, L. E. (2016b). Childhood maltreatment as a risk factor for COPD: findings from a population-based survey of Canadian adults. *International journal of chronic obstructive pulmonary disease*, *11*, 2641–2650. https://doi.org/10.2147/COPD.S107549

Souama, C., Lamers, F., Milaneschi, Y., Vinkers, C. H., Defina, S., Garvert, L., Stein, F., Woofenden, T., Brosch, K., Dannlowski, U., Galenkamp, H., De Graaf, R., Jaddoe, V. W. V., Lok, A., Van Rijn, B. B., Völzke, H., Cecil, C. a. M., Felix, J. F., Grabe, H. J., . . . Penninx, B. W. J. H. (2023). Depression, cardiometabolic disease, and their co-occurrence after childhood maltreatment: an individual participant data meta-analysis including over 200,000 participants. *BMC Medicine*, *21*(1). https://doi.org/10.1186/s12916-023-02769-y

Statistics Canada. (2023, September 21). *Mental Health and Access to Care Survey (MHACS)*. https://www23.statcan.gc.ca/imdb/p2SV.pl?Function=getSurvey&Id=1378635

Stephenson, E. (2023). Mental disorders and access to mental health care. *Insights on Canadian society. https://www150.statcan.gc.ca/n1/pub/75-006-x/2023001/article/00011-eng.htm*

Thompson, A. E., Anisimowicz, Y., Miedema, B., Hogg, W., Wodchis, W. P., & Aubrey-Bassler, K. (2016). The influence of gender and other patient characteristics on health care-seeking behaviour: a QUALICOPC study. *BMC family practice*, *17*, 38. https://doi.org/10.1186/s12875-016-0440-0

Walsh, C. A., MacMillan, H. L., Trocmé, N., Jamieson, E., & Boyle, M. H. (2008). Measurement of victimization in adolescence: Development and validation of the Childhood Experiences of Violence Questionnaire. *Child Abuse & Neglect*, *32*(11), 1037–1057. https://doi.org/10.1016/j.chiabu.2008.05.003

Wan, A., Bernstein, C. N., Graff, L. A., Patten, S. B., Sareen, J., Fisk, J. D., Bolton, J. M., Hitchon, C., Marriott, J. J., Marrie, R. A., & CIHR Team in Defining the Burden and Managing the Effects of Immune-mediated Inflammatory Disease (2022). Childhood Maltreatment and Psychiatric Comorbidity in Immune-Mediated Inflammatory Disorders. *Psychosomatic medicine*, *84*(1), 10–19. https://doi.org/10.1097/PSY.0000000000001025