Network structure induced bias in estimates of intrinsic generation times

Pratyush K. Kollepara[1], Chiara Poletto[2], and Joel C. Miller[1,3]

[1]Department of Mathematical and Physical Sciences, La Trobe University, Melbourne, Australia
[2]Department of Molecular Medicine, University of Padova, Padova, Italy
[3]Australian Centre for AI in Medical Innovation, La Trobe University, Melbourne, Australia

May 16, 2025

Abstract

The reproduction number is a critical measure of the severity of an infectious disease epidemic. The generation interval, defined as the time taken by an infector to create another infection from its time of infection, is crucial for estimating the basic reproduction number. However, the generation intervals observed, 'realised generation intervals', change over time depending on the dynamics of the epidemic. These time varying distributions are well understood for homogeneous populations, and can be used to infer the intrinsic generation interval distribution. For heterogeneous populations, the state-of-the-art method relies on the use of expensive network based or agent based simulations. We simplify this process by writing exact expressions for simple structured populations. We then use these expressions to demonstrate some previously unexplored biases in the estimation of the intrinsic generation times from the observed generation times, caused by the network structure assumed in the model and particularly the temporal structure of contacts.

1 Introduction

Generation time, the time a parent takes to produce an offspring since its birth, is an important quantity of interest in ecology. The concept has been used extensively in epidemiology, where it refers to the time taken for a newly infected individual to infect someone else. Knowing this interval distribution is necessary for planning contact tracing, quarantine, and isolation in case of an

outbreak [13]. It is also necessary to estimate $\mathcal{R}_0$, which quantifies the epidemic transmission potential, from the aggregated incidence of infections using the Wallinga-Lipsitch equation [15]. However, measuring this quantity and correctly interpreting it for outbreak analysis is not straightforward [17].

The Wallinga-Lipsitch equation requires the "intrinsic generation interval", which is defined as the time distribution of infectious contacts made by a primary case, or, equivalently, the time distribution of generated infections when the population is fully susceptible and mixes randomly [17]. The observed or realised generation intervals, obtained by contact tracing data, are different from the intrinsic ones, because of dynamical and population mixing effects, and also because of how generation intervals are defined and aggregated from available data. For instance, The forward generation interval distribution refers to the distribution of times it takes for a group of individuals infected at a given time to transmit the infection to others. On the other hand, the backward generation interval distribution takes the point of view of a cohort of infectees, infected at a given time, and is defined as the distribution of times to infect them [17].

The intrinsic generation interval, determined solely by the infector's infectiousness, remains constant throughout an outbreak. In contrast, forward and backward realized generation times vary. The functional relationship between intrinsic and realised generation time was mathematically studied for a randomly mixing population by the Champredon and Dushoff [17] within the renewal equation framework. The backward generation time increases monotonically, underestimating the intrinsic generation interval early on and overestimating it later. This bias arises from the incidence dynamics. During the growth phase, recent infections dominate the pool of potential infectors, skewing generation times shorter; during the decline, older infections dominate, lengthening the generation intervals. On the other hand, the forward-looking formulation of generation time avoids this dynamic bias. Still, its mean value shrinks right before the peak [17, 16, 12]. One reason for this contraction is the high rate at which susceptibles are being depleted near the peak. An infector may generate multiple transmissions throughout their infectious period and, if the number of susceptibles remains the same, then the transmission events are all equally likely to contribute to the generation intervals [17]. Conversely, if the availability of susceptibles shrinks significantly during the infectious period the later transmissions are less likely to lead to an infection and longer transmission intervals are less likely to be sampled. An alternative mechanism to understand the generation time contraction is the competition between infectors, racing to infect someone [16, 11]. When many infectors compete to infect the same susceptible individual, only the transmission by the fastest infector counts as an infection event and contributes to the generation time statistics, causing the generation interval to contract. The study of these dynamical mechanisms is essential to design techniques to fit the intrinsic generation time from contact tracing data [5, 14, 6].

Although most studies rely on the homogenous mixing assumption, contacts within households, schools, and workplaces are recurrent, heterogeneous, and spatially clustered [19, 3, 9, 10, 7, 8, 1, 2, 19, 4]. Previous studies investigating

clustering and recurrence showed that these enhance the competition between infectors and cause a stronger reduction of generation intervals — more than predicted by homogeneous-mixing models — which persists throughout the outbreak [19, 11]. Despite these findings, the effect of the contact network on the generation time dynamics remains under-studied, especially from an analytical perspective. Here, we use the edge-based compartmental modelling framework [18] to derive exact equations for realized generation intervals. The framework allows us to systematically address the role of contacts' heterogeneity - by comparing a homogeneous and a heterogeneous network - and recurrence - by comparing the cases in which contacts' identities continuously change or are fixed. Our theoretical framework, supported by simulations, provides insights into the dynamics of generation intervals and highlights differences between network topologies. Most importantly it points to potential sources of bias, that were left unexplored so far.

---

2 Methods

This section gives a brief overview of the Edge Based Compartmental Modelling (EBCM) framework [18] that is required for understanding the results in the next section. The EBCM framework can be used to solve the SIR model on configuration model networks. These networks are specified by a distribution of the degrees of nodes in the network but are limited to a tree-like structure (do not allow clustering or higher-order structures).

Real-world networks have a temporal nature: some contacts of an individual are fleeting (fast changing) while others are static (long-lasting). The models used here will consider versions of configuration model networks for each type of contact. The first one is known as the annealed or mean field configuration model where even though the number of contacts of an individual remains constant, the edges in the network rewire to new nodes constantly. The other is known as the quenched or static configuration model where the edges are static. In addition to these assumptions about the network structure, we also assume that both transmission and recovery from infection are described by a Poisson process. Detailed derivations of our results are presented in the Supplementary Methods section, and the results can be generalised to non-Poisson processes.

For both annealed and quenched networks, a standard compartmental modelling approach would divide the population up into sub-groups of homogeneous degree (number of contacts), resulting in at least as many differential equations as the number of sub-groups. The EBCM framework reduces such a large system of differential equations into two ordinary differential equations which can be solved numerically with ease.

The EBCM framework makes use of probability-generating functions

$$\psi(x) = \sum_i N_i \ x^i,$$

where N_i refers to the proportion or probability of nodes of degree class i. A variable θ is defined to be the probability that a randomly chosen partnership of a node has not transmitted to it yet. If a node of degree k is susceptible, then none of its k partners have transmitted to it. The probability for the node to be susceptible (given its degree k) is θ^k. The prevalence of susceptibles in the population can be computed using

$$S = \sum_k N_k \theta^k = \psi(\theta).$$

2.1 Annealed network

Homogeneous compartmental models use macroscopic variables such as the proportion of the population that is infected, susceptible, recovered etc. The probability a random interaction with another individual is with an infected individual is assumed to equal the proportion of the population that is infected. In a more general case where the number of partners differs among individuals, the probability a random contact is with an infected individual is the prevalence of half-edges in the population that are attached to an infected node. The variable $I(t)$ (the proportion of the population that is infected) is generalised to a variable $\pi_I(t)$ which is the proportion of half-edges that connect to infected nodes. Similarly, we generalize $\pi_S(t)$ and $\pi_R(t)$ which are the prevalence of half-edges with susceptible and recovered nodes respectively. We also define a new variable $\pi_i(t)$, which is the incidence of infected stubs, i.e. the instanteous rate at which half-edges with infected nodes are created by new infections.

We find that the probability a random half-edge connects to a susceptible node is

$$\pi_S = \frac{\sum_k k N_k \theta^k}{\sum_k k N_k} = \theta \frac{\psi'(\theta)}{\psi'(1)}$$

Knowing π_S requires $\theta(t)$. At the start of the epidemic, the initially susceptible nodes have not received any transmissions. So, using the initial condition $\theta(0) \approx 1$, the following differential equation (derived in ref [18]) can be solved to find $\theta(t)$:

$$\dot{\theta} = -\beta\theta + \beta \frac{\theta^2 \psi'(\theta)}{\psi'(1)} - \gamma\theta\ln\theta$$

2.2 Quenched network

In the quenched case, the variable $S(t)$ is generalised to a variable $\phi_S(t)$ which is the probability that the partner of a randomly selected node is susceptible (under the assumption that the randomly selected node has not transmitted to its partner).

$$\phi_S = \frac{\sum_k k N_k \theta^{k-1}}{\sum_k k N_k} = \frac{\psi'(\theta)}{\psi'(1)}$$

In this case, using a similar initial condition for θ as before, the following differential equation [18] is solved:

$$\dot{\theta} = -\beta\theta + \beta\frac{\psi'(\theta)}{\psi'(1)} - \gamma(1 - \theta)$$

3 Results

3.1 Annealed network

In a homogeneous population, the forward generation interval distribution can be calculated using the intrinsic generation interval distribution, $g(\tau)$, and the proportion or number of susceptible people, $S(t)$, with time [17].

$$f_H(t,\tau) = \frac{g(\tau)S(t+\tau)}{\int_0^\infty dx\ g(x)S(t+x)}$$

Here, $S(t)$ gives the probability for a random individual to come in contact with a susceptible individual at time t. In an annealed heterogeneous network this is no longer $S(t)$ and depends on the connectivity. We thus need to replace it with $\pi_S(t)$. Therefore, the forward generation interval distribution for an annealed configuration model network is

$$f_A(t,\tau) = \frac{g(\tau)\pi_S(t+\tau)}{\int_0^\infty dx\ g(x)\pi_S(t+x)}.$$

Turning our attention to the backward generation interval, Champredon and Dushoff's equation for the homogeneous case makes use of the incidence of new infections as a function of time [17]:

$$b_H(t,\tau) = \frac{g(\tau)i(t-\tau)}{\int_0^\infty dx\ g(x)i(t-x)}.$$

If we use the average incidence of stubs, instead of the incidence of infections, we obtain an exact expression for the backward generation interval distribution for an annealed heterogeneous network.

$$b_A(t,\tau) = \frac{g(\tau)\pi_i(t-\tau)}{\int_0^\infty dx\ g(x)\pi_i(t-x)}.$$

A general characteristic of the backward generation intervals is that they are stable in time when the incidence grows exponentially [17]. This remains valid in the heterogeneous network case, as can be seen by plugging in $\pi_i(t) = \pi_i(0)e^{\lambda t}$ into the above formula, leading to

$$b_A^{exp}(t,\tau) = (\lambda + \gamma)e^{-(\lambda+\gamma)\tau}$$

Similar to the forward generation interval, we define an effective reproduction number called the forward reproduction number, $\mathcal{R}_f$. It represents the expected number of infections created by an average infector who became infected at time t over its entire infectious period. For an annealed network,

$$\mathcal{R}_f^A(t) = \frac{\beta(\theta\psi''(\theta(t)) + \psi'(\theta(t)))}{\psi'(\theta(t))} \int_0^\infty d\tau \ e^{-\gamma\tau} \pi_S(t+\tau)$$

If a reproduction number is calculated empirically through observation of transmission chains, we would expect it to match with the forward reproduction number.

3.2 Quenched network

In an annealed network, a node continuously breaks its edges and re-connects to new edges. So, if a node infects another node, it does not compete with itself when transmitting again. Conversely, in a quenched network, a node can compete with itself — after the first transmission all later transmissions to the same neighbour have no effect. In the equations for the realised generation time we thus need to replace the intrinsic generation time $g(\tau)$ with $\tilde{G}(\tau) = e^{-(\beta+\gamma)\tau}$. In addition, the probability that a stub points to a susceptible node – which was $\pi_S(t)$ in the annealed case – is now given by $\phi_S(t)$. The modified version of the equation in [17] then becomes

$$f_Q(t,\tau) = \frac{\tilde{G}(\tau)\phi_S(t+\tau)}{\int_0^\infty dx \ \tilde{G}(x)\phi_S(t+x)}.$$

In the backward case, the incidence of infections, $i(t)$ is replaced with the average probability of incidence among the neighbours of a node, $\phi_i(t)$:

$$b_Q(t,\tau) = \frac{\tilde{G}(\tau)\phi_i(t-\tau)}{\int_0^\infty dx \ \tilde{G}(x)\phi_i(t-x)}$$

From this equation, we obtain that the backward generation intervals are stable when the epidemic grows exponentially, similar to the annealed case. The distribution in this regime is

$$b_Q^{exp}(t,\tau) = (\lambda + \beta + \gamma)e^{-(\lambda+\beta+\gamma)\tau}$$

Finally, the forward reproduction number can be derived for quenched network as follows

$$\mathcal{R}_f^Q(t) = \frac{\beta\psi''(\theta(t))}{\psi'(\theta(t))} \int_0^\infty d\tau \ e^{-(\beta+\gamma)\tau}\phi_S(t+\tau)$$

3.3 Impact of network properties on the realised generation time

We validate the exact results using event-based simulations [21]. Figures 1 and 2 show the comparison between analytical solutions and simulations for the annealed and quenched case, respectively. In each figure, we plot the epidemic

Figure 1: *Comparison of epidemics and their realised generation intervals using an event-based simulation method and the exact solution for annealed networks with two types of heterogeneous degree distributions (Left: the degree of a node can be 3, 5, or 7 with equal probabilities, Right: degree distribution is a truncated power law with maximum degree of 70 and exponent of -2). We find a good match between theory and simulations for the realised generation intervals and the forward reproduction number. The time-series are centred such that the prevalence peaks at time zero. The thin lines show trajectories from each of the 100 simulations, while the plot with markers shows the average. At the start and towards the end of the epidemic, significant noise is observed and the simulation deviates from the theory.*

profile, the forward and backward generation times and the forward reproduction number. The figures show the comparison for two different topologies that we call TMD and TPL70. TMD refers to a tri-modal degree distribution and TPL70 refers to a truncated power law degree distribution (See Figure captions more details). Both networks are heterogeneous, but the TPL70 network is a case of extreme heterogeneity. The forward generation intervals in the annealed case show a contraction qualitatively similar to what is predicted by the homogenous mixing equations [17]. This is, however, less evident when the network is extremely heterogeneous. In the quenched case forward generation times shrink substantially due to the intensified competition among infectors. While forward generation time varies substantially from one network to another, backwards generation time is quite robust to the network structure.

We can further analyse the impact of the network features by numerically solving the derived analytical expressions for different scenarios. This is significantly inexpensive compared to stochastic numerical simulations. We performed various numerical experiments and we summarise the results here:

Experiment (i) Structure: Annealed TPL70 network and homogeneous network with same mean degree. Disease parameters: Identical recovery rate, intrinsic generation interval ($\gamma = 1$) and $\mathcal{R}_0$. Result: Heterogeneity reduces the contraction in the forward generation interval. However, this could possibly be explained by the lower peak (hence lesser competition) in the heterogeneity (See 4.3).

Experiment (ii) Structure: Annealed TPL70 network and homogeneous network with same mean degree.
Disease parameters: Identical recovery rate, intrinsic generation interval ($\gamma = 1$). The transmission rate is selected such that the two epidemics have the same level of peak incidence of infections.
Result: Heterogeneity increases the contraction in the forward generation interval, despite the same maximum incidence of infections. This can be explained by noting that the peak of $\pi_i(t)$ is higher than $i(t)$. Furthermore, the backward generation intervals also differ significantly (See Figure 4.3).

Experiment (iii) Structure: Annealed TPL70 network Method: Compare distribution of realised intervals assuming homogeneity [17] (i.e. using $i(t)$ and $S(t)$) and not assuming homogeneity (i.e. using $\pi_i(t)$ and $\pi_S(t)$)
Result: Using the net incidence of infections underestimates the contraction in forward generation interval. In contrast, the backward generation intervals are relatively unchanged (See Figure 4.3).

Figure 2: *Comparison of epidemics and their realised generation intervals using an event-based simulation method and the exact solution for quenched networks with two types of heterogeneous degree distributions (Left: the degree of a node can be 3, 5, or 7 with equal probabilities, Right: degree distribution is a truncated power law with maximum degree of 70 and exponent of -2). We find a good match between theory and simulations for the realised generation intervals and the forward reproduction number. The time-series are centred such that the prevalence peaks at time zero. The thin lines show trajectories from each of the 100 simulations, while the plot with markers shows the average. At the start and towards the end of the epidemic, significant noise is observed and the simulation deviates from the theory for the truncated power law distribution.*

3.4 Implications for estimation of reproduction number

In this section, we will explore how errors could arise if disease parameters are inferred using existing methods to relate the observed generation interval and growth rate to the basic reproduction number.

If the intrinsic generation intervals and the exponential growth rate are known, the basic reproduction number can be estimated using the Wallinga-Lipsitch equation [15]. The growth rate can be estimated from incidence at the start of the epidemic. However, the intrinsic generation intervals are not observed. Instead we observe the realised generation intervals (forward or backward) which are influenced by the dynamics of the outbreak and the contact structure of the population.

So to fit model parameters and infer the basic reproduction number from the growth rate and the observed generation intervals, it is crucial to recognise the distinction between the realised generation intervals and the intrinsic generation intervals. In principle, the exact expressions for realised generation interval distributions can be used to infer the intrinsic generation interval distribution, which can be used to then estimate $\mathcal{R}_0$ [17, 5, 14, 6]. However, the Champredon and Dushoff's equation and further developments relies on the homogenous mixing assumption.

Our results from previous sectionss show that for a heterogeneous population, the conclusions will be incorrect if simple population aggregates such as prevalence of susceptibles or incidence of infections are used. We expect that the joint use of simple aggregate observations of an outbreak and realised generation intervals could bias the estimation of the basic reproduction number and the intrinsic generation intervals. To demonstrate this bias, we now look at an example case of outbreak analysis and two scenarios with different amounts of information available to the disease modeller.

Example Let us consider a pathogen causing an epidemic in different disconnected populations. There is a population in which the authorities have the capacity and resources to document the incidence of cases and also perform backward contact tracing (let us call it Population 1). There is another population where we do not have the capacity to perform contact tracing, but we can document case incidence (call it Population 2). Both populations are well described by a quenched and heterogeneous contact network. This is a simplification, as any real-world network will have a mixture of short and long time scale contacts. First, we describe the quantities of interest in the ground reality from the correct model. Then, we consider cases where an infectious disease modeller has some information about the contact structure and see how their modelling assumptions can affect model inference in both populations.

Using the ground reality model, we can write the basic reproduction numbers

in terms of the observable quantities:

$$\mathcal{R}_0^1 = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{1}$$

$$\mathcal{R}_0^2 = \frac{1}{1 - \lambda_2 T_{exp,2}} \tag{2}$$

where the λ is the exponential growth rate estimated from incidence and T_{exp} is the mean backward generation interval during the exponential growth phase, estimated from the contact tracing data. We can also write the average intrinsic generation time as a function of the same quantities. However, the expression depends also on the first and second moments of the degree distribution. See detailed methods for a detailed calculation.

Case A: The modeller does not know anything about the contact structure and assumes a well-mixed homogeneous model. The model has essentially two parameters, the transmission rate β and the recovery rate γ (which also specifies the intrinsic generation intervals). Using this model, the estimators for population 1 are:

$$\widehat{\gamma}_1 = \frac{1}{T_{exp,1}} - \lambda_1 \tag{3}$$

$$\widehat{\mathcal{R}_0^1} = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{4}$$

Comparing these estimates to the ground reality tells us that the basic reproduction number is unbiased but $\widehat{\gamma}_1$ is not as it does not properly account for the contact network properties – first and second moment of the degree distribution.

For population 2, nothing is known about the backward generation intervals, but the modeller has an estimate for the intrinsic generation intervals in population 1. Since by definition, intrinsic intervals would not vary among populations, the modeller might see it fit to treat $\widehat{\gamma}_1$ as an estimator for γ_2. This leads to an estimate of the reproduction number

$$\widehat{\mathcal{R}_0^2} = \frac{1 + (\lambda_2 - \lambda_1)T_{exp,1}}{1 - \lambda_1 T_{exp,1}} \neq \frac{1}{1 - \lambda_2 T_{exp,2}} \tag{5}$$

which is biased. We find that the bias in the estimator for population 2 is greater when the contact structure is different among the two populations. When the contact structure is identical, the estimator is unbiased when the growth rates are equal.

Case B: The modeller knows that the contacts are quenched and also has correct information of the average number of contacts $\langle K_1 \rangle$ and $\langle K_2 \rangle$ respectively. In the absence of more detailed information, they assume a homogeneous quenched network with degrees $k_1 = \langle K_1 \rangle$ and $k_2 = \langle K_2 \rangle$, respectively. The model has two other parameters, the transmission rate β and the recovery rate γ (which also specifies the intrinsic generation intervals). Using this model, the

estimators for population 1 are:

$$\widehat{\gamma}_1 = \frac{(k_1 - 2)(1/T_{exp} - \lambda) - \lambda}{k_1 - 1} \tag{6}$$

$$\widehat{\mathcal{R}_0^1} = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{7}$$

Comparison with the ground reality model shows that $\widehat{\gamma}_1$ is biased, as calculation does not correctly account for the second moment of the degree distribution, while $\widehat{\mathcal{R}_0^1}$ is not. As in the previous case, the modeller might see fit to treat $\widehat{\gamma}_1$ as an estimator for γ_2. This leads to an estimate of the reproduction number for population 2

$$\widehat{\mathcal{R}_0^2} = \frac{(\lambda_2 + \widehat{\gamma}_1)(k_2 - 1)}{\lambda_2 + \widehat{\gamma}_1(k_2 - 1)}$$

This estimator is biased, albeit behaves very differently than the biased estimator in Case A.

In summary, we find that if the incidence and backward contact tracing is available in a population, then even incorrect assumptions about the contact structure can yield an unbiased estimator for the basic reproduction number, even though the estimator for the intrinsic generation interval is biased. Using this estimated intrinsic generation interval in order to estimate the basic reproduction number in another population leads to biased estimates. See 3 for an illustration of the bias in $\widehat{\mathcal{R}_0^2}$.

While case A is realistic, case B is hypothetical since more information – i.e. the average number of contacts – is available and used in the analysis than what normally happens in reality. Paradoxically, case B presents a more severe bias. For Population 1, a truncated power law degree distribution is used with an exponent of -2 and maximum degree of 70. Population 2 has a similar structure but with a maximum degree of 30. The growth rate in Population 1 is $\lambda_1 = 0.5$ and recovery rate is $\gamma = 1$.

---

4 Discussion

In this article, we have generalised Champredon and Dushoff's theory [17] for realised generation intervals in a homogeneous and well-mixed population to heterogeneous networks, using the EBCM framework. This framework allows incorporating degree heterogeneity in a tree-like network. In structured populations, realised generation intervals have been investigated using network and agent-based simulations [19, 4]. These methods benefit from their ability to be modified quite easily – adding more coarse-grained structure or even including forms of structure beyond degree heterogeneity. However, they are slow and managing the noise reasonably would require a large number of simulations or a large population. In addition, they do not provide analytical insights. Some

Figure 3: *Comparison between the estimated and true value of the basic repro-
duction number in a heterogeneous quenched network when different homoge-
neous models are used and the intrinsic generation interval is estimated from a
different population (see 3.4 for a detailed explanation).*

analytical considerations were provided by a few works [20, 11]. Our exact
expressions, validated using simulations, present an advance in our ability to
predict generation intervals in simple heterogeneous contact structures and sets
up a foundation to obtain exact expressions for more complex heterogeneous
networks.

Generation intervals are useful in estimating the basic reproduction number
of an epidemic. Champredon and Dushoff showed how fitting the mean intrinsic
generation interval to the mean of the realised generation intervals (backward)
is incorrect [17]. They anticipate that heterogeneity may affect their results.
We confirm this, determine the manner by which it affects the results and more
importantly, through what mechanism. We found that the realised intervals
depend not on simple aggregate measures like net prevalence of susceptibles or
net incidence of infection, but on more involved measures such as the prevalence
of half-edges in the network which have a susceptible or a newly infected node
attached to them. Measurement of such quantities requires higher resolution
observation of the population, at the scale of groups which are homogeneous in
number of contacts.

We also use a series of hypothetical experiments to show that the estimates
of intrinsic intervals (inferred using realised generation intervals) are sensitive
to model assumptions. Despite this sensitivity, the estimates of $\mathcal{R}_0$ can be relied
upon as long as the estimated intrinsic generation intervals are used in the same
setting where they were estimated. If the estimated intrinsic intervals are used

in a different setting (say an epidemic caused by the same pathogen in a different setting where the contact/network structure differs) then the estimates of $\mathcal{R}_0$ will be biased. This practice of applying generation interval estimates or related parameter estimates from from one setting to an epidemic of the same pathogen in another setting is common [22, 23, 24, 25, 26, 27].

The forward reproduction number that was derived here is an effective reproduction number. Effective reproduction numbers are of two types: instantaneous or cohort-based (case reproduction number). The instantaneous reproduction number measures the transmission happening at certain time while the case reproduction number measures the transmission that is caused by a cohort of infectors who became infected at a certain time [28]. The forward reproduction number, $\mathcal{R}_f$, is equivalent to the case reproduction number [28, 29].

A key assumption throughout this work is that the recovery from infection is a Poisson process (the intrinsic generation interval distribution is exponential). While real-world diseases do not follow this assumption, the expressions for forward and backward generation intervals that we derived are fairly general. Real-world networks have many kinds of heterogeneities beyond the kind we have investigated here. They have clustering (triads that form a triangle), multiple layers and temporal variation. Here, we have only studied degree heterogeneity in two extreme cases of temporal variation – quenched (static partners) and annealed (fleeting contact with changing partners). In principle, a linear combination of the distributions derived here may be used to model contact networks with a more realistic temporal behaviour.

Data and Software

The software used in this study is available at
`https://github.com/praty-k/gi-networks`

Acknowledgements

This research was supported by use of the Nectar Research Cloud, a collaborative Australian research platform supported by the NCRIS-funded Australian Research Data Commons (ARDC). We thank Andrew Black and Eben Kenah for helpful conversations.

Funding: P.K.K.: La Trobe Graduate Research Scholarship and Full-Fee Research Scholarship; Ed Smith Applied Mathematics Fellowship.
C.P.: Cariparo Foundation through the program Starting Package; Department of Molecular Medicine through the program SID from BIRD funding.

Supplementary Methods

We use the framework of Edge Based Compartmental Models to derive exact expressions for the generation interval distribution [18]. We assume that the infectiousness, β, is constant, and that infectious individuals recover at a constant rate γ. Unless otherwise stated, the heterogeneous network refers to a configuration model network with a truncated power law degree distribution which has an exponent of -2, and integer degrees between 1 and 70. The mean degree of nodes in such a network is ~ 3 and the mean of square of degrees is ~ 43. These properties are used in calculating the basic reproduction number, growth rate etc. [18]

4.1   Annealed Network

The average probability of contacting an individual in state X who has k stubs (or is of degree k) is proportional to the product of probability of being in state X, of having k stubs, and the number of stubs k (because if a node has more stubs, then a random node is more likely to connect to it):

$$P(X,k) \sim X_k k, \text{and } \pi_X = \frac{\sum_k X_k k}{\sum_k N_k k}, \text{Thus, } \pi_S(t) = \frac{\sum_k S_k k}{\sum_k N_k k} = \frac{\sum_k N_k \theta^k k}{\sum_k N_k k} = \frac{\theta \psi'(\theta)}{\psi'(1)}$$

where θ is the probability that the neighbour of a randomly selected node has not transmitted to it yet. So, the number of susceptibles of degree k is equal to the number of individuals of degree k, times the probability that none of its k neighbours has transmitted to it. The probability with which a randomly selected individual would come in contact with the stub of a susceptible node is π_S, with the stub of an infected node is π_I, with the stub of a recovered node is π_R [18]. These three probabilities would sum up to unity.

Using the EBCM framework, we define new a quantity, $\pi_i(t)$, which is the probability of contacting the stub of a newly infected node at time t. This can also be described as the average incidence of infected stubs.

The depletion rate of susceptibles of degree k is given by:

$$\frac{dS_k}{dt} = -\beta k S_k(t) \pi_I(t)$$

This is also negative of the rate at which new infections are being created. At time t, the creation rate (i.e., the incidence) of new infections of degree k is

$$i_k(t) = \beta k S_k \pi_I(t)$$

The rate of contacting an infectious individual who was infected at time t and has k edges will be $k i_k(t)$. And the rate of contacting an individual infected at t will be the average $\frac{\sum_k k i_k}{\langle k \rangle}$. The incidence of infections of degree k is $-\frac{dS_k}{dt}$. Therefore, from equation (), the average incidence of stubs can be defined as

$$\pi_i = \beta \pi_I \frac{\sum_k k^2 S_k}{\langle k \rangle} = \beta \pi_I \frac{\sum_k k^2 N_k \theta^k}{\langle k \rangle} = \beta \pi_I \frac{\theta^2 \psi''(\theta) + \theta \psi'(\theta)}{\psi'(1)}$$

The incidence of infected stubs can also be obtained by considering the fact that the incidence of infected stubs is the same as the rate at which π_S decreases (since nodes, and hence their stubs, can only transition from state-S to state-I). Using the EBCM framework, an expression for $-\frac{d\pi_S}{dt}$ can be derived, which is equivalent to the expression for π_i derived above [18].

Forward Generation Interval Distribution In the homogeneous case, the number of new cases at time $t + \tau$ caused by an average person infected at time t is proportional to [17]

$$i_t(t + \tau) \sim g(\tau)S(t + \tau)i(t)$$

For a configuration model, the number of new cases at time $t + \tau$ with degree k caused by people with degree l infected at time t is proportional to

$$g(\tau)S_k(t + \tau)ki_l(t)l$$

So the incidence at time $t + \tau$ caused by the average person infected at time t is

$$i_t(t + \tau) \sim g(\tau)\sum_k kS_k(t + \tau)\sum_l li_l(t) = g(\tau)\pi_S(t + \tau)\sum_l li_l(t)$$

The distribution of forward generation intervals (τ) at time t is obtained by normalising the above derived incidence:

$$f_A(t, \tau) = \frac{g(\tau)\pi_S(t + \tau)}{\int_0^\infty dx\; g(x)\pi_S(t + x)}$$

Backward Generation Interval Distribution In the homogeneous case, the number of new cases at time t caused by an average person infected at time $t - \tau$ (i.e. τ units ago) is proportional to [17]

$$i_{t-\tau}(t) \sim g(\tau)S(t)i(t - \tau)$$

For a configuration model, the number of new cases at time t with degree k caused by people with degree l infected at time $t - \tau$ is proportional to

$$g(\tau)S_k(t)ki_l(t - \tau)l$$

So the incidence at time t caused by the average person infected at time $t - \tau$ is

$$i_{t-\tau}(t) \sim g(\tau)\sum_k kS_k(t)\sum_l li_l(t - \tau) \sim g(\tau)\pi_S(t)\pi_i(t - \tau)$$

Normalising this incidence derived above will yield the backward generation interval distribution at time t:

$$b_A(t, \tau) = \frac{g(\tau)\pi_i(t - \tau)}{\int_0^\infty dx\; g(x)\pi_i(t - x)}$$

4.2 Quenched Network

A quenched network differs from an annealed/well-mixed/mean-field network in that the partnerships are static in time. In a well-mixed network, a node does not compete with itself to infect a certain node since the edges rewire constantly. In the quenched case, if an infected node transmits to a (susceptible) neighbour twice, then only the first transmission counts as an infection event. The second transmission does not lead to an infection. Thus, the generation intervals are shorter compared to the well-mixed case even when there are no other competing infectors. When calculating the realised (forward and backward) generation intervals, we need to use an effective intrinsic generation interval which is equivalent to the distribution of first transmissions between a pair of infected and susceptible partners.

If the intrinsic generation intervals are given by

$$g(t) = \gamma e^{-\gamma t}$$

then the effective intrinsic generation interval is

$$\tilde{g}(t) = (\beta + \gamma)e^{-(\beta+\gamma)t}$$

This can be derived by considering the first transmission event between a pair of connected nodes. The rate at which the first transmission event occurs at time t is the product of the probabilities of the infector being infectious ($e^{-\gamma t}$), the infector not yet transmitting ($e^{-\beta t}$), the rate of transmission β. Normalising this product gives the rate at which first transmissions occur.

As the above derived equation shows, the first transmission event is a poisson process with rate $\beta + \gamma$. Instead of $\tilde{g}(t)$, what we will be using from here on is the probability that the first transmission has not occurred yet:

$$\tilde{G}(t) = e^{-(\beta+\gamma)t}$$

Forward Reproduction Number   For the quenched network, we will derive a forward reproduction number using the renewal equation. Fundamentally, the renewal equations relates the incidence at present to the incidence during the time that has passed. If the present time is t, then

$$\text{Incidence } (t) = \int_0^t \sum_k \text{Incidence of degree } k \text{ at time } (t - \tau)$$
$$\times \text{Rate of new infections produced by a single infector of}$$
$$\text{degree } k \text{ who was infected } \tau \text{ units ago}$$

Let us calculate the second term of the product. The number of new infections created at time $t + \tau$, by a node u, of degree k, infected at time t (τ units ago) can be computed by:

$$P(u \text{ is infectious at } t + \tau \text{ given it became infected at } t)$$
$$\times P(u \text{ didn't transmit until } t + \tau \text{given it became infected at } t)$$
$$\times (\text{No. of neighbours it could infect})$$
$$\times (\text{Rate at which } u \text{ transmits to a neighbour})$$
$$\times P(\text{a neighbour } v \text{ is susceptible at } t + \tau \text{ given } u \text{ was susceptible at time } t)$$
$$= e^{-(\beta+\gamma)\tau} \times (k - 1) \times \beta \times P(v_{t+\tau} = S \mid u_t = S)$$

$$P(v_{t+\tau} = S \mid u_t = S) = \frac{P(v_{t+\tau}=S \cap u_t=S)}{P(u_t=S)}$$

The denominator, $P(u_t = S)$ is $\theta^k(t)$. The numerator can be written as

$$P(v_{t+\tau} = S \cap u_t = S) = P(v_{t+\tau} = S \cap v \nrightarrow u \text{ by time } t)$$
$$\times P(k-1 \text{ neighbours } \nrightarrow u \text{ by time } t)$$
$$= \phi_S(t+\tau)\theta^{k-1}(t)$$

where $v \nrightarrow u$ denotes v did not transmit to u. Therefore,

$$P(v_{t+\tau} = S \mid u_t = S) = \frac{\phi_S(t+\tau)}{\theta(t)}$$

The number of new infections generated at time $t + \tau$, by a node of degree k who was infected at time t, is:

$$\beta(k-1)e^{-(\beta+\gamma)\tau}\frac{\phi_S(t+\tau)}{\theta(t)}$$

Next, let us try to find the incidence at time t in a degree class k. The probability of getting infected at time t given that a node u has degree k is

$$P(u \text{ is susceptible at } t)$$
$$\times \text{ Rate at which an infected neighbour transmits}$$
$$\times \text{ Number of neighbours who could transmit to } u$$
$$\times \text{Probability that a neighbour } v \text{ is infected and has not transmitted to } u$$
$$\text{given that } u \text{ is susceptible}$$
$$= \theta^k(t) \times \beta \times k \times P(v_t = I \cap v \nrightarrow u \mid u_t = S)$$

$$P(v_t = I \cap v \nrightarrow u \mid u_t = S) = \frac{P(v_t=I \cap v\nrightarrow u \cap u_t=S)}{P(u_t=S)}$$
$$= \frac{P(v_t=I \cap v\nrightarrow u) \times P(k-1 \text{ neighbours } \nrightarrow u \text{ by time } t)}{P(u_t=S)}$$
$$= \frac{\phi_I(t)}{\theta(t)}$$

Therefore, the probability of getting infected at time t given that the node has degree k is

$$\beta k \phi_I(t)\theta^{k-1}(t)$$

Total incidence at time t is therefore

$$N\beta\phi_I(t)\sum_k kN_k\theta^{k-1}(t) = N\beta\phi_I(t)\psi'(\theta(t))$$

where N is the population size.

Now, let us find the total number of new infections at time $t + \tau$ created by individuals who got infected at time t (τ units ago). We make use of the two quantities dervied above: incidence of degree k at time t and number of

infections created at time $t + \tau$ by a single infector of degree k who was infected at time t.

$$N \sum_k \beta k N_k \theta^{k-1}(t) \phi_I(t) \times \beta(k-1) e^{-(\beta+\gamma)\tau} \frac{\phi_S(t+\tau)}{\theta(t)}$$
$$= N\beta^2 \phi_I(t) \phi_S(t+\tau) e^{-(\beta+\gamma)\tau} \sum_k k(k-1) N_k \theta^{k-2}(t)$$
$$= N\beta^2 \phi_I(t) \phi_S(t+\tau) e^{-(\beta+\gamma)\tau} \psi''(\theta(t))$$

Dividing this by the total incidence at time t and integrating over τ gives the expected number of infections created over an infectious period by the average individual who was infected at time t.

$$\frac{\beta \psi''(\theta(t))}{\psi'(\theta(t))} \int_0^\infty d\tau \phi_S(t+\tau) e^{-(\beta+\gamma)\tau}$$

We call this quantity the forward reproduction number, in analogy to the forward generation interval. Note that the term in the product, $\frac{\psi''(\theta(t))}{\psi'(\theta(t))}$ depends only on θ and the degree distribution. It can be interpreted as the expected degree of a node that got infected at time t.

Forward Generation Interval Distribution The forward generation interval distribution at time t can be computed using the rates (as a function of time τ) at which new infections are produced by the infections which happened at t. This rate is proportional to the probability that a partner of a randomly selected node is susceptible, ϕ_S, and the probability that the selected node is still infectious in the future, given it becomes infected at present time, and the probability that it hasn't transmitted to this partner before. The distribution is obtained by normalising these rates against τ.

In a quenched network, the rate at which new infections are caused at time $t + \tau$, by infections which were generated at time t

$$i(t+\tau) \sim \tilde{G}(\tau) \phi_S(t+\tau)$$

because $\phi_S(t+\tau)$ is the probability that an infected neighbour of a susceptible node transmits and $\tilde{G}(\tau) = e^{-(\beta+\gamma)\tau}$ is the probability that a node infected τ time units ago would emit its first transmission. Therefore, the forward generation interval distribution will be given by

$$f_Q(t,\tau) = \frac{\tilde{G}(\tau)\phi_S(t+\tau)}{\int_0^\infty dx \ \tilde{G}(x)\phi_S(t+x)}$$

Backward Generation Interval Distribution The rate at which new infections are caused at time t due to infections created τ units ago (at time $t - \tau$) is

$$i_{t-\tau}(t) \sim \tilde{G}(\tau)\phi_i(t-\tau)$$

because $\phi_i(t - \tau)$ is the rate at which a susceptible neighbour of a node gets infected at time $t - \tau$ and enters the ϕ_I state (i.e. infected neighbour that has

not yet transmitted to our node of interest) and $g(\tau)$ is the probability density that a node infected τ units ago would emit its first transmission. Therefore, the backward generation interval can be obtained by normalising this rate

$$b_Q(t, \tau) = \frac{\tilde{G}(\tau)\phi_i(t - \tau)}{\int_0^\infty dx \ \tilde{G}(x)\phi_i(t - x)}$$

4.2.1 Estimation of reproduction number

Ground Reality: In Population I, the parameters are β_1, γ and K_1, which are the transmission rate, recovery rate (also specifies the intrinsic generation interval) and the random variable for the number of contacts. The growth rate is λ_1.

$$\lambda_1 = \beta_1 \frac{\langle K_1^2 \rangle}{\langle K_1 \rangle} - 2\beta_1 - \gamma \tag{8}$$

$$\mathcal{R}_0^1 = 1 + \frac{\lambda_1}{\beta_1 + \gamma} \tag{9}$$

Similarly, for Population 2,

$$\lambda_2 = \beta_2 \frac{\langle K_2^2 \rangle}{\langle K_2 \rangle} - 2\beta_2 - \gamma \tag{10}$$

$$\mathcal{R}_0^2 = 1 + \frac{\lambda_2}{\beta_2 + \gamma} \tag{11}$$

The recovery rate is identical for both populations because it defines the intrinsic generation intervals, which by definition, are not affected by changes in contact structure. Using the expression for backward generation intervals, the mean backward generation interval can be calculated when the epidemic is growing exponentially

$$T_{exp,1} = \frac{1}{\lambda_1 + \beta_1 + \gamma} \tag{12}$$

$$T_{exp,2} = \frac{1}{\lambda_2 + \beta_2 + \gamma} \tag{13}$$

The quantities of interest, which are the basic reproduction number and the intrinsic generation interval, can now be expressed in terms of the observable

variables

$$\mathcal{R}_0^1 = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{14}$$

$$\mathcal{R}_0^2 = \frac{1}{1 - \lambda_2 T_{exp,2}} \tag{15}$$

$$\gamma = \frac{\left(\frac{\langle K_1^2 \rangle}{\langle K_1 \rangle} - 2\right)\left(\frac{1}{T_{exp,1}} - \lambda_1\right) - \lambda_1}{\frac{\langle K_1^2 \rangle}{\langle K_1 \rangle} - 1} \tag{16}$$

$$= \frac{\left(\frac{\langle K_2^2 \rangle}{\langle K_2 \rangle} - 2\right)\left(\frac{1}{T_{exp,2}} - \lambda_2\right) - \lambda_2}{\frac{\langle K_2^2 \rangle}{\langle K_2 \rangle} - 1} \tag{17}$$

Case A: The modeller does not know anything about the contact structure and decides to use the simplest model available, the homogeneous and well-mixed SIR scheme to model these epidemics. This model has two parameters, the contact rate β and the recovery rate γ. The recovery rate also specifies the intrinsic generation interval distribution. The exponential growth rate in this model is $\beta - \gamma$ and this rate can be obtained from the incidence.

$$\lambda = \beta_1 - \gamma_1$$

Using the expression for backward generation intervals, under the assumption that incidence grows exponentially with rate λ, the mean backward generation interval, T_{exp} is

$$T_{exp} = \frac{1}{\lambda + \gamma_1}$$

Since, we have access to backward contact tracing, $T_{exp,1}$ is obtained from the data. Using these two equations in conjunction, the parameters β and γ can be estimated, along with the basic reproduction number.

$$\widehat{\beta}_1 = \frac{1}{T_{exp,1}} \tag{18}$$

$$\widehat{\gamma}_1 = \frac{1}{T_{exp,1}} - \lambda_1 = \beta_1 + \gamma \tag{19}$$

$$\widehat{\mathcal{R}_0^1} = 1 + \frac{\lambda_1}{\widehat{\gamma}_1} = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{20}$$

The estimated distribution of intrinsic generation intervals is incorrect, yet the estimate for the basic reproduction number is correct.

By definition, the intrinsic generation interval distribution depends solely on how the infectiousness of an infected person changes with time. It does not depend on contact structure or the availability of susceptible contacts. Therefore, the modeller might see it fit to use the estimated intrinsic interval distribution from Population 1 to estimate the basic reproduction number in Population 2

as they do not have contact tracing capacity there.

$$\widehat{\mathcal{R}_0^2} = 1 + \frac{\lambda_2}{\widehat{\gamma}_1} \tag{21}$$

$$\widehat{\mathcal{R}_0^2} = \frac{1 + (\lambda_2 - \lambda_1)T_{exp,1}}{1 - \lambda_1 T_{exp,1}} \neq \frac{1}{1 - \lambda_2 T_{exp,2}} \tag{22}$$

This estimate of the basic reproduction number is different from the true reproduction number unless the growth rates in the populations are equal $(\lambda_2 = \lambda_1)$, for which there is no a priori justification.

This example can also be framed in a different manner. Say there are two populations which are similar in their contact structure, but differ in their implementation of interventions.

Case B: The modeller knows that the population is well described by a quenched network and knows the average degrees of the two populations, so they assume a quenched homogeneous model. This model has three parameters, the contact rate β, the recovery rate γ and the degree of nodes, k (which is known). The exponential growth rate in this model is $\beta(k-2) - \gamma$ and this rate can be obtained from the incidence.

$$\lambda_1 = \beta_1(k_1 - 2) - \gamma_1$$

Using the expression for backward generation intervals, under the assumption that incidence grows exponentially with rate λ, the mean backward generation interval, $T_{exp,1}$ is

$$T_{exp,1} = \frac{1}{\lambda_1 + \gamma_1 + \beta_1}$$

Since, we have access to backward contact tracing, $T_{exp,1}$ is obtained from the data. Using these two equations in conjunction, γ_1 can be estimated, along with the basic reproduction number.

$$\widehat{\gamma}_1 = \frac{(k_1 - 2)(1/T_{exp} - \lambda) - \lambda}{k_1 - 1} \tag{23}$$

$$\widehat{\mathcal{R}_0^1} = 1 + \frac{\lambda_1}{\widehat{\gamma}_1 + \widehat{\beta}_1} = \frac{1}{1 - \lambda_1 T_{exp,1}} \tag{24}$$

The estimated distribution of intrinsic generation intervals is incorrect, but the estimate for the basic reproduction number is correct. The intrinsic generation interval can be estimated properly only when the degree of the homogeneous model matches the ratio of mean square degree and mean degree from the actual contact network.

As in the previous case, the estimated intrinsic generation intervals would be used to understand the dynamics in Population 2, which would lead to

$$\widehat{\mathcal{R}_0^2} = 1 + \frac{\lambda_2}{\widehat{\beta}_2 + \widehat{\gamma}_1} = \frac{(\lambda_2 + \widehat{\gamma}_1)(k_2 - 1)}{\lambda_2 + \widehat{\gamma}_1(k_2 - 1)} \tag{25}$$

This estimate of the basic reproduction number is different from the actual basic reproduction number in Population 2.

In this example, we do not fit the full time series of mean backward intervals to the models, instead we focus on a section of the time series where the behaviour is analytically tractable. We would guess that fitting the full time series would present more challenges and bias the estimators further.

Insert plot of $\widehat{\mathcal{R}_{0,II}}$ and $\mathcal{R}_{0,II}$ with β_2 on axis to show the estimation error.

4.3 Supplementary Results

Figure S1 *Epidemics and their realised generation intervals for two networks: a heterogeneous annealed network and a homogeneous annealed network. Parameters are selected such that basic reproduction number and intrinsic generation interval distribution are the same for both epidemics. Yet, the realised intervals look different. The heterogeneity clearly decreases the amount of contraction in the mean forward generation interval. This can be explained by look-*

ing at the incidence of stubs, $\pi_i(t)$. In the homogeneous case, the peak is higher than the heterogeneous epidemic. More competition among infectors shortens the generation time more. The starting time, $t = 0$, is selected such that the prevalence is 0.01%, and it can be seen that in the heterogeneous case, the mean generation interval is shorter at the start. This can be explained by the faster fall in the prevalence of susceptible stubs in the population, $\pi_S(t)$. The mean backward generation interval is quite similar at early times, and deviates at a later time. This can be explained by the same early exponential growth rate and different rates of exponential decline

Figure S2 *Epidemics and their realised generation intervals for two networks: a heterogeneous annealed network and the other a homogeneous annealed network, with identical mean degree. Parameters are selected such that the peak incidence and intrinsic generation interval distribution are the same for both epidemics. Yet, the realised intervals look very different.*

Figure S3 *An epidemic and its mean realised generation intervals on an annealed network with truncated power law degree distribution. The realised generation intervals calculated using the equations from [17] and the equations we derived are markedly different. The backward mean generation intervals are identical at the start of the epidemic. The homogeneous assumption underestimates the contraction in forward generation intervals.*

References

[1] Barrat, A., Cattuto, C., Colizza, V., Gesualdo, F., Isella, L., Pandolfi, E., Pinton, J.-F., Ravà, L., Rizzo, C., Romano, M., Stehlé, J., Tozzi, A. E., & Van den Broeck, W. (2013). Empirical temporal networks of face-to-face human interactions. *The European Physical Journal Special Topics*, 222(6), 1295–1309. https://doi.org/10.1140/epjst/e2013-01927-7

[2] Eames, K., Bansal, S., Frost, S., & Riley, S. (2015). Six challenges in measuring contact networks for use in modelling. *Epidemics*, 10, 72–77. https://doi.org/10.1016/j.epidem.2014.08.006

[3] Aleta, A., Martín-Corral, D., Bakker, M. A., Pastore y Piontti, A., Ajelli, M., Litvinova, M., Chinazzi, M., Dean, N. E., Halloran, M. E., Longini, I. M., Pentland, A., Vespignani, A., Moreno, Y., & Moro, E. (2022). Quantifying the importance and location of SARS-CoV-2 transmission events in large metropolitan areas. *Proceedings of the National Academy of Sciences*, 119(26), e2112182119. https://doi.org/10.1073/pnas.2112182119

[4] Pung, R., Russell, T. W., & Kucharski, A. J. (2024). Detecting changes in generation and serial intervals under varying pathogen biology, contact patterns and outbreak response. *PLOS Computational Biology*, 20(3), e1011967. https://doi.org/10.1371/journal.pcbi.1011967

[5] Park, S. W., Sun, K., Abbott, S., Sender, R., Bar-on, Y. M., Weitz, J. S., Funk, S., Grenfell, B. T., Backer, J. A., Wallinga, J., Viboud, C., & Dushoff, J. (2023). Inferring the differences in incubation-period and generation-interval distributions of the Delta and Omicron variants of SARS-CoV-2. *Proceedings of the National Academy of Sciences*, 120(22), e2221887120. https://doi.org/10.1073/pnas.2221887120

[6] Chen, D., Lau, Y.-C., Xu, X.-K., Wang, L., Du, Z., Tsang, T. K., Wu, P., Lau, E. H. Y., Wallinga, J., Cowling, B. J., & Ali, S. T. (2022). Inferring time-varying generation time, serial interval, and incubation period distributions for COVID-19. *Nature Communications*, 13(1), 7727. https://doi.org/10.1038/s41467-022-35496-8

[7] Loedy, N., Wallinga, J., Hens, N., & Torneri, A. (2024). Repetition in social contacts: Implications in modelling the transmission of respiratory infectious diseases in pre-pandemic and pandemic settings. Proceedings of the Royal Society B: Biological Sciences, 291(2027), 20241296. https://doi.org/10.1098/rspb.2024.1296

[8] Pung, R., Firth, J. A., Russell, T. W., Rogers, T., Lee, V. J., & Kucharski, A. J. (2024). Temporal contact patterns and the implications for predicting superspreaders and planning of targeted outbreak control. Journal of The Royal Society Interface, 21(221), 20240358. https://doi.org/10.1098/rsif.2024.0358

[9] Moreno López, J. A., Arregui García, B., Bentkowski, P., Bioglio, L., Pinotti, F., Boëlle, P.-Y., Barrat, A., Colizza, V., & Poletto, C. (2021). Anatomy of digital contact tracing: Role of age, transmission setting, adoption and case detection. Science Advances, 7, eabd8750. https://doi.org/10.1126/sciadv.abd8750

[10] Calmon, L., Colosi, E., Bassignana, G., Barrat, A., & Colizza, V. (2024). Preserving friendships in school contacts: An algorithm to construct synthetic temporal networks for epidemic modelling. PLOS Computational Biology, 20(12), e1012661. https://doi.org/10.1371/journal.pcbi.1012661

[11] Park, S. W., Champredon, D., & Dushoff, J. (2020). Inferring generation-interval distributions from contact-tracing data. Journal of The Royal Society Interface, 17(167), 20190719. https://doi.org/10.1098/rsif.2019.0719

[12] Svensson, Å. (2007). A note on generation times in epidemic models. Mathematical Biosciences, 208(1), 300–311. https://doi.org/10.1016/j.mbs.2006.10.010

[13] Kraemer, M. U. G., Pybus, O. G., Fraser, C., Cauchemez, S., Rambaut, A., & Cowling, B. J. (2021). Monitoring key epidemiological parameters of SARS-CoV-2 transmission. Nature Medicine, 27(11), 1854–1855. https://doi.org/10.1038/s41591-021-01545-w

[14] Manica, M., De Bellis, A., Guzzetta, G., Mancuso, P., Vicentini, M., Venturelli, F., Zerbini, A., Bisaccia, E., Litvinova, M., Menegale, F., Grané, C. M., Poletti, P., Marziano, V., Zardini, A., d'Andrea, V., Trentini, F., Bella, A., Riccardo, F., Pezzotti, P., Ajelli, M., Rossi, P. G., & Merler, S. (2022). Intrinsic generation time of the SARS-CoV-2 Omicron variant: An observational study of household transmission. The Lancet Regional Health – Europe, 19, 100446. https://doi.org/10.1016/j.lanepe.2022.100446

[15] Wallinga, J., & Lipsitch, M. (2007). How generation intervals shape the relationship between growth rates and reproductive numbers. Proceedings of the Royal Society B: Biological Sciences, 274(1609), 599–604. https://doi.org/10.1098/rspb.2006.3754

[16] Kenah, E., Lipsitch, M., & Robins, J. M. (2008). Generation interval contraction and epidemic data analysis. Mathematical Biosciences, 213(1), 71–79. https://doi.org/10.1016/j.mbs.2008.02.007

[17] Champredon, D., & Dushoff, J. (2015). Intrinsic and realized generation intervals in infectious-disease transmission. Proceedings of the Royal Society B: Biological Sciences, 282(1821), 20152026. https://doi.org/10.1098/rspb.2015.2026

[18] Miller, J. C., Slim, A. C., & Volz, E. M. (2011). Edge-based compartmental modelling for infectious disease spread. Journal of The Royal Society Interface, 9(70), 890–906. https://doi.org/10.1098/rsif.2011.0403

[19] Liu, Q.-H., Ajelli, M., Aleta, A., Merler, S., Moreno, Y., & Vespignani, A. (2018). Measurability of the epidemic reproduction number in data-driven contact networks. Proceedings of the National Academy of Sciences, 115(50), 12680–12685. https://doi.org/10.1073/pnas.1811115115

[20] Trapman, P., Ball, F., Dhersin, J.-S., Tran, V. C., Wallinga, J., & Britton, T. (2016). Inferring R 0 in emerging epidemics—The effect of common population structure is small. Journal of The Royal Society Interface, 13(121), 20160288. https://doi.org/10.1098/rsif.2016.0288

[21] Kiss, I. Z., Miller, J. C., & Simon, P. L. (2017). Mathematics of Epidemics on Networks: From Exact to Approximate Models (Vol. 46). Springer International Publishing. https://doi.org/10.1007/978-3-319-50806-1

[22] Saldaña, F., Daza-Torres, M. L., & Aguiar, M. (2023). Data-driven estimation of the instantaneous reproduction number and growth rates for the 2022 monkeypox outbreak in Europe. PLOS ONE, 18(9), e0290387. https://doi.org/10.1371/journal.pone.0290387

[23] Díaz-Brochero, C., & Cucunubá, Z. M. (2024). Epidemiological findings, estimates of the instantaneous reproduction number, and control strategies of the first Mpox outbreak in Latin America. Travel Medicine and Infectious Disease, 59, 102701. https://doi.org/10.1016/j.tmaid.2024.102701

[24] Shim, E., Tariq, A., Choi, W., Lee, Y., & Chowell, G. (2020). Transmission potential and severity of COVID-19 in South Korea. International Journal of Infectious Diseases, 93, 339–344. https://doi.org/10.1016/j.ijid.2020.03.031

[25] Rocklöv, J., Sjödin, H., & Wilder-Smith, A. (2020). COVID-19 outbreak on the Diamond Princess cruise ship: Estimating the epidemic potential and effectiveness of public health countermeasures. Journal of Travel Medicine, 27(3), taaa030. https://doi.org/10.1093/jtm/taaa030

[26] Munayco, C. V., Tariq, A., Rothenberg, R., Soto-Cabezas, G. G., Reyes, M. F., Valle, A., Rojas-Mezarina, L., Cabezas, C., Loayza, M., Chowell, G., Garro, D. C., Vasquez, K. M., Castro, E. S., Ordinola, I. S., Mimbela, J. M., Cornejo, K. M., Quijano, F. C., La Torre Rosillo, L., Ibarguen, L. O., . . . Ramos, W. (2020). Early transmission dynamics of COVID-19 in a southern hemisphere setting: Lima-Peru: February 29th–March 30th, 2020. Infectious Disease Modelling, 5, 338–345. https://doi.org/10.1016/j.idm.2020.05.001

[27] Salvatore, M., Basu, D., Ray, D., Kleinsasser, M., Purkayastha, S., Bhattacharyya, R., & Mukherjee, B. (2020). Comprehensive public health evaluation of lockdown as a non-pharmaceutical intervention on COVID-19 spread in India: National trends masking state-level variations. BMJ Open, 10(12), e041778. https://doi.org/10.1136/bmjopen-2020-041778

[28] Gostic, K. M., McGough, L., Baskerville, E. B., Abbott, S., Joshi, K., Tedijanto, C., Kahn, R., Niehus, R., Hay, J. A., De Salazar, P. M., Hellewell, J., Meakin, S., Munday, J. D., Bosse, N. I., Sherrat, K., Thompson, R. N., White, L. F., Huisman, J. S., Scire, J., . . . Cobey, S. (2020). Practical considerations for measuring the effective reproductive number, Rt. PLOS Computational Biology, 16(12), e1008409. https://doi.org/10.1371/journal.pcbi.1008409

[29] Fraser, C. (2007). Estimating Individual and Household Reproduction Numbers in an Emerging Epidemic. PLoS ONE, 2(8), e758. https://doi.org/10.1371/journal.pone.0000758