Title: Low Pass Whole Genome sequencing - a cost-effective new technique for the

identification of aneuploidies and copy number variations: a study of 1372 clinical

samples in an Indian cohort

Authors: Satish Kariyaiah[1], Rashmi Rasalkar[1], Meenakshi Bhat[2], Avinash Pradhan[1],

Deepika Rani Jena[1], Smitha C[1], Manjunath V[1], Lenika A[1], SriGanesh Jammula, Monisha

Morris[2], Ravi Gupta[1], Eswarachari Venkataswamy[1] Sakthivel Murugan S.M.[1], Ramprasad

V.L.[1] Priya Kadam[1]

Satish Kariyaiah and Rashmi Rasalkar are joint first authors

Affiliations: 1. Medgenome Labs Ltd, Bangalore, 2. Centre for Human Genetics, Bangalore

Conflict of Interest: No conflict of interest has been declared

Keywords: Low pass whole genome sequencing, copy number variations, aneuploidies,

prenatal genetic testing, postnatal genetic testing

Abstract

Background: Identification and confirmation of copy number variation is an important
aspect of genetic testing for several prenatal scenarios, such as abnormal maternal serum
screening, abnormal ultrasound findings, high risk results on non-invasive prenatal
screening as well as postnatal settings such as developmental delay, intellectual disabilities,
congenital anomalies and dysmorphism as well as in couples with recurrent miscarriages.
Low pass whole genome sequencing (lpWGS) followed by copy number variation (CNV)
analysis has shown identification of critical pathogenic/likely pathogenic variants
comprehensively with great accuracy and cost-effectiveness. We validated this technique
on retrospective routine clinical samples previously assessed by an orthogonal tests, either
chromosomal microarray or karyotyping or fluorescence in situ hybridization in a diagnostic
laboratory and evaluated its performance.

Material and Methods: The validation included 112 clinical samples which showed the
following events: 14 aneuploidies, 70 copy number variations of size range (50 kb - 116 Mb),
11 mosaics aneuplodies, 4 mosaics CNVs and 4 triploidies. These validation samples were
derived from 21 fetuses (6 chorionic villus biopsies and 15 amniotic fluid samples), 25
abortuses and 66 postnatal samples. The validation was performed at two different
resolutions - 1Mb and 50Kb. We then assessed the overall yield of the lpWGS assay on
1,260 clinical samples that comprised 801 fetuses, 346 products of conception/intrauterine
fetal demise/stillbirth and 113 postnatal cases.

Results: We obtained 100% concordance in detecting full aneuploidies and copy number
variations including mosaics and triploidy. The overall success rate of this assay that passed
all quality parameters was 99.8%. (1235/1238). The overall diagnostic yield for pathogenic
and likely pathogenic variants was 13.6% (168/1235), and specifically 7.3%, 28.8%, 15% for
fetuses, products of conception/intrauterine fetal demise/stillbirth and post natal samples
respectively. An additional 2.9% (36/1235) of overall samples showed CNVs that were
classified as variants of unknown significance.

Conclusion: Overall, low pass whole genome sequencing is a sensitive and robust technique
for identification of aneuploidies and copy number variations in a clinical setting.

Introduction

Chromosomal abnormalities in the form of aneuploidies and copy number variations (CNVs) as a likely cause of a specific phenotype are important in both prenatal and postnatal clinical settings Different techniques with increasing resolution and progressive technological advances have been employed over the last few decades in the identification of chromosome aneuploidies and CNVs. These include karyotyping which detectsCNVs with resolution of approximately 5Mb or more aneuploidies, polyploidies, balanced and unbalanced structural rearrangements, , and mosaicism[1], . Fluorescence in situ hybridization (*FISH*)[2], Multiplex ligation probe amplification (MLPA)[3], quantitative polymerase chain reaction[4] are targeted approaches for specific abnormalities. Chromosome arrays such as comparative genomic hybridization and single nucleotide polymorphism based array[5] have enabled analysis of a wider spectrum of abnormalities and with increased resolution. Chromosomal microarray had been established as the gold standard for identification of CNVs for structurally abnormal foetuses, stillbirths, products of conception and for children with intellectual disability and developmental delay by professional societies.[6,7] Overall, they have a proven greater yield of about 5-15% as compared to karyotyping[8]. However, these array techniques are chip- based and probe dependent, are expensive and have less comprehensive coverage. Emerging studies show that next generation sequencing with low coverage are accurate at CNV detection, low cost and with comprehensive coverage.[9,10]

Low-pass whole genome sequencing (lpWGS) as a method to identify aneuploidies and CNVs is relatively new but the utility has been clearly demonstrated in both prenatal and postnatal scenarios.[11,12,13] LpWGS refers to shallow whole genome sequencing (WGS) at a lower depth, typically around 0.5X to 5X coverage, as compared to the standard depth of 30X to 50X in traditional WGS. This reduction in coverage significantly decreases the cost and time required for analysis while still providing valuable genomic information. Although CMA is currently the first-line investigation for aneuploidies and CNV identificaton, the lpWGS technique developed on different NGS platforms along with sensitive CNV calling algorithms provides an alternative method to detect aneuploidies and CNVs accurately[11]. An increased yield with lpWGS has been demonstrated in a direct comparison between lpWGS

and chromosomal microarray in some studies.[14] Some of the advantages include lower
initial input DNA compared to CMA protocols, non-probe dependency, wider coverage of
genome, use of existing NGS equipment/platforms, lower costs and manpower
requirements for both wet lab and data-analysis procedures, and high throughput with a
shortened turnaround time.[12,13] A Chinese joint expert group in 2019 recommended the use
of lpWGS as first tier diagnostic testing in pregnant women for prenatal diagnosis, especially
couples with balanced chromosomal translocations, and/or with pregnancy loss.[15]
Consequently, this technique is being increasingly used in clinical practice in tertiary level
Chinese hospitals.[16] Recently updated guidelines by ACMG on the use of NGS techniques for
the identification of structural variants state that lpWGS provides an affordable CNV
detection method with a similar diagnostic yield and probable enhanced resolution than
microarrays.[17]

This study was aimed at validating the lpWGS technique against a standard tests and
evaluating its performance for use in detection of aneuploidies and CNVs. After the
validation phase, the study aimed at evaluating the diagnostic yield in 1260 clinical samples.

Materials and Methods

Validation cohort

This validation was performed in two phases to assess the ability of lpWGS to identifying
aneuploidies and CNVs. In the first phase CNVs greater than 1Mb were validated and in the
second phase we assessed the sensitivity of lpWG in detecting CNVs greater than 50Kb and
in triploidy identification. Samples with chromosomal mosiacism were included in both
phases.

Phase I (0.5X - 0.7X): We used 85 consented retrospective clinical samples with known
diagnostic results that included abberations ranging in size from 1.34 to 133 Mb. These
included 15 prenatal samples (4 chorionic villus biopsies, 11 amniotic fluid), 17 products of
conception and 53 postnatal samples (Figure 1). Of the total, 53 samples were previously
investigated by chromosomal microarray (CMA) while 11 were investigated by karyotyping
(KT), 15 by karyotyping and FISH for microdeletion syndromes, and 6 others were
investigated only by FISH for specific microdeletion syndromes, particularly 22q11.2
microdeletion (Table 1). We created a panel of normal reference using 68 clinical samples
with no known clinically relevant CNVs as seen on Affymetrix CytoScan™ Optima array. We
evaluated these 10 mosaic samples (Table 1, mosaics), of which consisted of 7 were
aneuploidy events and four CNVs with mosaic percentages ranging from 15 - 40% in phase 1
(Figure 1).
108
Phase II (3-5X Depth): To increase the resolution for identifying smaller CNVs (up to ~50Kb)
as well as mosaicism and triploidy, which require greater sequencing depth to capture
smaller variants, a total of 27 consented retrospective clincial samples were included (6
prenatal samples, 8 products of conception, and 13 postnatal samples). All these samples
were previously investigated by either Affymetrix CytoScan™ Optima array, CytoScan™ 750K
Array or CMA CytoScan™ XON Array and had CNVs of size range less than 1Mb and greater
than 50 Kb (range 47 - 660 Kb) (Table 1, Phase II, Figure 1).   To build a reference set, 42
samples without CNVs on CytoScan™ 750K Array were included.

We also included four triploidy samples, identified by MCC testing (16 STR markers/QF-PCR
for aneuploidy) and CMA by a inhouse triploidy detection algorithm in phase II**.** We also
included four mosaic aneuploidies with mosaic percentages from 30 - 65% (Table 1).

Library preparation and sequencing

DNA was extracted using both automated (QiaSymphony) and manual methods. QC-passed
samples were used for whole genome library preparation at a minimum of 50ng on the
Agilent Bravo platform using commercially available library prep kits. Prepared libraries
were sequenced on Illumina platforms (NovaSeq 6000 or X Plus). Around 2Gb data per
sample was generated in Phase I (~ 1 Mb resolution) and 15 Gb data for the samples used
for ~50 Kb resolution in Phase 2. The sequencing results generated ~16 million paired-end

reads of 150 bp per sample, corresponding to a 0.5X - 0.7X for >1Mb size and 3X - 5X

coverage for >50 Kb size.

Bioinformatics analysis and variant classification

The raw paired-end FASTQ files were processed using Cutadapt to remove adapter

sequences, reads shorter than 20 bp or with a quality score below 10 phred score. CNV

analysis was performed using both WisecondorX and Illumina's DRAGEN v4.2. WisecondorX

was employed solely for validation purposes.

For WisecondorX analysis, the trimmed reads were aligned to the GRCh38 reference

genome using the Sentieon BWA aligner (sentieon-genomics-201808). Coverage at a

defined bin size was computed from the aligned BAM files for each sample and converted

into a WisecondorX compatible NPZ format. All reference NPZ files were subsequently

merged to create a final reference NPZ file. Segmentation was then performed based on

reference ploidy, grouping bins with similar ploidy levels. Segments showing ploidy

deviations from the reference were identified and classified as copy number variations.

For analysis using DRAGEN, trimmed reads were directly employed for alignment and

CNV detection. The reads were aligned to the GRCh38 reference genome. CNV calling was

performed through a comparative coverage-based approach, where coverage across

predefined bin sizes was calculated. Subsequently, the data underwent bias correction,

addressing GC-content bias, resulting in the generation of a corrected target counts file.

Normalization was then carried out ensuring that all reference files were also GC-corrected

if the sample had undergone GC bias correction. Once normalized, segmentation of the data

was performed based on the processed signal. This was followed by CNV detection, where

any segment displaying a significant deviation from the normal ploidy level—established

using the reference—was classified as a CNV event.

To identify larger events, a bin size of 200 Kb was used, with analysis performed

against a model based on 68 reference samples. Similarly, for identifying events larger than

50 Kb, a finer bin size of 20 Kb was applied, utilizing a model constructed from 42 reference

samples.

Triploidy was identified by evaluating the proportion of mutant reads in heterozygous
SNPs.[18] To perform this analysis, high-quality heterozygous SNPs from the 1,000 Genomes
Project were utilized. For each site with sufficient depth, allele frequencies were calculated
based on reference and alternate bases across all samples. In triploidy, the allele frequency
at heterozygous positions is expected to be around 0.33 or 0.66, indicating a genetic gain.
The variants were then categorized into three distinct groups according to their copy
number states: 0.33 (with a range of 0.28–0.38), 0.66 (range of 0.62–0.72), and 0.5 (range of
0.45–0.55). Following this, the ratio between the number of sites with 0.33 and 0.66
frequencies, relative to those with 0.5, was computed. Upon comparing these ratios
between control and positive samples, a clear distinction emerged. Samples exhibiting a
ratio greater than 2.4 were classified as positive for triploidy.

Annotation and classification of all CNVs was performed using an in-house workflow
developed based on the ACMG and Clinical Genome Resource (ClinGen) guidelines.

Diagnostic yield of serial clinical samples by lpWGS in a clinical setting

We evaluated the assay yield from 1,260 clinically consented samples (prenatal - 801,
POC/Intrauterine deaths/stillbirths - 346, postnatal - 113). Demographic details such as
maternal age, gestational age and clinical indications for testing were recorded. All prenatal
samples were subjected to maternal cell contamination (MCC) analysis either directly or
combined with aneuploidy testing through quantitative fluorescence polymerase chain
reaction (QF-PCR). MCC and QF-PCR analysis also identified presence of triploidy. Briefly,
the Devyser Compact v3 QF-PCR kit was used in accordance with the manufacturer's
protocol. All PCR products were genotyped using a SeqStudio Genetic Analyzer (Thermo
Fisher Scientific) with GeneMappe software 6 (Thermo Fisher Scientific). Only MCC negative
samples were processed further. These samples were evaluated after the first phase of
validation and therefore the CNV resolution was limited to 1Mb.

Results

We validated the data in two phases, by evaluating the performance of lpWGS 112 samples

to detect aneuploidies and CNVs against previous orthogonal data from CMA, karyotype

and FISH. Then, we analysed the performace in 1260 consecutive clinical samples.

Validation results

Phase I – Identification of large CNV

All aneuploidies and CNVs greater than 1Mb, identified by orthogonal tests were identified by

lpWGS (Table 1).

lpWGS concordance with CMA - All 13 aneuploidies and 37 CNVs identified by previous testing of 53

samples by chrosomsomal microarray were also identified by lpWGS (Table 1). The CNV size ranged

from 1.22 – 116.51 Mb. There was a high degree of concordance between the event size detected

by CMA and lpWGS as seen in the scatter plot (Figure 2, I). The six samples that did not show CNVs

on CMA did not show any CNVs on lpWGS either. To evaluate reproducibility, we re-processed 13

samples three times, all of which showed full concordance (Supplementary Table 1).

lpWGS concordance with Karyotyping data – Of eleven samples initially tested by karyotyping, two

abnormal events, one each in V10 and V18 had been identified by karyotype. LpWGS identified and

further characterized these two events (Table 1), first, V10, a mosaic karyotype involving derivative

of Y chromosome was recognized as four copies of Chromosome Y, second, V18, karyotype had

shown 46,XY,?der(15), lpWGS revealed a duplication of 25.4 Mb at 4p16.3-p15.2, identifying the

probable chromosomal content within the derivative chromosome. Additionally, lpWGS identified a

CNV, duplication of size 1.7 Mb at cytoband region 17q11.2, in sample V39, where the karyotype

had been reported as normal. However, this was not expected to be detected on karyotype since it

was below the detection limit. These three events were further confirmed by chromosomal

microarray (Table 1). The remaining eight samples that had not show any abnormality on karyotype

with a detection limit of 5Mb, did not show any CNV on lpWGS of size greater than 5Mb, either.

lpWGS concordance with Karyotyping and FISH for 22q11.2 microdeletion data – Of the sixteen

samples previously evaluated by both karyotyping and FISH for 22q11.2 microdeletion syndrome, 3

samples, V24, V17 and V31, had shown abnormalities (Table 1). LpWGS assay identified 5 CNVs

among these three samples. Three CNVs were seen in one sample alone (V24). This case had a

karyotype of 46,XY,der(5)t(5;?)(p15;?). lpWGS identified, a deletion at 5p15.33-p14.2 and

duplications at 8q23.2-q24.3 and 6q16.3, with sizes of 23.3 Mb, 34.3 Mb, and 2 Mb,

respectively. Parental karyotyping was recommended due to suspected structural variation,
however, no further information was available as the case was lost to follow-up. These
events were further confirmed by running a CMA (Table1).

Second, sample V17 had shown a karyotype of 46,XY,der(6) and no deletion identified by FISH
probe 22q11.2 deletion or for 7q deletion. However, lpWGS identified a 36.5 Mb duplication at
cytoband 3q25.32-q29, clarifying the chromosomal component of the derivative chromosome while
no variant was seen in q arm of either chromosome 7 or 22. Lastly, in V31, where karyotype had
identified an deletion, 46,XY,del(10)p13) and where FISH was negative for 22q11.2, lpWGS identified
the precise breakpoints and size, at cytoband region 10p15.3-p14 and size of 11.4Mb respectively.
Both findings of lpWGS were again confirmed on CMA (Table 1). In the 13 remaining samples, where
no event was seen by karyotyping and FISH, lpWGS also did not identify any CNV.

lpWGS concordance by FISH alone - We evaluated 5 samples by lpWGS that were previously
evaluated by FISH, four by FISH for 22q11.2 deletion, and one for 15q11 deletion for Angelman
syndrome alone, based on clinical presentation. Two samples that had shown the 22q11.2
microdeletion by FISH were also identified by lpWGS where microdeletions in cytoband region
22q11.21 of sizes 4.4 (V44) and 2.6 Mb (V45) were identified. For a sample investigated by FISH for
Angelman syndrome, V37, no variation was detected on chromosome 15; however, lpWGS
identified a 1.4 Mb deletion at 17p11.2. The events seen in V37 and V45 was also confirmed by
chromosomal microarray (Table 1). However V44 could not be confirmed as the sample was
exhausted.

In two samples, that did not show any deletion by FISH for 22q11.2 deletion, showed two
CNVs in V15, a terminal gain and loss, at cytoband region 9q32-q34.3 and 3p26.3 - p26. of size 24
and 4.5 Mb respectively and in the other sample, V20, one CNV, a duplication at 4q35.1-q35.2 of
size 2.2Mb was detected. The events in V15 and V20 were confirmed on CMA (Table 1)

Concordance of mosaic events – Of the 10 mosaic samples analyzed in Phase I, the seven
mosaic aneuploidies that showed mosaicism between 15% - 35% demonstrated on CMA
data showed concordance with lpWGS in terms of relative size and coordinates of
chromosomal events. Sample V51, had exhibited two distinct events—a gain and a loss on
the X chromosome measuring 70.4 Mb and 76.5 Mb, respectively—showed variability in
size when analyzed by lpWGS, with the gain measured at 41 Mb and the loss at 9.2 Mb. Of

the mosaic CNVs, in three samples that showed about 40% mosaicism, of size range 15.9 –

78.6Mb were identified by lpWGS.

Overall, when CMA, karyotype, and FISH results were used as the reference, lpWGS

demonstrated, considering at least one CNV calling algorithm, 100% sensitivity for detecting

CNVs previously identified, highlighting the high consistency and robust performance of this

NGS platform.

Phase II – Identification of smaller CNVs, triploidy, mosaics

Of the 27 samples analyzed in Phase 2, all variations identified by CMA were successfully

detected by lpWGS. The DRAGEN^{TM} software, which demonstrated the highest

concordance with CMA data, was selected for subsequent analysis. The detected CNV size

ranged from 70 kb – 880 kb.

Identification of mosaic aneuploidies and CNVs In Phase 2, a different set of mosaic

aneuploid samples was analyzed using higher depth. The samples exhibited 30–62%

mosaicism, were readily identifiable by both CNV calling tools. The diverse sample set and

varied processing conditions allowed for an in-depth assessment of the method's accuracy

and consistency across multiple scenarios.

*Identification of Triploid*y -   All four cases exhibited a significantly higher ratio compared to the

controls and were detectable as Triploidy. One sample was reanalyzed for reproducibility, yielding

fully concordant results (Figure 2h).

Clinical Experience with 1260 Samples

Demographics and Indications for Testing

The demographic data and diagnostic yield are summarized in Table 2, while clinical

indications grouped into broad categories are presented in Figure 3. For the prenatal

samples, the most frequent indication for invasive testing was high risk on maternal serum

screening (45.4%), followed by ultrasound abnormalities (36.8%). In products of conception

(POCs), the leading indications were spontaneous abortion, missed abortion, or intrauterine

death (IUD) (42%) followed by ultrasound abnormalities (33.5%). In postnatal
cases, dysmorphism and congenital abnormalities being the most common.

Assay Success Rate and Technical Performance

Result by lpWGS assay was obtained in 99.8% (1235/1238) and therefore failure rate was
only 0.2%. Results were obtained in all postnatal samples, with assay failure observed in
only three prenatal samples (1 amniotic fluid sample and 2 POCs). The call rate and
distribution of normal and abnormal samples by category is shown in Figure 4.

A total of 22 samples (21 POCs and 1 CVS) exhibited identical maternal and fetal
genotypes (Figure 4). No fetal genotype was identified by maternal cell contamination
(MCC) or quantitative fluorescence PCR (QF-PCR) which might have been caused by
improper sample collection. These samples were excluded from further analysis

lpWGS Concordance with common aneuplodies identified by QF-PCR

Of the 1260 samples, 1147 were prenatal, of which 650 were booked for rapid aneuploidy
screening by QF-PCR. Of these, 51/639 (7.9%) aneuploides in chromosomes 21,18,13, and
sex chrosomsome aneuplodies were identified by QF-PCR which were also detected by
lpWGS. Also in 588/639 (92%) samples where QF-PCR did not pick up any anomaly in these
chromosomes showed normal two copies by lpWGS as well, giving 100% concordance for
common aneuploidy detection. Additional aneuplodies in chromosomes 3, 14, 16, 22 were
identified on lpWGS. The remaning 11 samples showed identical genotype to the mother
indicating absence of fetal DNA in the POC tissue and were therefore exluded from further
testing by lpWGS.

Diagnostic Yield

Of the 1,235 samples with identified results (204/1235, 16.5%) 131 were aneuploidies and
73 significant CNVs (Table 3). Of these 37 were pathogenic/likely pathogenic copy number
variations (CNVs) and 36 were classified as variants of uncertain significance (VOUS). The
overall diagnostic yield for aneuplodies and pathogenic/likely pathogenic CNVs was 13.6%,
specifically, x7.3% in fetuses, 28.8% in early miscarriages, intrauterine deaths, and

stillbirths, 15% in postnatal cases (Table 2). Variants classified as VOUS accounted

for 2.9% of all samples: specifically 2.6% in prenatal cases, 2.5% in POCs/IUDs/stillbirths,

6.2% in postnatal cases (Supplementary Table 2).

Aneuploidies - Aneuploidies accounted for 78% (131/168) of abnormalities. As expected,

common aneuploidies involving chromosomes 21, 18, 13, X, and Y were predominantly

identified in: 82.7% (48/58) of fetuses and 41.9% (39/93) of POCs/IUDs/stillbirths (Figure 2a-

c, Table 3, Figure 5). Rare autosomal aneuploidies accounted for 32.3% (30/93) and were

observed exclusively in POCs. Similarly, triploidies (12.9%) were identified only in POCs, as

expected.

Pathogenic/Likely pathogenic CNVs - Smaller CNVs constituted 22% (37/168) of all cases.

Smaller CNVs were detected in 12.9% (12/93) of POCs/IUDs/stillbirths (Figure 5). In

postnatal samples, pathogenic CNVs were more common (88.2%, 15/17), while common

aneuploidies were observed in only 12% of cases (Figure 5).. The pathogenic CNVs ranged in

size from 1 Mb to 81.4 Mb. Of these: 8.3% (3/36) involved two variants per sample, while

the remaining cases each showed a single variant. A total of 11.9% (20/168) of variants

were <5 Mb, sizes that would have been missed by conventional karyotyping. One sample

exhibited terminal gains and losses, suggesting a balanced translocation in the parents;

parental testing was recommended, however this case was lost to follow-up.

Identification of common microdeletion syndromes – Multiple well known microdeletion

syndromes were readily identified, including: 22q11.2 deletion syndrome (T11, T32, T35),

Angelman/Prader-Willi syndrome (T24, T27), Williams syndrome (T8,T13,T31,T36), Cri-du-

chat syndrome (T30), Jacobsen syndrome (T22,T33) These findings are detailed in (Figure

2d-g, Table 4).

Variant of uncertain significance (VOUS) - The size of the 36 VOUS ranged from 1.05 Mb to

9.9 Mb (Supplementary Table S2). Parental testing was conducted in 15 parents (5 couples),

identifying the variant in 2 parents, resulting in an inheritance rate of 40% (2/5 couples).

This additional information facilitated further refinement of variant classification as well as

genetic counselling for the family.

Discussion

We present the first large study on Indian clinical samples demonstrating the clinical-grade diagnostic utility of low-pass whole genome sequencing (lpWGS) for identification of aneuploidies and copy number variations (CNVs) in prenatal and postnatal samples. Our study achieved 100% detection rate of aneuploidies and CNVs previously detected by existing gold-standard technique such as chromosomal microarray (CMA). Additionally, we were able to identify CNVs below the resolution threshold of karyotyping, highlighting its superior resolution. Furthermore, we also implemented a method which led to successful identification of the chromosomal origin of imbalances in abnormal karyotypes, particularly in cases involving derivative chromosomes.

The rigorous validation process was divided into two-phases. The first phase was designed for detecting CNVs ≥1 Mb. Encouraged after obtaining 100% concordance we increased the resolution of LOD for CNV to 50 Kb in the second phase. This approach aligns with previous studies that reported reliable CNV detection at resolutions of 50–100 kb.[12,13] Importantly, our data showed no difference in resolution for CNV detection between prenatal and postnatal samples, underscoring the robustness of lpWGS across diverse sample types.

The depth of sequencing (0.5–0.7X) employed in our study was optimized for cost-effectiveness and scalability while maintaining high diagnostic accuracy. At this shallow sequencing depth, lpWGS effectively detected whole chromosome aneuploidies and small to large structural abnormalities. Similar studies have demonstrated the reliability of lpWGS at depths as low as 0.25X (Dong et al., 2017)[11], although depths below 0.5X may compromise the identification of triploidy and absence of heterozygosity.[10,19] By increasing the sequencing depth to 3-4X in phase two, we successfully identified triploidies using a SNP-based algorithm, further enhancing the clinical utility of lpWGS.

Comparisons between CNV data derived from microarrays and lpWGS showed high concordance in the estimated sizes of chromosome imbalances, with minor discrepancies in terms of chromosomal location attributable to differences in probe spacing across platforms. At a resolution of 1 Mb, our overall diagnostic yield for pathogenic/likely

pathogenic variants was 13.6%, comparable to the yield reported by Wang et al.[13] By
including VOUS, the yield increased to 16.3%, consistent with yields reported in prior
studies.[10,11,12] Diagnostic yields vary widely (2.83–53%) depending on sequencing depth,
resolution, and sample types, with the highest yields observed in POCs.

In line with expectations, aneuploidies were the predominant abnormalities detected
in fetuses (82.7%) and POC/stillbirth samples (41.9%), while smaller CNVs were more
prevalent in postnatal samples (88.2%). The technique reliably identified well-
known microdeletion syndromes, such as 22q11.2 deletion, Cri-du-chat syndrome,
Angelman/Prader-Willi syndrome, 1p36 deletion syndrome, and Wolf-Hirschhorn
syndrome.

We observed a VOUS rate of 2.9%, primarily in postnatal samples. It has been previously
suggested that lpWGS tends to detect more VOUS compared to CMA due to improved
resolution and uniform genome-wide coverage.[13] Previous studies have reported VOUS
rates ranging from 4–5.8% even by CMA. Therefore the rate observed in our study of 1260
samples by lpWGS was within the range reported in literature.[20] As with other genetic tests,
VOUS detection poses challenges for genetic counseling, particularly in prenatal settings.
Guidelines recommend careful interpretation with parental segregation testing, cautious
phenotypic correlation, and avoidance of reproductive decisions based solely on VOUS
findings.[21] Conversly, lpWGS may also identify novel clinically relevant CNVs that are
undetectable by CMA.

Detection of mosaicism presents another challenge in clinical diagnostics. Mosaic
aneuploidies are kown to occur in <1% of prenatal cases.[22] Our data showed reliable
mosaicism detection at levels 15%, however size variability was seen in one sample. That
sample probably had a complex karyotype involving chromosome X. Previous studies have
demonstrated mosaicism detection down to 5%.[9] Mosaicism detection accuracy remains
influenced by platform variability, sample quality, and maternal cell contamination. Clinical
interpretation of mosaicism, particularly at low levels (15–20%), requires caution, as it may
lead to diagnostic uncertainty and challenges in genetic counselling.[23]

Apart from the accuracy, our study also demonstrated that lpWGS has an extremely low

assay failure rate of 0.2%, significantly lower than that reported for the CMA (~4.6%).[10] This

low failure rate is particularly advantageous in prenatal settings, where re-sampling is often

impractical. Furthermore, lpWGS requires low input DNA (25–50 ng) compared to CMA (300

ng), making it suitable for resource-limited scenarios. Cost-effectiveness is another major

advantage of lpWGS assay. With lpWGS the cost of testing is reduced by 50% as compared

to CMA, our own calculations showed a 40-50% reduction in cost compared to low

resolution microarray.[10] As sequencing costs continue to decline, lpWGS is expected to

become increasingly accessible, particularly in low- and middle-income countries.[24]

One of the limitations of this study was that we did not evaluate additional rare variants

seen on lpWGS. Literature suggests that this technique may identify novel variants

undetectatble by microarrays and detected by lpWGS. Additionally, in phase two, we

restricted our scope to triploidies and did not analyze regions of homozygosity/uniparental

disomy, these are planned for subsequent phases at higher sequencing depth and

resolution.

Conclusion

In summary, lpWGS is a robust, sensitive, and scalable method for genome-wide

detection of aneuploidies and pathogenic CNVs. It's superior resolution, cost-effectiveness,

and ability to detect a wide range of abnormalities makes it an excellent alternative to CMA,

particularly in resource-limited clinical settings.

References

1. Smeets DF. Historical perspective of human cytogenetics: from microscope to

microarray. Clin Biochem. 2004;37:439-446.

2. Jia CW, Wang SY, Ma YM, Lan YL, Si YM, Yu L, Zhou LY. Fluorescence in situ

hybridization in uncultured amniocytes for detection of aneuploidy in 4210 prenatal

cases. Chin Med J (Engl). 2011;124:1164-1168.

3. Van Opstal D, Boter M, de Jong D, van den Berg C, Bruggenwirth HT, Wildschut HI, et

al. Rapid aneuploidy detection with multiplex ligation-dependent probe

amplification: a prospective study of 4000 amniotic fluid samples. Eur J Hum Genet.
2009;17:112-121.

4. Mann K, Hills A, Donaghue C, Thomas H, Ogilvie CM. Quantitative fluorescence PCR
analysis of >40,000 prenatal samples for the rapid diagnosis of trisomies 13, 18, 21
and monosomy X. Prenat Diagn. 2012;32:1197-1204.

5. Schaaf CP, Wiszniewska J, Beaudet AL. Copy number and SNP arrays in clinical
diagnostics. Annu Rev Genomics Hum Genet. 2011;12:25-51.

6. South S, Lee C, Lamb A, et al. ACMG Standards and Guidelines for constitutional
cytogenomic microarray analysis, including postnatal and prenatal applications:
revision 2013. Genet Med. 2013;15:901-909. doi:10.1038/gim.2013.129.

7. Miller DT, Adam MP, Aradhya S, Biesecker LG, Brothman AR, Carter NP, et al.
Consensus statement: Chromosomal microarray is a first-tier clinical diagnostic test
for individuals with developmental disabilities or congenital anomalies. Am J Hum
Genet. 2010;86:749-764.

8. Wapner RJ, Martin CL, Levy B, Ballif BC, Eng CM, Zachary JM, et al. Chromosomal
microarray versus karyotyping for prenatal diagnosis. N Engl J Med. 2012;367:2175-
2184.

9. Liang D, Peng Y, Lv W, Deng L, Zhang Y, Li H, et al. Copy number variation sequencing
for comprehensive diagnosis of chromosome disease syndromes. J Mol Diagn.
2014;16:519-526.

10. Chaubey A, Shenoy S, Mathur A, Ma Z, Valencia CA, Reddy Nallamilli BR, et al. Low-
pass genome sequencing: Validation and diagnostic utility from 409 clinical cases of
low-pass genome sequencing for the detection of copy number variants to replace
constitutional microarray. J Mol Diagn. 2020;22:823-840.

11. Dong Z, et al. Low-pass whole-genome sequencing in clinical cytogenetics: a
validated approach. Genet Med. 2016;18:940-948.

12. Wang J, Chen L, Zhou C, Wang L, Xie H, Xiao Y, et al. Prospective chromosome
analysis of 3429 amniocentesis samples in China using copy number variation
sequencing. Am J Obstet Gynecol. 2018;219(3):287.e1-287.e18.
doi:10.1016/j.ajog.2018.05.030.

13. Wang H, et al. Low-pass genome sequencing versus chromosomal microarray
analysis: implementation in prenatal diagnosis. Genet Med. 2020;22:500-510.

14. Clinical Genetics Group of Medical Genetics Branch Chinese Medical Association;
Professional Committee for Prenatal Diagnosis of Genetic Diseases Medical Genetics
Branch of Chinese Medical Association; Group of Genetic Disease Prevention and
Control Birth Defect Prevention and Control Committee of Chinese Society of
Preventive Medicine. Expert consensus on the application of low-depth whole
genome sequencing in prenatal diagnosis. Chin J Med Genet. 2019;36:293-296.

15. Zheng Y, Zhu B, Tan J, Guan Y, The Chinese Genomic Structural Variants Consortium,
Morton CC, et al. Experience of low-pass whole-genome sequencing-based copy
number variant analysis: a survey of Chinese tertiary hospitals. Diagnostics.
2022;12:1098. doi:10.3390/diagnostics12051098.

16. Raca G, Astbury C, Behlmann A, De Castro MJ, Hickey SE, Karaca E, et al. Points to
consider in the detection of germline structural variants using next-generation
sequencing: a statement of the American College of Medical Genetics and Genomics
(ACMG). Genet Med. 2023;25(2):100316. doi:10.1016/j.gim.2022.09.017.

17. Raman L, Dheedene A, De Smet M, Van Dorpe J, Menten B. WisecondorX: Improved
copy number detection for routine shallow whole-genome sequencing. Nucleic Acids
Res. 2018;47:1605-1614.

18. Geng Q, Cui X, Zhang Y, et al. Screening of triploid with low-coverage whole-genome
sequencing by a single-nucleotide polymorphism-based test in miscarriage tissue. J
Assist Reprod Genet. 2019;36:2525-2531. doi:10.1007/s10815-019-01588-6.

19. Mazzonetto PC, Villela D, da Costa SS, Krepischi ACV, Milanezi F, Migliavacca MP, et
al. Low-pass whole genome sequencing is a reliable and cost-effective approach for
copy number variant analysis in the clinical setting. Ann Hum Genet. 2024;88(2):113-
125. doi:10.1111/ahg.12532.

20. Mitrakos A, Kosma K, Makrythanasis P, Tzetis M. Prenatal chromosomal microarray
analysis: Does increased resolution equal increased yield? Genes. 2023;14:1519.
doi:10.3390/genes14081519.

21. Joynt ACM, Axford MM, Chad L, Costain G. Understanding genetic variants of

uncertain significance. Paediatr Child Health. 2021;27(1):10-11.

doi:10.1093/pch/pxab070.

22. Kang H, Wang L, Xie Y, Chen Y, Gao C, Li X, et al. Prenatal diagnosis of chromosomal

mosaicism in 18,369 cases of amniocentesis. Am J Perinatol. 2024;41(S 01):e2058-

e2068. doi:10.1055/s-0043-1770163.

23. Wallace SE, et al. Resources for genetics professionals — Mosaicism. In: Adam MP,

Feldman J, Mirzaa GM, et al., editors. GeneReviews® [Internet]. Seattle (WA):

University of Washington, Seattle; 1993-2024. Available

from: https://www.ncbi.nlm.nih.gov/books/NBK5854.

24. Srivastava S, Phadke SR. Low-pass genome sequencing: A good option for detecting

copy number variations. Genet Clin. 2022;15:April-June.

Table 1: List of retrospective clinical samples and its detail used for validation of lpWGS and comparison with Orthogonal test

Sl No.	Sample Type	Age Range | Gender	Orthogonal Test	Cytoband with alteration	Orthogonal Test			Lowpass Whole genome			Concordance check
					Start	End	Size (Mb)	Start	End	Size (Mb)	
						Phase I					
V1	POC	1st Tri	CMA	Trisomy 21	13634137	46677460	33.04	0	48129895	48.13	C
V2	CVS	-|NK	CMA	Trisomy 18	136227	80255845	80.12	0	78077248	78.08	C
V3	POC	-|NK	CMA	XXY; Trisomy 13; Trisomy 21	251880; 18862148; 13634137	156004066; 114342258; 46677460	155.75; 95.5; 33.0	518401; 18875554; 13850829	155783229; 114122514; 46484554	155.3; 95.25; 32.63	C
V4	POC	-|NK	CMA	XO; Trisomy 18	670673; 136227	156004066; 80255845	155.33; 80.1	1; 1	155270560; 80140651	155.2; 80.14	C†
V5	POC	-|NK	CMA	XO; Del:2q37.3; Dup:16p13.3p11.2	251880; 238580161; 35881	156004066; 241841231; 32836614	155.75; 3.26; 32.80	1 238530917; 1	155270560; 242032733; 31858519	155.2; 3.50; 31.86	C†
V6	POC	-|NK	CMA	XXY	2785592	155691858	152.91	1	155270560	155.2	C†
V7	POC	-|NK	CMA	XO	251880	156004066	155.75	1	155270560	155.3	C†
V8	POC	-|NK	CMA	XXX	2785592	155691858	152.91	518401	155783229	155.26	C
V9	POC	1st Tri|NK	CMA	Trisomy 12	64621	133201316	133.14	0	133851895	133.85	C
V10	DD	6-10Y|F*	KT; CMA	47,X,der(Y),der(Y)[32]/46,X,der(Y)[26]/45,X[2]; Dup Yp11.2q11.23	2782100	26277778	23.5	1	59373566	59.3	C
V11	PB	6-10 Y|M	CMA	Dup:Xq28	153063080	154164940	1.1	153029585	154403555	1.37	C
V12	AF	2nd Tri|NK	CMA	Del:Xp22.31	6568449	8188133	1.62	6567032	8126043	1.56	C
V13	PB	6-10 Y|M	CMA	Del:1p36.33p36.32	914087	4687603	3.77	1035549	4106654	3.07	C
V14	AF	2nd Tri|NK	CMA	Del:3q29	193054506	195291140	2.24	192701721	195307011	2.61	C
V15	DD	0-5 Y|M	F-22q; CMA	Negative; Del 3p26.3p26.1; Dup 9q32q34.3	20214; 114420869	4426396; 138125937	4.4; 23.7	0; 117178796	4560000; 141213431	4.5, 24	C
V16	PB	0-5 Y|M	CMA	Dup:3q13.2q13.31	112451525	115722398	3.27	112416860	115710247	3.29	C
V17	DD	0-5 Y|M	KT; F-22q, F-7q; CMA	46,XY, der(6); Negative; Negative; Dup 3q25.32q29	157312388	193491389	36.2	157001502	193501721	36.5	C
V18	DD	0-5 Y|M	KT; CMA	46, XY? der (15); Dup 4p16.3p15.2	68454	25612567	25.5	0	25640993	25.4	C
V19	PB	0-5 Y|F	CMA	Del:4q34.1q35.2	172079686	190036305	17.96	172151582	189871569	17.72	C
V20	DD	0-5 Y|NA	F-22q; CMA	Negative; Dup: 4q35.2	186249068	188459067	2.2	186981307	189181307	2.2	C
V21	PB	0-5 Y|F	CMA	Dup: 4q25; Del:17p11.2	111131505; 16880332	112863326; 20488761	1.73; 3.61	111140415; 16916072	112967231; 20229100	1.83; 3.31	C
V22	PB	0-5 Y|M	CMA	Del:5q35.2q35.3	175989093	178055505	2.07	176170924	177632806	1.46	C

V23	AF	-|NK	CMA	Del:5p15.33p15.2; Dup: 20p13p12.1	113462; 80929	13788173; 15514914	13.67; 15.43	0; 0	13810000; 15360000	13.81; 15.36	C
V24	DD	0-5 Y|M	KT; F-22q; CMA	46, XY, der (5)t(5;?)(p15;?); Negative; Del 5p15.33p14.3; Dup 8q23.2q24.3; Dup 6q16.3	113462; 110872286; 100875815	23297521; 145070385; 102907265	23.2; 34.2; 2	0; 111900277; 101330543	23313813; 146364022; 10333054	23.3, 34.3, 2	C
V25	AF	-|NK	CMA	Del:6q26q27; Dup:7q33q36.3	162350656; 133640234	170610394; 159327017	8.26; 25.68	162025576; 133113445	171115067; 159138663	9.09; 26.03	C
V26	PB	0-5 Y|M	CMA	Del:7q35q36.3	145083337	159327017	14.24	145209927	159193037	13.98	C
V27	AF	-|NK	CMA	Del:7p21.3p15.2	10611592	26483627	15.87	10660000	26460000	15.80	C
V28	PB	6-10 Y|M	CMA	Dup:8q24.21q24.3; Del:21q22.3	130350392; 46045960	145070385; 46673449	14.72; 0.63	130269822	143035595	12.77	C
V29	AF	-|NK	CMA	Del:9q22.1q22.31	87988200	92870072	4.88	90521336	95971336	5.45	C
V30	AF	2nd Tri|NK	CMA	Del:10p15.3p14	100048	7701527	7.60	100048	7701527	7.60	C
V31	DD	0-5 Y|F	KT; F-22q; CMA	46, XY, del(10)(p13); Negative; Del 10p15.3p14	54087	11273168	11.2	0	11460000	11.4	C
V32	POC	-|NK	CMA	Dup:11p15.5q23.3; Del:22q11.1q11.21	230616; 16408174	116744147; 20291006	116.51; 3.88	230616; 16408174	116744147; 20291006	116.51; 3.88	C
V33	PB	0-5 Y|M	CMA	Del:13q22.2q31.3	75664252	91427209	15.76	75526376	91390174	15.86	C
V34	POC	-|NK	CMA	Del:13q32.3q34; Dup:16p13.3p13.2	101067948; 35881	114342258; 8831829	13.27; 8.79	101770197; 5260000	115169878; 8910000	13.40; 3.65	C
V35	AF	2nd Tri|NK	CMA	Del:15q13.2q13.3	30820717	32622039	1.80	30885000	32485000	1.60	C
V36	PB	0-5 Y|M	CMA	Del:15q11.2q13.1	22582283	28697883	6.12	23571052	28127176	4.56	C
V37	DD	6-10 Y|M	ANGL; CMA	Negative; Del 17p12	14196248	15601354	1.4	14000000	15500000	1.4	C
V38	AF	2nd Tri|NK	CMA	Del:17q24.1q24.2	66060439	67467739	1.41	66032271	67400943	1.37	C
V39	DD	0-5 Y|F	KT; CMA	46,XX; Dup 17q11.2	30689342	31870304	1.2	29 007057	30707057	1.7	C
V40	PB	0-5 Y|M	CMA	Del:18q21.2q21.33	54720552	62003701	7.28	54761202	61988318	7.23	C
V41	POC	-|NK	CMA	Dup:19q13.2	39151229	40281520	1.13	38942109	40426354	1.48	C
V42	PB	50-55 Y|F	CMA	Del:21q22.3	45329955	46673449	1.34	45373861	46598733	1.22	C
V43	PB	6-10 Y|M	CMA	Del:21q11.2q21.3	13644166	25925065	12.28	13850829	25933790	12.08	C
V44	DD	26-30 Y|F	F-22q	Positive	-	-	-	18862659	23270380	4.4	C
V45	DD	0-5 Y|F	F-22q; CMA	Positive; Del 22q11.21	18950000	21108369	2.2	18862659	21470380	2.6	C
V46	PB	11-15 Y|M	CMA	Del:22q13.32q13.33	48199706	50759338	2.5	48199706	50759338	2.6	C
V47	PB	0-5 Y|M	CMA	Del:3p25.3; Dup:15q11.2	8617702; 22582283	9971426; 23060000	1.35; 0.48	8392526; 22852677	10442526; 23095635	2.05; 0.24	C
V48	PB	0-5 Y|F	CMA	Del:10q22.3; Del:15q11.2q13.1	79625681; 23102621	80213053; 28458904	0.59; 5.36	79551950; 23571052	80232513; 28127176	0.68; 4.56	C
V49	POC	-|NK	CMA	Mos. Trisomy 13[0.35]	18862148	114342258	95.5	18800001	114400000	95.6	2GB

V50	PB	0-5 Y|NK	CMA	Mos. Monosomy X[0.3]	2785592	155691858	152.91	200001	156000000	155.8	2GB
V51	POC	-|NK	CMA	Mos. Dup:Xp22.31q21.1[0.4]; Mos.Del:Xq21.1q28[0.4]	8186478; 78579045	78576179; 155233098	70.39; 76.65	0; 76708058	9150000; 155270560	42.2; 78.6	2GB
V52	POC	-|NK	CMA	Mos. Monosomy 19 [0.3]	260912	58445449	58.1	200001	58600000	58.4	2GB
V53	POC	-|NK	CMA	Mos. Trisomy 8 [0.20]	208049	145070385	144.8	200001	145000000	144.8	2GB
V54	CVS	-|NK	CMA	Mos. Trisomy 13 [0.15]	18862148	114342258	95.5	18800001	114400000	95.6	2GB
V55	AF	2nd Tri|NK	CMA	Mos. Trisomy 22[0.20]	16408174	50759338	34.3	15800001	50800000	35.0	2GB
V56	CVS	2nd Tri|NK	CMA	Mos. Trisomy 20 [0.30]; Trisomy 21	81022; 13634137	64282292; 46677460	64.2; 33.04	1; 13000001	64200000; 46800000	64.2; 33.8	2GB
V57	AF	2nd Tri|NK	CMA	Mos. Dup: 1q21.1q25.2[0.36]	143459799	178083001	34.6	143200001	178200000	35.0	2GB
V58	POC	-|NK	CMA	Mos. Del: 7q11.21q21.11[0.4]	62437390	77764079	15.3	10660000	26560000	15.9	2GB
				Phase II							
V59	PB	41-45 Y|M	CMA	Dup:Xp11.22	53522223	53604372	0.08	53531129	53596588	0.07	C
V60	PB	11-15 Y|M	CMA	Hom.Del:Xp11.23p11.22	50034443	50205615	0.17	50046551	50203420	0.16	C
V61	AF	-|NK	CMA	Hom.Del:Xp21.1	31613916	31761327	0.15	31618133	31746217	0.13	C
V62	PB	0-5 Y|M	CMA	Del: 6q25.3	156987387	157101227	0.113	157010931	157094882	0.1	C
V63	PB	0-5 Y|M	CMA	Del:2p16.3	47783201	47837567	0.05	47777481	47847637	0.07	C
V64	CVS	-|NK	CMA	Del:2q31.2	178716155	178979102	0.26	178697107	178972060	0.27	C
V65	POC	-|NK	CMA	Del:16p11.2	29555976	30215609	0.66	29600001	30200000	0.59	C
V66	PB	0-5 Y|F	CMA	Del:6p25.3	1593382	1701415	0.11	1585474	1692283	0.11	C
V67	PB	0-5 Y|M	CMA	Del:9q34.3	137547852	137657680	0.11	137525141	137651995	0.13	C
V68	CVS	-|NK	CMA	Del:10p13	14944633	15022198	0.08	14945159	15033707	0.09	C
V69	PB	16-20 Y|F	CMA	Del:14q23.1	58280647	58458765	0.18	58278240	58462794	0.18	C
V70	DD	6-10 Y|M	CMA	Del:14q22.1	50553477	50726580	0.17	50561349	50720465	0.16	C
V71	AF	-|NK	CMA	Del:15q11.2	22582283	23102621	0.52	22781078	23102105	0.32	C
V72	PB	0-5 Y|M	CMA	Del:15q25.2	82536429	82583477	0.05	82547192	82615826	0.07	C
V73	DD	-|NK	CMA	Del: 6q27	170279913	170605209	0.32	170267801	170601115	0.3	C
V74	PB	0-5 Y|M	CMA	Del:16p13.3	2081121	2235560	0.15	2092639	2228021	0.14	C
V75	PB	0-5 Y|M	CMA	Dup:17q21.31	46110932	46215310	0.10	46087375	46286901	0.20	C
V76	AF	-|NK	CMA	Del:22q11.21	20704600	21110475	0.41	20725934	21092778	0.37	C
V77	PB	6-10 Y|F	CMA	Del:22q13.33	50707646	50818468	0.11	50710333	50730443	0.11	C
V78	POC	-|NK	CMA	Mos.Trisomy 16 [0.55]	35881	90088654	90.05	40001	90120000	90.1	C

V79	POC	-|NK	CMA	Mos.Trisomy 13 [0.62]	18862148	114342258	95.48	18875194	114345127	95.5	C
V80	PB	0-5 Y |F	CMA	Mos.Trisomy 9 [0.5]	203862	138125937	137.92	100001	138160000	138.1	C
V81	POC	-|NK	CMA	Mos.Monosomy 21[0.3]	13634137	46677460	33.04	12940001	46680000	33.7	C
V82	POC	-|NK	MCC	Triploidy	-	-	-	-	-	-	C
V83	POC	-|NK	MCC	Triploidy	-	-	-	-	-	-	C
V84	POC	-|NK	MCC	Triploidy	-	-	-	-	-	-	C
V85	POC	-|NK	QF-PCR	Triploidy	-	-	-	-	-	-	C

CVS – Chorionic Villous Biopsy, PB – Peripheral Blood, POC – Products of Conception, DD – Direct DNA, AF – Amniotic Fluid, 1st Tri – First Trimester, 2nd Tri – Second Trimester, NK- Not Known, CMA – Chromosomal Microarray, MCC – Maternal Cell contamination, QF-PCR – Quantitative Fluorescence – Polymerase Chain Reaction

Note: Control samples were used to build the reference model, Phase I – 68 normal samples and Phase II – 42 normal samples, other samples where no event was detected by orthogonal testing - 6 CMA, 8 karyotyping, 13 karyotyping and FISH, samples are not shown in the Table 1, C - Concordance between CMA data and lpWGS by either Wiseconder/DRAGEN CNV calling tools.
[†]DRAGEN is capable of detecting events such as segmental loss or gain through its CNV calling workflow. However, it does not directly provide information on sex chromosome anomalies. To address this, we utilized our internal workflow to compute z-scores for the X chromosome by comparing the sample against control samples. Based on these z-scores, we predict whether the sample shows signs of conditions such as XO or XXY. Samples V47 and V48 in phase I had variants below the predetermined threshold.

Table 2: Demographic data and Diagnostic yield of the 1,260 samples evaluated by lpWGS

(n=1,260)	Samples	Gestational Age / Age	Maternal Age	Normal	Abnormal	
					P/LP	**VoUS**
Fetus	801 (63.6%)	18W 5D (11W 1D - 33W 3D)	30.2 (17-45)	90.1% (720/799)	7.3% (58/799)	2.6% (21/799)
POC/IUD/Still birth	346 (27.5%)	18W 0D (7W 3D - 34W)	29.6 (18-47)	68.7% (222/323)	28.8% (93/323)	2.5% (8/323)
Postnatal	113 (9.0%)	16Y 2M (10D - 49Y)	-	78.8% (89/113)	15.0% (17/113)	6.2% (7/113)
Overall	-	-	-	**83.5% (1031/1235)**	**13.6% (168/1235)**	**2.9%(36/1235)**

Table 3: Aneuploidies and pathogenic and likely pathogenic variants identified by LP_WGS
Triploidy was identified by MCC/QF PCR data. Subsequently algorithm was optimized to identify the Triploidy by increasing the depth of sequencing in validation phase II.

Nomenclature	Overall (n=168)	Fetus (n=58)	POC/IUD/Still birth (n=93)	Postnatal (n=17)
Common Aneuploidies				
Trisomy 21	38 (22.6%)	28 (48.3%)	8 (8.6%)	2 (11.8%)
Trisomy 18	16 (9.5%)	11 (19.0%)	5 (5.4%)	-
Trisomy 13	6 (3.6%)	3 (5.2%)	3 (3.2%)	-
Sex Chromosome Aberrations				
Turner syndrome	19 (11.3%)	1 (1.7%)	18 (19.4%)	-
Triple X syndrome	2 (1.2%)	2 (3.4%)	-	-
Klinefelter syndrome	2 (1.2%)	2 (3.4%)	-	-
Jacob's syndrome	2 (1.2%)	1 (1.7%)	1 (1.1%)	-
Others	3 (1.8%)	-	3 (3.2%)	-
Rare Autosomes Aberrations				
Trisomy 3	2 (1.2%)	-	2 (2.2%)	-
Trisomy 7	2 (1.2%)	-	2 (2.2%)	-
Trisomy 8	2 (1.2%)	-	2 (2.2%)	-
Trisomy 10	2 (1.2%)	-	2 (2.2%)	-
Trisomy 14	3 (1.8%)	-	3 (3.2%)	-
Trisomy 15	2 (1.2%)	-	2 (2.2%)	-
Trisomy 16	9 (5.4%)	-	9 (9.7%)	-
Trisomy 17	1 (0.6%)	-	1 (1.1%)	-
Trisomy 22	7 (4.2%)	-	7 (7.5%)	-
Other combinations				
Klinefelter syndrome +Trisomy 18	1 (0.6%)	-	1 (1.1%)	-
Triploidy	12 (7.1%)	-	12 (12.9%)	-
Aneuploidy (Including Triploidy)	131 (78.0%)	48 (82.8%)	81 (87.1%)	2 (11.8%)
Copy number variations (Pathogenic/Likely pathogenic)	37 (22.0%)	10 (17.2%)	12 (12.9%)	15 (88.2%)
<5Mb size	20 (20/1235,1.6%) (20/168,11.9%)	7 (7/1235,0.6%) (7/58, 12.1%)	6 (6/1235,0.5%) (6/93,6.5%)	7 (7/1235,0.6%) (7/17,41.2%)
>5 Mb size	17 (17/1235,1.4%)	3 (3/1235,0.2%)	6 (6/1235,0.5%)	8 (8/1235,0.6%)

Table 4: Summary of the smaller pathogenic copy number variation identified by the low pass whole genome sequencing.
1st Tri – First Trimester, 2nd Trimester – Second Trimester

SI No.	Sample Type	Age Range	Cytoband	Size (Mb)	Associated Syndrome
T1	Amniotic Fluid	1st Trimester	Del:17p12	1.50	Charcot-Marie-Tooth Disease Type 1A (CMT1A)
T2	Amniotic Fluid	-	Del:17q11.2	2.60	-
T3	Amniotic Fluid	2nd Trimester	Del:7q31.33-q33	10.50	-
T4	Amniotic Fluid	2nd Trimester	Del:17q12	1.75	Renal Cysts and Diabetes Syndrome (RCAD)
T5	Chorionic Villus Sample	1st Trimester	Mos.Del: Xq24	2.45	-
T6	Chorionic Villus Sample	-	Del:16p12.2-p11.2	8.00	Rubinstein-Taybi Syndrome
T7	Amniotic Fluid	2nd Trimester	Del: Xp22.31	1.85	Ichthyosis, X-linked (Steroid Sulfatase Deficiency)
T8	Amniotic Fluid	-	Mos.Del: 7q11.22-q11.23	3.75	Williams Syndrome
T9	Chorionic Villus Sample	2nd Trimester	Mos.Del: 13q32.3–q33.1	1.50	-
T10	Amniotic Fluid	2nd Trimester	Del: 12q21.1-q21.33	15.60	Noonan Syndrome (PTPN11 gene)
T11	Product of conception	-	Del: 22q11.21	1.41	DiGeorge Syndrome/Velocardiofacial Syndrome
T12	Product of conception	-	Del: 15q26.2-q26.3	7.80	-
T13	Product of conception	-	Del: 7q11.21-q11.23	13.96	Williams Syndrome
T14	Product of conception	-	Mos. Dup: 5q23.1-q23.3	14.00	Crohn's Disease (NOD2 gene)
T15	Product of conception	-	Del:8p23.3-p11.1, Mos.Dup:8q12.3 - q24.3	45.14,81.4	Langer-Giedion Syndrome; Burkitt Lymphoma
T16	Product of conception	1st Trimester	Del: 2q36.1-q36.3, Del: 11q22.3 - q23.1	6.2,1	Potocki-Shaffer Syndrome
T17	Product of conception	-	Del: 6p24.3-p23, Del: 7q31.32 - q31.33	5.8,1.3	Cornelia de Lange Syndrome
T18	Product of conception	-	Del: 2q37.2-q37.3	6.95	-
T19	Product of conception	-	Del: 14q12	1.60	-

T20	Product of conception	-	Del:1q21.1-q21.2	1.80	TAR Syndrome
T21	Product of conception	-	Dup: 17p12	1.40	-
T22	Peripheral Blood	0-5Y	Del: 11q23.3-q25	14.06	Jacobsen Syndrome
T23	Peripheral Blood	0-5 Y	Del: 11q23.3-q25	14.06	Jacobsen Syndrome
T24	Peripheral Blood	0-5 Y	Del: 15q11.2-q13.1	5.00	Prader-Willi Syndrome, Angelman Syndrome
T25	Peripheral Blood	26-30 Y	Del: 10q21.1	1.90	-
T26	Peripheral Blood	0-5 Y	Del: 1p32.1-p31.3	3.10	-
T27	Peripheral Blood	16-20 Y	Dup: 15q11.2-q13.1	7.29	Prader-Willi Syndrome, Angelman Syndrome
T28	Peripheral Blood	0-5 Y	Del: 10q11.22-q11.23	5.90	-
T29	Peripheral Blood	0-5 Y	Dup: 12p13.33-p13.31, Dup: 6q15	9.66,1.3	-
T30	Peripheral Blood	6-10 Y	Del: 5p15.33-p14.1	25.41	Cri-du-Chat Syndrome
T31	Peripheral Blood	6-10 Y	Del: 7q11.23	2.65	Williams Syndrome
T32	Peripheral Blood	6-10 Y	Del: 22q11.21	1.30	DiGeorge Syndrome, Velocardiofacial Syndrome
T33	Peripheral Blood	0-5 Y	Dup: 7p22.3-p21.3	8.36	Greig Cephalopolysyndactyly Syndrome
T34	Peripheral Blood	26-30 Y	Dup: 9p24.3-p13.1	38.71	Alagille Syndrome (JAG1 gene)
T35	Peripheral Blood	0-5 Y	Del: 22q11.21	2.20	DiGeorge Syndrome, Velocardiofacial Syndrome
T36	Peripheral Blood	0-5 Y	Del: 7q11.23	1.60	Williams syndrome
T37	Product of conception	-	Dup:13q31.3-q34, Del: 18q21.33 -q23	24.7;17.96	-

Figure 01: Validation flow chart divided in Phase 1 and Phase II

Figure 02: Commonly occurring Aneuploidies and events Trisomy 21(a) Trisomy 18 (b), XO/Turner syndrome (c), 7q11.23 del / Williams Beuren Syndrome (d), 15q11.2q13.1 del / Prader-Willi/Angelman Syndrome (e), 22q11.21 del / DiGeorge Syndrome (f), 5p15.33p13.3 del / Cri du Chat syndrome (g),Triploidy (h), scatterplot CMA vs lpWGS (i)

Figure 03: Distribution of Clinical indications of 1260 samples **A**. Fetus, **B**. POC, **C**. Postnatal

*Overall call rate is 99.8% (1235/1238), Failure rate 0.2%(3/1238) Abnormal (P/LP/VoUS)

Figure 04: Diagnostic yield for lowpass whole genome sequencing of 1260 samples.

Figure 05: Aneuploidies and pathogenic and likely pathogenic variants identified by lpWGS