Title: Observer: Creation of a Novel Multimodal Dataset for Outpatient Care Research

Kevin B. Johnson, MD, MS, FACMI, FAMIA, FAAP, FIAHSI, FAIMBE (corresponding)

David L. Cohen University Professor

Division of Biomedical Informatics, Department of Biostatistics, Epidemiology, and Informatics,

Perelman School of Medicine at the University of Pennsylvania

Department of Computer and Information Science, School of Engineering and Applied Science,

University of Pennsylvania, Philadelphia, USA

B202 Richards Building

3700 Hamilton Walk

University of Pennsylvania | Philadelphia, PA 19104-6116

+1 (215) 573-5885

Kevin.johnson1@pennmedicine.upenn.edu

Basam Alasaly, MS

Division of Biomedical Informatics, Department of Biostatistics, Epidemiology, and Informatics,

Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA

Kuk Jin Jang, PhD

Division of Biomedical Informatics, Department of Biostatistics, Epidemiology, and Informatics,

Perelman School of Medicine at the University of Pennsylvania

Department of Computer and Information Science, School of Engineering and Applied Science,

University of Pennsylvania, Philadelphia, USA

Eric Eaton, PhD

Department of Computer and Information Science, School of Engineering and Applied Science,

University of Pennsylvania, Philadelphia, USA

Sriharsha Mopidevi, MS

Division of Biomedical Informatics, Department of Biostatistics, Epidemiology, and Informatics,

Perelman School of Medicine, University of Pennsylvania, Philadelphia, USA

Ross Koppel, PhD, MA, FACMI, FIAHSI

Division of Biomedical Informatics, Department of Biostatistics, Epidemiology, and Informatics,

Perelman School of Medicine at the University of Pennsylvania, Philadelphia, USA

Professor of Sociology, School of Arts and Sciences, University of Pennsylvania, Philadelphia,

USA

rkoppel@sas.upenn.edu

MeSH Terms: Ambulatory Care, Artificial Intelligence, Data Curation, Primary Health Care,

Video Recording

Word Count – Abstract 242/250 words. Manuscript 3920 words.

Abstract

Objective: To support ambulatory care innovation, we created *Observer*, a multimodal dataset comprising videotaped outpatient visits, electronic health record (EHR) data and structured surveys. This paper describes the data-collection procedures and summarizes clinical and contextual features of the dataset.

Materials and Methods: A multistakeholder steering group shaped recruitment strategies, survey design, and privacy-preserving design choices. Consented patients and primary care providers (PCPs) were recorded using room-view and egocentric cameras. EHR data, metadata and audit logs were also captured. A custom de-identification pipeline, combining transcript redaction, voice masking, and facial blurring, ensured video and EHR HIPAA compliance.

Results: We report on the first 100 visits in this continually growing dataset. Thirteen PCPs from four clinics participated. Recording the first 100 visits required approaching 210 patients from which 129 consented (61%), with 29 patients missing their scheduled encounter after consenting. Visit lengths ranged from 5 to 100 minutes, covering preventive care to chronic disease management. Survey responses revealed high satisfaction: 4.24/5 (patients) and 3.94/5 (PCPs). Visit experience was unaffected by the presence of video recording technology.

Discussion: We demonstrate the feasibility of capturing rich, real-world primary care interactions using scalable, privacy-sensitive methods. Room layout and camera placement were key influences on recorded communication and are now added to the dataset. The *Observer* dataset enables future clinical AI research/development, communication studies, and informatics education among public and private user groups.

Conclusion: Observer is a new, sharable, real-world clinic encounter research and teaching resource with a representative sample of adult primary care data.

Background

The U.S. healthcare system—once a global exemplar—is now criticized for its high costs and relatively poor health outcomes.[1] While biomedical researchers and industry leaders have proposed incremental improvements to existing technological infrastructure—once seen as key to reform[2–4]--these efforts have proven inadequate for addressing the deeper systemic issues. Johnson refers to this as the "Era of Entanglement," a period marked by the intertwined demands of clinical care, regulation, and finance that have produced a fragmented, burdensome system.[5] As with the leap from horse-drawn carriages to motorized vehicles, transformative, rather than iterative, innovation is needed to meaningfully advance healthcare delivery.

The National Research Council defines this sort of thinking as radical change, where healthcare exploits "qualitatively new ways of conceptualizing and solving health and healthcare problems and revolutionary ways of addressing these problems."[4] However, these researchers will need to understand how healthcare works–they need access to data and observations that spur new hypotheses and ideas. The conundrum they face is how challenging it is in our current environment for non-medical faculty to obtain access to patient-care settings due to hospital and privacy policies.

Video as a data collection tool has been under-utilized in healthcare due to inherent barriers, such as confidentiality and privacy.[6–8] Earlier efforts to capture clinical visit knowledge recognized that tacit knowledge, non-verbal interpersonal interactions, and the necessity of understanding 'embodied cognition' all emphasize the need to video-capture interactions between physicians and their patients in a natural setting.[6,9–11] These data also enable the documentation of activities

in their entirety and complexity.[12] It builds on the work of others who have used video tapes and eye-tracking methods.[13–15] Video capture alone is an incomplete source of clinic visit detail, because of the increasing role of electronic health records, web applications, and other in-room technology.[16] Combining video data with a view of the activities taking place on the computer (EHR logs, clinical notes, and visit-related surveys) as well as a room layout showing the position of the technology with respect to the patient provides a multifaceted view, allowing for qualitative and quantitative analysis and a view supporting a holistic analysis of the encounter. This perspective is especially intriguing given current advances in multimodal AI.[17–23]

There are also concerns that the process of video recording may bias the content of the encounter, thereby lessening the value of these recordings due to the Hawthorne effect.[24] Interestingly, video recording has been shown to be less disruptive to natural participant behavior than human observers present during visits.[6] Another study examining physicians' consultation behavior found no evidence that awareness of video recording affects objective measures of their performance, supporting the use of video recording as a valuable tool for teaching and research purposes.[25]

To address the challenges of researcher access to clinic encounters, and to embrace the need for multimodal data, we embarked on creating the Observer dataset, a multimodal resource for multidisciplinary research. This paper describes those efforts and observational frameworks, initial studies enabled by the dataset, and initial observations.

Materials and Methods

The overarching goal of Observer is to provide a library of clinical exam rooms and the interactions among the patient, any other caregivers, the primary care provider (PCP), and technology.

Our goal is to provide a downloadable dataset to users. interested in research directly related to the clinical exam room. We anticipated that researchers, educators, and policymakers would be most interested in this resource. Ultimately, our plan is for Observer to be housed in a repository that will adhere to FAIR principles as defined below:

- *Findable* by providing metadata that clearly describes the data using a dashboard, being indexed in searchable repositories and using a digital object identifier to provide a persistent link to its location on the internet.

- *Accessible* by providing the website URL (https://observer.med.upenn.edu/) and data availability statement on a series of public sites, in addition to our lab site.

- *Interoperable* by mapping medical data to standard vocabularies, and using the OMOP table schema for Observer data.

- *Reusable* by providing clear usage licenses and documenting information provenance, including any sources for derivative metadata.

The remainder of this paper describes the dataset creation process and initial observations. Of note, this paper focuses on our first 100 recorded visits. We continue to record visits, with a goal of collecting 200 visits in adult primary care settings by the end of 2026.

Design Considerations

We began designing Observer by consulting with two steering groups of adult primary care stakeholders. The first group consisted of adult patients and family members. After a short presentation by the PI, this group reacted to a series of driving questions. They were all very supportive of the project and excited about how it might be used to improve their interactions with physicians and advanced practice nurses. They made the following recommendations to our recruitment and data collection plans:

1. Identify that we are Penn Medicine researchers working with their clinic team (and not, for example, billing or marketing personnel.)

2. Compensation for time and effort should be in the form of a $50 physical gift card, rather than in a form requiring a bank account or an email address.

3. Allow the patient to wear an egocentric camera. Our initial plan was to only ask the PCP to wear this camera, to provide eye tracking as well as a view of the EHR screen. This focus group suggested that seeing what the patient experiences would be invaluable.

4. Strongly consider obtain asking patients and family for visit feedback. Our initial plan was only to ask clinicians for feedback; patients immediately recognized that our plan did not include them, which might cause us to miss valuable insights from their perspective. We agreed and presented the idea of recording the patient watching and commenting about the recorded visit. Patients were concerned about how and where this would be conducted, and recommended we switch to a survey/structured interview after the visit.

A second set of focus groups consisted of clinic physicians and nursing staff. After a short presentation by the PI, the group discussion led to the following recommendations.

1. Consider the use of an after-visit clinician survey instead of being recorded watching and commenting on the visit. Clinicians were uncomfortable seeing themselves on the video.

2. To minimize clinic disruption, begin by recording only first visits of the day for each consented clinician, and progress to multiple visits in one clinic after staff and consented clinicians were comfortable with the processes.

3. To avoid the risk of new patients losing trust in the clinic team, confirm which patients we would approach until the consented clinicians and clinic managers were comfortable with the study processes.

Based on these conversations and a review of similar medicine-related data platforms[13–15,26–28], we constructed a data plan for Observer that holistically covered the activities of a typical ambulatory care visit. We subsequently obtained IRB approval for visit recording and data collection using the process described below, under broad consent for future use.

As noted, this dataset consists of ambulatory care visit videos linked to the EHR user action log, encounter summary, current and past diagnoses, problems, medications, allergies, and all visit-related correspondence from 24 hours before until 48 hours after the recorded visit. Each of these videos is accompanied by transcripts and final progress notes.

Our review of existing efforts to create ambulatory visit data sets was disappointing. In particular, none of these platforms supported the retention of multimodal PHI, which requires sophisticated tools for removing identifiers across modalities (facial recognition blur, voice alteration, metadata scrubbing) that go far beyond standard text-based de-identification[29,30] and extensive curation. Therefore we realized that to share these data, we would need to create a de-identification pipeline.

Data Model

To support interoperability with existing EHR-based data and less commonly available video-derived data, we developed a custom data model inspired by OMOP and MIMIC to structure our EHR-derived fields, visit metadata, and survey instruments.

Equipment Selection

We tested numerous cameras, microphone setups, and screen capture software tools under simulated conditions before selecting our current equipment for collecting encounter information. After tests and discussions with health system security, we ruled out using screen-capture software due to the difficulty in aligning those data with room video, challenges with some software capturing video from more than one application (such as from both the EHR and a web browser) and other equipment challenges, (e.g., bringing in a laptop with enterprise licenses for the PCP to use during the visit).

After testing various configurations, we selected the following:

- Room-view camera with microphone array and 360-degree room view, connected to a local zoom session, monitored by a research assistant outside the room.
- Camera worn on the head at eye level, providing both a microphone and a video from the wearer's perspective (egocentric view) offered to both the PCP and the patient.
- For telemedicine encounters, we selected Zoom, which supported secure downloads of video, and with which clinical teams and patients were familiar.
- To standardize and facilitate remote consent for patients, we initially used an established secure messaging platform.

- We also learned that linking EHR use to the line-by-line transcripts of the PCP-patient interactions is a valuable alignment for end users interested in understanding, for example, what is being discussed while the computer is being used.

Previous approaches to video-capturing clinical interactions, such as by Montague and Asan[31], relied on multiple fixed cameras positioned at different angles to capture gaze direction from external perspectives. We instead incorporated egocentric head-mounted cameras for both patients and physicians to capture first-person perspectives of what each participant actually sees during the visit. We also built on the previous work of others in this space, including Senathirajah and colleagues[13], whose scoping review confirmed that automated detection of eye gaze without eye tracking was feasible and adequate; and Khairat and colleagues[14], who found that user experience can be measured using surveys for physician satisfaction, system usability, and workload intensity. These latter observations motivated our physician survey design. More technical details may be found at https://observer.med.upenn.edu/.

Recruitment and Data Collection Process

Our study recruitment focused on patients and PCPs in adult primary care clinics throughout the University of Pennsylvania Health System, an integrated delivery system. We adopted the suggestion of our multistakeholder steering group and required that PCPs be enrolled first, followed by a convenience sample of their patients. Our final recruitment strategy is below, and summarized in Supplemental Figure 1. All recruitment materials and any changes made in our protocol are available at https://observer.med.upenn.edu/.

Primary Care Provider Enrollment: We were invited to present the goals of the project to primary care providers during their weekly team meetings. After the presentation and question-answering we provided informed consent forms (ICFs) to each PCP. The ICFs also included options for the degree of de-identification each PCP required for broad consent. Initially, consented PCPs were contacted to establish their preference for days and times best suited for video recording. This process was relaxed once PCPs and we became more comfortable with the study protocol.

Patient Enrollment: Our study team contacted any patients who were scheduled for visits within consented PCP's date and time preferences, and described the study and reviewed the consent form using an IRB-approved script. Patients who were willing to participate then received a link to the consent form via email or text message with an option to electronically sign the form, or were met by our team to review and sign the ICF before the visit. As offered to providers, patients also had the option to blur their face and/or obscure their voice with an additional option to blur any other body parts they specified. Patients were not recorded if ICFs were not signed before the patient was taken to the exam room. The University of Pennsylvania IRB approved this process.

Exam Room Preparation and Visit Data Collection: Before the visit, a research coordinator set up in-room cameras, then met the patient in the room to help them place and activate the egocentric camera. Any additional people accompanying the patient signed an informed consent to be recorded, but to have their image and audio altered. After the visit, the research coordinator collected all recording equipment and asked the patient to complete a short survey about the visit while still in the room (under 5 minutes on average). Once the surveys were completed, the research coordinator provided the patient with a gift card before escorting the patient out of the

exam room. Primary care providers were given an exit survey to complete and return to the research team. They received a gift card after returning the exit survey.

EHR Data Extraction: After the visit, a research coordinator completed collecting all data for the visit. Recorded videos were copied to a secure cloud platform (Microsoft Azure Databricks) for de-identification and trimming extraneous events, such as turning on the video recorders, which were captured by recorders already activated. Survey results were labeled with a visitID and stored electronically in a secure location. A request was made to our data and analytics core to extract all pertinent data from the EHR and to label these data with same visit ID. These data were provided to us without PHI and with dates shifted. We updated all data to reflect this new date and ID and stored these data in the same secure location as the video recordings and surveys. All other data derived from the visit video were managed by us, with labels provided by our honest broker. All other labels were removed before the data were available for review.

Privacy Preservation

To preserve privacy, we developed and implemented MedVidDeID[32], a modular pipeline designed to de-identify clinical audio-video (AV) data in compliance with HIPAA Safe Harbor standards.[33] This system integrates state-of-the-art, open-source tools into a six-stage workflow that treats audio and video modalities independently. First, we extract transcripts using WhisperX and de-identify them with a local instance of PHIlter. Audio is de-identified by redacting timestamped PHI segments and transforming voice characteristics to obscure speaker identity. Video frames are processed with YOLOv11 to detect and blur facial features using pose estimation and Kalman filtering. The de-identified audio and video are then recombined and subjected to manual quality control to ensure completeness of PHI removal. An analysis of our

first 11 hours of clinical encounters demonstrated a 91% video de-identification rate and reduced the need to conduct manual processing time by over 60%. This package enables scalable, secure sharing and analysis of AV clinical data while preserving patient privacy.

Results

Participant Recruitment

We recruited 13 PCPs from 4 clinic settings. We approached 210 patients scheduled to see these PCPs to participate in this study. A total of 129 patients agreed to participate (61%), from which we were able to capture a clinic visit for 100 (78%.) There were 29 patients who verbally agreed to participate but were not recorded due to illness, rescheduling, or logistical issues. Three patients participated more than once for separate visits. We contacted three patients whose primary language was non-English, of which two participated in the study (data not shown.) Figure 1 summarizes enrolled patient self-reported health and educational levels.

Table 1 and 2 summarize the results of our first 100 visits. Visit types, as extracted from EHR data represents the range of visit types for primary care.

Table 1 Visit characteristics.

Visit type	Count	Visit type	Count
Follow up	34	Check up	6
Annual Visit	20	Chronic disease management	4
Wellness visit	16	Physical	4
Return visit	7	Pre-op exam	2
New patient	6	Procedure	1

Figure 1 Patient characteristics. Data only available for the last 70 visits, due to the addition of these questions on the post-visit surveys.

PCPs were uniformly distributed across years in practice, ranging from 1-5 years to more than 20 years in practice. In response to the question, "How would you rate your comfort level with using technology in healthcare" most providers described themselves as either self-taught or needing instruction (see Table caption for definitions.) Visit length ranged from 5 - 100 minutes. PCPs rated their visit satisfaction as a weighted average of 3.95/5 across all visits, while their

patients rated visits as 4.24/5 on average, with occasionally discrepant scores provided by the

patient and clinician for the same visit (data not shown, but available on the Observer website.)

Overall, visit experience was unaffected by the presence of video recording technology. Of those

who perceived some impact, one was positive, where the PCP performing a physical exam

reminded a medical student that it was always important to examine the patient without a shirt

on. One other PCP reported negative impact without specifying details.

Table 2 Data about primary care provider (PCP) characteristics were obtained via self-reported post-visit surveys Possible answers to the self-reported technical competence question ("How would you rate your comfort level with using technology in healthcare?") ranged from being a novice ("initial help") to being comfortable with technology ("self taught") to being considered an expert ("Teacher").

PCP #	Total Visits	Visit Length Range	Mean PCP satisfaction with visit (1-5, SD)	Mean patient satisfaction (1-5, SD)	Years in Practice	Technical competence	# visits impacted by recording processes
1	12	16-39	3.9 (0.51)	5(0)	11-15	Initial help	0
2	12	16-44	3.8 (0.39)	5(0)	>20	Self-taught	0
3	8	20-43	4.0 (0)	4(0)	11-15	Self-taught	0
4	21	27-68	4.3 (0.56)	4.47 (0.72)	16-20	Initial help	1 (positively)
5	16	5-39	3.3 (1.65)	4.56 (0.51)	1-5	Teacher	0
6	14	9-33	4.0 (0)	4.29 (0.73)	6-10	Self-taught	0
7	6	26-64	4.2 (0.75)	4 (0.89)	6-10	Self-taught	4 (negatively)
8	3	67-100	3.7 (0.58)	3.67 (0.58)	>20	Initial help	0
9	4	27-49	4.0 (0)	4.2 (0.84)	1-5	Self-taught	0

10	1	23	4.0 (0)	5 (0)	16-20	Self-taught	0
11	1	31	5.0 (0)	4.5 (0.71)	6-10	Self-taught	0
12	1	27	5.0 (0)	4 (0)	11-15	Initial help	0
13	1	33	5.0 (0)	4 (0)	>20	Initial help	0
Total	100	5-100	3.94	4.24	n/a	n/a	

Observer Composition and Structure

At the time of this publication, the Observer dataset comprises 110 outpatient primary care visits, each recorded using a standardized protocol. For each visit Observer includes:

Room-view video recordings, captured with a stationary 360-degree camera equipped with a directional microphone array, positioned to provide a wide-angle perspective of the physical interaction between patient and clinician.

Egocentric videos, recorded using head-mounted cameras worn by the patient and the primary care provider. These recordings offer a first-person perspective of visual attention, body language, and interpersonal engagement. They also provide the equivalent of eye-tracking, enabling a view of how computer technology is used during visits.

EHR-derived audit logs, including timestamped records of all user interactions within the electronic health record during the visit and the surrounding 72-hour window (24 hours before through 48 hours after the encounter).

Structured metadata, covering reason for visit, patient demographics, visit type, clinician characteristics, room configuration, and survey responses.

Survey results from patients and clinicians after the visit, capturing perceptions of communication quality, comfort, and the perceived impact of recording technology.

Transcripts, automatically generated and manually verified, representing the verbal exchange between participants.

Clinical notes, including timestamped text entries from providers linked to specific patient visits, capturing detailed observations, diagnoses, and care plans, with metadata on note type and status.

Vital signs and measurements, recorded during visits, including quantitative data such as blood pressure, weight, height, pulse, and oxygen saturation, linked to the corresponding patient encounter.

To support scalable reuse, we provide standardized file formats (MP4 for video, CSV for logs and metadata, and JSON for annotations) and accompanying documentation that describes data provenance, schema definitions, and known quality attributes. All data packages are stored in a secure institutional archive, with download access granted through a verification process modeled after the MIMIC framework.

Clinic Encounter Visit Details

Figure 2 Summary of visit diagnoses.

Figure 2 summarizes the visit diagnoses covered in our first 100 recordings. Patients had

multiple active diagnoses with a frequency representative of a typical adult primary care setting.

Figure 3 provides a sample of the different room configurations. We observed that the physical

configuration of exam rooms and the EHR screens significantly affects the ability of PCPs and

patients to see each other face-to-face and to view the EHR.

Discussion

We report our experience creating the Observer dataset–a first-of-its-kind library of visit

recordings, EHR data, and many other forms of accompanying information. Previous efforts

such as the Activity Log Files Aggregation toolkit, while methodologically sound, lacked the

infrastructure for long-term data sharing and became inaccessible over time, requiring

subsequent studies to independently collect data, rather than build upon existing foundations; a

limitation that Observer aims to address. We anticipate that Observer will enable hypothesis

generation and testing, as well as an evolving dataset useful for model development and testing. Ethnographic researchers will be able to use Observer to understand compliance with healthcare delivery recommendations, observe care provider-patient interaction, and identify opportunities to improve both needed resources and cognitive support for both patients and PCPs.

Room configuration was surprisingly different in different clinics, and is likely to be an important covariate in studies of communication, as noted by Kumarapeli[34] and summarized in a systematic review.[35] For this reason, we added an image of the room configuration and metadata about this configuration to each visit's data set. These videos provide natural experiments to observe how screen and room arrangements impact the PCPs spend looking at EHR screens vs. looking at patients; and how PCPs use EHR data in dialogs and explanations.

In addition to hypothesis generation and testing,[3] these recordings allow a complete record of consultations to be viewed repeatedly for inter-observer reliability testing and open science.[36] After discussions with researchers at recent meetings, we believe that Observer can be useful for the following classes of problems because of the multimodal nature of the data. We provide a few examples in Table 3, below.

Table 3 Example Use Cases for the Observer Dataset and Their Required Data Modalities.

Use Case	Example	Video	Transcript	EHR data	Audit Log	Room Audio
Ethnographic Research	A researcher could compare visits with and without an automated scribe to measure differences in the frequency of direct patient-provider eye contact and the number of conversational interruptions.	x	x			x
Medical Education	A medical educator could task students with watching an Observer video and using a standardized checklist to evaluate the PCP's communication skills, then discussing the findings as a group.	x	x			x
Visit Summarization	A software engineer could train a model to generate a clinical summary from the visit transcript and then validate its accuracy by comparing the output against the physician's official EHR note.		x	x		x
Quantify Signs and Symptoms	A data scientist could develop a model that analyzes features (e.g., pitch, speaking rate, pauses, gait) to create a quantitative score for cognitive impairment, correlating it with relevant EHR data.	x	x	x		x

Cognitive Support Assessment	A human factors researcher could identify points where a PCP accesses a clinical guideline and analyzes subsequent dialogue to find how that information was raised in the patient conversation.	x	x		x	x
Administrative Burden Assessment	A health services researcher could quantify the amount of in-visit time wasted on administrative tasks (e.g., verbal discussion about next visit, time spent scheduling next visit) versus clinical assessment.		x	x	x	x
Visit summarization (NLP)	Fine-tuning a language model to generate patient-friendly visit summaries		x	x		
Cognitive health monitoring	Detecting speech markers of mild cognitive impairment across longitudinal visits		x	x		x
Predictive modeling (ML)	Predicting patient no-shows based on pre-visit behaviors and clinician communication patterns	x	x	x	x	x
New technology evaluation	Assessing usability of an AI-powered voice assistant for automated documentation in exam rooms	x	x	x	x	x

.

A key contribution of Observer dataset is the ability to synchronize observational data with process data extracted from EHR logs. These data can serve as powerful new sources for Audit and Feedback (A&F), as they help generate the actionable plans with specific advice for improvement that are known to make feedback most effective.[38] This dataset, along with its ability to generate specific, actionable feedback helps catalyze multimodal AI applications in a variety of contexts. For example, we are developing a model for identifying agenda topics using de-identified visit transcripts from Observer.[39] Another team is using multimodal data to develop feature-detection models that predict mild cognitive impairment. These data also support model fine tuning and provide examples for few-shot prompting of large language models. Other contexts will likely involve clinical decision support rules, providing real-time visit information, an almost infinite array of clinical care improvements, physician training, use and design of EHRs, the role of AI, patient facing chatbots, and exam room redesign.

Limitations and Challenges

Although we envision many uses for Observer, we recognize that any data resource of this type suffers from limitations that are either measurable or implied. For example, we have observed no systematic skew in recruiting physicians or patients; however, physician liability concerns could potentially introduce a selection bias in their participation, resulting in a biased sample of patients invited to participate. These potential biases necessitate vigilant monitoring and the implementation of measures to ensure a diverse and representative sample. The ethical implications of video recording and the potential risks to participant confidentiality and privacy must be meticulously addressed to maintain the integrity of the research and the trust of the participants. Balancing the benefits of rich, detailed data collection with the ethical

considerations and participants' comfort is crucial in leveraging representative video recordings for research purposes.[40,41]

The request for video recordings may have raised concerns among potential participants. Although we did not collect data from non-respondents, some individuals may have been reluctant to participate because of the possibility that a) the video recordings may be viewed by multiple individuals over time, b) unauthorized individuals may access video data due to improper storage, and c) an individual's identity could be more easily discerned from a video recording than written data. Although we emphasized privacy protections and included clear language about participants' right to withdraw at any time, some patients still expressed concerns that influenced their willingness to participate. We provided modest gift cards as a token of appreciation, and we allowed patients to select that level of privacy preservation they required to participate as a part of informed consent. These strategies seemed to support participation and may be useful for others collecting encounter video data.

Conclusion

Observer builds on previous work to create a data resource containing a completely de-identified view of ambulatory care visits—with multiple camera views, a visit transcript, an EHR audit log covering from 24 hours before the visit until 48 hours after the visit, and all pertinent EHR data used or accessible during the visit. Creating the Observer dataset required attention to a holistic view of a clinical encounter, as well as patient and PCP privacy using state-of-the art techniques. Next steps will focus on developing the dataset into a repository available for general use.

Data Availability

The Observer dataset is available at https://observer.med.upenn.edu/. Requests for access to its

data may be made by following the instructions on that website.

Code Availability

All code related to this project is available on GitHub, at https://github.com/kbjohnson-

penn/observer_frontend

Acknowledgements

We appreciate the thoughtful review of this manuscript by Dr. S.Trent Rosenbloom and Dr. John

Holmes, as well as the proofreading of this manuscript by Lauren Malloy.

Author Contributions

All authors contributed to the conception and design of this project. Mr. Mopidevi provided the

technical work leading to this manuscript and contributed to the writing of the Methods section.

Mr. Alasaly and Dr. Johnson analyzed the data reported in this manuscript. Drs. Koppel, Jang,

and Eaton contributed to the interpretation of the data reported in this manuscript. All authors

approve the submitted version of this manuscript.

Competing Interests

None

References

1. Blumenthal D, Shah, A, Gunja, MZ, Williams, II RD. *Mirror, Mirror 2024: A Portrait of the Failing U.S. Health System*. Accessed April 29, 2025. https://www.commonwealthfund.org/publications/fund-reports/2024/sep/mirror-mirror-2024

2. Stead WW. Rethinking electronic health records to better achieve quality and safety goals. *Annu Rev Med*. 2007;58:35-47. doi:10.1146/annurev.med.58.061705.144942

3. Johnson KB, Stead WW. Making Electronic Health Records Both SAFER and SMARTER. *JAMA*. 2022;328(6):523-524. doi:10.1001/jama.2022.12243

4. National Research Council (US) Committee on Engaging the Computer Science Research Community in Health Care Informatics. *Computational Technology for Effective Health Care: Immediate Steps and Strategic Directions*. (Stead WW, Lin HS, eds.). National Academies Press (US); 2009. Accessed June 13, 2022. http://www.ncbi.nlm.nih.gov/books/NBK20640/

5. Johnson KB, Neuss MJ, Detmer DE. Electronic health records and clinician burnout: A story of three eras. *J Am Med Inform Assoc*. 2021;28(5):967-973. doi:10.1093/jamia/ocaa274

6. Asan O, Montague E. Using video-based observation research methods in primary carehealth encounters to evaluate complex interactions. *Journal of Innovation in Health Informatics*. 2014;21(4):161-170. doi:10.14236/jhi.v21i4.72

7. Golembiewski EH, Espinoza Suarez NR, Maraboto Escarria AP, et al. Video-based observation research: A systematic review of studies in outpatient health care settings. *Patient Educ Couns*. 2023;106:42-67. doi:10.1016/j.pec.2022.09.017

8. Gilmore RO, Adolph KE, Millman DS, Gordon A. Transforming Education Research Through Open Video Data Sharing. *Adv Eng Educ*. 2016;5(2):http://advances.asee.org/wp-content/uploads/vol05/issue02/Papers/AEE-18-Gilmore.pdf.

9. Chan S. Using videos and multimodal discourse analysis to study how students learn a trade. *International Journal of Training Research*. 2013;11(1):69-78. doi:10.5172/ijtr.2013.11.1.69

10. Coleman T. Using video-recorded consultations for research in primary care: advantages and limitations. *Fam Pract*. 2000;17(5):422-427. doi:10.1093/fampra/17.5.422

11. Nassauer A, Legewie NM. Video Data Analysis: A Methodological Frame for a Novel Research Trend. *Sociological Methods & Research*. 2021;50(1):135-174. doi:10.1177/0049124118769093

12. Schaeffer JH. Videotape: New Techniques of Observation and Analysis in Anthropology. In: Hockings P, ed. *Principles of Visual Anthropology*. DE GRUYTER; 1995:255-284. doi:10.1515/9783110290691.255

13. Senathirajah Y, Borycki EM, Kushniruk A, Cato K, Wang J. Use of Eye-Tracking in Studies of EHR Usability - The Current State: A Scoping Review. *Stud Health Technol Inform*. 2019;264:1976-1977. doi:10.3233/SHTI190742

14. Khairat S, Coleman C, Ottmar P, Bice T, Carson SS. Evaluation of Physicians' Electronic Health Records Experience Using Actual and Perceived Measures. *Perspect Health Inf Manag*. 2022;19(1):1k.

15. Khairat S, Coleman C, Ottmar P, Jayachander DI, Bice T, Carson SS. Association of Electronic Health Record Use With Physician Fatigue and Efficiency. *JAMA Netw Open*. 2020;3(6):e207385. doi:10.1001/jamanetworkopen.2020.7385

16. Committee on Implementing High-Quality Primary Care, Board on Health Care Services, Health and Medicine Division, National Academies of Sciences, Engineering, and Medicine. *Implementing High-Quality Primary Care: Rebuilding the Foundation of Health Care*. (McCauley L, Phillips RL, Meisnere M, Robinson SK, eds.). National Academies Press; 2021:25983. doi:10.17226/25983

17. Bokade R, Navato A, Ouyang R, et al. A cross-disciplinary comparison of multimodal data fusion approaches and applications: Accelerating learning through trans-disciplinary information sharing. *Expert Systems with Applications*. 2021;165:113885. doi:10.1016/j.eswa.2020.113885

18. Bundy H, Gerhart J, Baek S, et al. Can the Administrative Loads of Physicians be Alleviated by AI-Facilitated Clinical Documentation? *J Gen Intern Med*. Published online June 27, 2024. doi:10.1007/s11606-024-08870-z

19. Reed M, Rampono B, Turner W, et al. A multicentre validation study of a smartphone application to screen hand arthritis. *BMC Musculoskelet Disord*. 2022;23(1):433. doi:10.1186/s12891-022-05376-9

20. Silverman D. *Interpreting Qualitative Data*. Seventh. Sage Publishing; 2024.

21. Classen DC, Longhurst C, Thomas EJ. Bending the patient safety curve: how much can AI help? *NPJ Digit Med*. 2023;6(1):2. doi:10.1038/s41746-022-00731-5

22. Acosta JN, Falcone GJ, Rajpurkar P, Topol EJ. Multimodal biomedical AI. *Nat Med*. 2022;28(9):1773-1784. doi:10.1038/s41591-022-01981-2

23. Xue C, Kowshik SS, Lteif D, et al. AI-based differential diagnosis of dementia etiologies on multimodal data. *Nat Med*. 2024;30(10):2977-2989. doi:10.1038/s41591-024-03118-z

24. Pringle M, Stewart-Evans C. Does awareness of being video recorded affect doctors' consultation behaviour? *Br J Gen Pract*. 1990;40(340):455-458.

25. Pringle M, Stewart-Evans C. Does awareness of being video recorded affect doctors' consultation behaviour? *Br J Gen Pract*. 1990;40(340):455-458.

26. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377-381. doi:10.1016/j.jbi.2008.08.010

27. Gilmore RO, Adolph KE, Millman DS. Curating identifiable data for sharing: The databrary project. In: *2016 New York Scientific Data Summit (NYSDS)*. IEEE; 2016:1-6. doi:10.1109/NYSDS.2016.7747817

28. Liberman, M, Cieri C. *The Evolution of the Linguistic Data Consortium.* Accessed January 22, 2024.

    https://scholar.google.com/scholar_url?url=https://www.ldc.upenn.edu/sites/www.ldc.upenn.edu/files/lrec1998-creation-distribution-use.pdf&hl=en&sa=X&ei=8ISuZf2RKcrJy9YPkL-OiAY&scisig=AFWwaeYm4hUhOlQ1bH2e8Q1mjVoG&oi=scholarr

29. Talby D. Delivering Regulatory-Grade, Automated, Multimodal Medical Data De-Identification. September 7, 2025. https://www.johnsnowlabs.com/delivering-regulatory-grade-automated-multimodal-medical-data-de-identification/

30. Anibal J, Huth H, Li M, et al. Voice EHR: introducing multimodal audio data for health. *Front Digit Health.* 2025;6. doi:10.3389/fdgth.2024.1448351

31. Montague E, Asan O. Dynamic modeling of patient and physician eye gaze to understand the effects of electronic health records on doctor-patient communication and attention. *Int J Med Inform.* 2014;83(3):225-234. doi:10.1016/j.ijmedinf.2013.11.003

32. Mopidevi VSSH, Jang KJ, Alasaly B, et al. MedVidDeID: Protecting Privacy in Clinical Encounter Video Recordings. Published online 2025. doi:10.2139/ssrn.5194782

33. *Guidance Regarding Methods for De-Identification of Protected Health Information in Accordance with the Health Insurance Portability and Accountability Act (HIPAA) Privacy Rule.* U.S. Department of Health and Human Services Accessed May 12, 2025. https://www.hhs.gov/hipaa/for-professionals/special-topics/de-identification/index.html

34. Kumarapeli P, de Lusignan S. Using the computer in the clinical consultation; setting the stage, reviewing, recording, and taking actions: multi-channel video study. *J Am Med Inform Assoc*. 2013;20(e1):e67-75. doi:10.1136/amiajnl-2012-001081

35. Patel MR, Vichich J, Lang I, Lin J, Zheng K. Developing an evidence base of best practices for integrating computerized systems into the exam room: a systematic review. *J Am Med Inform Assoc*. 2017;24(e1):e207-e215. doi:10.1093/jamia/ocw121

36. Adolph KE, Gilmore RO, Freeman C, Sanderson P, Millman D. Toward Open Behavioral Science. *Psychological Inquiry*. 2012;23(3):244-247. doi:10.1080/1047840X.2012.705133

37. Jamtvedt G, Young J, Kristoffersen D, Thomson O'Brien M, Oxman A. Audit and feedback: effects on professional practice and health care outcomes. In: The Cochrane Collaboration, ed. *The Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd; 2003:CD000259. doi:10.1002/14651858.CD000259

38. Ivers N, Yogasingam S, Lacroix M, et al. Audit and feedback: effects on professional practice. Cochrane Central Editorial Service, ed. *Cochrane Database of Systematic Reviews*. 2025;2025(3). doi:10.1002/14651858.cd000259.pub4

39. Jang KJ, Bhatti S, Pugh S, et al. Towards a Real-time Clinical Agenda Setting System for Enhancing Clinical Interactions in Primary Care Visits. *AAAI GenAI4Health Workshop*. Published online 2025. https://openreview.net/forum?id=Lu1dxr1bWV

40. Maldonado-Castellanos I, Barrios LM. Ethical Issues when Using Digital Platforms to Perform Interviews in Qualitative Health Research. *International Journal of Qualitative Methods*. 2023;22:16094069231165949. doi:10.1177/16094069231165949

41. Sartori L, Theodorou A. A sociotechnical perspective for the future of AI: narratives, inequalities, and human control. *Ethics Inf Technol*. 2022;24(1):4. doi:10.1007/s10676-022-09624-3