Timely Geographic Access to Blood Banking Facilities: A Pan-India Modeling Study

Short title: Blood Banking Accessibility in India

Gaurav Urs[1], Aamir Miyajiwala[1], Ankit Raj[1], Siddhesh Zadey[1,2,3,4]

1 Association for Socially Applicable Research (ASAR), Pune, Maharashtra, India
2 Global Emergency Medicine Innovation and Implementation (GEMINI) Research Center, Duke University, Durham, North Carolina, United States of America
3 Dr. D. Y. Patil Dental College and Hospital, Dr. D. Y. Patil Vidyapeeth, Pune, Maharashtra, India
4 Columbia University Mailman School of Public Health, New York City, New York, United States of America

Corresponding author: Siddhesh Zadey BSMS, MScGH; ASAR Office Address: D2 Sai Heritage, New DP Road, Aundh, Pune, Maharashtra, India 411007; Email: sidzadey@asarforindia.org

Article Type: Original Article
Word count: 3344
No. of figures: 6
No. of references: 31
Conflict of Interest: Siddhesh Zadey serves as the co-founding director of the Association for Socially Applicable Research (ASAR), Chair of the Asia Working Group, the G4 Alliance, Fellow at the Lancet Citizens' Commission on Reimagining India's Health System, and the Drafting Committee Member for Maharashtra State Mental Health Policy. The other authors declare no conflicts of interest.
Funding: None
Data Availability Statement: All data produced in the present study are available upon reasonable request to the authors.
Ethics approval and consent to participate: Not applicable since we did not use any human participants or animal subjects data. All data used in this manuscript were taken from public sources.
Consent for publication: Not applicable
Authors' contributions:
Study concept and design: Siddhesh Zadey
Acquisition, analysis, or interpretation of data: All authors
Drafting of the manuscript: Gaurav Urs, Siddhesh Zadey
Critical revision of the manuscript for important intellectual content: All authors
Statistical analysis: Aamir Miyajiwala, Ankit Raj, Siddhesh Zadey
Administrative, technical, or material support: Siddhesh Zadey
Study supervision: Siddhesh Zadey

Abstract

Background: Efficient blood transfusion planning requires knowing the distribution of and accessibility to blood banking facilities (BBFs). We mapped population-level, timely geographic access to BBFs across 729 districts across India.

Study Design and Methods: We conducted a pan-India geospatial modeling study using 2024 data from e-RaktKosh, a centralized blood management system. As outcomes, we estimated BBFs per million people, travel time to the nearest BBF, and access population coverage (APC), defined as the proportion of the population with access to the nearest BBF by motorized transport (60 minutes) and walking (30 minutes).

Results: In 2024, India had 5685 BBFs, averaging 4.16 facilities per million people. Median travel times to the nearest BBF were 35.25 minutes by motorized transport and 253.27 minutes by walking. For APCs, 93.86% of people were within 60 minutes by motorized transport of their nearest BBF, while 16.21% were within 30 minutes by walking. There were large differences in APCs between districts across states and notable rural-urban disparities.

Discussion: Access to blood banking facilities in India exhibits geographic disparities, with the northeastern and island Union Territories with prolonged median travel times and low APC values. Rural areas are disproportionately adversely affected. Future efforts should focus on using timely geographic access data for strategic infrastructure planning and resource allocation to enhance healthcare equity.

Keywords: access; blood banks; India; lower-middle-income countries; geospatial

1 Introduction

The World Health Organization (WHO) classifies blood and blood products as essential medicines and notes a significant gap in access to these critical resources across countries, adversely impacting access to blood in low- and middle-income countries.[1] Timely geographic access to blood and blood products is essential across acute specialties, including emergency, intensive care, surgical, obstetric, trauma, and anesthesia care. Beyond acute specialties, such access is vital for managing chronic hemato-oncological conditions that require regular transfusions.[2] Hence, establishing efficient and equitable blood banking systems is arguably the most important way to meet the above-mentioned healthcare needs.[3]

Timely geographic access to blood relies, partly, on spatial proximity to BBFs. Prolonged travel times result in delays in accessing blood transfusions, potentially increasing the risk of adverse consequences, especially for patients in critical and chronic conditions or those needing emergent or urgent care.[4] Hence, access time to BBFs is vital for the equitable utilization of blood banking services.[5]

In low- and middle-income countries (LMICs), blood supply primarily relies on blood banks and storage facilities. LMICs are known to face barriers concerning availability, safety, and infrastructure.[5,6] While studies in LMICs have focused on these challenges, efforts to evaluate timely geographic access to BBFs are limited and localized to smaller geographies, regardless of the high priority.[7,8]

India, the most populous country with over 1.4 billion residents, faces formidable challenges in ensuring access to a large and heterogeneous blood banking system regulated under the National Blood Transfusion Council (NBTC).[9] Previously, studies have evaluated the performance of Indian BBFs for blood volume availability, accreditation, safety, infrastructure, regulations, etc., using systematic composite indices.[10,11] However, they did not include timely geographic access, which is the primary focus of this study.

We aimed to assess the various measures of geographic access to BBFs in India. First, we mapped the BBF geolocations and estimated BBF density per million people. Second, we analyzed the travel times to the nearest BBFs by different modes of transport. Third, we assessed the proportion of people who can access BBFs within 30 and 60-minute intervals, which we defined as population access coverage (APC). We present estimates for the above outcomes of geographic accessibility at national, state, and district levels with rural and urban disaggregations. Finally, we also assessed the state-level correlations between APC and previously published BBF performance indices, as a step toward concurrent validation.

2 Methods

2.1 Data sources

We obtained geographic locations for 5685 blood facilities across 729 districts, within 28 states and eight union territories (UTs) from e-RaktKosh, a centralized blood management system maintained by the Centre for Development of Advanced Computing (CDAC).[12] We used Awesome Tables API to geocode the locations. A series of visualizations helped

identify outliers and anomalies, which were then corrected manually using Google Maps by verifying their addresses.

Population counts were sourced from high-resolution (1 sq. km.) United Nations-adjusted projections for 2020 from the WorldPop dataset.[13] We obtained friction surface 2019 rasters for motorized transport and walking from the Malaria Atlas Project (MAP), an international scientific effort, primarily concerned with mapping the global response to malaria.[14] We used these rasters to calculate travel times to BBF from each square kilometer (1 km²) to BBFs.[15]

We obtained the administrative boundaries of India from publicly available and validated shapefiles.[16] A previously published global raster with ordinal catchment area (CA) categories based on population densities and proximity to high-density urban areas provides standardized rural-urban regions.[17] We classified CA categories >7 as rural areas.

2.2 Outcomes

We studied three outcomes. First, BBF density was examined as the number of BBFs per million people to understand the geographic distribution of BBFs relative to the underlying population residing in a given geography. Second, the median travel time to the nearest BBF was investigated. Travel time estimates for each 1km² raster were used, and the median and interquartile range (IQR) values were reported at aggregate group levels. Finally, the access population coverage (APC) was defined as the percentage of the population with timely access to the nearest BBF. We used 60 minutes as the timely access threshold for travel by motorized transport and 30 minutes for walking. The APC integrated both the population and timeliness aspects of access.

2.3 Analysis

We manually cleaned the BBF addresses to improve machine readability before geocoding, performed using the Google Maps application programming interface (API) and the 'Awesome Table' add-on for Google Sheets. Any BBFs not captured were manually geocoded in Google Maps. The locations with the most relevant address string were selected from those that returned multiple sets of coordinates. Geo-coordinate similarity was used to identify and remove duplicates, and points beyond India's latitude and longitude limits were excluded.

The WorldPop 2020 and rural-urban CA rasters were analyzed to estimate population distribution at various geographic levels. The global multi-category CA raster (1 km² resolution at the equator, with each pixel representing an agglomeration category) was first cropped to fit within India's national boundaries. The raster was then divided into two layers: rural (CA agglomeration categories >7) and urban (CA agglomeration categories ≤7). We then overlaid the WorldPop 2020 population raster for India (also 1 km² resolution) on these binarized rural-urban CA rasters to get rural and urban population rasters. To ensure compatibility, the CA raster was resampled to match the population raster's extent, resolution, and origin. In the rural population raster, rural area pixels were assigned a value of 1, while urban areas were marked as 'NA', and vice versa in the urban population raster. Residential population in each pixel was calculated by multiplying the population count by the

corresponding category value (1 for rural areas in the rural raster). The total population was derived by summing the rural and urban populations. This was done for all geographic levels: subnational (districts and states) and national (all India). It is important to note that not all districts had both rural and urban areas since some districts were entirely rural. In this analysis, 728 districts had rural areas while 691 districts had urban areas.

We calculated the travel times based on the shortest time taken to traverse the MAP friction surface from each 1 km² pixel to its nearest BBF location.[18] Two accessibility rasters were generated based on the modes of transport: one for walking and the other for motorized transport, providing travel times in minutes assuming optimal speeds. We estimated the median and interquartile range (IQR) for the travel times at different geographic levels, i.e., district, state, and national levels.

For APC calculations, we used timely access thresholds (60 minutes for motorized transport and 30 minutes for walking) to create binary accessibility rasters, where pixels below the threshold were coded 1, and those beyond were marked as 'NA'. The accessibility raster was then aligned and overlaid onto the population raster to determine the population with timely access. We calculated the sum of pixels coded '1' to get the population with timely access at different geographic levels, i.e., district, state, and national levels. Finally, the proportion of the population with timely access (i.e., the APC) was calculated by dividing the population with access in a region by its residential population for rural and urban disaggregations.

We conducted the above geospatial analyses in R software (version 4.3.2). Overall, they are similar to our recent work on population-level timely geographic access to medical college hospitals, COVID testing centers, primary and community healthcare centers, surgical care facilities, and palliative care centers in India.[19–23]

Previously, we analyzed the blood banking performance at the state and district levels using the 2016 data from the NBTC.[11] In that analysis, we constructed a composite index made up of 49 state-level variables across seven themes, including accreditation, ownership, regulation, infrastructure, safety, volume, and workforce. As an initial concurrent validation (i.e., sanity check) for the timely access work presented here, we assessed the ecological relationships between the APC values for BBFs and the previously described BB performance index using Spearman correlations. We conducted the analysis using the SR Plot tool.[24] We classified the strength of correlation as follows: very strong (0.7 to 0.9), strong (0.4 to 0.7), moderate (0.3 to 0.4), weak (0.2 to 0.3), and negligible (0.1 to 0.2), and used the conventional 5% alpha threshold for determining statistical significance.[24,25] We used state-level data instead of district-level data for this analysis since district boundaries have undergone significant changes between 2016 and 2024.

3 Results
3.1 Density of BBFs
In 2024, India had 5685 BBFs. Among states/UT, Tamil Nadu had the highest number of BBFs (642), while the UT of Lakshadweep had just one BBF **(Figure 1A)**. The national BBF

density was 4.12 facilities per million people. The BBF density ranged from 1.34 in Bihar to 25.34 in Lakshadweep. Thirteen (36%) states had BBF density below the national value. Among 729 districts, Bangalore in Karnataka had the highest number of BBFs (104), while 108 districts had only one BBF each. Regarding BBF density, Serchhip in Meghalaya had the highest BBF density (40.54), while Suburban Mumbai in Maharashtra had the lowest (0.19) (**Figure 1B**). Three hundred and sixty-four districts had BBF density lower than the national value. Generally, districts in Tamil Nadu, Kerala, and South Karnataka had higher densities, while those in northern and eastern states of Bihar, Uttar Pradesh, West Bengal, and Jharkhand had lower BBF densities.

3.2 Travel time to the nearest BBF
3.2.1 Motorized Transport
By motorized transport, the national median travel time to the nearest BBF was 35.25 minutes (IQR: 20.20–62.40). For urban areas, the median travel time was 6.66 minutes (IQR: 2.37–20.22), about 5 times lower than that for rural areas: 35.91 minutes (IQR: 20.99–63.05) (**Figure 2A**). Among states/UTs, Lakshadweep had the shortest median travel time to the nearest BBF at 1.40 minutes (IQR: 0.70–1.77), over 25 times faster than the national median time. Ladakh had the longest median travel time at 1781.16 minutes (IQR: 1125.50–2342.05). Across districts, Mumbai in Maharashtra recorded the shortest median travel time of 0.92 minutes (IQR: 0.52–1.35), about 38 times faster than the national value. Anjaw in Arunachal Pradesh had the longest median travel time of 1434.51 minutes (IQR: 991.29–1913.09), over 40 times longer than the national value.

3.2.2 Walking
By walking, the national median travel time to the nearest BBF was 253.27 minutes (IQR: 153.11–392.38). For urban areas, the median travel time was 66.65 minutes (IQR: 25.83–188.33), about 4 times faster than that for rural areas: 256.59 minutes (IQR: 157.87–394.67) (**Figure 2B**). Among states/UTs, Lakshadweep had the shortest median travel time to a BBF by walking, at 9.63 minutes (IQR: 4.81–13.0), over 26 times faster than the national value. Across districts, Lakshadweep in Lakshadweep Islands recorded the shortest median travel time at 9.63 minutes (IQR: 4.81–13.01), over 26 times faster than the national median time. The North & Middle Andaman region of the Andaman and Nicobar Islands had the longest median travel time of 2331.40 minutes (IQR: 1617.12–2995.20), over 9 times longer than the national value.

3.3 Access Population Coverage (APC)
3.3.1 Motorized Transport
Nationally, 1,291,542,656 people (93.86% of the Indian population) had access to BBFs within 60 minutes by motorized transport. Among the states/UTs, New Delhi and Chandigarh had the highest APC at 100%, while Andaman and Nicobar Islands had the lowest value of 48.56% (**Figure 3A**). Forty-nine districts had an APC value of 100%, while three districts had an APC of 0% (**Figure 3B**).

Nationally, 99.44% of the population residing in urban areas and 91.83% of those in rural areas could access their nearest BBF within 60 minutes. Urban areas of five states/UTs had an APC of 100%, while only two rural areas achieved the same. At the district level, 396 urban and 45 rural areas had 100% APC, while three rural and two urban areas had an APC value of 0% **(Figure 4)**.

3.3.2 Walking
Nationally, 223,158,576 people (16.21% of the Indian population) had access to BBFs within 30 minutes by walking. Among the states/union territories, New Delhi led with an APC of 69.05%, while Ladakh had the lowest value at 2.76% **(Figure 5A)**. Among districts, Shahdara, New Delhi had an APC of 97.94%, while 78 districts had an APC value of 0% **(Figure 5B)**.

Nationally, 53.02% of the population residing in urban areas and only 2.63% of rural areas had access to BBFs within 30 minutes by walking. Among states/UTs, rural Puducherry had the highest APC at 13.52%, while rural areas in two states/UTs had an APC of 0%. Urban areas in Mizoram recorded the highest APC of 91.76% while those in Bihar noted the lowest APC of 21.9%. At the district level, only the urban Lower Subansiri in Arunachal Pradesh had an APC of 100%, while urban areas in 30 districts had 0% APC. There were 48 rural areas across districts without access to BBFs within 30 minutes, while three areas had 100% APC **(Figure 6)**.

3.4 Correlations with Blood Banking Performance Index
In a sample of 28 states and eight UTs, between the blood banking performance index and APC values, we found a moderate positive correlation for APC within 60 minutes by motorized transport (r=0.39, p<0.001) and a strong positive correlation for APC within 30 minutes by walking (r=0.42, p=0.04). Both these were statistically significant at the alpha threshold of 0.05.

4 Discussion
4.1 Summary of findings
In 2024, India had 5685 blood banking facilities (BBFs) for about 1.4 billion. The national density was 4.16 BBFs per million people, ranging from 1.34 to 12.94 BBFs per million people across states/UTs. The median walking time to the nearest BBF was 425.74 minutes, while motorized transport took 113.69 minutes. Nationally, 93.86% of the population could reach their nearest BBF in 60 minutes by motorized transport and 16.22% of the population could reach their nearest BBF within 30 minutes by walking. This meant that 88,521,598 people did not have access to a BBF within 60 minutes by motorized transport and 1,208,079,465 did not have access to the nearest BBF within 30 minutes by walking. Urban areas had almost universal access, whereas rural areas had lower coverage. We also observed heterogeneities in access outcomes across states/UTs and districts. At the state level, the APCs were correlated with the blood banking performance index, pointing to preliminary evidence for concurrent validity.

4.2 Previous studies and relevance of findings

In this study, we measured travel times over distances, as the duration to reach a facility depends on various factors beyond distance. Further, APC integrates population with travel time to offer a more comprehensive view of geographical accessibility to care. Previously, another modeling study focusing more on blood availability also assessed timely access in specific regions of the northern and northeastern Indian states.[26] However, a pan-India assessment providing district-level values with rural-urban disaggregations was missing.

Our study noted disparities between rural and urban areas. Rural areas had lower BBF densities, longer travel times, and lower APC values in comparison to urban areas in India. These findings echo the observations of the Association of Rural Surgeons of India (ARSI) that highlighted the disparity in blood supply distribution between rural and urban areas, noting that while over two-thirds of the population resided in rural areas, most blood banks were concentrated in urban centers.[5]

4.3 Policy implications

Our findings help determine whether the lack of access to BBFs is due to either a shortage of BBFs, challenges in transportation to reach them, or both.

We noted disparities between rural and urban areas. Rural areas with low BBF density, such as Bihar, West Bengal, Uttar Pradesh, and Jharkhand, should prioritize establishing new BBFs to meet population demands. Rural regions of Arunachal Pradesh, Assam, and parts of Uttar Pradesh face challenges in travel times mainly due to challenging terrain and poor road infrastructure. In these areas, deploying mobile blood banks and establishing satellite blood banks can help expand services and increase blood collection coverage. Furthermore, partnerships with non-governmental organizations and local community health workers can bridge coverage gaps.[27]

In urban areas, despite having a higher number of BBFs, access remains limited due to high population density and unequal distribution of blood banks, which are often concentrated in centers.[28] This makes it difficult for people in peripheral regions to access blood banks in emergencies, especially in metropolitan cities like Mumbai, New Delhi, and Kolkata.[29] Moreover, the public-private divide, the cost of blood and blood products, and emergency transport delays due to traffic congestion can further exacerbate the lack of timely access.[19] To address these challenges, decentralizing blood bank functions, promoting real-time inventory systems (e.g., e-RaktKosh), establishing blood transport services, and educating the public about access points can help improve access population coverage.[12,30]

In states with longer travel times and where building BBFs might be challenging, such as Ladakh, Arunachal Pradesh, and Himachal Pradesh, the focus should shift to alternative solutions like walking blood banks, immediate autologous transfusion, and donor blood drop strategies.[27] Rigorous mapping of 'blood deserts', i.e., regions lacking access to blood and blood products and associated healthcare services, can help inform local policy changes and implementation strategies.[31]

4.4 Strengths and limitations

To evaluate timely geographic access to BBFs in India, we employed a novel method using multiple outcome measures. Our pan-India study comprehensively analyzed the densities of BBFs per million people, travel times to the nearest BBF by two modes of transport (walking and motorized), and the access population coverage that combines population and timeliness measures for 5685 BBFs across 729 districts. A novel feature of the study is the assessment of access across rural and urban areas. We also add to the broader global health data inventory in terms of the geolocation data for the Indian BBFs. The study focuses on timely geographic access beyond the common blood donation index, which is critical from a health systems perspective. These study methods can be adopted for similar analyses in other low- and middle-income countries.

Our study has several limitations. First, our analysis is subject to the assumptions and limitations of the original data sources, including e-RaktKosh, Worldpop, and MAP. Second, the APC estimates do not consider whether individuals have access to or own motor vehicles. Third, our study calculates travel times and APCs based on walking and motorized land transport, without factoring in other modes of travel, such as waterways, that might be relevant to islands and settlements near larger rivers. Fourth, we did not examine differences in access based on demographic factors such as age, gender, or socioeconomic status. Finally, the correlation between the blood banking performance index and APC values is ecological and can have aggregation bias. Even so, these findings are novel and pave the way for future studies that provide a more nuanced and comprehensive picture of geographical access to BBFs

5 Conclusion

In this study, we analyzed geographic access to blood bank facilities through multiple outcome measures that looked at population, travel times, and a combination of the two. Our study quantified the differential patterns in timely geographical access to BBFs across states/UTs and districts and disparities across rural and urban areas. Future policies must focus on the equitable distribution of facilities in terms of travel times. Future research should focus on utilizing geospatial tools to optimize resource allocation and improve accessibility to BBFs.

Acknowledgments

We thank the Health Informatics Group managing e-Raktkosh for sharing the blood banking facility list with us.

References

1   World Health Organization. Availability, safety and quality of blood products (WHA63.12). https://www.who.int/publications/i/item/WHA63.12 (accessed March 24, 2024).

2   Cabalar I, Le TH, Silber A, *et al.* The role of blood testing in prevention, diagnosis, and

management of chronic diseases: A review. *Am J Med Sci* 2024; published online April 16. DOI:10.1016/j.amjms.2024.04.009.

World Health Organisation. Blood safety and availability. World Health Organisation. 2023; published online June 2. https://www.who.int/news-room/fact-sheets/detail/blood-safety-and-availability (accessed June 12, 2023).

Gammon RR, Rosenbaum L, Cooke R, *et al.* Maintaining adequate donations and a sustainable blood supply: Lessons learned. *Transfusion* 2021; 61: 294–302.

Jenny HE, Saluja S, Sood R, *et al.* Access to safe blood in low-income and middle-income countries: lessons from India. *BMJ Glob Health* 2017; 2: e000167.

Kralievits KE, Raykar NP, Greenberg SLM, Meara JG. The global blood supply: a literature review. *Lancet* 2015; 385 Suppl 2: S28.

Custer B, Zou S, Glynn SA, *et al.* Addressing gaps in international blood availability and transfusion safety in low- and middle-income countries: a NHLBI workshop. *Transfusion* 2018; 58: 1307–17.

Kundu S, Munoz-Valencia A, Virk SK, *et al.* Defining blood deserts and access to blood products for 660 million people: A geospatial analysis of eight states in northern india. 2024. DOI:10.2139/ssrn.4724384.

Mammen JJ, Asirvatham ES, Lakshmanan J, *et al.* The clinical demand and supply of blood in India: A National level estimation study. *PLoS ONE* 2022; 17: e0265951.

Asirvatham ES, Mammen JJ, Lakshman J, *et al.* Assessment of performance of blood banks in india: A national level cross sectional study. *Indian J Hematol Blood Transfus* 2021; 37: 640–7.

Urs GM, Krishna Kumar P, Kamath Y, Zadey S. Indexing blood banking performance in India: A retrospective cross-sectional analysis of states and districts. *Dialogues in Health* 2023; 3: 100160.

G O I. e-RaktKosh:Centralized Blood Bank Management System. eRaktKosh. 2016; published online April 7. https://www.eraktkosh.in/BLDAHIMS/bloodbank/transactions/bbpublicindex.html# (accessed June 13, 2023).

School of Geography and Environmental Science, University of Southampton; Department of Geography and Geosciences, University of Louisville; Departement de Geographie, Universite de Namur, Center for International Earth Science Information Network (CIESIN), Columbia University (2018). WorldPop. 2020; published online June 22. https://dx.doi.org/10.5258/SOTON/WP00671 (accessed July 5, 2024).

MAP. Malaria Atlas Project - MAP. Welcome to the Malaria Atlas Project - MAP. https://malariaatlas.org/ (accessed March 17, 2021).

European Commission. Global Accessibility Map. Global Accessibility Map. https://forobs.jrc.ec.europa.eu/gam/description (accessed July 14, 2024).

Sabde Y, Diwan V, De Costa A, Mahadik VK. Mapping the rapid expansion of India's medical education sector: planning for the future. *BMC Med Educ* 2014; 14: 266.

Cattaneo A, Nelson A, McMenomy T. Global mapping of urban-rural catchment areas reveals unequal access to services. *Proc Natl Acad Sci USA* 2021; 118. DOI:10.1073/pnas.2011990118.

Weiss DJ, Nelson A, Vargas-Ruiz CA, *et al.* Global maps of travel time to healthcare facilities. *Nat Med* 2020; 26: 1835–8.

Thakkar H, Reddy C, Passi VR, *et al.* Assessing Population-level Accessibility to Medical College Hospitals in India: A Geospatial Modeling Study. *Cold Spring Harbor Laboratory* 2024. DOI:10.1101/2024.08.11.24311839.

Patil S, Miyajiwala A, Arora A, *et al.* Geospatial accessibility to primary and community health centres in india. *medRxiv* 2025; published online May 13. DOI:10.1101/2025.05.13.25327502.

Sharma P, Thakkar H, Patil A, *et al.* Access to palliative care in India: situational analysis and modeling of access from public healthcare centers. *Res Sq* 2025; published online April 29. DOI:10.21203/rs.3.rs-6535976/v1.

Poddar S, Gorkar A, Passi VR, *et al.* Geographical Access to COVID-19 Testing Centers in India. COVID-19 Testing Centers Accessibility India. 2025. DOI:10.2139/ssrn.5085689.

Jadhav T, Vissoci JRN, Zadey S. Measuring timely geographical access to surgical care in India: a geospatial modelling study. *Lancet Glob Health* 2022; 10 Suppl 1: S29.

Xiong M, Xu Y, Zhao Y, *et al.* Quantitative analysis of artificial intelligence on liver cancer: A bibliometric analysis. *Front Oncol* 2023; 13: 990306.

Akoglu H. User's guide to correlation coefficients. *Turk J Emerg Med* 2018; 18: 91–3.

Kundu S, Munoz Valencia A, Virk SK, *et al.* Defining blood deserts and access to blood products for 660 million people: a geospatial analysis of eight states in Northern India. *BMJ Glob Health* 2024; 9. DOI:10.1136/bmjgh-2024-015637.

Kundu S, Gerk A, Glatter R, Poenaru D. Innovative transfusion strategies for blood deserts in disaster settings. *BMJ Glob Health* 2024; 9. DOI:10.1136/bmjgh-2024-016854.

Raykar NP, Raguveer V, Abdella YE, *et al.* Innovative blood transfusion strategies to address global blood deserts: a consensus statement from the Blood Delivery via Emerging Strategies for Emergency Remote Transfusion (Blood DESERT) Coalition. *Lancet Glob Health* 2024; 12: e522–9.

Mammen JJ, Asirvatham ES, Sarman CJ, Ranjan V, Charles B. A review of legal, regulatory, and policy aspects of blood transfusion services in India: Issues, challenges, and opportunities. *Asian J Transfus Sci* 2021; 15: 204–11.

Organization WH. Guidance On Ensuring A Sufficient Supply Of Safe Blood And Blood Components During Emergencies. World Health Organization, 2023.

The Lancet Haematology. Blood deserts: a vision to tackle blood inaccessibility. *Lancet Haematol* 2024; 11: e241.

Figures

Figure 1: A) Geographic locations and B) density per million people of blood banking facilities across 729 districts.

Figure 2: Median travel time (in minutes) to the nearest blood banking facility by A)
motorized transport and B) walking across 36 states/Union Territories. The dotted
lines note the national values.

Figure 3: Access population coverage within 60 minutes by motorized transport for A) states/Union Territories and B) districts.

Figure 4: Access population coverage within 60 minutes by motorized transport in rural and urban areas across districts.

Figure 5: Access population coverage within 30 minutes by walking across A) states/Union Territories and B) districts.

Figure 6: Access population coverage within 30 minutes by walking in rural and urban areas across districts.