Rapidly Progressing Glaucoma: Clinical, Structural, and Socioeconomic Drivers of Treatment Escalation

Lok Hin Lee[1], DPhil; Yangyiran Xie[1], BS; Annabelle Pan[2], MD; Chris Bradley[2], PhD; Jithin Yohannan[2, 3], MD, MPH

1 Vanderbilt School of Medicine, Vanderbilt University, Nashville, Tennessee, United States of America

2 Wilmer Eye Institute, Johns Hopkins School of Medicine, Baltimore, Maryland, United States of America

3 Malone Center for Engineering in Healthcare, Johns Hopkins University, Maryland, United States of America

Meeting Presentation: This material has not been presented previously.

Financial Support: Supported by grants from the National Institute of Health 1K23EY032204-03 and Research to Prevent Blindness Unrestricted Grant.

Proprietary Interest Statement: Jithin Yohannan is a consultant for Abbvie, Topcon and Ivantis/Alcon. Chris Bradley is a consultant for Radius XR. The authors have no conflicting proprietary or financial interests related to this manuscript.

Running Head: Treatment Escalation for Rapidly Progressing Glaucoma

Key Words: Glaucoma, Visual Field Progression

Corresponding Author and address for reprints:

Jithin Yohannan, MD, MPH

Wilmer Eye Institute

1800 Orleans Street

Baltimore, MD 21287

jyohann1@jhmi.edu

Abbreviations and Acronyms

•   ADI – Area Deprivation Index

•   CI – Confidence Interval

•   CPT – Current Procedural Terminology

•   dB – Decibel

•   ICD – International Classification of Diseases

•   IOP – Intraocular Pressure

•   logMAR – Logarithm of the Minimum Angle of Resolution

•   MD – Mean Deviation

•   MIGS – Minimally Invasive Glaucoma Surgery

•   OCT – Optical Coherence Tomography

•   OR – Odds Ratio

•   RNFL – Retinal Nerve Fiber Layer

•   SITA – Swedish Interactive Thresholding Algorithm

•   SVI – Social Vulnerability Index

•   VA – Visual Acuity

•   VF – Visual Field

•   ZIP – Zone Improvement Plan (ZIP Code)

Abstract

Purpose:

To evaluate clinical and sociodemographic factors associated with selecting treatments in glaucoma

patients with rapid visual field (VF) progression.

Design:

Retrospective cohort study

Participants:

2,583 eyes from 1,514 adults with 5 or more 24-2 visual fields over five years and at least one optical

coherence tomography (OCT) scan.

Methods:

Rapid progressors were defined by mean deviation (MD) slopes worse than -1 dB/year. Demographic

(age, gender, race), clinical (intraocular pressure (IOP), VF metrics, OCT measures), and socioeconomic

(social vulnerability index, or SVI) variables were collected. Patients were categorized based on the most

intensive treatment received in the first seven years: medical management, minimally invasive procedures

(e.g., minimally invasive glaucoma surgery or laser), or aggressive procedures (e.g., filtering surgery or

external ciliodestruction). Logistic regression was performed to identify demographic, clinical, and

socioeconomic factors associated with treatment intensity.

Main Outcome Measures:

Odds of treatment selection based on rapid VF progression

Results:

Rapid progressors had significantly higher odds of receiving aggressive procedures (odds ratio [OR] 3.35,

95% confidence interval [CI] 2.12–5.30, p < 0.001) and any procedure (OR 2.74, 95% CI 1.87–4.01,

p < 0.001), yet only 15% of rapid progressors underwent aggressive procedures in the first seven years.

Among rapid progressors, elevated IOP, worse MD, smaller rim area, and thinner retinal nerve fiber layer

predicted aggressive or minimally invasive procedures, while lower IOP and milder damage reduced the

likelihood. Higher SVI was associated with a reduced likelihood of receiving minimally invasive

procedures among rapid progressors (OR 0.006, 95% CI 0.000–0.63, p < 0.04).

Conclusion:

Although rapid progression was a strong predictor of aggressive procedures, fewer than one in five

underwent aggressive IOP-lowering interventions. Baseline IOP and structural severity appeared to

supersede VF progression in clinical decisions. Patients in areas of higher socioeconomic vulnerability

were also less likely to receive less invasive procedures. Better integrating rates of functional decline and

addressing socioeconomic barriers may help optimize care for rapidly progressing glaucoma patients.

Introduction:

Glaucoma remains one of the foremost causes of irreversible blindness worldwide and is anticipated to affect an expanding proportion of the aging population[1, 2]. Because the disease often progresses silently in its early stages, prompt detection and intervention are critical for preserving visual function and preventing long-term disability. Visual field (VF) testing is commonly employed to monitor disease progression, with the rate of change in mean deviation (MD) serving as a key measure of decline in visual function. Multiple studies suggest that approximately 5-10% of patients with glaucoma exhibit rapid progression, most commonly defined as eyes experiencing MD worsening exceeding -1 dB/year[3, 4, 5].

A variety of clinical factors have been linked to faster rates of progression. Older age, elevated intraocular pressure (IOP), worse baseline MD, disc hemorrhage, and pseudoexfoliation have all been implicated in more rapid progression[6, 7].

Beyond traditional ocular and systemic risk factors, social determinants of health have emerged as critical considerations in glaucoma care. Composite indices such as the Social Vulnerability Index (SVI) and Area Deprivation Index (ADI) incorporate multiple metrics—educational attainment, income, and housing stability, among others — have been used in previous studies on glaucoma to gauge community-level disadvantage[8, 9, 10]. Although higher levels of SVI or ADI may correlate with delayed access to treatment, more advanced disease at presentation and faster rates of glaucoma worsening[11], some studies also suggest that individuals residing in high-SVI (more disadvantaged) areas might eventually obtain more aggressive interventions, reflecting complex and sometimes counterintuitive relationships between socioeconomic status and healthcare utilization[12].

Within this complex landscape, treatment approaches can vary widely. Although aggressive procedures, such as tradition filtering surgery, has been shown to slow VF loss in glaucomatous patients[13], there remain no clear criteria for escalating beyond medical, laser, or minimally invasive surgical interventions,

and clinical practices differ widely. Understanding the drivers behind treatment decisions for aggressive

surgery versus less invasive therapies, especially for rapidly progressing patients, can, therefore, assist in

developing best practices and identifying ways to improve care.

Here, we aim to identify glaucoma patients who demonstrate rapid progression and examine the ocular,

clinical, and socioeconomic factors that shape treatment strategies. Specifically, we explore whether these

rapidly progressing individuals are likely to undergo aggressive procedures, such as filtering surgery, or

whether they remain on medical treatment or receive less invasive procedures. We aim to clarify the

underlying contributors to treatment escalation and to highlight potential gaps in care for patients who,

despite substantial functional decline, may not receive robust intervention.

Methods:

Study Population

This study was conducted in accordance with the tenets of the Declaration of Helsinki and was approved

by the Johns Hopkins University School of Medicine Institutional Review Board. As this was a

retrospective study using data collected in routine clinical practice, the need for informed consent was

waived by the Institutional Review Board.

We included all adult patient eyes with glaucoma or glaucoma suspect diagnosis based on ICD H40

diagnosis at the Wilmer Eye Institute between 2013 and 2024 with at least five years of follow-up from

the initial VF and at least five reliable 24-2 VFs within the first five years. All VFs were measured using

Swedish Interactive Thresholding Algorithm (SITA)-Standard, SITA-Fast or SITA-Faster test strategies.

Reliable VFs were defined as having <15% false positives and <25% false negatives where available for

eyes with suspect, mild, or moderate glaucoma status or <50% false negatives for eyes with severe

glaucoma[14].

Each eye was also required to have at least one reliable Zeiss Cirrus Optical Coherence Tomography

(OCT) scan within one year of the initial VF to evaluate the impact of OCT-derived metrics on rapid

progression. Reliable OCTs were defined as having a minimum signal strength of six and superior,

inferior, and average retinal nerve fiber layer (RNFL) thicknesses greater than 30um. RNFL thickness less

than 30um is likely due to errors in registration or artifact, as it is approximately three standard deviations

below the floor RNFL value on a Zeiss Cirrus OCT scan[15].

Eyes were classified as having mild, moderate, or severe glaucoma based on their initial VF using

Hodapp-Parish Anderson criteria for MD: mild (MD > -6dB), moderate (MD ≤ -6 dB and MD > -12 dB)

and severe (MD ≤ -12 dB). A linear regression model was fit to the MD values for each eye to estimate

the rate of MD progression. Patients were classified as rapid progressors if the rate of MD progression

within the first five years was less than -1 dB/year and classified as non-rapid progressors otherwise.

Visual acuities for patients were measured using a Snellen chart at the first visit and converted to the

logarithm of the minimum angle of resolution (logMAR). In line with previous literature, we converted

"Count Fingers" to logMAR 1.85, "Hand Motion" to logMAR 2.30, "Light Perception" to logMAR 2.70

and "No Light Perception" to logMAR 3.00[16].

Patients were classified as having received either aggressive procedures or minimally invasive procedures

if they had a corresponding Current Procedural Terminology (CPT) code recorded within seven years of

their first visual field assessment and glaucoma diagnosis. This would be during or after the period of

rapid progression as previously defined for all included patients. Treatment codes for aggressive

procedures included filtering surgeries or external ciliodestruction, whereas minimally invasive procedure

codes included codes which corresponded to Minimally Invasive Glaucoma Surgery (MIGS) or laser

procedures. Patients were assumed to be treated medically otherwise. A comprehensive list of CPT codes

for aggressive and minimally invasive procedures is provided in online-only supplemental material.

Socioeconomic Vulnerability Index

We used the Centers for Disease Control and Prevention and Agency for Toxic Substances and Disease

Registry Social Vulnerability Index 2022 to estimate social vulnerability. This dataset provides an

estimation of the overall socioeconomic vulnerability index (SVI) at a census tract level based on 1)

socioeconomic status, 2) household characteristics, 3) racial and minority status, and 4) housing type and

transportation. SVI ranges from 0 to 1 with a higher value representing a higher level (worse) of

socioeconomic vulnerability. Patient ZIP codes were mapped to census tracts using the U.S. Department

of Housing and Urban Development's ZIP Code Crosswalk Files to assign an SVI to each patient

Descriptive Statistical Analysis

First, we assessed whether there were any statistically significant differences between rapid progressors

and non-rapid progressors for age, gender, SVI, race, and a variety of baseline measures from the first

clinical visit, including MD, average RNFL thickness, cup-to-disc ratio, rim area, intraocular pressure,

and visual acuity. The Wilcoxon rank sum test was used for continuous variables — all the continuous

variables violated assumptions of normality ($p < 0.05$) based on a Shapiro-Wilk test — while the Chi-

squared test was used for categorical variables.

We then structured our analysis to compare treatment patterns across two key dimensions: (1) progression

status (rapid vs. non-rapid progressors), and (2) treatment intensities (aggressive, minimally invasive or

medical) to create six distinct clinical groups for comparison (all possible combinations of progression

status and treatment intensity). We used the Kruskal-Wallis test for continuous variables and the Chi-

squared test used for categorical variables to assess differences across all six clinical groups. If any

significant differences were found, post-hoc pairwise comparisons with Bonferroni correction were

performed to determine which groups maintained significant differences.

Modelling the impact of clinical, demographic and SVI variables on procedure selection

We utilized a multivariable logistic regression model to assess the extent to which clinical and demographic variables predict the likelihood of receiving aggressive procedures, minimally invasive procedures, or "any procedure" (i.e., aggressive or minimally invasive procedures). Separate analyses were conducted for rapid and non-rapid progressors to identify factors significantly associated with procedure selection decisions in each group. A sensitivity analysis was performed across all models by varying the intervention timeframe, adjusting the time frame for treatment receipt from seven years to alternative thresholds of three and five years. All statistical analysis was conducted using Python 3.9.0 with the Statsmodels 0.14.4 and Scipy 1.14.1 libraries.

Results:

181   Table 1. Demographics.

	Overall	Rapid Progressors (n = 182, 7% of all eyes)				Non-Rapid Progressors (n = 2401, 92.9% of all eyes)				P-Value (Comparison between overall rapid and non-rapid progressors)	P-Value (Comparison between treatment groups)
		All Treatments	Aggressive Procedure	Minimally Invasive Procedure	Medical Treatment	All Treatments	Aggressive Procedure	Minimally Invasive Procedure	Medical Treatment		
Patients (# eyes)	2583	182 (100%)	28 (15.4%)	10 (5.5%)	144 (79.1%)	2401 (100%)	102 (4.2%)	100 (4.2%)	2199 (91.6%)		
Biological sex, n (%)										0.96	0.69
Female	1403 (54.32%)	98 (53.85%)	13 (46.43%)	6 (60.00%)	79 (54.86%)	1305 (54.35%)	58 (56.86%)	61 (61.00%)	1186 (53.93%)		
Male	1180 (45.68%)	84 (46.15%)	15 (53.57%)	4 (40.00%)	65 (45.14%)	1096 (45.65%)	44 (43.14%)	39 (39.00%)	1013 (46.07%)		
Age, mean (years)	74.42 [66.52, 80.61]	76.69 [67.48, 82.87]	77.76 [73.29, 84.38]	81.61 [71.74, 82.88]	76.30 [67.18, 82.51]	74.30 [66.43, 80.51]	75.58 [70.11, 80.79]	73.28 [66.11, 80.31]	74.28 [66.37, 80.50]	0.01	0.02
Race, n (%)										0.08	0.04
Asian	187 (7.24%)	14 (7.69%)	2 (7.14%)	1 (10.00%)	11 (7.64%)	173 (7.21%)	6 (5.88%)	5 (5.00%)	162 (7.37%)		
Black or African American	836 (32.37%)	76 (41.76%)	12 (42.86%)	3 (30.00%)	61 (42.36%)	760 (31.65%)	41 (40.20%)	33 (33.00%)	686 (31.20%)		
White or Caucasian	1418 (54.90%)	84 (46.15%)	14 (50.00%)	6 (60.00%)	64 (44.44%)	1334 (55.56%)	46 (45.10%)	53 (53.00%)	1235 (56.16%)		
Other	97 (3.76%)	7 (3.85%)	0 (0.00%)	0 (0.00%)	7 (4.86%)	90 (3.75%)	7 (6.86%)	9 (9.00%)	74 (3.37%)		
American Indian/Alaska Native	6 (0.23%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	6 (0.25%)	0 (0.00%)	0 (0.00%)	6 (0.27%)		
Other Pacific Islander	8 (0.31%)	1 (0.55%)	0 (0.00%)	0 (0.00%)	1 (0.69%)	7 (0.29%)	2 (1.96%)	0 (0.00%)	5 (0.23%)		
Not Disclosed/Unknown	31 (1.20%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	0 (0.00%)	31 (1.29%)	0 (0.00%)	0 (0.00%)	31 (1.41%)		
SVI	0.35 [0.22, 0.51]	0.36 [0.23, 0.54]	0.35 [0.30, 0.55]	0.23 [0.21, 0.34]	0.36 [0.23, 0.55]	0.35 [0.22, 0.50]	0.42 [0.25, 0.56]	0.34 [0.19, 0.56]	0.34 [0.22, 0.50]	0.15	0.06
VF Characteristics											
MD at Baseline	-2.54 [-5.38, -0.78]	-4.33 [-7.94, -2.01]	-9.19 [-13.59, -5.18]	-3.38 [-8.24, -0.98]	-3.83 [-6.63, -1.81]	-2.46 [-5.19, -0.71]	-6.49 [-12.18, -2.64]	-2.63 [-4.81, -0.61]	-2.34 [-4.92, -0.67]	0	0
Slope of MD	-0.05 [-0.37, 0.25]	-1.43 [-1.98, -1.17]	-1.41 [-1.90, -1.22]	-1.32 [-1.52, -1.16]	-1.43 [-2.03, -1.17]	-0.00 [-0.28, 0.28]	-0.27 [-0.55, 0.10]	-0.09 [-0.41, 0.23]	0.01 [-0.26, 0.29]	0	0
Number of Visits in five year duration	6.00 [5.00, 7.00]	6.00 [5.00, 8.00]	6.00 [5.00, 8.00]	6.50 [5.00, 7.00]	6.00 [5.00, 8.00]	6.00 [5.00, 7.00]	6.00 [6.00, 7.75]	6.00 [5.00, 7.00]	6.00 [5.00, 7.00]	0	0
Clinical Characteristics											
IOP (mmHg)	17.00 [14.00, 20.00]	17.00 [14.00, 21.00]	18.00 [15.50, 22.25]	18.00 [14.25, 20.00]	16.50 [14.00, 20.00]	17.00 [14.00, 20.00]	18.00 [15.00, 22.00]	18.00 [15.75, 21.00]	17.00 [14.00, 20.00]	0.76	0.01
Visual Acuity (logMAR)	0.10 [0.00, 0.18]	0.18 [0.10, 0.30]	0.24 [0.10, 0.48]	0.10 [0.10, 0.18]	0.18 [0.00, 0.30]	0.10 [0.00, 0.18]	0.10 [0.00, 0.30]	0.10 [0.00, 0.30]	0.10 [0.00, 0.18]	0	0
Glaucoma Severity										0	0
Mild Glaucoma	2031 (78.63%)	117 (64.29%)	8 (28.57%)	7 (70.00%)	102 (70.83%)	1914 (79.72%)	47 (46.08%)	84 (84.00%)	1783 (81.08%)		
Moderate Glaucoma	357 (13.82%)	42 (23.08%)	10 (35.71%)	2 (20.00%)	30 (20.83%)	315 (13.12%)	28 (27.45%)	9 (9.00%)	278 (12.64%)		
Severe Glaucoma	195 (7.55%)	23 (12.64%)	10 (35.71%)	1 (10.00%)	12 (8.33%)	172 (7.16%)	27 (26.47%)	7 (7.00%)	138 (6.28%)		
OCT Characteristics											
Average RNFL Thickness (µm)	78.43 [68.72, 86.96]	74.24 [63.67, 83.98]	67.21 [56.93, 73.74]	67.03 [62.10, 83.90]	76.07 [64.91, 84.36]	78.67 [69.21, 87.32]	67.50 [60.13, 77.81]	79.07 [72.27, 86.37]	79.04 [69.98, 87.62]	0	0
Average Cup Disc Ratio	0.67 [0.57, 0.74]	0.72 [0.59, 0.79]	0.77 [0.65, 0.83]	0.72 [0.64, 0.80]	0.71 [0.59, 0.78]	0.66 [0.57, 0.74]	0.74 [0.64, 0.81]	0.66 [0.56, 0.72]	0.66 [0.56, 0.73]	0	0
RIM Area (mm²)	1.00 [0.82, 1.20]	0.94 [0.72, 1.20]	0.70 [0.57, 0.93]	0.89 [0.65, 1.06]	0.98 [0.77, 1.25]	1.00 [0.83, 1.20]	0.75 [0.66, 1.03]	1.04 [0.87, 1.21]	1.01 [0.84, 1.21]	0.01	0

*Continuous variables are presented as median [IQR].

A total of 2,583 eyes from 1,514 patients were included in this study. Table 1 lists key demographic and

clinical characteristics of rapid (n = 182 eyes, 7.0%) and non-rapid progressors (n = 2,401 eyes, 93.0%)

stratified by treatment received in the first seven years.

We first examined differences in demographic and social vulnerability characteristics between rapid and

non-rapid glaucoma progressors, emphasizing factors potentially influencing treatment decisions (Table

1). Rapid progressors differed significantly from non-rapid progressors in racial distribution (p = 0.04),

with a notably higher proportion of African-derived patients among rapid progressors (41.76% vs.

31.65%). Rapid progressors were also significantly older compared to non-rapid progressors (median age

76.7 vs. 74.3 years, p = 0.01). However, no significant differences were observed in biological sex (p =

0.69) or Social Vulnerability Index (SVI) (see Table 1 for full details).

Regarding baseline clinical characteristics, rapid progressors demonstrated significantly worse glaucoma

severity distribution (p < 0.001), characterized by greater proportions of moderate (23.08% vs. 13.12%)

and severe glaucoma cases (12.64% vs. 7.16%). Rapid progressors and non-rapid progressors had the

same median number of visits (6.0), though the distribution was significantly different (p < 0.001).

Subsequent analysis of the distributions shows that the non-rapid progressor group had a more

pronounced tail than the rapid-progressor group (higher kurtosis), reflecting more outliers in visit

frequencies. Additionally, rapid progressors had significantly worse baseline visual field mean deviation

(MD: -4.33 vs. -2.46 dB, p < 0.001), poorer visual acuity (logMAR: 0.18 vs. 0.10, p < 0.001), and worse

optical coherence tomography (OCT) measures, including thinner retinal nerve fiber layer (RNFL)

thickness (74.2 µm vs. 78.7 µm, p < 0.001) and larger cup-to-disc ratios (0.72 vs. 0.66, p < 0.001). No

significant differences were identified in intraocular pressure (IOP) (see Table 1 for full details).

When we examined differences between the six progression-treatments groups (rapid/non-rapid

progressors and by treatment type), we found significant differences in all measured variables except

biological sex (p=0.69) and SVI (p=0.06). Subsequent pairwise analyses identified significant differences

in MD slope for each treatment type between rapid vs. non-rapid progressors (all p<0.001). We also find

that MD slope did not differ pairwise between all rapid progressor treatment groups, but differed between

all non-rapid progressor treatment groups. Treatment patterns followed disease severity at initial

presentation. Rapid progressors receiving aggressive procedures had significantly worse baseline MD

(−9.19 [−13.59, −5.18] dB) compared to all other groups (range of medians: −2.34 to −6.49 dB, all p <

0.001). Among non-rapid progressors, those with a worse baseline MD were more likely to receive

aggressive procedures (p < 0.001). Rapid progressors managed medically required significantly more

follow-up visits than non-rapid medical patients (p < 0.001), indicating closer monitoring despite

conservative management. Similarly, among non-rapid progressors, patients receiving aggressive

procedures attended more visits compared to minimally invasive or medical groups (both p ≤ 0.016).

Rapid progressors who received aggressive procedures had significantly worse visual acuity (logMAR

0.24 vs. 0.10; p=0.004) and thinner RNFL thickness (medians 67.21 µm vs. 79.07 µm; p≤0.003)

compared to non-rapid progressors who received minimally invasive procedures or medical treatment,

highlighting worse baseline visual impairment and structural damage. Likewise, among non-rapid

progressors, aggressive procedures remained associated with poorer visual acuity and thinner RNFL

compared with minimally invasive or medical treatments (all p < 0.001), indicating that initial severity is

associated with aggressive procedures regardless of rapid progression status.

Table 2. Logistic regression for predictors of intervention (aggressive procedures, minimally

invasive procedures, or any procedure) in all patients.

Variable	Aggressive Procedures Odds Ratio (95% CI)	P-value	Minimally Invasive Procedures Odds Ratio (95% CI)	P-value	Any Procedure Odds Ratio (95% CI)	P-value
Age (per 1 year increase)	1.010 (0.994–1.026)	0.216	1.007 (0.991–1.023)	0.399	1.009 (0.997–1.021)	0.125
Gender (Male vs. Female)	0.751 (0.525–1.073)	0.116	0.748 (0.517–1.082)	0.124	0.758 (0.580–0.990)	0.042*
Race (Asian vs. White)	0.881 (0.401–1.936)	0.752	0.749 (0.316–1.778)	0.513	0.827 (0.457–1.499)	0.532

Race (Black or African American vs. White)	1.138 (0.730–1.774)	0.568	1.035 (0.648–1.654)	0.885	1.104 (0.789–1.544)	0.563
Race (Other vs. White)	1.395 (0.595–3.273)	0.444	2.536 (1.246–5.164)	0.010*	2.070 (1.163–3.685)	0.013*
SVI (per unit increase)	3.382 (1.223–9.350)	0.019*	0.494 (0.166–1.470)	0.205	1.280 (0.591–2.772)	0.532
Rapid Progressor Status	3.354 (2.122–5.302)	0.000*	1.546 (0.840–2.846)	0.162	2.742 (1.873–4.013)	0.000*
MD at First Visit (per dB decrease)	1.094 (1.059–1.130)	0.000*	0.984 (0.940–1.030)	0.481	1.059 (1.031–1.088)	0.000*
IOP (per mmHg increase)	1.037 (1.006–1.069)	0.017*	1.052 (1.019–1.085)	0.002*	1.049 (1.025–1.073)	0.000*
VA (per unit logMAR increase)	2.138 (1.230–3.714)	0.007*	0.763 (0.329–1.765)	0.527	1.506 (0.930–2.437)	0.096
Average Cup Disc Ratio (per unit increase)	0.212 (0.033–1.355)	0.101	0.613 (0.123–3.063)	0.551	0.397 (0.113–1.397)	0.150
Average RNFL Thickness (per µm increase)	0.982 (0.965–0.999)	0.033*	0.991 (0.975–1.009)	0.328	0.987 (0.974–0.999)	0.037*
Rim Area (per mm² increase)	0.142 (0.052–0.390)	0.000*	1.009 (0.413–2.462)	0.985	0.403 (0.200–0.811)	0.011*

* Females serve as the reference group for gender, and white patients serve as the reference group for race. All procedure groups

are compared with those not in the procedure as a reference group. Asterisks indicate statistically significant p-values ($p < 0.05$)

Next, we assessed which demographic, clinical, and socioeconomic factors were associated with different

intervention types (aggressive, minimally invasive, or any procedure) using logistic regression, first

across all patients (Table 2), then rapid progressors (Table 3), and finally non-rapid progressors (Table 4).

In the overall patient cohort (Table 2), male sex modestly reduced the odds of receiving any procedure

(odds ratio [OR]: 0.76, 95% confidence interval [CI]: 0.58–0.99, p = 0.042). Patients identified as a

race other than White, Black, or Asian had significantly increased odds of undergoing minimally invasive

procedures (OR: 2.54, 95% CI: 1.25–5.16, p = 0.010) and any procedure (OR: 2.07, 95% CI: 1.16–

3.69, p = 0.013). Higher (worse) social vulnerability as measured by SVI was also associated with

increased odds of aggressive procedures (OR: 3.38 per unit increase, 95% CI: 1.22–9.35, p = 0.019). In

contrast, age, Asian race, and Black or African American race did not significantly influence procedural

selection.

Among clinical predictors, rapid progression strongly increased the odds of receiving aggressive

procedures (OR: 3.35, 95% CI: 2.12–5.30, p < 0.001) and any procedure (OR: 2.74, 95% CI: 1.87–

4.01, p < 0.001). Worse visual field damage at baseline, as indicated by more negative MD, also

predicted aggressive procedures (OR: 1.09 per dB decrease, 95% CI: 1.06–1.13, p < 0.001) and any

procedure (OR: 1.06 per dB decrease, 95% CI: 1.03–1.09, p < 0.001). Elevated intraocular pressure

(IOP) at the first clinical visit was a consistent predictor across all intervention types—aggressive (OR:

1.04 per mmHg increase, 95% CI: 1.01–1.07, p = 0.017), minimally invasive (OR: 1.05 per mmHg

increase, 95% CI: 1.02–1.09, p = 0.002), and any procedure (OR: 1.05 per mmHg increase, 95% CI:

1.03–1.07, p < 0.001). Worse visual acuity (higher logMAR) significantly increased the likelihood of

aggressive interventions (OR: 2.14 per unit increase, 95% CI: 1.23–3.71, p = 0.007). On structural

evaluation, thinner average retinal nerve fiber layer (RNFL) thickness (OR: 0.98 per μm increase, 95%

CI: 0.97–1.00, p = 0.033) and larger rim area (OR: 0.14 per mm² increase, 95% CI: 0.05–0.39,

p < 0.001) were significantly associated with lower odds of aggressive procedures. Average cup-to-disc

ratio did not significantly influence procedural likelihood.

Table 3. Logistic regression for predictors of intervention (aggressive procedures, minimally

invasive procedures, or any procedure) in rapid progressors.

Variable	Aggressive Procedures Odds Ratio (95% CI)	Aggressive Procedures P-value	Minimally Invasive Procedures Odds Ratio (95% CI)	Minimally Invasive Procedures P-value	Any Procedure Odds Ratio (95% CI)	Any Procedure P-value
Age (per 1 year increase)	1.005 (0.964–1.047)	0.826	1.025 (0.964–1.090)	0.428	1.007 (0.973–1.043)	0.672
Gender (Male vs. Female)	1.453 (0.600–3.516)	0.407	0.770 (0.209–2.837)	0.694	1.183 (0.552–2.534)	0.665
Race (Asian vs. White)	1.300 (0.222–7.605)	0.771	0.768 (0.078–7.515)	0.820	1.118 (0.257–4.858)	0.882
Race (Black or African American vs. White)	1.156 (0.353–3.791)	0.811	1.722 (0.368–8.048)	0.490	1.346 (0.496–3.652)	0.560
SVI (per unit increase)	6.667 (0.441–100.787)	0.171	0.006 (0.000–0.625)	0.031*	0.751 (0.073–7.753)	0.810
MD at First Visit (per dB decrease)	1.136 (1.040–1.242)	0.005*	0.982 (0.860–1.122)	0.794	1.099 (1.016–1.188)	0.018*
IOP (per mmHg increase)	1.095 (1.011–1.186)	0.026*	1.019 (0.900–1.154)	0.761	1.089 (1.017–1.168)	0.015*

VA (per unit logMAR increase)	1.554 (0.469–5.149)	0.471	0.034 (0.000–2.930)	0.137	0.965 (0.325–2.868)	0.949
Average Cup Disc Ratio (per unit increase)	0.050 (0.000–6.231)	0.224	1.103 (0.003–413.396)	0.974	0.152 (0.003–8.208)	0.355
Average RNFL Thickness (per µm increase)	0.999 (0.962–1.038)	0.965	1.019 (0.964–1.077)	0.515	1.005 (0.973–1.038)	0.750
Rim Area (per mm² increase)	0.050 (0.003–0.749)	0.030*	0.141 (0.005–3.910)	0.248	0.062 (0.007–0.564)	0.014*

* Females serve as the reference group for gender, and white patients serve as the reference group for race. All procedure groups

are compared with those not in the procedure as a reference group. Asterisks indicate statistically significant p-values ($p < 0.05$)

Among rapid progressors (Table 3), we found that key predictors from the overall patient cohort remained

consistent. Elevated IOP at the first clinical visit continued to significantly increase the likelihood of

receiving aggressive procedures (OR 1.10 per mmHg increase, 95% CI: [1.01, 1.19], $p < 0.03$) and any

procedure (OR 1.09 per mmHg increase, 95% CI: [1.02, 1.17], $p < 0.02$). Worse initial MD significantly

correlated with higher odds of aggressive procedures (OR 1.14 per dB decrease, 95% CI: 1.04–1.24,

$p = 0.005$) and any procedure (OR 1.10 per dB decrease, 95% CI: 1.02–1.19, $p = 0.018$). Smaller rim

area significantly increased the odds of aggressive procedures (OR 0.05 per mm² increase, 95% CI:

0.003–0.75, $p = 0.030$) and any procedure (OR 0.06 per mm² increase, 95% CI: 0.007–0.564,

$p = 0.014$).

Higher SVI significantly reduced the likelihood of minimally invasive procedures (OR 0.006 per unit

increase, 95% CI: 0.000–0.63, $p = 0.031$), suggesting socioeconomic factors might limit access to

minimally invasive options among rapid progressors. Higher SVI also tended to be associated with

aggressive procedures in this cohort, although this did not reach statistical significance (OR 6.67 per unit

increase, 95% CI: 0.441–100.787, $p = 0.171$). Demographic factors such as age, sex, and race, as well

as baseline visual acuity and average cup-to-disc ratio did not reach statistical significance.

Table 4. Logistic regression for predictors of intervention (aggressive procedures, minimally

invasive procedures, or any procedure) in non-rapid progressors.

Variable	Aggressive Procedures Odds Ratio (95% CI)	Aggressive Procedures P-value	Minimally Invasive Procedures Odds Ratio (95% CI)	Minimally Invasive Procedures P-value	Any Procedure Odds Ratio (95% CI)	Any Procedure P-value
Age (per 1 year increase)	1.011 (0.993–1.029)	0.234	1.005 (0.989–1.023)	0.533	1.009 (0.996–1.021)	0.183
Gender (Male vs. Female)	0.659 (0.442–0.982)	0.041*	0.756 (0.512–1.115)	0.159	0.713 (0.534–0.952)	0.022*
Race (Asian vs. White)	0.794 (0.327–1.927)	0.610	0.644 (0.253–1.641)	0.356	0.729 (0.379–1.401)	0.343
Race (Black or African American vs. White)	1.082 (0.670–1.745)	0.748	0.893 (0.551–1.448)	0.646	0.988 (0.693–1.408)	0.946
SVI (per unit increase)	2.805 (0.925–8.510)	0.069*	0.713 (0.230–2.213)	0.558	1.326 (0.580–3.027)	0.504
MD at First Visit (per dB decrease)	1.088 (1.050–1.129)	0.000*	0.984 (0.937–1.034)	0.524	1.055 (1.024–1.086)	0.000*
IOP (per mmHg increase)	1.030 (0.995–1.066)	0.091*	1.055 (1.021–1.091)	0.001**	1.046 (1.020–1.073)	0.000*
VA (per unit logMAR increase)	2.204 (1.157–4.198)	0.016**	1.114 (0.459–2.700)	0.812	1.717 (1.001–2.944)	0.050*
Average Cup Disc Ratio (per unit increase)	0.197 (0.033–1.172)	0.074	0.819 (0.175–3.840)	0.800	0.516 (0.149–1.783)	0.296
Average RNFL Thickness (per μm increase)	0.978 (0.959–0.997)	0.024*	0.991 (0.973–1.009)	0.316	0.984 (0.971–0.998)	0.025*
Rim Area (per mm² increase)	0.169 (0.055–0.521)	0.002*	1.301 (0.520–3.254)	0.574	0.560 (0.265–1.181)	0.128

* Females serve as the reference group for gender, and white patients serve as the reference group for race. All procedure groups

are compared with those not in the procedure as a reference group. Asterisks indicate statistically significant p-values (p < 0.05).

Among non-rapid progressors (Table 4), predictors of intervention were also similar to the overall cohort

but varied in magnitude. Male sex was significantly associated with lower odds of aggressive procedures

(OR 0.659, 95% CI: 0.442–0.982, p = 0.041) and any procedure (OR 0.713, 95% CI: 0.534–0.952,

p = 0.022). Higher initial IOP significantly predicted minimally invasive procedures (OR 1.055 per

mmHg increase, 95% CI: 1.021–1.091, p = 0.001) and any procedure (OR 1.046 per mmHg increase,

95% CI: 1.020–1.073, p < 0.001). Better baseline MD (less negative) was significantly associated with

lower odds of aggressive procedures (OR 1.088 per dB decrease, 95% CI: 1.050–1.129, p < 0.001) and

any procedure (OR 1.055 per dB decrease, 95% CI: 1.024–1.086, p < 0.001). Smaller optic disc rim

area (OR 0.169 per mm² increase, 95% CI: 0.055–0.521, p = 0.002) and thinner RNFL thickness (OR

0.978 per μm increase, 95% CI: 0.959–0.997, p□=□0.024) significantly increased odds for aggressive

procedures, reinforcing their clinical significance. No significant differences were observed for age, race,

average cup-to-disc ratio, SVI, or visual acuity.

Sensitivity Analysis

When shortening the timeframe for treatment after initial clinical diagnosis from seven years to five and

three years, some predictors changed in their statistical significance. Male sex and socioeconomic

vulnerability were significant predictors in the seven-year analysis for both aggressive procedures and

minimally invasive procedures (e.g., seven-year odds ratio for male gender for either procedure = 0.76,

p□= 0.04). However, they often lost statistical significance in the three-year models (e.g., three-year odds

ratio =□0.76, p□=□0.153).

In contrast, elevated baseline IOP showed a consistent positive association with the likelihood of

aggressive and minimally invasive procedures. For example, each additional mmHg of IOP increased the

odds of aggressive procedures by roughly 4–5% in both 3-year (OR□=□1.04, p□<□0.02) and 5-year

models (OR□=□1.05, p□<□0.01), and maintained similar associations at 7 years (OR□=□1.04,

p□<□0.02). Therefore, despite narrowing the treatment window, baseline IOP and key functional and

structural measures of disease severity (worse MD, smaller rim area) remained the most robust predictors

of both early and later interventions.

Finally, the primary predictors of aggressive procedures, particularly disease severity, remained consistent

across all models and stayed statistically significant across all timeframes. For instance, a better mean

deviation at the first visit continued to be a strong predictor of lower odds of aggressive procedures (e.g.,

OR = 1.085, 95% CI: 1.041–1.131 at 3 years, p = 0.0001; OR = 1.091, 95% CI: 1.050–1.131 at 5 years, p

= 0.000004; OR = 1.099, 95% CI: 1.064–1.136 at 7 years, p < 0.001). Similarly, the optic disc rim area

consistently demonstrated large effect sizes for predicting aggressive procedures (e.g., OR□≈□0.09–0.22

across models, p □<□0.02), indicating that patients with smaller rim areas were more likely to undergo

early and more invasive procedures.

Discussion

In this study, we evaluated the clinical and socioeconomic determinants of treatment decisions in

glaucoma patients with different visual field progression rates. Despite finding that rapid progression was

associated with greater odds of receiving either aggressive procedures (OR □3.35) or any procedure

(OR □2.74), only 20% of rapidly progressive patients underwent any non-medical procedure, of which

15% were aggressive IOP lowering procedures within seven years of their initial visit. Among patients

who progressed rapidly, aggressive procedures were more commonly pursued among those who had

higher baseline intraocular pressures and worse functional or structural parameters. Furthermore, a higher

social vulnerability index was associated with a lower likelihood of receiving less aggressive therapy,

which highlights the potential role of socioeconomic factors in shaping treatment decisions.

While previous studies[3,4] have reported a 5-12% rapid progression in glaucomatous patients, to our

knowledge, this is the first study that specifically reports treatment patterns in rapid progressors and how

these may vary from non-rapid progressors. In our cohort of 2,583 eyes from 1,514 patients, only 15% of

rapidly progressing eyes ultimately underwent aggressive procedures within seven years of diagnosis. We

also found that there was no significant pairwise difference for MD slope between treatment groups for

rapid progressors – in other words, if there was a faster rate of MD worsening, it was not associated with

more aggressive treatment within the rapid progressor cohort. This relatively low percentage and similar

MD slopes between treatment categories for rapid progressors is clinically significant and may not be

intuitive, as rapid progression typically raises clinical concerns warranting aggressive interventions to

preserve visual function. Potential reasons for this observed gap include clinician reliance on baseline

disease severity markers (e.g., intraocular pressure, structural OCT findings) over functional progression

rates, patient hesitancy or refusal of invasive procedures, perceived surgical risks, socioeconomic barriers

limiting access to interventions, or clinician preference for conservative management due to concerns

about complications. Specifically, in the overall cohort and rapid progressors, eyes with a higher IOP,

worse MD, smaller optic disc rim area or thinner RNFL at presentation were significantly more likely to

undergo aggressive procedures or any procedure. When we restricted the analysis to rapid progressors

alone, the association between structural and functional damage as a predictor of aggressive procedures

remained robust, and for certain variables such as baseline MD and IOP, the odds ratios were even greater

in this subgroup. Our data suggest that rapid progression, as defined retrospectively from at least five VF

tests, may not always align with initial clinical impressions or decisions which may be based on baseline

disease severity or IOP. Since clinicians may not have access to longitudinal VF progression data when

initially deciding on treatment escalation, our results underscore the potential value of incorporating

standardized and timely measures of VF loss rate earlier in clinical decision-making processes to better

identify and manage patients who may rapidly progress or incorporating models which can forecast rapid

progression risk from early data points[17, 18]. This observation highlights a potential gap in aggressive

intervention among patients demonstrating rapid functional decline, as most rapidly progressing patients

did not receive definitive aggressive procedures. Future studies could prospectively assess how earlier

identification and consideration of rapid VF progression rates could influence treatment escalation

decisions in clinical practice.

Across the entire population, patients with worse (higher) SVI were more likely to undergo aggressive

procedures, and patients with worse (higher) SVI who were rapid progressors had a markedly reduced

likelihood of receiving minimally invasive procedures (OR = 0.006, p = 0.031). This may reflect more

advanced disease at presentation or reduced access to healthcare in earlier stages. Prior research has

consistently found that higher neighborhood-level socioeconomic strongly correlates with delayed

diagnosis or inconsistent follow-up, which may lead to more urgent need for definitive procedures once

severe damage is detected[8, 9, 10, 12]. Although we adjusted our analysis for baseline disease severity, our

results also highlight a significant association between socioeconomic vulnerability and the types of

glaucoma interventions patients received, underscoring potential disparities in access or utilization of

different treatment modalities. This suggests potential unmeasured confounding variables, from disease

severity factors associated with SVI as well as a possible direct causal impact of SVI on treatment

selection. These findings align with previous literature demonstrating disparities in access to minimally

invasive procedures among different demographic groups[19].

In line with previous literature, we find that demographic characteristics also shaped treatment

approaches[8,12,20]. Male patients were less likely to undergo either aggressive or minimally invasive

procedures in the overall cohort (OR = 0.758, p = 0.042), and less likely to receive aggressive procedures

or any procedure in non-rapid progressors (OR = 0.659 for aggressive procedures; OR = 0.714 for any

procedure with both p < 0.05). Additionally, individuals who did not identify as Asian, African American,

or Caucasian were more frequently treated with minimally invasive procedures, suggesting that racial and

ethnic identity may be related to different treatment preferences, referral patterns, or provider perceptions.

There are several limitations to this study. This was a retrospective, single-center study in an academic

setting, which may limit the generalizability of the results to other practice environments or diverse

patient populations. Treatment decisions and outcomes were drawn from clinical data analyzed

retrospectively, and unmeasured confounders such as patient preferences, medication adherence, and

clinician and clinic-specific practice patterns may have influenced both the selection of therapy and

observed progression rates, especially when considering the effect of socioeconomic status. We also

relied on administrative CPT and ICD-10 codes to capture both procedures and diagnoses, which may

have possible misclassifications, and repeated interventions in the seven-year follow-up period post initial

diagnoses were not explicitly accounted for. Furthermore, the SVI used in this study provides a census-

tract level estimate of socioeconomic status which is necessarily a coarse measure, and may therefore not

have fully encapsulated the individual level social determinants of health that can affect treatment access

and decision-making. Finally, our inclusion criteria of requiring at least five reliable visual fields within

five years and one reliable OCT within one year of first visual field may have excluded patients with

more severe disease, limited follow-up or suboptimal testing, which may have introduced selection bias

into our results, as such patients may have rapidly progressed had they not had a procedure earlier in their

disease course.

386 Conclusion

In conclusion, this study demonstrates that although rapid visual field progression was associated with

greater likelihood of receiving aggressive or minimally invasive treatments, fewer than one in five rapidly

progressing eyes received aggressive intervention within seven years of initial presentation. Furthermore,

our data suggest that socioeconomic factors may play a pivotal role, especially in the reduced uptake of

minimally invasive treatment in patients with a worse SVI, highlighting the need to address disparities in

healthcare access and utilization. Future research efforts should focus on refining risk stratification tools,

potentially integrating the current or forecasted rate of VF progression more prominently into treatment

algorithms, and mitigating socioeconomic barriers so we can optimize outcomes for patients with rapidly

progressing glaucoma.

References

1. Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol*. 2006;90(3):262-267.

2. Tham Y-C, Li X, Wong TY, Quigley HA, Aung T, Cheng C-Y. Global prevalence of glaucoma and projections of glaucoma burden through 2040. *Ophthalmology*. 2014;121(11):2081-2090.

3. Jackson AB, Martin KR, Coote MA, et al. Fast progressors in glaucoma: prevalence based on global and central visual field loss. *Ophthalmology*. 2023;130(5):462-468.

4. Huang X, Poursoroush A, Sun J, et al. Identifying factors associated with fast visual field progression in patients with ocular hypertension based on unsupervised machine learning. *J Glaucoma*. 2024;33(11):815-822.

5. Chauhan BC, Garway-Heath DF, Goñi FJ, et al. Practical recommendations for measuring rates of visual field change in glaucoma. *Br J Ophthalmol*. 2008;92(4):569-573.

6. Kim JH, Rabiolo A, Morales E, et al. Risk factors for fast visual field progression in glaucoma. *Am J Ophthalmol*. 2019;207:268-278.

7. Prata TS, De Moraes CGV, Teng CC, et al. Factors affecting rates of visual field progression in glaucoma patients with optic disc hemorrhage. *Ophthalmology*. 2010;117(1):24-29.

8. Shaheen A, Medeiros FA, Swaminathan SS. Association between greater social vulnerability and delayed glaucoma surgery. *Am J Ophthalmol*. 2024;268:123-135.

9. Tseng VL, Kitayama K, Yu F, Coleman AL. Disparities in glaucoma surgery: a review of current evidence and future directions for improvement. *Transl Vis Sci Technol*. 2023;12(9):2.

10. Tseng VL, Kitayama K, Yu F, Pan D, Coleman AL. Social vulnerability, prevalence of glaucoma, and incidence of glaucoma surgery in the California Medicare population. *Ophthalmol Glaucoma*. 2023;6(6):616-625.

11. Almidani L, Bradley C, Herbert P, et al. The impact of social vulnerability on structural and

functional glaucoma severity, worsening, and variability. *Ophthalmol Glaucoma.* 2024;7(4):380-

390.

12. Musa I, Bansal S, Kaleem MA. Barriers to care in the treatment of glaucoma: socioeconomic

elements that impact the diagnosis, treatment, and outcomes in glaucoma patients. *Curr*

Ophthalmol Rep. 2022;10(3):85-90.

13. Folgar FA, De Moraes CGV, Prata TS, et al. Glaucoma surgery decreases the rates of localized

and global visual field progression. *Am J Ophthalmol.* 2010;149(2):258-264.

14. Yohannan J, Cheng M, Da J, et al. Evidence-based criteria for determining peripapillary OCT

reliability. *Ophthalmology.* 2020;127(2):167-176.

15. Yohannan J, Wang J, Brown J, et al. Evidence-based criteria for assessment of visual field

reliability. *Ophthalmology.* 2017;124(11):1612-1620.

16. Schulze-Bonsel, K., Feltgen, N., Burau, H., Hansen, L., & Bach, M. (2006). Visual acuities "hand

motion" and "counting fingers" can be quantified with the freiburg visual acuity test.

Investigative ophthalmology & visual science, *47*(3), 1236-1240.

17. Herbert P, Hou K, Bradley C, et al. Forecasting risk of future rapid glaucoma worsening using

early visual field, OCT, and clinical data. *Ophthalmol Glaucoma.* 2023;6(5):466-473.

18. Shuldiner SR, Boland MV, Ramulu PY, et al. Predicting eyes at risk for rapid glaucoma

progression based on an initial visual field test using machine learning. *PLoS One.*

2021;16(4):e0249856.

19. Vasu P, Hall RP, Wagner IV, et al. Racial disparities in microinvasive glaucoma surgery for

management of primary open-angle glaucoma: a propensity-matched cohort study. *Am J*

Ophthalmol. 2025;271:96-103.

20. Allison K, Hodges B, Shahid MM, Feng C. Racial and gender disparities for glaucoma treatment

rates in Upstate New York. *J Clin Med.* 2024;13(23):7225.

451