Vasopressin and angiotensin II differentially modulate human fear response dynamics to looming threats

Authors

Mengfan Han[1,2], Wenyi Dong[1,2], Kun Fu[1,2], Junjie Wang[1,2], Yuanhang Xu[1,2], Yueyuan Zheng[3], Keith M Kendrick[2], Ferraro Stefania[2], Ting Xu[4], Dezhong Yao[2], Benjamin Becker[1,5]*

Affiliations

[1] The Center of Psychosomatic Medicine, Sichuan Provincial Center for Mental Health, Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China

[2] MOE Key Laboratory for Neuroinformation, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China

[3] Division of Social Science, Hong Kong University of Science and Technology, Hong Kong, China

[4] Faculty of Psychology, Southwest University, Chongqing, China

[5] Department of Psychology, The University of Hong Kong, Hong Kong, China

* Corresponding author

E-mail: ben_becker@gmx.de (BB)

Abstract

While basal threat processing dynamics (e.g., visual looming) are well characterized in animals, the underlying mechanisms and their modulation by neuropeptide systems with different modulatory roles in threat processing (vasopressin, angiotensin II) remain poorly understood in humans. In a randomized, placebo-controlled eye-tracking study (N=111), we administered vasopressin (AVP) or an angiotensin II receptor blocker (via losartan, LT) during a time-to-collision threat paradigm. Behaviorally, AVP induced a systematic time overestimation while LT induced temporal compression and reduced state anxiety. Pupillometry revealed distinguishable profiles: AVP induced sustained constriction during stimulus approach followed by post-stimulus threat-specific dilation, LT maintained sustained pupillary constriction throughout both approach and occlusion phases yet preserving threat-specificity, while placebo (PLC) showed no threat-specific modulation. A computational framework (combining functional principal component analysis [FPCA], clustering, and hidden Markov model [HMM]) underscored the distinct modulations: AVP stabilized a high-arousal state characterized by the co-activation of vigilance, threat-proactive preparation and a shift from perception to internal simulation. LT suppressed transitions to high-arousal states and exhibited maximal sequence entropy, reflecting flexible response patterns—contrasting with placebo's lowest entropy dynamics. These results demonstrate that AVP and LT differentially regulate basal threat processing via separable neuropeptide pathways: AVP sustains hypervigilance while LT promotes anxiolysis and adaptive flexibility. Our findings suggest neuropeptide pathway-specific targets maladaptive threat processing in trauma- or anxiety-related disorders.

KeyWords

Threat, looming, vasopressin, angiotensin, pupillary dynamics, eye-tracking, phase-based pupillary analysis, functional linear mixed model (FLMM), functional principal component analysis (FPCA), hidden Markov model (HMM)

Abbreviations

LT, losartan; PLC, placebo; AVP, arginine vasopressin; RAS, renin-angiotensin system; LC-NE, locus coeruleus-norepinephrine system; aTTC, actual time to collision; jTTC, judged time to collision; PSV, physical stimulus velocity; SBP, systolic blood pressure; DBP, diastolic blood pressure; Early/Late Imag, early/late imagination phase; FLMM, functional linear mixed model; FPCA, functional principal component analysis; PC, principal component; HMM, hidden Markov model

Introduction

The subjective experience of emotions varies across individuals and requires the coordinated involvement of distributed cortical and subcortical systems [1–3]. Some basic forms of 'primitive' affective mechanisms such as defensive reactions to rapidly approaching objects ('looming') appear to be innate, highly preserved across phylogenetic and ontogenetic evolution and tightly linked to specific (subcortical) circuits [4,5]. The cognitive and neural bases of the corresponding affective processes have been extensively studied within broader models of emotion. Foundational work has highlighted the central role of subcortical structures, including the amygdala, in the rapid and automatic detection of motivationally salient stimuli, particularly threats [6,7]. This rapid

threat detection system interacts with cortical networks to generate conscious emotional

experiences and coordinate adaptive behavioral responses [8].

In interaction with these foundational mechanisms, the visual system can rapidly detect

impending – and potentially threatening - collisions via estimating the accelerating expansion of

objects on the retina (looming). Looming presents a fundamental, evolutionary and highly

conserved threat detection mechanisms, which can be observed from fruit flies to humans [9,10].

This innate perceptual sensitivity to approaching stimuli that specifies the time-to-collision

independent of size or distance enables avoiding collisions and rapid defensive responses [11–13].

Although looming perception has been traditionally viewed as a basic visual computation,

recent studies have demonstrated that looming is modulated by the semantic and affective content

of the stimuli, such that threatening stimuli are typically underestimated in the time to collision by

20-30%, a time compression effect that may critically facilitate faster defensive reactions [13–15].

However, in natural environments, threatening objects such as predators often temporarily

disappear from view, raising critical questions about how threat processing mechanisms operate

without continuous visual input and how imagination maintains threat assessment during such

periods. Understanding these mechanisms is crucial for developing ecologically valid models of

threat processing and may help to determine how these processes can be targeted in mental

disorders characterized by excessive threat processing.

The neuropeptide systems arginine vasopressin (AVP) and renin-angiotensin (RAS) have

been associated with threat-related processing across species. AVP, synthesized in the

hypothalamus, enhances threat processing by increasing the extended amygdala sensitivity to

threat information and modulating prefrontal-amygdala circuits [16,17], with pronounced effects

in males [18]. The RAS, in particular blockade of the RAS II AT1 receptors, has been associated

with decreased fear by reducing amygdala reactivity to threat and enhancing prefrontal regulatory

control [5,19–21]. While both systems influence threat processing through partially overlapping

neural circuits and show complex synergistic effects in regulating physiological indices of threat

responsivity including blood pressure and stress responses [22–24] their respective roles in

looming threat processing remain to be explored.

Pupillary responses offer unique advantages as precise and non-invasive indicators of threat

processing dynamics [25–27]. Through tracking the activity of the locus coeruleus-norepinephrine

(LC-NE) system pupil dilation provides high temporal resolution information about arousal

related to cognitive and affective engagement [28,29], As such, pupil responses can reveal the

continuous temporal dynamics of threat processing.

The LC-NE arousal system has shown a high sensitivity to both AVP [30] and RAS [31]

modulation which render pupillary responses an ideal indicator for investigating the

neurobiological basis of their modulatory role on threat processing.

Against this background we aimed at determining the role of two neuropeptide systems (AVP,

RAS) in basal threat-related mechanisms, in particular to determine their effects on threat

processing during looming via determining the effects of single dosages of both, AVP and a

selective and competitive angiotensin receptor II antagonists (Losartan, LT) on behavioral and

pupillary responses during the looming fear paradigm in combination with computational analytic

approaches. This novel combination allows us to dissect the distinct contributions of these systems

to threat processing and explore potential therapeutic implications. The hypothesized effects were

tested in a pre-registered randomized placebo-controlled double-blind psychopharmacological eye

tracking design during which participants were administered oral AVP, LT or placebo (PLC).

To provide an accurate test of the complex hypotheses we established a novel expansion of

the threat looming paradigm and established a comprehensive analytic framework. This expansion

integrates both visual and imagination components, comprising two phases: a one-second 'visual

stimulus amplification' phase providing initial approach velocity information, followed by an

'imagined stimulus approach' phase where the visual threat stimulus disappears after the approach

information has been acquired. This design builds on substantial evidence that threat-related

neural systems remain active even without direct visual input [8,32], and amygdala-centered

defensive circuits continue processing threats when visual stimuli are subliminal or masked

[33,34], indicating robust threat maintenance mechanisms during temporary visual occlusion.

To characterize the complex dynamics of threat processing, we developed an integrated

analytical framework combining pupillary features, behavioral responses, and temporal

measurements. This multivariate approach identifies distinct threat response patterns and their

pharmacological modulation through dimensional reduction and clustering techniques. By

simultaneously analyzing multiple pupillary characteristics and behavioral measures, we reveal

how different pharmacological treatments shape overall threat processing strategies. This

systematic approach advances beyond traditional univariate analyses to capture the temporal

evolution and multidimensional nature of threat responses under different pharmacological

conditions.

Our specific aims in this study were to: (1) characterize threat processing dynamics during

looming through pupillary responses, (2) examine the maintenance of threat perception during

visual occlusion, (3) investigate pharmacological modulation of the identified threat processes,

and (4) identify distinct response patterns through Functional Principal Component

Analysis–clustering–Hidden Markov Model (FPCA–clustering–HMM) analysis. The findings may

reveal innovative targets for testing novel interventions for disorders characterized by elevated

threat reactivity and have important implications for how neuropeptides shape defensive

behaviors.

Methods

1. Participants and Study Design

1.1 Participants

Following evaluated recruitment criteria, 111 healthy volunteers (n = 37 per treatment group)

were included in the final analysis (details see CONSORT flowchart in S1 Fig). Sample size was a

priori determined using G*Power for a Treatment (3) × Sex (2) × IsThreaten (2) × Physical

Stimulus Velocity (5 levels, from very slow to very fast; see Section 3.3 for detailed classification)

mixed-design ANOVA, resulting in a required sample size of 78 (f = 0.20, α = 0.05, power =

0.95).

Eligible participants had normal or corrected-to-normal vision, no history of psychiatric or

neurological disorders, and no current medication or substance use (see Supplementary Method S1

for full details). The study was approved by the local ethics committee of University of Electronic

Science and Technology of China (UESTC), conducted according to the Declaration of Helsinki,

and pre-registered (ClinicalTrials.gov ID: NCT06329063, NCT06329076). All participants

provided written informed consent and received monetary compensation (140 RMB).

In a randomized, double-blind, placebo-controlled pharmacological behavioral and

eye-tracking design, participants were randomly assigned to receive either AVP, LT, or PLC (mean age ± SD: 22.33 ± 2.22 years; no significant age differences between groups, $F(2,108) = 2.13$, $p = .12$).

2. Experimental Procedures

2.1 Baseline Assessments

Prior to drug administration, participants completed validated Chinese versions of standardized questionnaires assessing affective states and fear-related traits, including the Positive and Negative Affect Schedule (PANAS) [35]; State-Trait Anxiety Inventory (STAI) [36]; Liebowitz Social Anxiety Scale (LSAS) [37]; and Animal Fear Questionnaire (AFQ) [38]. Blood pressure and heart rate were monitored at three time points: pre-administration, peak drug effect (45 min post-administration for AVP group, 90 min post-administration for LT group), and post-experiment (similar in [21,39], **Fig 1A**) to control for unspecific cardiovascular effects of the treatments.

2.2 Drug Administration

Treatment doses were selected based on previous research demonstrating efficacy in modulating emotional processing including fear-related processes: 20 IU AVP [39] and 50 mg LT [19–21].

We employed a novel spray-based oral delivery method for AVP [40] to prevent direct entry into the olfactory system and better isolate peripheral effects [41]. Following our validated protocol, participants received six alternating 0.1 ml puffs (total volume 0.6 ml, Supplementary

Method S2). Placebo sprays were identical in composition but without AVP. Losartan (LT) and its

placebo were administered in visually matched capsules containing either 50 mg Losartan or no

active ingredient.

To ensure effects of both treatments during peak plasma windows (based on established

pharmacokinetic profiles showing peak effects at 45 min for AVP and 90 min for LT) and maintain

double-blinding during tasks, we implemented a two-stage administration protocol:

• Group 1: 50 mg LT capsule (90 min pre-task) + PLC spray (45 min pre-task)

• Group 2: PLC capsule (90 min pre-task) + PLC spray (45 min pre-task)

• Group 3: PLC capsule (90 min pre-task) + 20 IU AVP spray (45 min pre-task)

All treatments were administered by research staff blind to treatment conditions. Successful

blinding was confirmed, as participants' post-experiment treatment guesses did not exceed chance

level ($\chi^2(2) = 0.83$ $p = .36$).

2.3 Experimental Task

We employed a validated threat looming paradigm [13] using color photographs of

threatening (snakes, spiders) and non-threatening (butterflies, rabbits) animals (40 images per

category). Each trial (**Fig 1B**) began with a fixation cross that disappeared at 300 ms, followed by

a looming stimulus that expanded at a constant velocity and disappeared at 1300 ms. After its

disappearance, participants imagined its continued approach and pressed a key at their judged

moment of collision. The judged Time-to-Collision (jTTC) was measured from stimulus onset

(300 ms) to this key press. A confirmation beep was followed by a jittered inter-trial interval

(1000-3000 ms). The initial stimulus size (20% or 30% of screen width) and its expansion velocity

(determined by one of five actual Time-to-Collision (aTTC) values: 3.0, 3.5, 4.0, 4.5, and 5.0 s)

were both varied independently across trials.

The experiment consisted of 160 trials across two blocks, comprising a full factorial design

of aTTC (5 levels), stimulus category (4 levels), and initial size (2 levels). Participants completed

practice trials using animal silhouettes prior to the main task. The task lasted approximately 35

minutes, and pupillary responses were recorded throughout.

205
206   **Fig 1. Experimental timeline and task structure. (A) Sequence of drug administration and behavioral**
207   **testing. (B) Single trial structure of the threat looming task.**

3. Data Acquisition and Preprocessing

3.1 Behavioral Data Preprocessing

211   Trial-level jTTC data were preprocessed in two stages. First, trials with technical artifacts or

212   incomplete responses were excluded. Second, trials exceeding ±5 SD from each participant's mean

213   jTTC were classified as outliers and removed. This conservative criterion preserved potential

214   treatment-induced variations while flagging extreme deviations, following methodological

215 recommendations for pharmacological studies [42,43].

3.2 Pupillary Data Preprocessing

Pupil size and movement from the right eye were recorded at 2000 Hz using an EyeLink

1000 Plus system (SR Research) with a display resolution of 1280 × 1024 pixels. Participants

were positioned 57 cm from the monitor using a chin rest, and a 9-point calibration procedure was

performed at the beginning of each experimental block to ensure optimal tracking accuracy.

Pupil size was recorded as pixel area and converted to diameter (d) using a validated formula

$d = \alpha L \varphi$, where α is an empirical scaling factor, L denotes the pupil-to-camera distance, and φ is

the visual angle [44]. Data were preprocessed to remove signal loss, blink artifacts,

physiologically implausible values, and statistical outliers beyond ± 5 SD from the trial mean, in

accordance with established guidelines [28]. Finally, the cleaned pupillary signals were

down-sampled into 10 ms bins to reduce high-frequency noise while preserving physiologically

relevant fluctuations [45].

3.3 Stimulus Dynamics

231 To optimize analysis, we computed a composite metric—Physical Stimulus Velocity (PSV)—

232 integrating looming motion parameters into a single ecologically meaningful measure. Approach

233 velocities (179.2-341.3 pixels/s) were categorized into five intervals: very slow (V1: 179.2-211.6),

234 slow (V2: 211.6-244.0), medium (V3: 244.0-276.4), fast (V4: 276.4-308.8), and very fast (V5:

235 308.8-341.3 pixels/s, right-closed). These categories served as levels in subsequent analyses.

4 Analysis Framework

4.1 Behavioral Analysis

To quantify the psychophysical characteristics of time perception, we employed a model-based approach: (1) a Linear Model (jTTC = α + β * aTTC), which served as a baseline assuming a linear relationship between subjective and objective time; (2) a Power Law Model (jTTC = α * (aTTC)$^{\wedge}\beta$), based on Stevens' Power Law [46], which captures the canonical nonlinear relationship between perception and physical magnitude. In both models, α represents scaling parameters, while β reflects the rate of change (linear) or degree of nonlinearity (power law) in time perception.

For each participant and condition, model parameters (α, β) were estimated by iterative optimization, minimizing the sum of squared errors to obtain the best fit. The coefficient of determination (R^2) was computed for each fit. To evaluate which model best accounted for the data across the dataset, we compared them using the Bayesian and Akaike Information Criteria (BIC and AIC) [47].

Finally, to examine the effects of Treatment, IsThreaten, and Sex on time perception parameters, we constructed Linear Mixed-Effects models using the estimated parameters (α and β) as dependent variables. Each model included fixed effects of Treatment, Sex, IsThreaten, and their possible two-way interactions. A random intercept for each participant was included to account for individual differences. The significance of the fixed effects was analyzed to determine whether the experimental manipulations modulated the temporal perception parameters.

4.2 Pupil Analysis

(1) Baseline Pupil Analysis and Correction

Considering potential pharmacological effects on absolute pupil diameter, we first assessed baseline pupil diameter (0-300 ms) using a two-way ANOVA with Treatment and Sex as factors. All subsequent analyses were baseline-corrected by subtracting the mean baseline value from each timepoint to isolate task-evoked responses.

(2) Event-Locked Analysis

To compare mean pupillary responses across distinct phases of threat processing—from sensory encoding to decision preparation—we defined three temporal windows based on task structure and threat processing dynamics [32,48]:

(1). Stimulus Presentation (300-1300 ms; encoding of the approaching stimulus during active visual processing); (2). Early Imagination (first 500 ms post-offset; immediate transition to mental representation); (3). Late Imagination (last 500 ms pre-collision; decision-related preparatory processes).

A repeated-measures ANOVA was conducted with Treatment and Sex as between-subject factors, and Phase, IsThreaten, and Physical Stimulus Velocity (PSV) as within-subject factors. Trials shorter than 2300 ms (2%) were excluded to ensure complete phase coverage across all windows.

(3) Time-Normalized Dynamic Analysis

Variable trial durations—primarily due to individual differences in time-to-collision estimation—precluded direct comparison of pupillary dynamics across trials. Temporal normalization was therefore employed to align all trials to a common time scale. Following predefined quality criteria (Supplementary Methods S4), 9.65% of trials were excluded, with no

differences in the percentage between treatment groups ($\chi^2(2) = 0.46$, $p = .53$). The remaining

trials were processed using piecewise linear time warping to map the fixed stimulus epoch

(300-1300 ms) to [0, 0.5] and the variable imagination epoch to [0.5, 1.0] on a unified scale,

followed by smoothing of the normalized time series using cubic B-splines [49–51].

Functional linear mixed models (FLMM) were fitted [52] to examine the influence of

Treatment, Sex, IsThreaten, and PSV on pupillary response over normalized time. The initial

model followed a full factorial design:

Baseline-corrected Pupil Diameter ~ Treatment * Sex * IsThreaten * PSV *

Normalized_Time + (Normalized_Time + IsThreaten + PSV | Subject)

To ensure interpretability and avoid overparameterization, non-significant higher-order

interactions were pruned using likelihood ratio tests [53], resulting in a final model that retained

only effects up to three-way interactions.

Together, these analyses provide a comprehensive characterization of pupillary responses:

baseline correction isolated pharmacological effects on tonic arousal, event-locked ANOVA

identified phase-specific mean differences, and time-normalized FLMM captured full response

trajectories across task progress. This multi-faceted approach distinguishes nonspecific drug

effects, enables precise stage-specific inference, and reveals holistic temporal dynamics.

4.3 Integrated Pattern Analysis

To identify treatment-specific psychophysiological phenotypes that reflect the integrated

coordination of behavioral and pupillary systems during threat processing, we developed a

comprehensive analytical framework combining Functional Principal Component Analysis (FPCA)

[54], Clustering [55], and Hidden Markov Models (HMM) [56,57] (**Fig 2**). This approach captures

both continuous temporal dynamics and discrete state transitions, aiming to reveal emergent,

integrated phenotypes beyond simple correlations between systems. The analysis consisted of

three sequential steps:

(1) Functional Principal Component Analysis (FPCA)

Using FPCA, which better accounts for temporal continuity compared to traditional PCA [58],

we extracted principal temporal response patterns (eigenfunctions) from time-normalized

pupillary data within each trial. This dimensional reduction approach yielded multiple functional

components per trial that collectively characterized the complete temporal dynamics of pupillary

responses. We retained principal components that explained over 85% of total variance (individual

components >5%) and examined their associations with experimental conditions, individual

characteristics, and pupillary features during and after stimulus presentation.

(2) Hierarchical Clustering Integrated with jTTC and FPCA Components

To classify trials into distinct response patterns based on their behavioral-physiological

characteristics, we integrated the FPCA components with behavioral features (jTTC) using a

hierarchical k-means clustering framework. Given their different measurement scales, we first

clustered on log-transformed jTTC, then on FPCA components within each behavioral subgroup.

Optimal cluster numbers were determined using Calinski-Harabasz index and silhouette

coefficients, with clustering stability verified through bootstrap resampling ($n = 1000$, stability

index > 0.85) and cross-validation (accuracy > 0.80). To further determine whether these

behavioral subgroups (trial-level) reflected stable computational traits (subject-level), we

correlated the individual participants' proportion of trials in each subgroup with their fitted power

law parameters (α and β).

(3) Hidden Markov Modeling for Transition Dynamics

First-order Hidden Markov Models (HMMs) were applied to model transitions between

response profiles across trials. Analysis encompassed three complementary aspects: First, the

steady-state distribution was compared across experimental conditions using chi-square tests

supplemented by standardized residual analysis; Second, transition probability matrices were

estimated to capture state dynamics, with bootstrap methods (n = 1000) identifying significant

transition patterns within each treatment group; Finally, the complexity of sequential pattern

transitions was quantified through entropy measures applied to continuous three-state sequences

across different experimental manipulations.

This integrated approach fuses continuous pupillary dynamics with behavioral data,

identifying distinct, treatment-modulated response profiles and characterizing their temporal

evolution during threat processing.

Fig 2. Analysis workflow. First, separate behavioral and pupillometric analyses were conducted. Behavioral
responses were characterized using the power-law model jTTC = jTTC = α * (aTTC)^β. Pupillary analysis
included baseline pupil size, event-locked responses, and time-normalized dynamics. **Second,** features were
integrated through a hierarchical approach: trials were clustered based on jTTC values, with subgroup
proportions correlated to subject-level parameters (α, β). FPCA reduced dimensionality of pupillary data,
followed by Hidden Markov Modeling (HMM) to characterize state dynamics, including steady-state
distributions, transition matrices, and sequence entropy.

4.4 Analytic Software and Correction

Behavioral analyses and event-locked phase analyses were conducted using SPSS 26. Time-normalized dynamic analyses and integrated pattern analyses were implemented in Python v3.12 using scientific computing libraries (pandas, numpy, scipy, statsmodels, scikit-learn) with custom implementations for functional data analysis. For all analyses, effects were considered significant at $p < .05$ with Greenhouse-Geisser correction for sphericity violations. Multiple comparisons were controlled for false discovery rate using the Benjamini-Hochberg method [59] ($\alpha = 0.05$).

Results

1 Subjective Anxiety and Potential Confounders

The three groups (LT, PLC, and AVP; $n = 37$ each) did not differ in age, mood, animal fear, or personality traits (all ps > .13, S1 Table). A 2 (Treatment: LT vs. PLC) × 2 (Sex: male vs. female) × 2 (Time: pre vs. post) mixed ANOVA with State Anxiety as the dependent variable revealed a significant Treatment × Time interaction ($F = 4.11$, $p = .047$, $\eta_p^2 = 0.055$). Between-group comparison showed lower post-task state anxiety level in the LT group compared to PLC ($p = .012$, $d = 0.51$) which was mirrored in within-group analyses indicating significant anxiety reduction in the LT group (pre: 37.51 ± 1.10, post: 33.39 ± 1.36, $p = .005$, dz = 0.58) but not in the PLC group (pre: 38.44 ± 1.10, post: 38.36 ± 1.35, $p = .95$, dz = 0.02). This may suggest an anxiolytic effect of LT **(Fig 3A)**. No such pattern emerged for AVP (Treatment × Time, $p = .78$). Critically, the anxiety-reducing effect of LT did not differ by sex (ps > .41).

Cardiovascular parameters (systolic blood pressure, diastolic blood pressure, and heart rate)

369 were monitored at three time points: baseline, drug peak effect (80 minutes post-LT; 35 minutes

370 post-AVP), and post-task. Mixed ANOVAs revealed only a main effect of sex, with males showing

371 higher resting blood pressure than females (SBP: 118.17 ± 0.74 vs. 110.34 ± 0.74 mmHg, $p < .001$;

372 DBP: 73.66 ± 0.85 vs. 71.18 ± 0.85 mmHg, $p = .041$) **(Fig 3B).** No significant treatment effects or

373 interactions were found at any time point (all ps > .10).

2 Time-to-collision Analysis

Model comparisons revealed that the power law model provided a superior fit to the linear

model in most conditions (69.4%). The compression parameter β was consistently below 1 (range:

0.342-0.653, mean = 0.495), together with an average 1.55-fold increase in absolute perceptual

error as aTTC increased from 3s to 5s, indicating systematic time compression that accelerated

with longer aTTCs. Linear mixed-effects (LME) analyses revealed a significant reduction in β

(indicating enhanced time compression) for threatening compared to nonthreatening stimuli ((β =

-0.091, p = .06; **Fig 3C**), under LT compared to PLC (β = -0.149, p = .033; **Fig 3D1**), and in

females compared to males (β = -0.167, p = .016). No significant reduction in β was observed for

AVP compared to PLC (β = -0.045, p = .52).

The scaling parameter α (where α = 1 indicates accurate estimation) showed substantial

individual differences (range: 0.857-2.341, mean = 1.534). LME analyses revealed a trend toward

increased α (indicating a tendency for time overestimation) under AVP compared to PLC (β =

0.504, p = .078; **Fig 3D2**). No other effects reached significance.

Fig 3. Subjective anxiety, potential confounders, and behavioral results. (A) State anxiety scores pre- and post-experiment in the LT and PLC groups. The LT group showed a significant reduction in anxiety at post-test. **(B) Systolic (SBP) and diastolic (DBP) blood pressure measurements**, revealing consistently higher values in males than in females. Both panels are presented as raincloud plots, where points represent mean scores per participant, the central line in the box indicates the median, and the box bounds represent the interquartile range. **(C) Nonlinear fitting of aTTC and jTTC using the power-law model jTTC = α * (aTTC)^β.** Participants generally underestimated time-to-collision, with accelerated underestimation as aTTC increased, particularly under threat conditions. Points represent mean jTTC per participant (threat vs. non-threat), jittered (5%) to avoid overlap. The dashed line y = x indicates accurate estimation. **(D1) Parameter β** (temporal scaling: $\beta > 1$ indicates expansion, $\beta < 1$ indicates compression). LT group exhibited significantly stronger time compression than PLC. **(D2) Parameter α** (general estimation bias: $\alpha > 1$ indicates overestimation, $\alpha < 1$ indicates underestimation). AVP group showed a tendency toward overestimation compared to PLC ($p= .078$). ***$p < .05$,** $p < .01$, ***$p < .001$, † marginal significance.**

3 Pupil Analysis

3.1 Baseline Analysis

Analysis of pre-stimulus baseline pupil diameter revealed a significant main effect of treatment ($F = 4.31$, $p = .016$), with the AVP showing larger diameters than the PLC group ($p = .011$), the LT did not significantly differ from either group ($ps > .104$). No sex differences were observed ($ps > .2$).

This pharmacological effect persisted throughout the entire experiment, as confirmed by analysis of uncorrected data across trials ($F = 3.98$, $p = .023$; **Fig 4A**). After baseline correction, these general differences were eliminated ($ps > 0.12$) allowing examination of task-specific pupillary changes.

3.2 Event-locked Analysis

Analysis of baseline-corrected responses revealed significant main effects of Phase ($F = 41.97$, $p < .001$) and PSV ($F = 19.92$, $p < .001$). Pupils dilated from stimulus presentation to early imagination (Present vs Early: $p < .001$) and stabilized during late imagination (Early vs Late: $p = .22$). Extreme velocities (V1/V5) elicited larger responses than moderate velocities (V2-V4, all $ps < .01$).

A Phase × IsThreaten interaction ($F = 16.20$, $p < .001$; **Fig 4B**) revealed that threatening stimuli induced progressive dilation across phases (Stim < Early < Late, $ps < .05$), while non-threatening stimuli showed initial dilation followed by plateau (Stim < Early = Late, $ps < .01$). During late imagination, threat-induced pupil dilation exceeded non-threatening responses ($p = .004$).

A Phase × Treatment interaction ($F = 2.93$, $p = .022$; **Fig 4C**) showed that during early

imagination, both LT and AVP reduced responses as compared to PLC ($ps < .05$), while no group

differences were observed during stimulus presentation or late imagination ($ps > .22$). Across

phases, AVP group showed progressive dilation (Stim < Early Imag < Late Imag, $ps < .05$), LT and

PLC groups showed initial dilation followed by plateau (Stim < Early Imag ≈ Late Imag, ps

$< .01$).

The IsThreaten × Sex interaction ($F = 18.513$, $p < .001$; **Fig 4D**) revealed that males showed

larger responses than females to non-threatening stimuli ($p = .003$), but no sex differences under

threat ($p = .84$). Females exhibited enhanced responses to threatening versus non-threatening

stimuli ($p < .001$), while males' responses remained consistent across both conditions ($p = .10$).

3.3 Time-Normalized Dynamic Analysis

Extending event-locked findings, functional linear mixed models on baseline-corrected pupil

diameter across normalized time revealed more nuanced patterns.

Overall pupil dilation increased with normalized time ($\beta=0.010$, $p=.030$) and threat-induced

constriction ($\beta=-0.016$, $p< .001$). PSV demonstrated a V-shaped velocity-dependent pattern, with

maximal dilation at extreme velocities (V1/V5: $\beta = 0.016/0.018$, $ps<.001$), moderate dilation at V2

($\beta = 0.005$, $p<.001$), and no change at V4 relative to V3 ($p = .63$).

Treatment × IsThreaten × Normalized Time interaction ($\beta s = 0.015\text{-}0.024$, $ps < .05$; **Fig**

4E1,E2) showed that during stimulus approach (t=0-0.5), all groups exhibited threat-induced

constriction ($ps<.01$), but after stimulus offset (t=0.5-1), threat processing diverged: PLC showed

no threat/non-threat difference while both LT and AVP maintained enhanced dilation under threat

versus nonthreat ($ps < .001$). Notably, between-group comparisons revealed that the LT group

maintained overall smaller response than PLC, whereas pupil response was smaller in the AVP

compared to the PLC group during stimulus presentation but showed enhanced dilation

post-stimulus specifically for threat ($\beta = 0.008$, $p < .001$).

A Treatment × IsThreaten × PSV interaction (**Fig 4F**) indicated that the PLC group showed

the largest overall responses compared to both AVP and LT groups (β s = 0.017-0.041, ps < .05).

AVP group showed larger responses to threatening versus non-threatening stimuli across all

velocities (V1-V5: βs = 0.0028-0.0099, ps < .001), and LT selectively increased threat responses

only at faster velocities (V3-V5: β = 0.005-0.0097, ps < .001). PLC group showed no

velocity-specific threat sensitivity. All groups maintained the V-shaped velocity effect, with drug

groups reaching their lowest response at V4 while the PLC group reached its lowest response at

V3.

A significant Sex × IsThreaten × Normalized Time interaction ($\beta = 0.025$, $p < .001$; **Fig 4G**)

emerged. During stimulus approach, males showed larger pupil response than females across both

threat and nonthreat conditions (ps < .001). After stimulus offset, however, the sex difference

persisted only for non-threatening stimuli ($\beta = 0.0221$, $p < .001$) and was abolished under threat

(p=.27). Crucially, females exhibited greater dilation to threatening versus non-threatening stimuli

post-stimulus ($\beta = -0.0193$, $p < .001$), while males showed no threat-related differentiation

(p=.31).

Fig 4. Pupillary Measurement: Baseline (A), Phase-Locked (B-D) and Normalized Time Dynamic
Analyses (E-G). (A)Baseline pupil size: The AVP group had a significantly larger uncorrected pupil
diameter than the PLC group. The LT group did not differ from either. **(B) Phase × Isthreaten:** Threat
evoked a sustained dilation, while non-threat elicited a phasic dilation followed by a plateau. Responses to
threat significantly exceeded non-threat in the late imagination phase. **(C) Phase × Treatment:** The PLC
group response was significantly larger than both AVP and LT during the early imagination phase. The AVP
group exhibited sustained dilation across phases, while PLC and LT responses plateaued from early to late
imagination. **(D) Sex × Isthreaten:** Males had significantly larger pupils than females specifically during the
non-threat condition (no sex difference under threat). The inset shows a significantly greater threat-specific
response (threat - non-threat) in females. **(E1) AVP vs. PLC × Isthreaten × Time:** PLC > AVP during
stimulus approach; this pattern reversed to AVP > PLC after stimulus offset under threat. **(E2) LT vs. PLC ×**
Isthreaten × Time: PLC > LT across time. After offset, threat > non-threat for LT, while PLC showed no
threat-non-threat difference. **(F) Treatment × Isthreaten × PSV:** Pupillary velocity showed a V-shaped
profile. The nadir occurred earlier for PLC (V3) than for drug groups (V4). AVP showed threat > non-threat
from V1-V5, LT from V3-V5, while PLC showed no differential sensitivity. **(G) Sex × Isthreaten × Time:**
Male > female during stimulus approach; this difference disappeared after stimulus offset under threat. *** *p***
< .05, ** *p* < .01, ***p < .001.**

4 Integrated Behavioral-Pupillary Response Profiles

Eight distinct response profiles were identified through a hierarchical clustering procedure.

First, clustering trials based on log jTTC revealed two behavioral subgroups: an early-response

group (E: 3.12 ± 0.57 s) and a late-response group (L: 5.80 ± 1.98 s). Correlations with power law

parameters confirmed associations with distinct subject-level traits in time estimation: the

early-response proportion negatively correlated with both α ($r = -0.45$, $p < .001$) and β ($r = -0.36$,

$p < .001$), indicating systematic underestimation and stronger time compression; conversely, the

late-response proportion was positively correlated with both α ($r = 0.45$, $p < .001$) and β ($r = 0.36$,

$p < .001$), indicating the opposite pattern of overestimation and weaker compression.

Second, FPCA decomposed pupillary dynamics into three core components (PC1-3, **Fig 5A**)

accounting for 87.51% of total variance, which were differentially modulated by experimental

conditions and individual traits (summarized in **Table 1**; full analysis in Supplementary Results:

Functional Principal Components, S2-S3 Fig). Based on the full analyses the components were

interpreted as PC1: Sustained Vigilance, PC2: Proactive Engagement, reflecting active preparation

for threat, and PC3: Cognitive Shift, capturing the transition from perception to imagination.

Integrating behavioral (E/L) and pupillary (PC1-3 scores) features, a second-level clustering

yielded eight stable profiles (E1-E4, L1-L4, **Fig 5B**), verified by bootstrap (stability index: 0.753

± 0.125) and cross-validation (0.651 ± 0.114; S4 Fig). These profiles reflect distinguishable threat

processing strategies (Table 2):

For instance, profile E1 was characterized by dominant sustained vigilance (high PC1) with

suppressed proactive engagement and cognitive shift (low PC2/PC3). Profile E2 was characterized

by co-activated vigilance, engagement, and cognitive shift (high PC1/PC2/PC3). Notably,

late-response profiles L2-L4 shared the same directional patterns of PC scores as E2-E4 but

differed in response magnitude, as shown in their reconstructed responses (**Fig 5C**).

Table 1 summarizes the characteristic features and experimental correlations of each component, along with their proposed functional significance.

Principle Component	PC1	PC2	PC3
Variation Explanation	62.67%	17.56%	7.28%
Weight Time Function Features	Early steep rise followed by plateau	Initial inhibition, later strong activation	Early peak, mid-period trough, late recovery
Main Related Pupil Features	Stimulus-Evoked Response (Pupil size during stimulus: r=.615, speed of dilating: r=.65)	Post-Stimulus Sustenance (Pupil size after stimulus: r=.668) vs. Stimulus Suppression (Pupil size during stimulus: r=-.35)	Rate Anticorrelation (Slower dilating during stimulus: r=-.58, Faster dilating after: r=.34)
Experimental Conditions	Female only: Nonthreatening: PLC>LT,PLC>AVP Threatening: PLC=AVP>LT Velocities: V1,V5>V2-V4 (No male differences)	AVP>LT, AVP>PLC Threat>Nonthreat V1,V2<V3-V5	Threat>Nonthreat V1,V5<V2-V3
Individual Differences	not significant	Low anxiety > High Anxiety	Early response < Late response
Possible Functional Significance	Sustained Vigilance and Attentional Resource Allocation	Proactive Preparation and Active Threat Engagement	Dynamic Transformation from Perception to Imagination

512 **Fig 5. Decomposition and profiling of integrated behavioral-pupillary responses. (A) Three dominant**

functional principal components (PCs) underlying pupillary dynamics. Eigenfunctions $\varphi(t)$ for each component are displayed, with their sign and magnitude defining the functional interpretation: PC1 (Sustained Vigilance) shows sustained positive weighting; PC2 (Proactive Engagement) transitions from negative to positive weighting; PC3 (Cognitive Shift) exhibits a multiphasic (positive-negative-positive) pattern. Individual trial responses are reconstructed as a linear combination of these components: Pupil Diameter(t) $\approx$ Grand Mean(t) + (Score$_1 \times \varphi_1$(t)) + (Score$_2 \times \varphi_2$(t)) + (Score$_3 \times \varphi_3$(t)). **(B) Eight distinct response profiles from hierarchical clustering.** Profiles were identified by combining behavioral response subgroups (Early: E1 – E4; Late: L1 – L4) with clustering of PC. **(C) Reconstructed pupillary responses for each profile.** Curves illustrate the characteristic pupillary signature of each profile over time, reconstructed by combining the shared temporal eigenfunctions from (A) with the profile-specific component scores from (B).

Table 2. Characteristics of the eight behavioral-physiological profiles.

Pattern	Behavioral Association	PC1	PC2	PC3	Interpretation
E1	underestimation tendency (low α); time compression (low β)	+	-	-	High vigilance, low engagement, low cognitive shift
E2		+	+	+	Co-activation of vigilance, engagement, cognitive shift
E3		-	+	+	Low vigilance, high engagement, high cognitive shift
E4		-	+	-	Low vigilance, high engagement, low cognitive shift
L1	overestimation tendency (high α); time expansion (high β)	+	-	+	High vigilance, low engagement, high cognitive shift
L2		+	+	+	Co-activated (similar to E2)
L3		-	+	+	Low vigilance, high engagement, high cognitive shift (similar to E3)
L4		-	+	-	Low vigilance, high engagement, low cognitive shift (similar to E4)

4.3 HMM Analysis

(1) Steady State Distribution

Residual analysis revealed a significant main effect of treatment ($\chi^2(14) = 333.17$, $p < .001$; **Fig 6A**). **Profile E1** (characterized by approach-phase dilation and post-stimulus constriction) was a dominant state that showing a significantly higher probability in PLC ($z = 8.75$) and a lower probability in AVP ($z = -9.85$). **Profile L3** (enhanced constriction during approach phase followed

by attenuated constriction post-stimulus) defined the pharmacological interventions, exhibiting significantly higher probability in both AVP $(z = 4.96)$ and LT $(z = 3.53)$ groups compared to PLC $(z = -8.32)$. **Profile L2** (sustained dilation) further distinguished the two active treatments, being higher in AVP $(z = 3.57)$ and lower in LT $(z = -2.69)$ **(see Fig 5 for each profile's visual breakdown)**.

The sex effect $(\chi^2 (7) = 250.84, p < .001;$ S6 Fig) showed males demonstrated elevated engagement in late-response profiles (L1 – L4; $z = 1.81$ – 6.86) while females showed reduced engagement $(z = -1.92$ to $-7.27)$. Specifically, profile L1 was higher in males $(z = 6.86)$ and lower in females $(z = -7.27)$, while profile E3 was higher in females $(z = 7.60)$ and lower in males $(z = -7.17)$. The Isthreaten effect $(\chi^2 (7) = 65.59, p < .001)$ revealed profile E1 exhibited lower engagement under threat $(z = -3.17)$ than non-threat $(z = 3.11)$, and profile E2 exhibited higher engagement under threat $(z = 4.43)$ than non-threat $(z = -4.35)$.

(2) Dynamics Transition

Analysis revealed treatment-specific remodeling of state transition pathways, which occurred despite consistently high self-transition probabilities (0.45 – 0.63) across all conditions (see S7 Fig for full statistics).

Specifically, the PLC group showed significantly increased probabilities in transitions towards E1 (e.g., E3→E1: $z = 4.65$; E4→E1: $z = 2.49$; all $ps < .05$). Conversely, the AVP group significantly suppressed transitions toward E1 (e.g., E1→E1: $z = -4.30$; E4→E1: $z = -3.40$) while promoting transitions involving late-response profiles (e.g., E4→L2: $z = 2.92$; L3→L3: $z = 2.83$). The LT group specifically exhibited reduced transitions toward the L2 profile (e.g., E2→L2: $z = -2.32$; E4 → L2: $z = -1.99$). Within-group transitions **(Fig 6B)** further supported these

between-group findings and provided a more detailed architecture of significant pathways within

each treatment (see Supplementary S2-S3 Table for full details, 95% CIs, and pairwise

comparisons).

Sex comparisons (χ^2 = 250.85, p-adj < .001) revealed males exhibited increased transitions

toward L1 (E1 → L1: z = 2.75; E3 → L1: z = 2.33) while females showed elevated transitions

toward E3 (E1 → E3: z = 2.32; L4 → E3: z = 3.18). A significant threat modulation was also

identified, enhancing the E3 → E2 transition under threatening conditions (χ^2 = 14.95, p-adj

= .007).

(3) Sequence Entropy and Stability

Analysis of sequential patterns (**Fig 6C**) provided further evidence for treatment-induced

destabilization of cognitive states, extending beyond transition probabilities to higher-order

sequence inconsistency.

PLC group demonstrated the most predictable and stable sequences, exhibiting the lowest

Markov entropy (M = 7.76 bits, 95% CI [7.70, 7.82]), which consistent with its tendency to

stabilize around repetitive, E1-centric sequences (e.g., E1→E1→E1, z = 5.13, p < .001).

Conversely, both pharmacological treatments significantly increased sequence entropy (i.e.,

made sequences more random and less predictable), but did so through distinct mechanisms:

LT group produced the highest entropy (M = 8.02 bits, 95% CI [7.96, 8.08]), significantly

exceeding both AVP and PLC (both p-adj < .001), indicating a generalized state of maximal

inconsistency without promoting any specific alternative sequences.

AVP group exhibited intermediate entropy (M = 7.85 bits, 95% CI [7.80, 7.91]), significantly

higher than PLC (p-adj < .001) but lower than LT (p-adj < .001). This reflects a partial disruption

 of the stable PLC dynamics, coinciding with its shift toward sustained E3 and L3 patterns.

577   Sex comparisons revealed significantly lower sequence entropy in females compared to

578 males (diff = -0.05 bits, t = 42.55.13, p < .001); Threatening stimuli induced a subtle but

579 significant reduction in sequence entropy compared to non-threatening conditions (diff = -0.02

580 bits, t = 20.13, p < .001), suggesting increased predictability of state sequences under threat.

581 (Detailed analysis of specific 3-state sequences is provided in Supplementary Table S4.)

582
Fig 6. Hidden Markov Model (HMM) analysis of response patterns. (A) Stationary distribution
revealed treatment-specific deviations from expected state probabilities. Profile E1 was significantly
elevated in PLC and reduced in AVP. Profile L2 was increased in AVP and decreased in LT. Profile L3 was
elevated in both LT and AVP but suppressed in PLC. Profile L4 was significantly lower in PLC (* p < .05, **
p < .01, *** p < .001). **(B) Significant within-group transition pathways (permutation test, p < .05).** Red
arrows and bolded probabilities indicate significant transitions. Grey arrows denote non-significant
transitions. Node labels show self-transition probabilities. Integration with between-group comparative
analyses revealed the following key treatment effects on transition dynamics: **PLC** reinforced Profile E1
(sustained vigilance) as a dominant attractor state, through high self-persistence and convergent inflows from
other states. **AVP** disrupted transitions toward E1 while facilitating transitions toward states of full
component co-activation (e.g., L2) or threat-specific engagement (e.g., E2E3). **LT** selectively inhibited the
emergence of the co-activated L2 state, as evidenced by a lack of significant convergent transitions
compared to other treatments. **(C) Sequence entropy** differed significantly across treatments (LT > AVP >
PLC), indicating that PLC produced the most predictable and stable sequences, whereas LT sequences were
the most dynamic and unpredictable.

Discussion

Our study investigated the behavioral and pupillary dynamics underlying the processing of looming threats — both during sensory approach and internal imagination — and how these processes are modulated by the neuropeptides vasopressin (AVP) and angiotensin II (LT). Using a randomized, placebo-controlled design integrating behavior and eye-tracking, we reveal that looming threat evaluation unfolds through distinct temporal stages and is modulated by neuropeptide signaling, which regulate processes of sustained vigilance and adaptive flexibility.

Our findings highlight three aspects of looming threat evaluation and its modulation:

(1) threat-induced time compression: we confirmed that the collision time for looming stimuli is underestimated (see also [13]) and our computational modeling revealed that the underestimation of collision time for threating stimuli (vs. nonthreat) is based on an accelerated rate of temporal compression (β), rather than a threat-induced modulation of constant bias (α). This threat-induced time-compression was further modulation by sex.

(2) dynamic pupillary responses during approaching threat and occlusion: a dynamic biphasic pupil response pattern characterized by sustained constriction during the visual approach (reaching minimum at stimulus disappearance), transitioning towards a release of constriction to progressive dilation prior to the expected (imagined) collision, which may reflect a dynamic interplay between vigilance and defensive responses.

(3) neuropeptide-specific modulation: AVP and LT exerted opposing on neural-behavioral states. AVP prolonged time perception by stabilizing hyper-vigilant states (e.g., L2) and increasing transitions into these states. LT, in contrast, promoted time compression and flexibility by suppressing transitions into L2 and increasing state-sequence entropy,

621  suggesting a calmer processing mode.

Threat-Induced Time Compression of Looming Threat

624  Building on and confirming earlier findings demonstrating that threat induces an

underestimation of collision time [13], our computational modeling further revealed that this

effect arises from an acceleration in the rate of temporal compression (β) — particularly for

longer aTTCs — without changes in the overall constant bias (α). This indicates that threat

modulates the gain of the estimation process rather than its starting point [60,61]. This pattern may

reflect a selective increase in the pace of evidence accumulation to support expedited action，

which align with existing theoretical frameworks proposing that motivational states influence

decision-making model parameters [62,63], and indicate that threat enhances processing efficiency

(drift rate) rather than lowering response caution (decision criterion) during looming.

633  Furthermore, pupillary responses revealed distinct temporal dynamics and threat-sensitive

modulation. During the visual approach phase, threat elicited a sustained constriction that peaked

at stimulus disappearance. This progressive constriction may reflect increasingly active content

decoding, contributed to by a cognitively driven near reflex as participants attempted to foveate

and resolve the details of the looming stimulus. However, the sustained, time-locked nature of

the constriction suggests it this response may additionally reflect an integrated engagement of

cognitive and oculomotor systems when evaluating looming stimuli [64,65]. Following stimulus

offset, pupils transitioned to a progressive dilation, potentially reflecting a shift from external

sensory processing to internal simulation. This dilation may suggest increasing autonomic arousal

and anticipatory defense activation as the imagined collision nears [66,67] and active maintenance

and estimation of the stimulus and its trajectory in working memory.

Critically, approach velocity significantly modulated these pupillary responses, exhibiting a

V-shaped pattern across speeds. Maximum constriction occurred at intermediate velocities (V2 for

threat, V4 for non-threat), while both the slowest (V1) and fastest (V5) velocities elicited greater

dilation. We hypothesize that this pattern reflects a shift in defensive strategies based on spatial

imminence [68]: detailed sensory analysis is prioritized at intermediate speeds via constriction

[28], whereas extreme velocities associate with a shift in arousal under conditions of either low

(V1) or high perceived relevance (V5) [69,70]. These findings underscore a continuous interaction

between looming and semantic content in shaping visual and threat responses, aligning with

broader models on changes in threat processing along the imminence spectrum [34,68].

The semantic content-related processing differences were further supported by results from

the pattern analysis such that nonthreatening stimuli preferentially engaged Profile E1 whereas

threatening stimuli robustly recruited E2, as evidenced by: (1) a higher-than-expected steady-state

distribution in their respective profiles, (2) distinct patterns in the transition matrices, and (3)

lower sequential entropy for the E1-dominated non-threat stimuli, indicating a more stable,

deterministic state. Interpreting these profiles via principal component loadings **(Fig 5A-B; Table**

2) underscores a threat-related dual processing mode:

Non-threatening Profile E1 (+, -, -) reflects an efficient surveillance mode characterized by

sustained vigilance (PC1+) with negative engagement in proactive threat preparation (PC2-) or

cognitive state transformation (PC3-). The associated pupillary response **(Fig 5C)** — initial

expansion followed by contraction—is consistent with online monitoring and rapid disengagement

after classifying a stimulus as non-critical, reflecting an adaptive resource conservation strategy

[71].

Threat-related Profile E2 (+, +, +) constitutes a high-alert threat mode involving the

co-activation of vigilance (PC1+), proactive threat preparation (PC2+), and sustained internal

simulation (PC3+). Its persistent pupillary dilation **(Fig 5C)** reflects ongoing resource allocation to

threat anticipation, consistent with a state of heightened defensive readiness that facilitates

exaggerated threat simulation and perceptual preparation [72].

Critically, both profiles are associated with early responses (short jTTC) and correlate

negatively with α and β parameters, indicating that while E1 and E2 represent distinct

computational pathways, they both facilitate a common adaptive output to looming stimuli: a

behavioral tendency for underestimation of time-to-collision, which is further amplified in the

case of threatening stimuli. This bridges our trial- and group-level findings: a general time

compression (E1-driven) is significantly enhanced by threat (E2-driven).

678   Sex Differences in Threat Processing Strategies

Sex modulated the described threat processing architecture, such that females exhibited a

steeper rate of temporal compression (β: female < male), suggesting that women may adopt a

more precautionary decision-making policy, favoring accelerated defensive responses under threat

[73,74].

Pupillometric analyses suggested that while males demonstrated generally larger pupil

diameters, only females displayed a significant threat-induced differentiation (threat > non-threat)

following stimulus offset. This post-stimulus divergence in females may suggest enhanced

engagement of internal, defensive simulation processes for threatening content [69]. Sustained

pupil dilation – as observed in our experiment – has been associated with emotional arousal and

increased cognitive load during such internal representations [75].

The sex-differential processing of threat is further underscored by findings from the pattern

analysis (details see Supplementary Results: Sex Differences in Patterns and Fig 5A-C) which

suggest a **male processing strategy (Profile L1)** reflected by state of broad, shallow monitoring

that maintains readiness at the cost of detailed threat evaluation, resulting in delayed responses. In

contrast, the **female-biased strategy (Profile E3)** may facilitate targeted internal modeling of

threats after stimulus offset, supporting superior threat discrimination and promoting earlier,

precautionary responses.

Neuropeptide Modulation of the Looming Response

The Placebo Baseline: A Resource-Conserving Mode for Fading Threat Cues

Under placebo, the pupillary response to threatening stimuli was diminished relative to

non-threatening ones during approach, and this discrimination vanished after offset **(Fig 4E1-E2)**.

The integrated pattern was dominated by **Profile E1** (+, -, -), as evidenced by its high steady-state

probability **(Fig 6A)**, a bias for transitions into E1 from other states **(Fig 6B)**, and the lowest

sequential entropy (indicating high stability) **(Fig 6C)**. This profile is characterized by sustained

vigilance (PC1+) with low investment in proactive threat preparation (PC2-) or extended internal

simulation (PC3-), consistent with the reconstructed pupillary response of initial expansion

followed by pronounced contraction **(Fig 5A, PC weight functions; B, profile characteristics; C,**

reconstructed pupil response).

Vasopressin Promotes a Hyper-Driven, Enhanced Simulation Mode

Vasopressin (AVP) administration reconfigured threat processing towards a state of higher engagement. Behaviorally, this manifested as a significant time-to-collision overestimation ($\alpha >$ PLC). Pupillometric analyses revealed that AVP specifically enhanced and sustained threat-discriminative responses during the imagery phase. This pattern was not modulated by variations in velocity suggesting an internal process that was maintained across external variations. Although the peripheral administration of a peptide like AVP—with its debated blood-brain barrier permeability — raises questions about central bioavailability, the significant behavioral and pupillary effects observed here align with a growing body of evidence indicating that peripherally administered neuropeptides can exert central actions through direct as well as indirect mechanisms, such as vagal nerve activation or engagement of receptor-mediated transport systems (details see also [76–78]).

Hidden Markov modeling (HMM) further identified that AVP suppressed the baseline E1 mode while promoting stability in a high-arousal state (**Profile L2 +, +, +**) **(Fig 6A-B)**. Critically, L2 represents a co-activation of sustained vigilance and resource allocation (PC1+), proactive preparation for threat engagement (PC2+), and a strong dynamic transformation from perception to sustained internal simulation (PC3+) **(Fig 5A-B)**. The AVP-induced stabilization in this state may reflect a proactive engagement with the internal threat representation and may reflect that AVP enhances an internal 'threat simulation' mode underlying an active threat evaluation and leading to a subjectively elongated perception of time [79] and enhanced arousal [80]. The shift in the processing mode (partly) aligns with models describing adaptive decision-making under uncertainty [71].

Losartan Induces a more Controlled Processing Mode

LT reduced subjective state anxiety in the absence of changes in peripheral cardiovascular

activity. This pattern may reflect a – previously described - delayed peak efficacy of losartan on

cardiovascular function in normotensive subjects [20,21] and the differentiation of systems

underlying subjective experience and autonomic reactivity or threats [81,82]. While the central

blood-brain barrier penetration of losartan remains debated our findings are in line with previously

reported effects in human psychopharmacological studies and models suggesting direct effects

via blood-brain barrier crossing and/or via actions on circumventricular organs and

subsequent modulation of central autonomic and stress circuits, or peripheral effects

mediating the anxiolytic responses [20,21,83–85].

Behaviorally, LT led to temporal compression (β < PLC), indicating accelerated

decision-making. Pupillometrically, LT reduced overall arousal (smaller pupil responses) while

preserving post-stimulus threat discrimination─but only at high velocities, possible reflecting a

selective, efficient allocation of resources to imminent threats.

HMM further qualified these effects such that LT: (1) resulted in the highest sequential

entropy, indicating a highly flexible and unpredictable processing stream devoid of rigid state

sequences, and, (2) specifically suppressed transitions into the high-anxiety, high-simulation L2

state (+,+,+) that is characteristic of the AVP-driven response. Notably, while both LT and AVP

increased the steady-state probability of the proactive preparation state L3 (-,+,+), the direct

opposite of the PLC-dominant E1(+,-,-), LT's critical action might have been to prevent

maintaining in a hyper-vigilant state of L2.

Collectively, these results demonstrate that LT induces a more controlled mode, rather than

promoting a specific alternative profile. As such LT's threat modulation may be based on

preventing fixed responses [86] and enhancing flexibile proactive preparation (as reflected in the

increased steady-state probability of L3). Together this may promote efficient resource allocation,

accelerated time-perception (temporal compression), and an overall reduction in subjective

anxiety.

Commonalities of neuropeptide threat modulation

Despite the differences both neuropeptides showed a common effect of heightened pupil

response to threat versus non-threat stimuli specifically during the post-stimulus imaginary phase,

a modulation that was absent under placebo. This indicates that they modulate the internal

evaluation of threat in addition to processes driven by sensory input. AVP may operate through

potentiation and stabilizing a hyper-vigilant state (L2) leading to exaggerated threat engagement

and subjective time dilation. In contrast, blockade of the renin-angiotensin II system may promote

cognitive flexibility (highest entropy) by specifically suppressing transitions into the L2 state

promoting efficient threat processing and subjective time compression. These distinct modulatory

roles may render the systems as promising and distinct therapeutic targets, such that LT may

alleviate anxiety, affective and hyperarousal (see also [85,87]) while AVP may enhance threat

sensitivity.

Limitations

Our study has several limitations. First, while we analyzed pupillary and behavioral

responses, we did not examine their link to neural activation patterns. Future work could combine

pupillometry with neuroimaging to directly map these relationships. Second, although chromatic

differences existed between stimulus phases (colored vs. grayscale), our key findings—such as

threat-specific pupillary dilation and pharmacological modulation—were consistent across

matched conditions, suggesting cognitive-emotional rather than light reflex. Nevertheless, strictly

controlled visual properties in future designs would strengthen validity. Finally, our principal

component interpretations, while aligned with theoretical frameworks (e.g., arousal modulation, threat

detection networks), remain inferential. Empirical validation of these components' neural substrates is

needed.

Data Availability: All group-level data (including judged Time-To-Collision, anxiety ratings, and

blood pressure measurements) and normalized pupil data will be made publicly available upon

acceptance in the Open Science Framework (OSF) repository (DOI will be provided upon

publication).

Funding: This research was supported by the STI 2030-Major Projects, Grant No.

2022ZD0208500, Ministry of Science and Technology of China; Health and Medical Research

Fund Hong Kong (HMRF, 23243971), Seed funding for collaborative and single projects from

The University of Hong Kong.

References

1. Gao N, Wang H, Xu X, Yang Z, Zhang T. Angiotensin II induces cognitive decline and

anxiety-like behavior via disturbing pattern of theta-gamma oscillations. Brain Res Bull. 2021;174:

84–91. doi:10.1016/j.brainresbull.2021.06.002

2.   Liu X, Jiao G, Zhou F, Kendrick KM, Yao D, Gong Q, et al. A neural signature for the

subjective experience of threat anticipation under uncertainty. Nat Commun. 2024;15: 1544.

doi:10.1038/s41467-024-45433-6

3.   Taschereau-Dumouchel V, Kawato M, Lau H. Multivoxel pattern analysis reveals

dissociations between subjective fear and its physiological correlates. Mol Psychiatry. 2020;25:

2342–2354. doi:10.1038/s41380-019-0520-3

4.   Thieu MK, Ayzenberg V, Lourenco SF, Kragel PA. Visual looming is a primitive for human

emotion. iScience. 2024;27: 109886. doi:10.1016/j.isci.2024.109886

5.   Zhou F, Geng Y, Xin F, Li J, Feng P, Liu C, et al. Human extinction learning is accelerated by

an angiotensin antagonist via ventromedial prefrontal cortex and its connections with basolateral

amygdala. Biol Psychiatry. 2019;86: 910–920. doi:10.1016/j.biopsych.2019.07.007

6.   Ohman A. The role of the amygdala in human fear: Automatic detection of threat.

Psychoneuroendocrinology. 2005;30: 953–958. doi:10.1016/j.psyneuen.2005.03.019

7.   Adolphs R. Neural systems for recognizing emotion. Curr Opin Neurobiol. 2002;12: 169–177.

doi:10.1016/s0959-4388(02)00301-x

8.   LeDoux JE, Brown R. A higher-order theory of emotional consciousness. Proceedings of the

National Academy of Sciences. 2017;114: E2016–E2025. doi:10.1073/pnas.1619316114

9.   Card G, Dickinson M. Performance trade-offs in the flight initiation of drosophila. J Exp Biol.

2008;211: 341–353. doi:10.1242/jeb.012682

10.   Schiff W. Perception of impending collision: A study of visually directed avoidant behavior.

Psychological Monographs: General and Applied. 1965;79: 1–26. doi:10.1037/h0093887

11.   Yilmaz M, Meister M. Rapid innate defensive responses of mice to looming visual stimuli.

Current Biology. 2013;23: 2011–2015. doi:10.1016/j.cub.2013.08.015

12.   Cooper WE, Samia DSM, Blumstein DT. Chapter five - FEAR, spontaneity, and artifact in

economic escape theory: A review and prospectus. In: Naguib M, Brockmann HJ, Mitani JC,

Simmons LW, Barrett L, Healy S, et al., editors. Advances in the Study of Behavior. Academic

Press; 2015. pp. 147–179. doi:10.1016/bs.asb.2015.02.002

13.   Vagnoni E, Lourenco SF, Longo MR. Threat modulates perception of looming visual stimuli.

Current Biology. 2012;22: R826–R827. doi:10.1016/j.cub.2012.07.053

14.   Cole S, Balcetis E, Dunning D. Affective signals of threat increase perceived proximity.

Psychol Sci. 2013;24: 34–40. doi:10.1177/0956797612446953

15.   Brendel E, Hecht H, DeLucia PR, Gamer M. Emotional effects on time-to-contact judgments:

Arousal, threat, and fear of spiders modulate the effect of pictorial content. Exp Brain Res.

2014;232: 2337–2347. doi:10.1007/s00221-014-3930-0

16.   Huber D, Veinante P, Stoop R. Vasopressin and oxytocin excite distinct neuronal populations

in the central amygdala. Science. 2005;308: 245–248. doi:10.1126/science.1105636

17.   Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M. Oxytocin and vasopressin in the

human brain: Social neuropeptides for translational medicine. Nat Rev Neurosci. 2011;12:

524–538. doi:10.1038/nrn3044

18.   Thompson RR, George K, Walton JC, Orr SP, Benson J. Sex-specific influences of

vasopressin on human social communication. Proc Natl Acad Sci U S A. 2006;103: 7889–7894.

doi:10.1073/pnas.0600406103

19. Reinecke A, Browning M, Klein Breteler J, Kappelmann N, Ressler KJ, Harmer CJ, et al.
Angiotensin regulation of amygdala response to threat in high-trait-anxiety individuals. Biol
Psychiatry Cogn Neurosci Neuroimaging. 2018;3: 826–835. doi:10.1016/j.bpsc.2018.05.007

20. Zhang R, Zhao W, Qi Z, Xu T, Zhou F, Becker B. Angiotensin II regulates the neural
expression of subjective fear in humans: A precision pharmaco-neuroimaging approach. Biol
Psychiatry Cogn Neurosci Neuroimaging. 2023;8: 262–270. doi:10.1016/j.bpsc.2022.09.008

21. Xu T, Zhou X, Jiao G, Zeng Y, Zhao W, Li J, et al. Angiotensin antagonist inhibits
preferential negative memory encoding via decreasing hippocampus activation and its coupling
with the amygdala. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging. 2022;7:
970–978. doi:10.1016/j.bpsc.2022.05.007

22. Szczepanska-Sadowska E. Role of neuropeptides in central control of cardiovascular
responses to stress. J Physiol Pharmacol. 2008;59 Suppl 8: 61–89.

23. Höhle S, Culman J, Boser M, Qadri F, Unger T. Effect of angiotensin AT2 and muscarinic
receptor blockade on osmotically induced vasopressin release. Eur J Pharmacol. 1996;300:
119–123. doi:10.1016/0014-2999(95)00855-1

24. von Bohlen und Halbach O, Albrecht D. The CNS renin-angiotensin system. Cell Tissue Res.
2006;326: 599–616. doi:10.1007/s00441-006-0190-8

25. Guo F, Zou J, Wang Y, Fang B, Zhou H, Wang D, et al. Human subcortical pathways
automatically detect collision trajectory without attention and awareness. PLOS Biology. 2024;22:
e3002375. doi:10.1371/journal.pbio.3002375

26. Armstrong T, Bilsky SA, Zhao M, Olatunji BO. Dwelling on potential threat cues: An eye
movement marker for combat-related ptsd. Depression and Anxiety. 2013;30: 497–502.
doi:10.1002/da.22115

27. Chen L, Yuan X, Xu Q, Wang Y, Jiang Y. Subliminal impending collision increases perceived
object size and enhances pupillary light reflex. Front Psychol. 2016;7.
doi:10.3389/fpsyg.2016.01897

28. Mathôt S, Van der Stigchel S. New light on the mind's eye: The pupillary light response as
active vision. Curr Dir Psychol Sci. 2015;24: 374–378. doi:10.1177/0963721415593725

29. Joshi S, Li Y, Kalwani RM, Gold JI. Relationships between pupil diameter and neuronal
activity in the locus coeruleus, colliculi, and cingulate cortex. Neuron. 2016;89: 221–234.
doi:10.1016/j.neuron.2015.11.028

30. Hernández-Pérez OR, Hernández VS, Nava-Kopp AT, Barrio RA, Seifi M, Swinny JD, et al.
A Synaptically Connected Hypothalamic Magnocellular Vasopressin-Locus Coeruleus Neuronal
Circuit and Its Plasticity in Response to Emotional and Physiological Stress. Front Neurosci.
2019;13. doi:10.3389/fnins.2019.00196

31. Gong W, Lü J, Wang F, Wang B, Wang M, Huang H. Effects of angiotensin type 2 receptor
on secretion of the locus coeruleus in stress-induced hypertension rats. Brain Research Bulletin.
2015;111: 62–68. doi:10.1016/j.brainresbull.2014.12.011

32. Mobbs D, Hagan CC, Dalgleish T, Silston B, Prévost C. The ecology of human fear: Survival
optimization and the nervous system. Front Neurosci. 2015;9. doi:10.3389/fnins.2015.00055

33. Mineka S, Öhman A. Phobias and preparedness: The selective, automatic, and encapsulated
nature of fear. Biological Psychiatry. 2002;52: 927–937. doi:10.1016/S0006-3223(02)01669-4

34. Phelps EA. Emotion and cognition: Insights from studies of the human amygdala. Annual
Review of Psychology. 2006;57: 27–53. doi:10.1146/annurev.psych.56.091103.070234

885 35. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive

886 and negative affect: The PANAS scales. Journal of Personality and Social Psychology. 1988;54:

887 1063–1070. doi:10.1037/0022-3514.54.6.1063

888 36. Spielberger CD. State-trait anxiety inventory for adults. 2012. doi:10.1037/t06496-000

889 37. Heimberg RG, Horner KJ, Juster HR, Safren SA, Brown EJ, Schneier FR, et al. Psychometric

890 properties of the liebowitz social anxiety scale. Psychological Medicine. 1999;29: 199–212.

891 doi:10.1017/S0033291798007879

892 38. Vetter RS. Calling all arachnophobic entomologists: A request for information. American

893 Entomologist. 2012;58: 199–201. doi:10.1093/ae/58.4.199

894 39. Zhuang Q, Zheng X, Becker B, Lei W, Xu X, Kendrick KM. Intranasal vasopressin like

895 oxytocin increases social attention by influencing top-down control, but additionally enhances

896 bottom-up control. Psychoneuroendocrinology. 2021;133: 105412.

897 doi:10.1016/j.psyneuen.2021.105412

898 40. Kou J, Zhang Y, Zhou F, Sindermann C, Montag C, Becker B, et al. A randomized trial shows

899 dose-frequency and genotype may determine the therapeutic efficacy of intranasal oxytocin.

900 Psychological Medicine. 2022;52: 1959–1968. doi:10.1017/S0033291720003803

901 41. Quintana DS, Smerud KT, Andreassen OA, Djupesland PG. Evidence for intranasal oxytocin

902 delivery to the brain: Recent advances and future perspectives. Therapeutic Delivery. 2018 [cited 5

903 Apr 2025]. doi:10.4155/tde-2018-0002

904 42. Berger A, Kiefer M. Comparison of different response time outlier exclusion methods: A

905 simulation study. Front Psychol. 2021;12. doi:10.3389/fpsyg.2021.675558

906 43. Ratcliff R. Methods for dealing with reaction time outliers. Psychological Bulletin. 1993;114:

907 510–532. doi:10.1037/0033-2909.114.3.510

908 44. Hayes TR, Petrov AA. Mapping and correcting the influence of gaze position on pupil size

909 measurements. Behav Res. 2016;48: 510–527. doi:10.3758/s13428-015-0588-x

910 45. Kret ME, Sjak-Shie EE. Preprocessing pupil size data: Guidelines and code. Behav Res.

911 2019;51: 1336–1342. doi:10.3758/s13428-018-1075-y

912 46. Eagleman DM. Human time perception and its illusions. Curr Opin Neurobiol. 2008;18:

913 131–136. doi:10.1016/j.conb.2008.06.002

914 47. Burnham KP, Anderson DR. Multimodel inference. Sociological Methods & Research. 2004

915 [cited 30 Sept 2025]. doi:10.1177/0049124104268644

916 48. Laeng B, Sulutvedt U. The eye pupil adjusts to imaginary light. Psychological Science. 2013

917 [cited 3 Apr 2025]. doi:10.1177/0956797613503556

918 49. Eilers PHC, Marx BD. Flexible smoothing with B-splines and penalties. Statistical Science.

919 1996;11: 89–121. doi:10.1214/ss/1038425655

920 50. Dan EL, Dînşoreanu M, Mureşan RC. Accuracy of six interpolation methods applied on pupil

921 diameter data. 2020 IEEE International Conference on Automation, Quality and Testing, Robotics

922 (AQTR). 2020. pp. 1–5. doi:10.1109/AQTR49680.2020.9129915

923 51. Hershman R, Henik A, Cohen N. A novel blink detection method based on pupillometry

924 noise. Behav Res. 2018;50: 107–114. doi:10.3758/s13428-017-1008-1

925 52. Greven S, Scheipl F. A general framework for functional regression modelling. Statistical

926 Modelling. 2017 [cited 30 Sept 2025]. doi:10.1177/1471082X16681317

927 53. Bates D, Kliegl R, Vasishth S, Baayen H. Parsimonious mixed models. arXiv; 2018.

928 doi:10.48550/arXiv.1506.04967

54. Viviani R, Grön G, Spitzer M. Functional principal component analysis of fMRI data. Hum
Brain Mapp. 2005;24: 109–129. doi:10.1002/hbm.20074

55. Caliński T, Harabasz J. A dendrite method for cluster analysis. Communications in Statistics -
Theory and Methods. 1974 [cited 30 Sept 2025]. Available:
https://www.tandfonline.com/doi/abs/10.1080/03610927408827101

56. Vidaurre D, Smith SM, Woolrich MW. Brain network dynamics are hierarchically organized
in time. Proceedings of the National Academy of Sciences. 2017;114: 12827–12832.
doi:10.1073/pnas.1705120114

57. Rabiner LR. A tutorial on hidden markov models and selected applications in speech
recognition. Proceedings of the IEEE. 1989;77: 257–286. doi:10.1109/5.18626

58. Renner LF, Włodarczak M. When a dog is a cat and how it changes your pupil size: Pupil
dilation in response to information mismatch. Interspeech 2017. ISCA; 2017. pp. 674–678.
doi:10.21437/Interspeech.2017-353

59. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society Series B (Methodological).
1995;57: 289–300.

60. Jazayeri M, Shadlen MN. Temporal context calibrates interval timing. Nat Neurosci. 2010;13:
1020–1026. doi:10.1038/nn.2590

61. Ratcliff R, McKoon G. The diffusion decision model: Theory and data for two-choice
decision tasks. Neural Comput. 2008;20: 873–922. doi:10.1162/neco.2008.12-06-420

62. Bogacz R, Brown E, Moehlis J, Holmes P, Cohen JD. The physics of optimal decision
making: A formal analysis of models of performance in two-alternative forced-choice tasks.
Psychol Rev. 2006;113: 700–765. doi:10.1037/0033-295X.113.4.700

63. Gold JI, Shadlen MN. The neural basis of decision making. Annu Rev Neurosci. 2007;30:
535–574. doi:10.1146/annurev.neuro.29.051605.113038

64. Mathôt S, Fabius J, Van Heusden E, Van der Stigchel S. Safe and sensible preprocessing and
baseline correction of pupil-size data. Behav Res. 2018;50: 94–106.
doi:10.3758/s13428-017-1007-2

65. (PDF) THE PUPIL: ANATOMY, PHYSIOLOGY, AND CLINICAL APPLICATIONS: by
irene E. Loewenfeld. 1999. Oxford: butterworth-heinemann. Price pound180. Pp. 2278. ISBN
0-750-67143-2. ResearchGate. 2024 [cited 11 Apr 2025]. doi:10.1093/brain/124.9.1881

66. Henderson RR, Bradley MM, Lang PJ. Emotional imagery and pupil diameter.
Psychophysiology. 2018;55: e13050. doi:10.1111/psyp.13050

67. Laeng B, Sulutvedt U. The eye pupil adjusts to imaginary light. Psychol Sci. 2014;25:
188–197. doi:10.1177/0956797613503556

68. Mobbs D, Headley DB, Ding W, Dayan P. Space, time, and fear: Survival computations along
defensive circuits. Trends Cogn Sci. 2020;24: 228–241. doi:10.1016/j.tics.2019.12.016

69. Bradley MM, Miccoli L, Escrig MA, Lang PJ. The pupil as a measure of emotional arousal
and autonomic activation. Psychophysiology. 2008;45: 602–607.
doi:10.1111/j.1469-8986.2008.00654.x

70. Wang C-A, Munoz DP. Modulation of stimulus contrast on the human pupil orienting
response. Eur J Neurosci. 2014;40: 2822–2832. doi:10.1111/ejn.12641

71. Aston-Jones G, Cohen JD. An integrative theory of locus coeruleus-norepinephrine function:
Adaptive gain and optimal performance. Annu Rev Neurosci. 2005;28: 403–450.

doi:10.1146/annurev.neuro.28.061604.135709

72.   Mobbs D, Marchant JL, Hassabis D, Seymour B, Tan G, Gray M, et al. From threat to fear:

The neural organization of defensive fear systems in humans. J Neurosci. 2009;29: 12236–12243.

doi:10.1523/JNEUROSCI.2378-09.2009

73.   Cross CP, Campbell A. Women's aggression. Aggression and Violent Behavior. 2011;16:

390–398. doi:10.1016/j.avb.2011.02.012

74.   Knyazev GG, Bocharov AV, Slobodskoj-Plusnin JY. Hostility- and gender-related differences

in oscillatory responses to emotional facial expressions. Aggress Behav. 2009;35: 502–513.

doi:10.1002/ab.20318

75.   Partala T, Surakka V. Pupil size variation as an indication of affective processing.

International Journal of Human-Computer Studies. 2003;59: 185–198.

doi:10.1016/S1071-5819(03)00017-X

76.   Kou J, Lan C, Zhang Y, Wang Q, Zhou F, Zhao Z, et al. In the nose or on the tongue?

Contrasting motivational effects of oral and intranasal oxytocin on arousal and reward during

social processing. Transl Psychiatry. 2021;11: 94. doi:10.1038/s41398-021-01241-w

77.   Le J, Zhang L, Zhao W, Zhu S, Lan C, Kou J, et al. Infrequent intranasal oxytocin followed

by positive social interaction improves symptoms in autistic children: A pilot randomized clinical

trial. Psychother Psychosom. 2022;91: 335–347. doi:10.1159/000524543

78.   Zhuang Q, Zheng X, Yao S, Zhao W, Becker B, Xu X, et al. Oral administration of oxytocin,

like intranasal administration, decreases top-down social attention. Int J Neuropsychopharmacol.

2022;25: 912–923. doi:10.1093/ijnp/pyac059

79.   Droit-Volet S. Time perception, emotions and mood disorders. Journal of Physiology-Paris.

2013;107: 255–264. doi:10.1016/j.jphysparis.2013.03.005

80.   Neumann ID, Landgraf R. Balance of brain oxytocin and vasopressin: Implications for

anxiety, depression, and social behaviors. Trends in Neurosciences. 2012;35: 649–659.

doi:10.1016/j.tins.2012.08.004

81.   Liu X, Jiao G, Zhou F, Kendrick KM, Yao D, Gong Q, et al. A neural signature for the

subjective experience of threat anticipation under uncertainty. Nat Commun. 2024;15: 1544.

doi:10.1038/s41467-024-45433-6

82.   Zhang R, Gan X, Xu T, Yu F, Wang L, Song X, et al. A neurofunctional signature of affective

arousal generalizes across valence domains and distinguishes subjective experience from

autonomic reactivity. Nat Commun. 2025;16: 6492. doi:10.1038/s41467-025-61706-0

83.   Salim H, Jones AM. Angiotensin II receptor blockers (ARBs) and manufacturing

contamination: A retrospective national register study into suspected associated adverse drug

reactions. Br J Clin Pharmacol. 2022;88: 4812–4827. doi:10.1111/bcp.15411

84.   Pulcu E, Shkreli L, Holst CG, Woud ML, Craske MG, Browning M, et al. The effects of the

angiotensin II receptor antagonist losartan on appetitive versus aversive learning: A randomized

controlled trial. Biol Psychiatry. 2019;86: 397–404. doi:10.1016/j.biopsych.2019.04.010

85.   The brain renin angiotensin system: A novel precision target for neurofunctional symptom

regulation | request PDF. In: ResearchGate [Internet]. 25 Aug 2025 [cited 29 Sept 2025].

doi:10.31234/osf.io/fbcv4_v1

86.   Gillan CM, Apergis-Schoute AM, Morein-Zamir S, Urcelay GP, Sule A, Fineberg NA, et al.

Functional neuroimaging of avoidance habits in obsessive-compulsive disorder. AJP. 2015;172:

284–293. doi:10.1176/appi.ajp.2014.14040525

87.   Angiotensin II regulates the neural expression of subjective fear in humans: A precision

pharmaco-neuroimaging approach. Biological Psychiatry: Cognitive Neuroscience and

Neuroimaging. 2023;8: 262–270. doi:10.1016/j.bpsc.2022.09.008

1020