Title: Yoga Effect on Quality-of-Life Study Among Patients with Idiopathic Pulmonary

Fibrosis (YES-IPF)

Authors and affiliation: Suha Kadura[1], Soumik Purkayastha[2,6], Josh Benditt[1], Amit

Anand[3], Bridget Collins[1], Miguele De Quadros[4], Mafara Hobson[4], MJ Biswas[4],

Lawrence Ho[1], Cathie Spino[5], Ganesh Raghu*[1]

1- Division of Pulmonary, Critical Care and Sleep Medicine, Center for Interstitial Lung

Diseases, Department of Medicine, University of Washington, Seattle, WA, USA

2- Department of Biostatistics, University of Pittsburgh, Pittsburgh, PA, USA.

3- Division of Pulmonary, Critical Care and Sleep Medicine, Beth Israel Deaconess

Medical Center, Boston, MA, USA.

4- Independent certified yoga instructors

5- Department of Biostatistics, University of Michigan, Ann Arbor, MI, USA.

6- Center for Health Equity Research and Promotion, Pittsburgh, PA, USA

***Corresponding Author:** Ganesh Raghu, MD**,** University of Washington, Campus Box

356175, Seattle, WA 98195 USA**.** Email: graghu@uw.edu

Running title: Yoga Effect on Quality-of-Life in IPF Patients;

Descriptor Number: 9.23 Interstitial Lung Disease;

Total words: 2990

Abstract words: 246

Summary: We examine if the implementation of a modified yoga program improves

patient-reported outcomes in patients with idiopathic pulmonary fibrosis. Our study

reveals significant post-yoga improvements in the yoga group compared to the control

group across different patient-reported outcomes. Such outcomes are meaningful

endpoints in patients with idiopathic pulmonary fibrosis and serve as a critical endpoint

in related clinical trials.

Contributions:

1. *Suha Kadura:* Data gathering, interpretation of result, writing first draft, revised

versions and collated inputs from all authors of the manuscript.

2. *Soumik Purkayastha:* Data analyses, interpretation of result, writing first draft,

revised versions and collated inputs from all authors of the manuscript.

3. Josh Benditt and Amit Anand: Developed the protocol and study design for the

modified yoga program /exercises used in the trial; reviewed the data analyzed,

interpreted results.

4. Josh Benditt, Amit Anand, Miguele De Quadros, Mafara Hobson, and MJ Biswas:

Certified yoga instructors for patients.

5. Bridget Collins: Enrolled consenting participants in the study, participated in study

design, interpretation of results, reviewing manuscript and provided input.

6. *Lawrence Ho:* Enrolled and encouraged patients to participate in the trial.

7. *Cathie Spino:* Reviewed data analyses and interpretation, reviewed manuscript

and provided input.

8. *Ganesh Raghu:* designed the trial and study protocol, enrolled consenting

participants in the study, interpretation of results, supervised the conduct of trial

and execution of the trial; manuscript writing.

All authors reviewed the drafts of manuscript, provided input, contributed to

writing the manuscript and approved the final manuscript for submission.

Data availability statement: In accordance with our protocol registered at

ClinicalTrials.gov (NCT02848625), individual participant data from this study will not be

made available publicly. This approach was established to protect participant privacy

and confidentiality as stated in our informed consent process. Summary data supporting

the findings of this study are available within the article and its supplementary materials.

Researchers interested in collaborative analyses may contact the corresponding author

with specific research questions and proposed methodologies for consideration.

Funding Statement: No external funding or financial support was received for the

conduct of this research, data analysis, manuscript preparation, or submission.

ABSTRACT

Rationale: Patients with idiopathic pulmonary fibrosis have poor health-related quality

of life.

Objective: Determine whether a modified yoga program in patients with idiopathic

pulmonary fibrosis improves quality of life compared to usual care.

Methods: Randomized controlled, non-blinded, pilot clinical trial lasting 12 weeks with 2

arms involving 63 adults with idiopathic pulmonary fibrosis. The yoga program-arm

consisted of interventions such as seated postures, breathing and meditation exercises

designed by authors for idiopathic pulmonary fibrosis patients. The control arm

continued with usual standard of care. Analysis of covariance was performed; no

multiplicity adjustments were made on account of this being a pilot study.

Measurements: The primary outcomes were week-12 scores in seven patient-reported

instruments, each with sub-domains, totaling 21 outcomes.

Results: 60 of the 63 participants (32 randomized to yoga; 31 receiving standard of

care) completed the study (one death in each arm and one withdrawal in yoga group).

Analysis of covariance for week-12 scores, adjusting for baseline scores and

confounders, revealed significant treatment effects favoring yoga in the L-IPF cough

domain (-9.29 points, 95% CI -18.37 to -0.20; p=0.045), in the L-IPF total score (-7.11,

95% CI -13.15 to -1.06; p=0.022), and in the R-scale-PF cough domain (-1.18, 95% CI -

2.27 to -0.10; p=0.034) in the study population.

Conclusion: Patients with idiopathic pulmonary fibrosis participating in a yoga program

demonstrated significant improvement in quality of life assessed by the cough and total

scores of L-IPF and the cough score of R-scale-PF than those receiving usual care.

Abstract word count: 246

Key words: Idiopathic Pulmonary Fibrosis, Yoga, Health-Related Quality-of-Life,

Patient-Reported Outcome Measures, Cough.

TEXT

1. Introduction

Idiopathic pulmonary fibrosis carries significant mortality and symptom burden, with

limited therapeutic options. Despite the discovery of antifibrotic agents that slow the rate

of decline of forced vital capacity, patients living with idiopathic pulmonary fibrosis suffer

from persistent respiratory symptoms and diminished quality of life. There are no

available pharmacologic therapies for idiopathic pulmonary fibrosis that have been

shown to impact patient-reported outcomes, such as symptoms or health-related

quality-of-life. Though biomarkers such as forced vital capacity or image patterns may

identify effects of interventions on biological pathways of the disease process, they

likely do not fully encompass treatment impact on how patients 'feel, function, and

survive,' and only weakly correlate with patient-reported symptom measures [2–4].

Identification of treatment strategies for idiopathic pulmonary fibrosis with the aim of

enhancing patients' quality of life is a top priority.

Yoga is a mind-body practice that has become an increasingly popular and effective

method to treat and manage symptoms of chronic diseases, and is the most commonly

used complementary health approach in the United States [5]. According to a 2017

national survey, about one in seven U.S. adults practiced yoga in the past 12 months,

and the percentage of people practicing yoga grew from 2007 to 2017 [6]. A mounting

body of literature demonstrates the benefits of yoga not only in components of general

wellness including stress, mental/emotional health, and sleep quality, but has also been

shown to improve symptoms and quality of life in various chronic diseases including

Chronic Obstructive Pulmonary Disease, asthma, cancer, heart failure, arthritis, and

chronic pain [7–12].

Although classical yoga includes other elements, yoga as commonly practiced in the

United States typically emphasizes standard components such as breathing exercises

(pranayama), postures (asanas), and meditation (dhyana). Central to asanas and

pranayama is diaphragmatic breathing, which may lead to improved chest expansion

and decreased work of breathing. Studies evaluating the benefits of yoga in chronic

respiratory diseases (largely asthma and Chronic Obstructive Pulmonary Disease) have

shown improvement in variables such as forced expiratory volume in 1 second, tidal

volume and minute ventilation, as well as increase in measures such as alveolar gas

exchange and improved submaximal and maximal oxygen consumption, in addition to

quality of life [13–16].

We sought to assess the effect of a standardized virtual, modified yoga program

(henceforth, treatment) designed specifically for individuals with idiopathic pulmonary

fibrosis at our center, with the effect on health-related quality of life as a primary

objective. We hypothesized that a 12-week modified yoga program, designed

specifically for individuals with idiopathic pulmonary fibrosis, would improve health-

related quality of life compared to usual care.

2. Methods

2.1 Study design: This study was a non-blinded, single center, randomized pilot trial

conducted by the Center for Interstitial Lung Diseases at the University of Washington

Medical Center. It was approved by the Institutional Review Board at the University of

Washington and registered with ClinicalTrials.gov (NCT02848625). See Sections 2 to 6

of Online Data Supplement for more details.

2.2 Participants: Consenting adults with idiopathic pulmonary fibrosis meeting criteria

established in 2018 by the American Thoracic Society, European Respiratory Society,

Japanese Respiratory Society, and Latin American Thoracic Society at time of screening

were eligible [17]. Exclusion criteria included comorbidities interfering with yoga (e.g.,

musculoskeletal discomfort), evaluation for lung transplant or hospice care, participation

in yoga and/or pulmonary rehabilitation outside of the study at screening and during the

study period. All patients were otherwise allowed to receive usual care for idiopathic

pulmonary fibrosis including treatment with *nintedanib* or *pirfenidone* and supplemental

oxygen when necessary, during the study period.

2.3 Randomization and masking: Eligible participants were randomly allocated in a

1:1 ratio to either the treatment arm (following the modified yoga program), or the

control arm (receiving standard of care and following usual activities) with stratification

by supplemental oxygen need prior to enrollment to account for disease severity. The

randomization schedule was prepared using computer-generated permuted random

blocks of size 4 for each stratum. The sequence of allocation was hidden from the

investigators to prevent selection bias. Participants, investigators, and providers were

not blinded to intervention; see Figure E1 of the Online Data Supplement.

2.4 Procedures: The modified yoga program consisted of seated postures, breathing,

and meditation exercises designed by 5 certified yoga instructors tailored for fibrosis

patients (see Table E1 and Section 9 of Online Data Supplement). Chair yoga, as used

in our study, is a gentler yoga alternative for older adults. Classical yoga with more

strenuous movement that may involve poses on the floor, significant joint and muscle

stretching was considered not to be appropriate for the typical idiopathic pulmonary

fibrosis patient population. Chair yoga is quite popular and many studios and training

courses for its practice are available, with prior studies demonstrating benefit in elderly

patients [18].

The protocol was intended to be in-person, though was amended due to the COVID-19

pandemic to reduce exposure risk, with participants in the treatment arm participating in

group sessions conducted virtually via Zoom in individuals' homes (5-6 per group).

Sessions lasted one hour, twice/week, for 12 consecutive weeks, led by a certified yoga

instructor. The same instructor led each group's sessions. Participants received a

secure link to a pre-recorded demo video to practice during their free time.

Participants in the treatment group received a journal at the beginning of the program

and were encouraged to write in it at the end of each class (see Section 8 of Online

Data Supplement). Journal entries were used in the qualitative assessment of the

program. Attention control conditions were not implemented for the control group due to

lack of feasibility and resource limitations during the pandemic. All patients were

evaluated in person at baseline and after 12 weeks when lung function and 6-minute

walk tests were performed, and patient-reported outcome measurements were

collected.

2.5 Outcomes: The *primary outcomes* measured were total/domain-specific scores

from seven validated health-related quality of life tools collected at baseline and at the

end of the 12-week study period. These included the Living with Idiopathic Pulmonary

Fibrosis symptoms module (L-IPF symptoms), a recently developed and validated visual

analog scale (VAS) for IPF called Raghu scale for pulmonary fibrosis (R-Scale-PF),

King's Brief Interstitial Lung Disease questionnaire (K-BILD), EuroQol-Five-Dimensional

Five-Level questionnaire (EQ-5D-5L), Hospital Anxiety and Depression (HAD) Scale,

Patient-Reported Outcomes Measurement Information System Sleep Disturbance and

Sleep-Related Impairment (PROMIS SD and SRI), and Epworth Sleepiness Scale

(ESS) [19–26].

Secondary outcomes included changes from baseline to 12 weeks in absolute and

percent predicted forced vital capacity (FVC), diffusing capacity of the lungs for carbon

monoxide (DLCO), and 6-minute walk distance (6MWD).

2.6 Questionnaires for Patient Reported Outcomes

L-IPF questionnaire symptoms module: The L-IPF symptoms module contains 15 items,

with the total score derived as the average of the dyspnea (7 items), cough (5 items),

and energy/fatigue (3 items) domains, each with a 24-hour recall period [19]. Each item is

rated on a 5-point scale (0-4), with higher scores connoting greater impairment. The

minimum clinically important difference estimates have been reported as 6 to 7 for the

dyspnea domain score and 4 to 5 points for the cough domain score, though these

estimates were investigated in a population of patients with progressive interstitial lung

disease, other than idiopathic pulmonary fibrosis, which does not specifically apply to

our study population [27].

R-scale-PF questionnaire: The R-scale-PF is a visual analog scale assessing 5 domains

(cough, shortness of breath, fatigue, depressed mood, and overall sense of well-being).

Each domain score ranges from 0 to 10 with 0.5 increments. The total score ranges

from 0 to 50 with lower numbers indicating better health-related quality of life [20]. A

minimum clinically important difference estimate has not yet been defined in fibrosis

patients.

Details for the remaining utilized questionnaires are provided in Section 7 of Online

Data Supplement.

2.7 Statistical analysis

Study Design and Analysis Population: This pilot trial utilized patient-reported outcomes

as primary outcomes, with sample size determined by feasibility rather than power

calculations, as well-defined data on clinically important differences were not available

for the fibrosis population in the implemented instruments. The primary analysis set was

the modified intention-to-treat population defined as all randomized participants with a

week 12 visit

Primary Analysis: Between-group comparisons for outcome scores were conducted

using analysis of covariance (ANCOVA), adjusting for baseline questionnaire scores,

age, sex, BMI, and supplementary oxygen use. Any of the outcomes that demonstrated

significant improvement in the study population was considered a favorable outcome. All

analyses were conducted in R (version 4.4.1). Given this is a pilot trial, p-values were

not adjusted for multiple comparisons. Missing data were handled via multiple

imputation with chained equations. Sensitivity analyses using complete cases were

performed.

3. Results

Between January 1-June 10, 2021, 63 consecutive, eligible patients provided consent.

Thirty-two were randomly assigned to yoga and 31 to the control group. Thirty (94%) of

32 participants in the yoga group and 30 (97%) of 31 participants in the control group

completed 12 weeks of assessment (Figure 1) to form the study population. One

participant from each study arm was missing at least one week 12 questionnaires.

DLCO data were not available for one participant at baseline and four at week 12, one

did not have week 12 FVC data, and two participants were missing 6MWD at baseline

and 4 at week 12.

Baseline demographics are presented in Table 1. Most participants were male (41 [68%]

of 60), with mean age of 73.4 [SD 8.21] years. Median disease duration was 3.2 years.

Most characteristics are comparable between the treatment groups, with no clinically

meaningful differences. Participants had mild to moderate impairment in lung function at

baseline (mean FVC% predicted was 74.8 [SD 21.44] and % predicted DLCO was 54.0

[18.27]). Twenty-two (37%) participants used supplemental oxygen. Fifty-one (85%)

participants were on background antifibrotic therapy with either *nintedanib* or

pirfenidone. Comorbid conditions included obstructive sleep apnea in 28 (47%)

participants, emphysema in 5 (8%), gastroesophageal reflux disease (confirmed by pH

probe) in 25 (42%), and pulmonary hypertension (confirmed by right heart

catheterization) in 11 (18%).

In Table 2, adjusted comparisons between yoga and control groups for primary

outcomes using multiple imputation are reported. For the L-IPF cough domain, the yoga

group showed improvement from baseline score (mean 27.83 [SD = 25.69]) to week 12

(mean 18.10 [21.23]), while the control group remained stable (baseline: 24.83 [22.95];

week 12: 25.00 [22.09]). The imputation-based analysis showed a treatment effect that

was statistically significant, favoring yoga (adjusted treatment effect: -9.29; 95% CI: [-

18.37, -0.20]; p=0.045). The yoga group showed improvement from baseline total L-IPF

score (mean 30.84 [SD = 18.77]) to week 12 (mean 23.37 [16.49]), while the control

group remained stable (baseline: 26.49 [19.60]; week 12: 27.16 [19.92]). The

imputation-based analysis showed a treatment effect that was statistically significant,

favoring yoga (adjusted treatment effect: -7.11; 95% CI: [-13.15, -1.06]; p=0.022). In the

R-scale-PF cough domain, the yoga group demonstrated significant reduction in cough

(baseline: 3.20 [SD 2.64]; week 12: 2.07 [2.31]), whereas the control group showed a

slight worsening (baseline: 2.87 [2.21]; week 12: 3.14 [2.56]). The imputation-based

analyses demonstrated significant treatment effects (-1.18; 95% CI: [-2.27, -0.10];

p=0.034).

Figure 2 illustrates individual patient trajectories from baseline to week 12 across three

pulmonary fibrosis outcome measures through a waterfall plot.

In Table E2 of Online Data Supplement, unadjusted and adjusted comparison between

the yoga and control groups for secondary outcomes are shown using the multiple

imputation approach. There were no significant differences in any of the secondary

outcomes over 12 weeks. Findings from the sensitivity analyses are presented in Table

E3 of Online Data Supplement.

A qualitative content analysis was performed through participants' journals. Central

themes identified leading to sense of improvement in overall well-being included social

connection, managing fibrosis symptoms, changed mindset (Figure E2 of Online Data

Supplement). Regarding social connection, participants reported feeling a sense of

community and support during the program. Participants felt less self-conscious about

experiencing symptoms (such as cough) and enjoyed speaking with others about their

experiences and being able to monitor their progress alongside others who shared their

diagnosis. Several participants reported incorporating more deliberate and controlled

breathing techniques into daily activities, with improved knowledge and understanding

of the connection between mind and body. From a mindset standpoint, participants

reported feeling an increase in confidence in their ability and motivation to increase

physical activity and push themselves despite their debilitating diagnosis.

4. Discussion

This pilot trial provides evidence that participants receiving 12-week modified yoga program experienced improvements in health-related quality-of-life outcomes, particularly in the cough domain scores of both the L-IPF and R-Scale-PF questionnaires, compared to those who continued usual daily activities. The strength of this conclusion is bolstered by consistency of results showing greater improvements in multiple quality of life instruments—including HADS, K-BILD, total R-scale-PF score, PROMIS SRI and ESS, following intervention in the yoga group than the control group.

To our knowledge, this is the first idiopathic pulmonary fibrosis clinical trial reporting improvement in outcomes in the intervention group as primary endpoints. As emphasized by a recent landmark international symposium involving patients, academic investigators, and regulatory representatives, endpoints in idiopathic pulmonary fibrosis clinical trials should resonate more closely with the tangible patient experience [2]. The integration of adequately validated outcomes is an important step in this direction. The statistically significant improvement in the L-IPF cough domain score (and numeric improvement in KBILD score) in the yoga arm participants in our study is clinically meaningful to patients, as both of these questionnaires were developed with input from idiopathic pulmonary fibrosis patients and have undergone extensive validity and reliability testing [2,21,28]. In a patient population with progressive interstitial lung disease, other than idiopathic pulmonary fibrosis, Swigris et al. provided estimates of 4 to 5 points in the L-PF cough domain score as a clinically meaningful threshold [27]. Other scale tools, such as cough visual analogue scale, carry prognostic and quality of life

value in fibrosis and are being increasingly used in interventional trials in idiopathic

pulmonary fibrosis [32,33].

Previous studies have demonstrated a clear benefit of yoga in other chronic respiratory

diseases on validated health-related quality-of-life scores, including asthma and COPD

[8,9,34,35]. Our study is the first to evaluate the effect of yoga on patients with idiopathic

pulmonary fibrosis with positive findings despite the barrier of virtual rather than in-

person sessions due to the COVID-19 pandemic, particularly since previous data

suggests participant preference for in-person yoga classes [36]. Two unpublished results

showed mixed results regarding improvements in quality of life scores following

participation in yoga, although both were designed as feasibility/safety studies with less

than 20 participants each (one study in idiopathic pulmonary fibrosis patients and

another in patients with fibrotic ILD of any cause) [37,38]. We did not find statistically

significant improvement in anxiety and depression scores measured by the HADS in

contrast to a previous study that evaluated a general group of chronic lung disease

patients awaiting lung transplant [39].

Although numerically greater improvements in the ESS and PROMIS SRI scores were

observed in the yoga group, our study did not show statistically significant benefit in

sleep quality despite prior reports showing improvement in sleep quality and insomnia

with yoga participation in other patient populations [40]. There is a complicated

relationship with sleep quality in idiopathic pulmonary fibrosis patients, in whom

comorbid sleep related breathing disorders such as OSA and insomnia are common

and likely worsen the quality of life and may even increase mortality in patients with

idiopathic pulmonary fibrosis [41–43]. Another factor reported to be potentially contributing

to increased arousal index in idiopathic pulmonary fibrosis is comorbid

gastroesophageal reflux disease [41]. Further prospective studies are needed to better

evaluate the effects of yoga on overall sleep quality in idiopathic pulmonary fibrosis

patients.

Possibilities of the potential mechanism for improved quality of life in fibrosis patients

participating in yoga include improved posture, allowing more efficient use of abdominal

and diaphragmatic muscles. Various yoga exercises such as deep diaphragmatic

breathing can improve vagal tone, which may help diminish enhanced cough reflex

sensitivity as previously described in patients with idiopathic pulmonary fibrosis [44].

The results of this study must be interpreted in the context of several limitations amidst

its strengths. First, this was a single center, pilot trial designed to assess feasibility with

the aim of informing future studies with adequate sample size calculations and powered

to detect significant differences in trial outcomes between groups. This was an

exploratory pilot trial designed primarily to assess feasibility and generate hypotheses

for future studies. We acknowledge that with multiple endpoints (21 in total) and no

adjustment for multiple comparisons, our findings are vulnerable to Type I error.

However, the consistency of findings across multiple cough-related measures

strengthens our confidence in these specific results. Future confirmatory trials should

pre-specify primary endpoints and include appropriate statistical adjustments. A second

limitation is the relatively small sample size and study duration of only 12 weeks. Third,

since a blinded trial is not possible, it is difficult to distinguish between yoga-specific and

nonspecific benefits (e.g. expectation bias, social connection during class, interpersonal

skills of the yoga instructor). Finally, future trials should include well-validated measures

of cough such as The Leicester Cough Questionnaire [45,46].

In summary, this is the first idiopathic pulmonary fibrosis clinical trial demonstrating

improvement in patient-reported outcomes as primary endpoints, and the first study to

evaluate the effect of yoga practice in this patient population. Besides demonstrating

safety and feasibility of a yoga program that patients can engage in from the comfort of

their homes, the study demonstrated improved health-related quality of life measures in

patients with idiopathic pulmonary fibrosis following participation in a yoga program

supervised by instructors. These results are clinically significant given the critical need

for therapeutic options that improve outcomes that are meaningful to patients suffering

with this disease. A larger, multi-center trial is warranted to further establish improved

health-related quality of life via participation in a yoga program tailored for patients with

idiopathic pulmonary fibrosis, in keeping with an emphasis on patient-centered

outcomes.

REFERENCES

1.   Raghu G, Remy-Jardin M, Richeldi L, et al. Idiopathic Pulmonary Fibrosis (an

Update) and Progressive Pulmonary Fibrosis in Adults: An Official

ATS/ERS/JRS/ALAT Clinical Practice Guideline. *Am J Respir Crit Care Med*.

2022;205(9):e18-e47. doi:10.1164/rccm.202202-0399ST

2.   Raghu G, Ghazipura M, Fleming TR, et al. Meaningful Endpoints for Idiopathic

Pulmonary Fibrosis (IPF) Clinical Trials: Emphasis on 'Feels, Functions, Survives'.

Report of a Collaborative Discussion in a Symposium with Direct Engagement from

Representatives of Patients, Investigators, the National Institutes of Health, a

Patient Advocacy Organization, and a Regulatory Agency. *Am J Respir Crit Care*

Med. 2024;209(6):647-669. doi:10.1164/rccm.202312-2213SO

3.   Swigris JJ, Han M, Vij R, et al. The UCSD shortness of breath questionnaire has

longitudinal construct validity in idiopathic pulmonary fibrosis. *Respir Med*.

2012;106(10):1447-1455. doi:10.1016/j.rmed.2012.06.018

4.   Aronson KI, Martin-Schwarze AM, Swigris JJ, et al. Validity and Reliability of the

Fatigue Severity Scale in a Real-World Interstitial Lung Disease Cohort. *Am J Respir*

Crit Care Med. 2023;208(2):188-195. doi:10.1164/rccm.202208-1504OC

5.   *2016 Yoga in America Study Conducted by Yoga Journal & Yoga Alliance*.

https://www.yogaalliance.org/2016YogaInAmericaStudy

6. Clarke TC, Barnes PM, Black LI, Stussman BJ, Nahin RL. Use of Yoga, Meditation,

and Chiropractors Among U.S. Adults Aged 18 and Over. *NCHS Data Brief*.

2018;(325):1-8.

7. Selvan P, Hriso C, Mitchell J, Newberg A. Systematic review of yoga for symptom

management during conventional treatment of breast cancer patients. *Complement*

Ther Clin Pract. 2022;48:101581. doi:10.1016/j.ctcp.2022.101581

8. Holland AE, Hill CJ, Jones AY, McDonald CF. Breathing exercises for chronic

obstructive pulmonary disease. Cochrane Airways Group, ed. *Cochrane Database*

of Systematic Reviews. Published online October 17, 2012.

doi:10.1002/14651858.CD008250.pub2

9. Yang ZY, Zhong HB, Mao C, et al. Yoga for asthma. *Cochrane Database Syst Rev*.

2016;4(4):CD010346. doi:10.1002/14651858.CD010346.pub2

10. Andrea Cortés-Ladino C, Augusto Arias-Ortiz W, Porras-Ramírez A. Effectiveness of

Yoga and Acupuncture in Rheumatoid Arthritis: A Systematic Review and Meta-

Analysis. *Evid Based Complement Alternat Med*. 2023;2023:9098442.

doi:10.1155/2023/9098442

11. Chow SL, Bozkurt B, Baker WL, et al. Complementary and Alternative Medicines in

the Management of Heart Failure: A Scientific Statement From the American Heart

Association. *Circulation*. 2023;147(2):e4-e30. doi:10.1161/CIR.0000000000001110

12. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute,

and Chronic Low Back Pain: A Clinical Practice Guideline From the American

College of Physicians. *Ann Intern Med.* 2017;166(7):514-530. doi:10.7326/M16-

2367

13. Vitacca M, Clini E, Bianchi L, Ambrosino N. Acute effects of deep diaphragmatic

breathing in COPD patients with chronic respiratory insufficiency. *Eur Respir J.*

1998;11(2):408-415. doi:10.1183/09031936.98.11020408

14. Goyeche JR, Ago Y, Ikemi Y. Asthma: The yoga perspective. Part I. The

somatopsychic imbalance in asthma: towards a holistic therapy. *J Asthma Res.*

1980;17(3):111-121. doi:10.3109/02770908009105671

15. Satish V, Rao RM, Manjunath NK, et al. Yoga versus physical exercise for cardio-

respiratory fitness in adolescent school children: a randomized controlled trial. *Int J*

Adolesc Med Health. 2018;32(3). doi:10.1515/ijamh-2017-0154

16. Shaw I, Shaw BS, Brown GA. Role of diaphragmatic breathing and aerobic exercise

in improving pulmonary function and maximal oxygen consumption in asthmatics.

Science & Sports. 2010;25(3):139-145. doi:10.1016/j.scispo.2009.10.003

17. Raghu G, Remy-Jardin M, Myers JL, et al. Diagnosis of Idiopathic Pulmonary

Fibrosis. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. *Am J Respir*

Crit Care Med. 2018;198(5):e44-e68. doi:10.1164/rccm.201807-1255ST

18. Park J, McCaffrey R, Newman D, Liehr P, Ouslander JG. A Pilot Randomized

Controlled Trial of the Effects of Chair Yoga on Pain and Physical Function Among

Community-Dwelling Older Adults With Lower Extremity Osteoarthritis. *J American*

Geriatrics Society. 2017;65(3):592-597. doi:10.1111/jgs.14717

19. Swigris JJ, Andrae DA, Churney T, et al. Development and Initial Validation Analyses

of the Living with Idiopathic Pulmonary Fibrosis Questionnaire. *Am J Respir Crit*

Care Med. 2020;202(12):1689-1697. doi:10.1164/rccm.202002-0415OC

20. Scallan C, Strand L, Hayes J, et al. R-scale for pulmonary fibrosis: a simple, visual

tool for the assessment of health-related quality of life. *Eur Respir J.*

2022;59(1):2100917. doi:10.1183/13993003.00917-2021

21. Patel AS, Siegert RJ, Brignall K, et al. The development and validation of the King's

Brief Interstitial Lung Disease (K-BILD) health status questionnaire. *Thorax.*

2012;67(9):804-810. doi:10.1136/thoraxjnl-2012-201581

22. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr*

Scand. 1983;67(6):361-370. doi:10.1111/j.1600-0447.1983.tb09716.x

23. Johns MW. A new method for measuring daytime sleepiness: the Epworth

sleepiness scale. *Sleep.* 1991;14(6):540-545. doi:10.1093/sleep/14.6.540

24. EuroQol Group. EuroQol--a new facility for the measurement of health-related

quality of life. *Health Policy.* 1990;16(3):199-208. doi:10.1016/0168-8510(90)90421-

9

25. Buysse DJ, Yu L, Moul DE, et al. Development and validation of patient-reported

outcome measures for sleep disturbance and sleep-related impairments. *Sleep.*

2010;33(6):781-792. doi:10.1093/sleep/33.6.781

26. Yu L, Buysse DJ, Germain A, et al. Development of short forms from the PROMIS^{TM}

sleep disturbance and Sleep-Related Impairment item banks. *Behav Sleep Med.*

2011;10(1):6-24. doi:10.1080/15402002.2012.636266

27. Swigris JJ, Bushnell DM, Rohr K, Mueller H, Baldwin M, Inoue Y. Responsiveness

and meaningful change thresholds of the Living with Pulmonary Fibrosis (L-PF)

questionnaire Dyspnoea and Cough scores in patients with progressive fibrosing

interstitial lung diseases. *BMJ Open Respir Res.* 2022;9(1):e001167.

doi:10.1136/bmjresp-2021-001167

28. Swigris J, Cutts K, Male N, Baldwin M, Rohr KB, Bushnell DM. The Living with

Pulmonary Fibrosis questionnaire in progressive fibrosing interstitial lung disease.

ERJ Open Res. 2021;7(2):00145-02020. doi:10.1183/23120541.00145-2020

29. Nolan CM, Birring SS. PROMising developments in IPF patient-reported outcome

measures. *Eur Respir J.* 2022;59(1):2102312. doi:10.1183/13993003.02312-2021

30. Moor CC, Mostard RLM, Grutters JC, Bresser P, Wijsenbeek MS. The use of online

visual analogue scales in idiopathic pulmonary fibrosis. *Eur Respir J.*

2022;59(1):2101531. doi:10.1183/13993003.01531-2021

31. Swigris JJ, Pryor JB, Aronson KI, Guess TA, Solomon JJ. Interstitial Lung Disease

Patients' Global Impressions of Symptoms, Severity Ratings, and Meaningfulness of

Changes Over Time. *Ann Am Thorac Soc.* Published online August 12, 2024.

doi:10.1513/AnnalsATS.202405-457OC

32. Khor YH, Johannson KA, Marcoux V, et al. Epidemiology and Prognostic

Significance of Cough in Fibrotic Interstitial Lung Disease. *Am J Respir Crit Care*

Med. Published online March 27, 2024:rccm.202311-2101OC.

doi:10.1164/rccm.202311-2101OC

33. Wu Z, Smith DJF, Yazbeck L, et al. Cough Severity Visual Analog Scale Assesses

Cough Burden and Predicts Survival in Idiopathic Pulmonary Fibrosis. *Am J Respir*

Crit Care Med. 2024;209(9):1165-1167. doi:10.1164/rccm.202311-2169LE

34. Anshu null, Singh N, Deka S, et al. The effect of yoga on pulmonary function in

patients with asthma: A meta-analysis. *Complement Ther Clin Pract*.

2023;50:101682. doi:10.1016/j.ctcp.2022.101682

35. Cai Y, Ren X, Wang J, Ma B, Chen O. Effects of Breathing Exercises in Patients

With Chronic Obstructive Pulmonary Disease: A Network Meta-analysis. *Arch Phys*

Med Rehabil. 2024;105(3):558-570. doi:10.1016/j.apmr.2023.04.014

36. Brinsley J, Smout M, Davison K. Satisfaction with Online Versus In-Person Yoga

During COVID-19. *J Altern Complement Med*. 2021;27(10):893-896.

doi:10.1089/acm.2021.0062

37. Fell CD, Jessa I, Hatley S, Crawford L, Culos-Reed N. Yoga For Idiopathic

Pulmonary Fibrosis: A Pilot Study. In: *A42. NOVEL PHENOTYPES, OUTCOME*

MEASURES, TREATMENT STRATEGIES, AND COMORBID ILLNESS. American

Thoracic Society; 2011:A1527-A1527. doi:10.1164/ajrccm-

conference.2011.183.1_MeetingAbstracts.A1527

38. Kreuter M, Wasmuth L, Kotzerke M, et al. Yoga in patients with fibrosing interstitial

lung diseases – a feasibility trial. In: *Idiopathic Interstitial Pneumonias*. European

Respiratory Society; 2020:773. doi:10.1183/13993003.congress-2020.773

39. Santana MJ, S-Parrilla J, Mirus J, Loadman M, Lien DC, Feeny D. An assessment of

the effects of Iyengar yoga practice on the health-related quality of life of patients

with chronic respiratory diseases: a pilot study. *Can Respir J.* 2013;20(2):e17-23.

doi:10.1155/2013/265406

40. Wang WL, Chen KH, Pan YC, Yang SN, Chan YY. The effect of yoga on sleep

quality and insomnia in women with sleep problems: a systematic review and meta-

analysis. *BMC Psychiatry.* 2020;20(1):195. doi:10.1186/s12888-020-02566-4

41. Hagmeyer L, Herkenrath SD, Treml M, Pietzke-Calcagnile A, Anduleit N, Randerath

W. Sleep-related breathing disorders in idiopathic pulmonary fibrosis are frequent

and may be associated with pulmonary vascular involvement. *Sleep Breath.*

2023;27(3):961-971. doi:10.1007/s11325-022-02686-z

42. Kolilekas L, Manali E, Vlami KA, et al. Sleep Oxygen Desaturation Predicts Survival

in Idiopathic Pulmonary Fibrosis. *Journal of Clinical Sleep Medicine.*

2013;09(06):593-601. doi:10.5664/jcsm.2758

43. Troy LK, Young IH, Lau EMT, et al. Nocturnal hypoxaemia is associated with

adverse outcomes in interstitial lung disease. *Respirology.* 2019;24(10):996-1004.

doi:10.1111/resp.13549

44. van Manen MJG, Birring SS, Vancheri C, et al. Cough in idiopathic pulmonary

fibrosis. *Eur Respir Rev.* 2016;25(141):278-286. doi:10.1183/16000617.0090-2015

45. Birring SS. Development of a symptom specific health status measure for patients

with chronic cough: Leicester Cough Questionnaire (LCQ). *Thorax.* 2003;58(4):339-

343. doi:10.1136/thorax.58.4.339

46. Key AL, Holt K, Hamilton A, Smith JA, Earis JE. Objective cough frequency in

Idiopathic Pulmonary Fibrosis. *Cough.* 2010;6(1):4. doi:10.1186/1745-9974-6-4

47. Stanojevic S, Kaminsky DA, Miller MR, et al. ERS/ATS technical standard on

interpretive strategies for routine lung function tests. *Eur Respir J.*

2022;60(1):2101499. doi:10.1183/13993003.01499-2021

FIGURE LEGENDS

1. Figure 1: CONSORT diagram.

2. Figure 2: Waterfall plot to illustrate individual patient changes from baseline to

week 12 across three pulmonary fibrosis outcome measures. Panel (A) shows

changes in the cough domain score using the Living with Idiopathic Pulmonary

Fibrosis (L-IPF) scale; (B) presents changes in the total domain score via the L-

IPF scale, and (C) displays changes in the cough domain score using the Raghu

Scale for Pulmonary Fibrosis (R-Scale-PF).

 TABLES

522 **Table 1.** Baseline characteristics of the study population.

	Control group (N=30)	Treatment group (N=30)	Overall (N=60)
Age (years)			
Mean (SD)	74.3 (7.41)	72.5 (8.97)	73.4 (8.21)
Body Mass Index (kg/m^2)			
Mean (SD)	26.5 (4.66)	27.6 (4.55)	27.1 (4.60)
Sex, n (%)			
Female	10 (33%)	9 (30%)	19 (32%)
Male	20 (67%)	21 (70%)	41 (68%)
Ethnicity, n (%)			
White	27 (90%)	27 (90%)	54 (90%)
Other	3 (10%)	3 (10%)	6 (10%)
Current/former smoker? n (%)			
No	19 (63%)	16 (53%)	35 (58%)
Yes	11 (37%)	14 (47%)	25 (42%)
Time since diagnosis (years)			
Median (Interquartile range)	3.6 (2.93)	3.0 (4.39)	3.2 (3.67)
[25th percentile – 75th percentile]	[2.31 – 5.24]	[1.92 – 6.30]	[2.21 – 5.88]
Surgical lung biopsy done? n (%)			
No	17 (57%)	19 (63%)	36 (60%)
Yes	13 (43%)	11 (37%)	24 (40%)
FVC (L)			
Mean (SD)	2.7 (0.77)	2.9 (0.86)	2.8 (0.82)
Median [25th percentile – 75th percentile]	2.7 [2.02 – 2.96]	3.1 [2.22 – 3.70]	2.8 [2.08 – 3.37]
FVC % predicted			
Mean (SD)	70.0 (21.84)	79.6 (20.28)	74.8 (21.44)
Median [25th percentile – 75th percentile]	70.5 [58.75 – 84.25]	77.5 [66.75 – 94.75]	73.0 [65.00 – 91.25]
DLCO (mmol/min/kPa)			
Mean (SD)	12.4 (3.41)	13.3 (4.42)	12.8 (3.93)
Missing	0 (0%)	1 (3.3%)	1 (1.7%)
DLCO % predicted value [*]			
Mean (SD)	52.5 (15.38)	56.0 (20.97)	54.0 (18.27)
Missing	0 (0%)	1 (3.3%)	1 (1.7%)
6MW distance (feet)			
Mean (SD)	1311 (405.7)	1269 (403.6)	1290 (401.7)
Missing	1 (3.3%)	1 (3.3%)	2 (3.3%)
Supplemental oxygen use, n (%) [†]			

	Control group (N=30)	Treatment group (N=30)	Overall (N=60)
Supplemental O2 not required	20 (67%)	18 (60%)	38 (63%)
Supplemental O2 required	10 (33%)	12 (40%)	22 (37%)
Obstructive sleep apnea reported, n (%)			
No	19 (63%)	13 (43%)	32 (53%)
Yes, without PAP therapy	4 (13%)	10 (33%)	14 (23%)
Yes, with PAP therapy	7 (23%)	7 (23%)	14 (23%)
Emphysema reported, n (%)			
No	29 (97%)	26 (87%)	55 (92%)
Yes	1 (3%)	4 (13%)	5 (8%)
Antifibrotics, n (%)			
No	3 (10%)	6 (20%)	9 (15%)
Yes	27 (90%)	24 (80%)	51 (85%)
Gastroesophageal reflux disease, n (%)			
No/Unknown	13 (43%)	15 (50%)	28 (47%)
Yes, confirmed by pH probe	14 (47%)	11 (37%)	25 (42%)
Yes, not confirmed by pH probe	3 (10%)	4 (13%)	7 (12%)
Pulmonary hypertension, n (%)			
No/Unknown	22 (73%)	16 (53%)	38 (63%)
Yes, confirmed by right heart catheterization	4 (13%)	7 (23%)	11 (18%)
Yes, not confirmed by right heart cath.	4 (13%)	7 (23%)	11 (18%)
Notes: [*] Percent predicted FVC and DLCO calculated using GLI reference equations [47]. [†] Supplemental oxygen use is a stratification factor.			

523   **Table 2. Changes from baseline to week 12 and adjusted treatment effect of modified yoga**

524   **program in primary outcomes for the study population.**

	Baseline*	Week 12*	Change† (Week 12 – Baseline)	Adjusted Treatment Effect of Yoga‡	
				Point estimate (95% CI)	p-value
Living with Idiopathic Pulmonary Fibrosis (L-IPF) (Cough)					
MYP	27.83 (25.69)	18.10 (21.23)	-9.31 (21.03)	-9.29 (-18.37, -0.20)	0.045
Control	24.83 (22.95)	25.00 (22.09)	0.17 (19.14)		
Living with Idiopathic Pulmonary Fibrosis (L-IPF) (Dyspnea)					
MYP	24.14 (21.81)	21.27 (20.02)	-2.61 (12.09)	-2.86 (-8.98, 3.27)	0.353
Control	21.27 (20.02)	22.86 (22.48)	0.99 (11.03)		
Living with Idiopathic Pulmonary Fibrosis (L-IPF) (Overall)					
MYP	30.84 (18.77)	23.37 (16.49)	-7.18 (13.57)	-7.11 (-13.15, -1.06)	0.022
Control	26.49 (19.60)	27.16 (19.92)	0.66 (10.70)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Cough)					
MYP	3.20 (2s.64)	2.07 (2.31)	-0.97 (2.29)	-1.18 (-2.27, -0.10)	0.034
Control	2.87 (2.21)	3.14 (2.56)	0.28 (2.05)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Shortness of Breath)					
MYP	4.13 (2.62)	4.00 (2.58)	0.00 (1.71)	-0.16 (-1.12, 0.80)	0.741
Control	3.27 (2.48)	3.66 (2.96)	0.38 (1.90)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Fatigue)					
MYP	4.43 (2.45)	4.31 (2.88)	0.00 (2.20)	0.09 (-1.05, 1.24)	0.869
Control	3.67 (2.66)	3.83 (3.08)	0.14 (2.08)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Mood)					
MYP	2.63 (2.61)	1.90 (2.54)	-0.79 (2.46)	-0.21 (-1.37, 0.96)	0.722
Control	1.63 (1.92)	1.48 (2.11)	-0.07 (2.09)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Wellbeing)					
MYP	3.03 (1.88)	3.21 (2.35)	0.07 (2.46)	-0.57 (-1.81, 0.67)	0.362
Control	2.80 (2.04)	3.48 (2.95)	0.72 (2.17)		
Raghu Scale for Pulmonary Fibrosis (R-scale-PF) (Overall)					
MYP	17.43 (9.63)	15.48 (9.88)	-1.69 (8.65)	-2.27 (-6.68, 2.14)	0.306
Control	14.23 (8.84)	15.59 (10.93)	1.45 (7.60)		
King's Brief Interstitial Lung Disease (K-BILD) (Psychological)					
MYP	36.40 (10.52)	38.45 (9.10)	1.90 (8.51)	0.03 (-4.28, 4.35)	0.988
Control	40.20 (10.04)	40.17 (10.18)	-0.03 (9.95)		
King's Brief Interstitial Lung Disease (K-BILD) (Breathlessness and Activities)					
MYP	24.03 (8.72)	24.21 (7.89)	0.14 (4.76)	0.06 (-2.12, 2.25)	0.953
Control	25.57 (8.59)	25.63 (7.89)	0.07 (4.62)		
King's Brief Interstitial Lung Disease (K-BILD) (Chest Symptoms)					
MYP	11.27 (2.21)	11.93 (2.2)	0.62 (1.90)	0.24 (-0.61, 1.08)	0.578
Control	11.70 (2.31)	12.00 (2.12)	0.30 (1.58)		
King's Brief Interstitial Lung Disease (K-BILD) (Overall)					
Yoga	71.73 (17.79)	74.59 (17.27)	2.62 (11.73)	0.66 (-5.73, 7.04)	0.837
Control	77.57 (18.33)	77.80 (17.64)	0.23 (13.94)		
EuroQol-5D-5L (EQ-5D-5L) Visual Analogue Scale (VAS)					
MYP	70.53 (18.77)	70.76 (16.71)	0.90 (13.27)	2.60 (-3.56, 8.76)	0.400
Control	71.57 (21.88)	68.43 (21.66)	-3.13 (12.40)		

EuroQol-5D-5L (EQ-5D-5L) Index					
MYP	0.81 (0.11)	0.80 (0.16)	-0.01 (0.11)	0.00 (-0.05, 0.04)	0.853
Control	0.82 (0.10)	0.82 (0.12)	0.00 (0.07)		
Hospital Anxiety and Depression Scale (HADS) (Anxiety)					
MYP	4.80 (4.13)	3.90 (3.72)	-0.90 (2.87)	-0.10 (-1.28, 1.07)	0.859
Control	3.23 (2.49)	2.93 (2.66)	-0.17 (2.19)		
Hospital Anxiety and Depression Scale (HADS) (Depression)					
MYP	4.57 (3.82)	3.76 (3.52)	-0.90 (3.07)	-0.88 (-2.21, 0.46)	0.193
Control	4.17 (2.90)	4.28 (3.5)	0.17 (2.16)		
Hospital Anxiety and Depression Scale (HADS) (Overall)					
MYP	9.37 (7.45)	7.66 (6.68)	-1.79 (4.97)	-1.09 (-3.32, 1.14)	0.331
Control	7.40 (4.80)	7.17 (5.51)	-0.03 (3.59)		
Patient-Reported Outcomes Measurement Information System (PROMIS™) Sleep Disturbance (SD)					
MYP	45.29 (9.52)	45.80 (8.73)	0.89 (8.60)	2.45 (-1.03, 5.93)	0.163
Control	48.62 (9.48)	46.29 (9.6)	-2.13 (5.76)		
Patient-Reported Outcomes Measurement Information System (PROMIS™) Sleep-Related Impairment (SRI)					
MYP	47.32 (8.78)	45.77 (8.08)	-1.10 (7.62)	-1.82 (-5.63, 1.99)	0.342
Control	44.63 (9.61)	45.92 (10.66)	1.58 (7.66)		
Epworth Sleepiness Scale (ESS)					
MYP	7.03 (3.94)	6.03 (4.40)	-0.90 (2.92)	-0.70 (-2.44, 1.05)	0.427
Control	4.63 (2.92)	4.90 (3.57)	0.38 (3.16)		

Notes:

[*] Week 12 is the end of the intervention period. Mean (standard deviation) values were computed for all patients with complete data at baseline and week 12.

[†] We perform an unadjusted comparison of the change (week 12 – baseline) of each of the primary outcomes between the two study arms (treatment and control).

[‡] A linear regression model was used treatment differences at week 12 for outcomes, adjusting for age, sex, BMI, supplementary oxygen usage, and the baseline value of the specific outcome measure under consideration. To handle missing data, multiple imputation was performed using the chained equations method. Pooled estimates, confidence intervals (CIs), and p-values were obtained using Rubin's rules (N=60) with * denoting statistically significant findings at level 0.05.

 FIGURES

527
528 **Figure 1:** CONSORT diagram.

Figure 2:

Waterfall plot to illustrate individual patient changes from baseline to week 12 across

three pulmonary fibrosis outcome measures. Panel (A) shows changes in the cough

domain score using the Living with Idiopathic Pulmonary Fibrosis (L-IPF) scale; (B)

presents changes in the total domain score via the L-IPF scale, and (C) displays

changes in the cough domain score using the Raghu Scale for Pulmonary Fibrosis (R-

Scale-PF).