Unlocking uterine biology at home: a validated system for DNA, RNA, and microbial

analysis from menstrual effluence

Stephen Gire[1]*, Xitong Li*[1]*, Claire Toth*[1]*, Meet Doshi*[3]*, Sachin Kumar Gupta*[3]*, Scott Parker*[2]*,*

Debbie Boles[2]*, Ridhi Tariyal*[1]

1. NextGen Jane, Inc. Oakland, California

2. Labcorp, Research Triangle Park, North Carolina

3. Elucidata, San Francisco, California

Corresponding Authors:

Stephen Gire: stephen@nextgenjane.com

Ridhi Tariyal: ridhi@nextgenjane.com

ABSTRACT

Introduction: Access to accurate, non-invasive diagnostics remains a critical unmet need in women's health. Menstrual effluence, containing endometrial tissue, immune cells, and microbial communities, represents a clinically relevant specimen for genomic and molecular pathology applications, yet has historically been underutilized due to concerns about sample integrity and variability.

Methods: We developed and validated a standardized, at-home tampon-based collection system designed to preserve nucleic acids at ambient temperature for clinical-grade analyses. 1,067 tampon samples from 328 participants underwent, RNA sequencing and metatranscriptomic profiling to assess specimen transcript integrity, diagnostic fidelity, and microbial composition over time. 12 patients were exome sequenced using matched menstrual effluence and whole blood to assess assay concordance between sample types.

Results: RNA extracted from menstrual effluence maintained stability for up to 14 days without refrigeration, achieving sufficient yield and quality for sequencing in >97% of samples. Variant detection via exome sequencing demonstrated 100% concordance among overlapping single nucleotide variants between menstrual fluid and matched venous blood, confirming clinical equivalency for genetic testing. Transcriptomic analyses revealed cycle-dependent variation in key reproductive and immune markers, while metatranscriptomic profiling identified shifts in microbial communities consistent with known reproductive tract dysbiosis.

Conclusions: Standardized at-home collection of menstrual effluence provides a clinically actionable platform that supports remote specimen acquisition without compromising molecular assay fidelity, offering a scalable solution to improve access to carrier screening, reproductive health assessment, and infectious disease monitoring in clinical practice.

INTRODUCTION

Women's health has been underserved by non-specific diagnostics and limited therapeutic

options. Despite high disease burden, conditions like endometriosis, fibroids, and recurrent

miscarriage are diagnosed and treated with tools that lack precision, leading to significant delays

in care[1]. Broader reliance on non-specific clinical pathways has led to a one-size-fits-all approach

to care[2–4]. For instance, gonadotropin-releasing hormone (GnRH) analogs—despite inducing

premature menopause and causing severe side effects such as bone loss and mood

disturbances[5] – remain a first-line treatment for endometriosis[6] and fibroid-related heavy

bleeding[7]. CA125, a non-specific marker of inflammation, is monitored across a wide range of

conditions including ovarian cancer[8], heart failure[9], miscarriage risk[10], and endometriosis[11],

despite limited diagnostic specificity[12]. Without condition-specific diagnostics, targeted therapies

have remained underdeveloped, perpetuating inadequate care across puberty, pregnancy, and

menopause.

At-home biospecimen collection systems offer a scalable way to improve diagnostics and enable

precision medicine in women's health. Remote collection of biological samples allows clinical data

to be captured outside of brick-and-mortar healthcare settings, improving access and enabling

earlier decision-making[13]. For research, these systems offer opportunities to build rich,

longitudinal, high-dimensional datasets that are essential for studying reproductive biology at

scale. However, few such datasets exist, and those available are poorly harmonized[14], limiting

their cross-study comparability and value for biomarker discovery.

Menstrual effluence is a rich but underutilized biospecimen that closely mirrors the endometrial

environment and presents novel opportunities for diagnostic development. Menstrual fluid is

expelled naturally and contains a mix of shed endometrium, interstitial fluid, blood, mucosa, and

epithelial cells from the cervix and vagina, along with reproductive tract microbiota[15,16]. Organoids

derived from menstrual effluence demonstrate similar transcriptomic profiles and hormone

responsiveness as those from invasive endometrial biopsies[17]. Importantly, menstrual effluence

contains a unique population of immune cells enriched for innate and adaptive subsets involved

in tissue remodeling and immune tolerance[18], unlike peripheral blood. The sample's heterogeneity

captures a broader snapshot of uterine and vaginal health than traditional biopsies or swabs.

Despite its promise, menstrual effluence has historically been viewed as technically difficult to

work with. Approximately 30% of cells in menstrual fluid are non-viable[15,16], and tissue aggregates

and mucosal debris complicate laboratory workflows. Additionally, cellular lysis releases

nucleases that can degrade nucleic acids, compromising downstream molecular assays. Sample

composition also varies by collection method[19] – e.g., pad vs. tampon vs. menstrual cup –

introducing variability and contributing to skepticism about its reliability as a biospecimen.

However, these challenges are surmountable with proper pre-analytical protocols.

New advances have transformed how menstrual effluence is viewed in diagnostics and research.

Recent studies have generated high-resolution datasets from menstrual fluid and uterine tissue

that offer insight into reproductive pathophysiology[15,16,20]. Transcriptomic data from menstrual

effluence have revealed dynamic biological processes such as epithelial-to-mesenchymal

transition relevant to endometriosis progression[21]. Endotoxins detected in menstrual samples

have been linked to adverse pregnancy outcomes[22], and menstrual immune profiles have shown

potential for diagnosing inflammatory and infectious conditions[23]. These discoveries demonstrate

that menstrual effluence is not only clinically relevant but may offer information inaccessible via

other sample types.

We developed a standardized, tampon-based system to enable high-quality menstrual fluid

collection at home. To address pre-analytical variability and improve sample stability, we built a

self-contained tampon collection kit that also captures metadata such as cycle day and bleeding

volume, which serve as important co-variables in transcriptomic analyses. Prior research has

shown that aligning sample timing with menstrual phase improves diagnostic accuracy in

cardiovascular[24] and mental health applications[25]. The NextGen Jane® (NGJ) platform enables

consistent sample collection, better supports hormonal context interpretation, and allows for

widespread use in both clinical and research settings.

We validated our tampon-based sample collection system through large-scale deployment and

demonstrated its feasibility and diagnostic utility in three clinically relevant applications through

DNA exome sequencing (carrier screening) and unbiased RNA sequencing of human and

microbial cell types (molecular phenotyping and microbial surveillance). We have collected 2,025

RNA-seq and metatranscriptomic datasets from 1,067 tampons submitted by 328 U.S.

participants. All samples were home-collected and shipped at ambient temperature to a

centralized lab in Oakland, CA, without loss of data quality – demonstrating the viability of

decentralized sampling for molecular diagnostics.

MATERIALS AND METHODS

A thorough accounting of materials and methods can be found in Supplemental Methods.docx

Patient Recruitment and Metadata Collection

Patients were recruited under IRB-approved protocols (20192619, 20191947, 20233438).

Consent was obtained from participants who agreed to tampon-based specimen collection and

completed a survey on clinical history, menstrual regimen, and demographics. Data were

organized in four tiers: patient-level (race, ethnicity), cycle-level (age, geography, birth control),

tampon/kit-level (flow type, cycle day), and sample-level (RNA concentration, library size).

Sample accession and collection dates were recorded to assess sample stability over time. Flow

rate was calculated from the weight of the tampon before and after collection. Metadata can be

found in Supplemental_RNASeq.xlsx. Patient IDs C01-, H01, E01), Kit IDs (NAPS) and Sample

IDs (ESGs), are randomized and anonymized ID's that cannot be tied back to patients outside

the research team.

Sample Collection and Processing

120   Participants used organic cotton tampons during multiple days of menstruation, wearing them for

121   4 hours. After collection, the tampon was sealed in a collection jar with 20mL of preservation

122   buffer (Norgen Biotek). Samples were sent back to the lab via USPS. Upon receipt, samples were

123   processed, including extrusion, centrifugation, and aliquoting into cryovials for storage at -80°C.

Hemoglobin Content Spectrophotometry

126   The tampon sample was diluted 1:4 with a preservation buffer and analyzed using a UV/Vis

127   spectrophotometer to measure absorbance at 550nm.

Nucleic Acid Extraction and Quality Assessment

130   RNA and DNA were extracted using the Norgen Column-Based RNA Extraction and MagMax

131   mirVana Total RNA Isolation methods. For RNA, DNase treatment was applied, followed by clean-

132   up with RNA XP clean beads. DNA was extracted using the MagMax-96 DNA Multi-Sample Kit.

133   The extracted nucleic acids were quantified using a Qubit 4.0 fluorometer.

Exome Sequencing and Carrier Screening

136   DNA exome sequencing was outsourced to Labcorp for analysis under their established

137   protocols. SNVs and sample metadata can be found in Supplemental_DNASeq.xlsx

RNA Sequencing and Analysis

RNA libraries were prepared using the Zymo-Seq RiboFree Total RNA Library Kit, followed by

sequencing on an Illumina NextSeq2000. Data processing included adapter trimming, alignment

to the hg38 human genome, and gene count normalization using STAR Aligner and

FeatureCount. RNA degradation effects were accounted for using the DegNorm program.

Sequencing metadata and gene expression values can be found in Supplemental_RNASeq.xlsx.

Tampon and menstrual cup samples were compared to publicly available RNA sequencing data

on GTeX to calculate Spearman correlations between tissue types. Samples used for this analysis

can be found in Supplemental_GTeX.xlsx. Differential expression was performed in XLSTAT.

Microbiome Analysis

Metatranscriptomic reads were aligned to microbial databases using Kraken2 to generate raw

read counts. For microbiome diversity, species-level and genus-level counts were aggregated to

genus level and analyzed using log-transformation and CPM normalization. Sequencing

metadata and gene expression values can be found in Supplemental_RNASeq.xlsx

Metadata and Statistical Analysis

Metadata, including clinical phenotypes and sample characteristics, were used in ANOVA to

assess statistical significance across various groups. Both pvalue and cohen's f for effect size

were calculated for all analyses using XLSTAT. Analysis focused on bleeding phenotypes, flow

rates, and other collection-related variables.

RESULTS

A tampon-based at-home collection system can reproducibly collect menstrual effluence

for downstream molecular analyses under real-world mailing and storage conditions.

The NGJ tampon collection system was designed for ease-of-use, integration into daily routines,

and standardization to ensure reliable biological preservation. Participants use the provided

TOTM-brand organic, low-absorbency tampons, minimizing synthetic additives and heavy metals.

Tampons are worn for four hours – reducing toxic shock risk and facilitates sample processing.

The cardboard applicator reduces contact with the absorbent core, limiting commensal

contamination. A nitrile glove is included for removal and placement into the jar. Once sealed, the

cap pulls the string into the jar, preventing leaks, and a shuttle punctures a foil seal to release the

preservation buffer, supersaturating the tampon. The sample is bagged, intake questions

completed and returned via standard mail. User-tested instructions guide participants through the

collection process. A picture of the collection kit and cover of the instructions for use can be found

in Figure 1.

Demographic Data of Sample Population – We analyzed menstrual samples collected on cycle

days 1–3 with complete phenotypic and technical data (Supplementary Table 1). Inclusion criteria

ensured samples were menstrual (not off-cycle vaginal tampons), had complete metadata (flow

type, collection time, bleeding phenotype), and passed sequencing quality checks (strandedness,

ribosomal rRNA content and uniquely mapped reads). This resulted in 1,135 sequences from 601

tampons across 276 participants.

Table 1 summarizes participant demographics and key flow characteristics. 75% (207

participants) identified as Non-Hispanic White, with the remaining 69 participants identifying as

Asian, Black, or more than one race. Of 365 cycles, 287 were from individuals not using birth

control. Enrolled participants were asked to report the **flow phenotype** (overall period heaviness),

characterized as heavy, medium or light, corresponding to their subjective experience of their

entire menstrual cycle at the time of collection. **Flow type** (heaviness on the collection day) was

also reported by participants. These measures differ – e.g., a heavy-flow tampon from a light

bleeder may differ significantly from one of a heavy bleeder.

Menstrual effluence preserved in stabilization buffer maintains sufficient RNA integrity for

sequencing for at least 14 days at ambient temperature.

Home-based sample collection enables large-scale, longitudinal studies across diverse locations.

We report nucleic acid yields from our tampon collection system and transcript degradation index

(DI), similar to the transcript integrity number (TIN)[26,27]. For sequencing, 11% of samples failed at

least once. Among initially failed samples, 79% were successfully resequenced, yielding a final

estimated sample failure rate of 2.3%. Approximately 60 million reads are generated per sample

(median = 60 million ± 23 million reads), which include both human and microbial transcripts.

Nucleic acid extracted from a 600ul aliquot of sample yields enough RNA for downstream

sequencing. After lysate release from tampons, an average of 17 mL of total volume

(mean = 17.13 ± 2.73) was recovered, providing sufficient material for downstream assays. A

500 ng RNA threshold – required for Zymo RiboFree sequencing – was routinely achieved.

Although this threshold is reachable via deep-well extraction, optimizing magnetic bead protocols

significantly improved nucleic acid recovery (Table 2), enabling use of standard plate formats.

Using 600 µL of lysate, 500 ng RNA was obtained in 93% of samples, with a median RNA yield

of 1.5 µg.

We assessed transcript degradation during ambient-temperature storage by measuring the DI of

the 25th percentile of genes using DegNorm[26]. RNA degradation causes 3′ coverage bias in RNA-

seq data. The DI [26] quantifies this degradation at the gene level, with higher scores reflecting

greater 3′ bias. Based on sample collection and processing dates, samples spent an average of

6.7 ± 3.52 days in buffer. No significant change in 5′–3′ coverage bias (DI_25) between days 1

and 21 (Figure 2a), indicating stability at typical shipping conditions. We further assessed the

impact of menstrual flow rate on RNA quality. Flow rate had a strong effect on DI_25: participants

with slower flows (<1 grams/hr) had higher degradation, whereas faster flow rates (1–2 grams/hr

or >2 grams/hr) correlated with reduced degradation (p=0.002; cohen's f = 0.230 Figure 2b).

Extended collection times in slow-flow individuals may exacerbate degradation.

No seasonal effects on DI_25 were observed (Figure 2b). We did detect significantly lower overall

degradation in samples collected in the Midwest region of the United States (p = 0.023; cohen's f

= 0.127) (Figure 2d). While not statistically significant, samples from the Midwest also show an

increase in mean flow rate compared to other regions (Midwest mean= 1.423 g/hr; Other Region

mean = 1.149 g/hr; p = 0.090; cohen's f = 0.110). This could explain the decrease in DI_25 scores

for this region. A total of 11 kits were successfully sequenced at least three times by separate

technicians, showing a median coefficient of variation between replicates of 0.149 (supplemental

data).

From 600 µL lysate, we consistently recovered sufficient RNA and DNA for sequencing. RNA

remains stable for at least two weeks in buffer at ambient temperatures, showing minimal

transcript degradation (see Figure 2a,b) – making this system ideal for at-home collection. With

17 mL total lysate per tampon (equivalent volume to extract and sequence the sample 28 times),

ample material remains for molecular assays and biobanking. Collectively, we have biobanked

2,013 tampon-derived samples.

DNA from tampon-collected menstrual effluence yields variant calls that are concordant

with those obtained from matched blood-based exome sequencing.

At-home, longitudinal collection of menstrual effluence offers accessible prenatal and antenatal

screening, especially in gynecological deserts or where remote sampling enhances clinical care[28].

To assess concordance of single nucleotide variants (SNVs) between menstrual effluence and

venous blood, both sample types were collected from 12 participants and exome sequenced via

Labcorp's Inheritest.

Percentage of overlapping SNVs between blood and menstrual samples was high

(mean = 0.983 ± 0.0089), which is similar to other studies showing high concordance of SNVs

between different sample types from the same patient[29]. On average, menstrual effluence

detected 143 more SNVs than venous blood. Non-overlapping SNVs had lower coverage than

overlapping SNVs, likely explaining the difference (median coverage depth for overlapping

SNVs = 103 vs. non-overlapping SNVs = 40). Four participants carried pathogenic variants

detected in both sample types: GJB2:p.Gly12fs (hearing loss), CFTR:p.Ile507_Phe508delinsille

(cystic fibrosis), CAPN3:p.Arg489Gln (limb-girdle muscular dystrophy), and HEXA:p.Arg499Hi

(Tay-Sachs), with allele frequencies ranging from 38% to 62% (Figure 3).

The transcriptomic profile of menstrual effluence represents a unique composite of

uterine, vaginal, and immune-derived cells distinct from isolated reproductive tissues.

To assess tissue similarity, we compared RNA expression profiles of menstrual effluence

collected via tampon (MB) and menstrual cup (MC) to GTEx data. Reproductive tissues (ovary,

vagina, fallopian tubes, cervix, uterus), whole blood, and small intestine (as outgroup) were

included. Spearman correlation was used to evaluate global similarity. Differential expression was

performed between menstrual effluence and GTEx whole blood samples to determine overlapping

genes between menstrual and reproductive tissues. The top 25 overexpressed genes in

menstrual effluence show some overlap with reproductive tissues in GTEx (Figure 4b),

demonstrating both shared and unique gene programs between the tissues. MB and MC samples

were highly correlated (Spearman's r = 0.97), with key differences in keratinized epithelium

expression (Supplemental Figure 1a), likely due to vaginal wall cell inclusion in tampon samples[19].

Reproductive tissues showed low correlation (r = 0.23–0.32), comparable to small intestine or

whole blood (r = 0.45; Figure 4a). This suggests that menstrual effluence shares only a limited

transcriptional overlap with reproductive tissue biopsies, similar to the non-reproductive tissues

included in the analysis. Differences in cycle phase and sequencing technology of GTEx data

may also contribute to this low correlation. The mixture of blood, uterine, and vaginal tissue could

represent shared gene programs between multiple tissue types leading to low correlations to any

one tissue.

Gene expression signatures in menstrual effluence vary according to user-reported hormonal use, flow rate, and cycle timing, enabling molecular phenotyping.

Endometrial gene expression provides valuable insight into menstrual physiology and molecular
phenotypes, spanning objective metrics like flow rate (grams/hour) to subjective variables such
as self-reported cycle day and flow type. To identify potential molecular correlates, we curated 89
endometrial-enriched genes from the Human Protein Atlas. Of these, 64 were detected in our
sequencing data, with 30 genes showing JaneScore-normalized expression ≥2. Notable
examples include canonical endometrial markers such as *PAEP*, *ESR1*, and several matrix
metalloproteinases (*MMP10*, *MMP11*). The genes not present or at low expression may represent
inherent differences between menstrual effluence and endometrium or be due to varying
approaches in RNA sequencing.

Flow rate varies by birth control and subjective flow phenotype. Patients using any form of birth
control exhibited reduced menstrual flow, as determined from 569 samples with reported data
(Figure 5a). When flow rate was combined with spectrophotometric hemoglobin measurements
at 550 nm, self-identified heavy bleeders (n = 83) showed significantly darker, more hemoglobin-
rich effluence ($p < 0.0001$; cohen's f=0.352); Figure 5b). This single-tampon assay presents a
scalable alternative to the burdensome alkaline hematin method, which requires collection of all
products used over an entire cycle.

Genes enriched in the endometrium correlate with cycle day and flow type. Expression of *GNLY*,
a broad-spectrum antimicrobial protein, varied significantly by cycle day ($p < 0.0001$; cohen's
f=0.394; Figure 5c), indicating immune modulation across menstruation. Furthermore, self-
reported flow heaviness (Heavy, Medium, Light, Spotting) was strongly associated with *PAEP*
expression in tampon samples ($p < 0.0001$; cohen's f= 0.469; Figure 5d).

Menstrual effluence contains transcriptionally active microbial signatures that vary with menstrual day and may include clinically relevant pathogens.

Menstrual samples show a time dependent dysbiosis in microbial transcription with an increase in BV causing bacteria on days 2 and 3 of menstruation. RNA transcriptional activity is distinct from the presence (relative abundance) of microbial DNA. We performed genus-level metatranscriptomic analysis using Kraken2 on RNA reads. We identified high transcriptional activity from several bacterial genera commonly associated with vaginal and uterine microbiota, including *Prevotella*, *Gardnerella*, *Corynebacterium*, *Finegoldia*, and *Anaerococcus* (Figure 6a). These genera are often implicated in microbial dysbiosis and reproductive tract infections. *Lactobacillus* species – typically associated with a healthy reproductive tract – were found to inversely correlate with the abundance of other bacterial taxa. In samples where *Lactobacillus* activity was high, the activity of other bacteria was generally suppressed. However, many samples exhibited low *Lactobacillus* levels (sum of all *Lactobacillus* species below 50% in 746 out of 1,135 sequences). alongside elevated expression of potentially pathogenic or opportunistic species. Interestingly, total *Lactobacillus* transcriptional activity decreased significantly on menstrual days 2 and 3 (Figure 6b), suggesting a temporal shift in microbial composition during menstruation. These findings present an opportunity to explore and understand typical microbial transitions during a single menstrual cycle and map how vaginal microbiome returns to equilibrium after a menstrual disruption.

DISCUSSION

This study presents a standardized, at-home tampon-based system for the collection and stabilization of menstrual effluence, offering a scalable approach for molecular diagnostics and research in women's health. The system preserves RNA and DNA for at least two weeks without refrigeration, supporting applications such as genetic carrier screening, RNA and exome

sequencing, and microbial profiling. Its simplicity and suitability for at-home sampling help

overcome persistent barriers to diagnostic access, especially in underserved populations.

Through the NGJ platform, patients seeking to get pregnant could access their genetic screening

results and uterine microbiome insight from a single sample collected at home.

A key contribution of this work is the ability to maintain nucleic acid integrity in a biologically

complex, degradation-prone matrix. Menstrual effluence contains a heterogeneous mixture of

blood, endometrial tissue, vaginal secretions, and immune cells, which contribute to enzymatic

degradation. By limiting tampon wear to four hours, using chemically inert, low-absorbency cotton

tampons, and stabilizing samples in a high-performance buffer, we preserved high-quality RNA

and DNA for downstream molecular applications. Higher menstrual flow rates correlated with

improved RNA quality, likely due to greater blood content and reduced degradation time.

Post-collection experiments further underscored the importance of immediate stabilization, as

RNA integrity declined significantly in samples exposed to 37°C for just 30 minutes before

preservation (Supplementary Figure 1). These findings provide actionable insights for real-world

implementation of mail-in collection systems.

From a systems biology perspective, our findings highlight menstruation as a dynamic and

immunologically active window into reproductive health. Transcriptomic profiles varied across the

cycle and by hormonal use, with genes such as *PAEP*, *ESR1*, *GNLY*, and *MMP10* reflecting

hormonal withdrawal, immune activation, and tissue remodeling. These changes are difficult to

capture via biopsy or swabs, underscoring the value of menstrual effluence for temporal molecular

profiling that should be accounted for in research and clinical findings[30–32].

The high concordance between venous blood and menstrual effluence in exome sequencing

supports its use as a non-invasive alternative for carrier screening. Variants linked to conditions

such as cystic fibrosis and Tay-Sachs were concordantly detected in both sample types,

demonstrating that menstrual effluence can expand access to genomic testing.

Our system also links subjective menstrual experiences with objective molecular measurements.

Variables such as flow heaviness and cycle day correlated with RNA gene expression patterns,

suggesting tampon-derived metrics could serve as proxies for subjective reporting or hematin-

based methods in heavy menstrual bleeding classification.

Beyond transcriptomics, metatranscriptomic profiling revealed day-specific shifts in microbial

communities. Decreases in *Lactobacillus* and increases in *Gardnerella* and *Prevotella* during

menstruation suggest a transient dysbiosis. Coupling host transcriptional responses with

microbial activity enables nuanced assessment of conditions like bacterial vaginosis or chronic

endometritis, which affect a significant proportion of infertile individuals[33]. Immune activation

markers may help determine whether antibiotic treatment is warranted in asymptomatic cases[34,35].

Our system allows for this dual measurement in a single, non-invasive specimen. RNA

transcriptional profiles of the menstrual microbial community expand the potential of menstrual

effluence as a non-invasive monitoring tool for chronic infections, sexually transmitted infections,

or dysbiosis-associated conditions like pelvic inflammatory disease.

The reproducibility and standardization of this collection method also make it well-suited for

research at scale. By capturing molecular profiles tied to menstrual phenotypes, this platform can

enable the development of classifiers for endometriosis, fibroids, and other gynecologic

conditions. Future work will endeavor to utilize this system to better understand the molecular

characteristics of menstrual effluence and how variation in cycle data within patients and between

patients may influence normal or abnormal menstruation.

CONCLUSION

In conclusion, the development and validation of a standardized tampon-based system for menstrual effluence collection marks a significant advancement in women's health diagnostics. Our findings demonstrate that menstrual effluence, a sample previously considered suboptimal for molecular analysis, can be preserved effectively for RNA and DNA extraction, enabling genetic screening, molecular phenotyping, and pathogen surveillance. This system offers unique opportunities for at-home, longitudinal monitoring of reproductive health, facilitating both clinical care and research applications.

The ability to link menstrual biomarkers with clinical outcomes, such as genetic carrier status or microbial dysbiosis, provides new avenues for early detection and personalized treatment strategies. By providing reliable, reproducible molecular data across diverse menstrual phases and bleeding phenotypes, this approach enhances the potential for diagnostic tools that address the unique health needs of women. Future work will focus on refining the system's integration into clinical practice, developing molecular classifiers for reproductive diseases like endometriosis, and expanding the applications of menstrual effluence in broader health research.

Conflicts of Interest

Ridhi Tariyal, Stephen Gire, Xitong Li, and Claire Toth are or were employed by NextGen Jane, Inc. which manufactures the tampon collection kit. The other authors did not report any potential conflicts of interest.

Funding Sources

This work was funded in part through an SBIR FastTrack grant from NICHD. Unique Federal Award Identification Number R44HD103159.

Acknowledgements

403
We would like to thank Stephen Palmer, Ruth Kulicke, Matt Stremlau, Ryan Phan, and Pardis

Sabeti for their helpful edits. We would like to thank the LabCorp team that helped coordinate

exome sequencing efforts: Marcia Eisenberg, Christina Smith, and Megann Vaughn. We would

like to thank Abhishek Jha and Yogesh Lakhotia for bioinformatic support. Margaret Eisen ,Julia

Carr and Aparna Kola coordinated all clinical collection efforts for this study. The analyses in this

paper were partly funded through an SBIR Fast Track grant with the National Institute of Child

Health and Human Development *5R44HD103159*.

REFERENCES

1. Cromeens MG, Carey ET, Robinson WR, Knafl K, Thoyre S. Timing, delays and pathways to diagnosis of endometriosis: a scoping review protocol. BMJ Open. BMJ; 2021 Jun 24;11(6):e049390. PMCID: PMC8231065

2. Domar A, Vassena R, Dixon M, Costa M, Vegni E, Collura B, Markert M, Samuelsen C, Guiglotto J, Roitmann E, Boivin J. Barriers and factors associated with significant delays to initial consultation and treatment for infertile patients and partners of infertile patients. Reprod Biomed Online. Elsevier BV; 2021 Dec;43(6):1126–1136. PMID: 34756644

3. Fryer J, Mason-Jones AJ, Woodward A. Understanding diagnostic delay for endometriosis: a scoping review [Internet]. Obstetrics and Gynecology. medRxiv; 2024. Available from: https://www.medrxiv.org/content/10.1101/2024.01.08.24300988v1.full

4. Regensteiner JG, McNeil M, Faubion SS, Bairey-Merz CN, Gulati M, Joffe H, Redberg RF, Rosen SE, Reusch JE, Klein W. Barriers and solutions in women's health research and clinical care: a call to action. Lancet Reg Health Am. Elsevier; 2025 Apr;44:101037. PMCID: PMC11937288

5. Gonadotropin releasing hormone (GnRH) analogues. LiverTox: Clinical and Research Information on Drug-Induced Liver Injury. Bethesda (MD): National Institute of Diabetes and Digestive and Kidney Diseases; 2012. PMID: 31643199

6. Saleh FL, Taylor HS. Clinical applications of gonadotropin-releasing hormone analogues: a broad impact on reproductive medicine. F S Rep. Elsevier BV; 2023 Jun;4(2 Suppl):83–87. PMCID: PMC10201293

7. Madueke-Laveaux OS, Ciebiera M, Al-Hendy A. GnRH analogs for the treatment of heavy menstrual bleeding associated with uterine fibroids. F S Rep. Elsevier BV; 2023 Jun;4(2 Suppl):46–50. PMCID: PMC10201292

8. Charkhchi P, Cybulski C, Gronwald J, Wong FO, Narod SA, Akbari MR. CA125 and ovarian cancer: A comprehensive review. Cancers (Basel). MDPI AG; 2020 Dec 11;12(12):3730. PMCID: PMC7763876

9. Pan C, Zhou M, Jian Y, Zeng Y, Wang M, Chen F. CA125: An increasingly promising biomarker of heart failure. Curr Pharm Des. Bentham Science Publishers Ltd.; 2021;27(37):3871–3880. PMID: 33461452

10. Al Mohamady M, Fattah GA, Elkattan E, Bayoumy R, Hamed DA. Correlation of serum CA-125 and progesterone levels with ultrasound markers in the prediction of pregnancy outcome in threatened miscarriage. Int J Fertil Steril. 2016 Jan;9(4):506–511. PMCID: PMC4793171

11. Chen Y, Pan M, Zuo Y, Yang B, Wang S. Research progress of CA125 in endometriosis: Teaching an old dog new tricks. gocm. BMJ; 2022 Dec;2(4):191–198.

12. Akinwunmi BO, Babic A, Vitonis AF, Cramer DW, Titus L, Tworoger SS, Terry KL. Chronic medical conditions and CA125 levels among women without ovarian cancer. Cancer Epidemiol Biomarkers Prev. American Association for Cancer Research (AACR); 2018 Dec;27(12):1483–1490. PMCID: PMC6279537

13. Rivo J, Page TF, Arrieta A, Amofah SA, McCann S, Kassaye H, Rodriguez A, Williams ML. The impact of comprehensive pre-visit preparation on patient engagement and quality of care in a population of underserved patients with diabetes: Evidence from the care management medical home center model. Popul Health Manag. Mary Ann Liebert Inc; 2016 Jun;19(3):171–177. PMID: 26440513

14. Vargas E, García-Moreno E, Aghajanova L, Salumets A, Horcajadas JA, Esteban FJ, Altmäe S. The mid-secretory endometrial transcriptomic landscape in endometriosis: a meta-analysis. Hum Reprod Open. Oxford University Press (OUP); 2022 Apr 4;2022(2):hoac016. PMCID: PMC9022214

15. Schwalie PC, Bafligil C, Russeil J, Zachara M, Biocanin M, Alpern D, Aasna E, Deplancke B, Canny G, Goncalves A. Single-cell characterization of menstrual fluid at homeostasis and in endometriosis [Internet]. medRxiv. 2024. p. 2024.05.06.24306766. Available from: https://www.medrxiv.org/content/10.1101/2024.05.06.24306766v1.full-text

16. Dobay A, Haas C, Fucile G, Downey N, Morrison HG, Kratzer A, Arora N. Microbiome-based body fluid identification of samples exposed to indoor conditions. Forensic Sci Int Genet. Elsevier BV; 2019 May;40:105–113. PMID: 30785061

17. Cindrova-Davies T, Zhao X, Elder K, Jones CJP, Moffett A, Burton GJ, Turco MY. Menstrual flow as a non-invasive source of endometrial organoids. Commun Biol. Springer Science and Business

Media LLC; 2021 Jun 17;4(1):651. PMCID: PMC8211845

18.   van der Molen RG, Schutten JHF, van Cranenbroek B, ter Meer M, Donckers J, Scholten RR, van
der Heijden OWH, Spaanderman MEA, Joosten I. Menstrual blood closely resembles the uterine
immune micro-environment and is clearly distinct from peripheral blood. Hum Reprod. Oxford
University Press (OUP); 2014 Feb;29(2):303–314. PMID: 24249743

19.   Vlasac IM, Stolrow HG, Thayer ZM, Christensen BC, Rivera L. DNA-based cell typing in menstrual
effluent identifies cell type variation by sample collection method: toward noninvasive biomarker
development for women's health. Epigenetics. Informa UK Limited; 2025 Dec;20(1):2453275. PMID:
39825876

20.   Marečková M, Garcia-Alonso L, Moullet M, Lorenzi V, Petryszak R, Sancho-Serra C, Oszlanczi A,
Icoresi Mazzeo C, Wong FCK, Kelava I, Hoffman S, Krassowski M, Garbutt K, Gaitskell K,
Yancheva S, Woon EV, Male V, Granne I, Hellner K, Mahbubani KT, Saeb-Parsy K, Lotfollahi M,
Prigmore E, Southcombe J, Dragovic RA, Becker CM, Zondervan KT, Vento-Tormo R. An integrated
single-cell reference atlas of the human endometrium. Nat Genet [Internet]. 2024 Aug 28; Available
from: http://dx.doi.org/10.1038/s41588-024-01873-w PMID: 39198675

21.   Demir AY, Demol H, Puype M, de Goeij AFPM, Dunselman GAJ, Herrler A, Evers JLH,
Vandekerckhove J, Groothuis PG. Proteome analysis of human mesothelial cells during epithelial to
mesenchymal transitions induced by shed menstrual effluent. Proteomics. Wiley Online Library;
2004 Sep 24;4(9):2608–2623. PMID: 15352236

22.   Kamiyama S, Teruya Y, Nohara M, Kanazawa K. Impact of detection of bacterial endotoxin in
menstrual effluent on the pregnancy rate in in vitro fertilization and embryo transfer. Fertil Steril.
Elsevier BV; 2004 Oct;82(4):788–792. PMID: 15482745

23.   Naseri S, Rosenberg-Hasson Y, Maecker HT, Avrutsky MI, Blumenthal PD. A cross-sectional study
comparing the inflammatory profile of menstrual effluent vs. peripheral blood. Health Sci Rep. Wiley
Online Library; 2023 Jan 5;6(1):e1038. PMCID: PMC9813904

24.   Gursoy AY, Caglar GS, Kiseli M, Pabuccu E, Candar T, Demirtas S. CRP at early follicular phase of
menstrual cycle can cause misinterpretation for cardiovascular risk assessment. Interv Med Appl
Sci. Akademiai Kiado Zrt.; 2015 Dec;7(4):143–146. PMCID: PMC4701164

25.   Handy AB, Greenfield SF, Yonkers KA, Payne LA. Psychiatric symptoms across the menstrual cycle
in adult women: A comprehensive review. Harv Rev Psychiatry. Ovid Technologies (Wolters Kluwer
Health); 2022;30(2):100–117. PMCID: PMC8906247

26.   Xiong B, Yang Y, Fineis FR, Wang JP. DegNorm: normalization of generalized transcript
degradation improves accuracy in RNA-seq analysis. Genome Biol. 2019 Apr 16;20(1):75. PMCID:
PMC6466807

27.   Wang L, Nie J, Sicotte H, Li Y, Eckel-Passow JE, Dasari S, Vedell PT, Barman P, Wang L,
Weinshiboum R, Jen J, Huang H, Kohli M, Kocher JPA. Measure transcript integrity using RNA-seq
data. BMC Bioinformatics. Springer Science and Business Media LLC; 2016 Feb 3;17(1):58.
PMCID: PMC4739097

28.   Byström JW, Vikström L, Rosendal E, Gröning R, Gwon YD, Nilsson E, Sharma A, Espaillat A,
Hanke L, McInerney G, Puhar A, Cava F, Karlsson Hedestam GB, Thunberg T, Monsen T, Elgh F,
Evander M, Johansson AF, Överby AK, Ahlm C, Normark J, Forsell MN. At-home sampling to meet
geographical challenges for serological assessment of SARS-CoV-2 exposure in a rural region of
northern Sweden, March to May 2021: a retrospective cohort study. Euro Surveill [Internet].
European Centre for Disease Control and Prevention (ECDC); 2023 Mar;28(13). Available from:
http://dx.doi.org/10.2807/1560-7917.ES.2023.28.13.2200432 PMCID: PMC10064644

29.   Zhu Q, Hu Q, Shepherd L, Wang J, Wei L, Morrison CD, Conroy JM, Glenn ST, Davis W, Kwan ML,
Ergas IJ, Roh JM, Kushi LH, Ambrosone CB, Liu S, Yao S. The impact of DNA input amount and
DNA source on the performance of whole-exome sequencing in cancer epidemiology. Cancer
Epidemiol Biomarkers Prev. American Association for Cancer Research (AACR); 2015
Aug;24(8):1207–1213. PMCID: PMC4526319

30.   Schisterman EF, Mumford SL, Sjaarda LA. Failure to consider the menstrual cycle phase may cause
misinterpretation of clinical and research findings of cardiometabolic biomarkers in premenopausal
women. Epidemiol Rev. Oxford University Press (OUP); 2014;36(1):71–82. PMCID: PMC3873842

31.   Lamon S, Le Carré J, Petito G, Duong HP, Luthi F, Hiam D, Léger B. The effect of the menstrual
cycle on the circulating microRNA pool in human plasma: a pilot study. Hum Reprod. 2023 Jan
5;38(1):46–56. PMCID: PMC9825267

32. Rocks D, Cham H, Kundakovic M. Why the estrous cycle matters for neuroscience. Biol Sex Differ.
2022 Oct 28;13(1):62. PMCID: PMC9615204
33. Hmaidan S, Holt E, Finer Z, Session DR, Roach M. The use of endometrial culture for targeted
treatment of endometritis in patients experiencing infertility and recurrent pregnancy loss. Fertil
Steril. Elsevier BV; 2022 Oct;118(4):e208.
34. Muzny CA, Schwebke JR. Asymptomatic bacterial vaginosis: To treat or not to treat? Curr Infect Dis
Rep [Internet]. Springer Science and Business Media LLC; 2020 Dec;22(12). Available from:
http://dx.doi.org/10.1007/s11908-020-00740-z PMCID: PMC8015387
35. Ng BK, Chuah JN, Cheah FC, Mohamed Ismail NA, Tan GC, Wong KK, Lim PS. Maternal and fetal
outcomes of pregnant women with bacterial vaginosis. Front Surg. 2023 Feb 13;10:1084867.
PMCID: PMC9968788
36. Uhlen M, Oksvold P, Fagerberg L, Lundberg E, Jonasson K, Forsberg M, Zwahlen M, Kampf C,
Wester K, Hober S, Wernerus H, Björling L, Ponten F. Towards a knowledge-based Human Protein
Atlas. Nat Biotechnol. Springer Science and Business Media LLC; 2010 Dec;28(12):1248–1250.
PMID: 21139605
37. The human proteome in endometrium - The Human Protein Atlas [Internet]. [cited 2025 Apr 16].
Available from: https://www.proteinatlas.org/humanproteome/tissue/endometrium

Table 1: Patient, Cycle, and Tampon level data collection summary

A. Patient Level Data for 276 Patients		B. Cycle Level Data for 365 Cycles		C. Tampon/kit level Data for 601 Tampons	
Age		**Age**		**Season of Collection**	
<35	166	<35	218	Spring	223
35-40	74	35-40	99	Summer	157
>40	36	>40	48	Autumn	135
Ethnicity		**Birth Control**		Winter	86
Hipanic of Latino	33	No birth control	287	**Flow Type**	
Non-Hispanic	235	Progestin	28	Heavy	294
Other/NA	8	Estrogen	2	Medium	183
Race		Estrogen and Progestin	41	Light	113
Asian	23	Other	7	Spotting	11
Black or African American	21	**Bleeding Phenotype**		**Cycle day**	
More than on race	21	Heavy	170	1	172
White	207	Medium	154	2	284
American Indian or Alaska Native	1	Light	41	3	145
Other/NA	3				
Region of Collection					
Midwest	42				
Northeast	46				
South	91				
West	97				

Data was collected at the **a.** patient level, as well as the **b.** cycle and **c.** tampon level. Each patient

may have multiple collections within a cycle, and between many monthly cycles. During this time

of collection, patients age, change birth control, or may change their bleeding phenotype. We

have collected data on these three hierarchical levels to reflect these changes. This dataset

includes 601 tampons collected in 365 monthly cycles from 276 participants.

Table 2: Nucleic acid recovery statistics

	Nucleic acid recover per 100ul sample lysate		
	number of extractions	median ng RNA per 100ul lysate	average stdev between replicates
	ng nucleic acid		
RNA - MagBead	508	251.67	28.43
RNA - Column	627	78.20	28.14
DNA - MagBead	12	80.80	NA

Total amount of nucleic acid (in nanograms) per 100ul of lysate was calculated by multiplying the

elution volume by the concentration of nucleic acid and then divided by the total volume of sample

lysate extracted. Data was calculated only on kits with at least 2 replicates between column-based

and magnetic bead-based extraction methods for RNA in order to calculate standard deviation of

extracted RNA at the kit level. DNA samples were not extracted in replicates and so standard

deviation was not calculated.

Figure 1: NextGen Jane tampon collection kit

NextGen Jane collection kit. **a.** Nitrile glove included for tampon removal. **b.** Included tampon with

applicator. **c.** Cap that seals tampon in jar and releases preservative onto tampon. **d.** Basic

phenotypic data about current collection is recorded on the inside flap of the box. These include

the flow type of the tampon being collected, the day of the cycle, the date, time in and time out for

collection, as well as a list of any medications taken during the collection **e.** Collection jar includes

a sealed buffer cup with preservative. **f.** Biohazard bag with absorbent material for shipping. **g.**

Tampon instructions for use. **h.** Front cover of the tampon collection instructions for use.

Figure 2: Degradation Index and sequence library size performance metrics

Sequencing metrics for monitoring degradation of RNA transcripts in menstrual collections. Higher

DI scores correlate to more degraded samples. **a.** DI_25 score variance grouped by days sample

left in buffer during transit show no significant changes by day (p = 0.107; cohen's f = 0.101).

Samples were grouped days 1-3, 4-6, 7-9, and 10 or greater. **b.** DI_25 score plotted by rate of

menstrual flow (grams of fluid collected divided by the hours of collections. Samples are bucketed

in 0 to 1 grams of fluid lost per hour, 1 to 2 grams lost per hour, and over 2 grams lost per hour.

There was a significant decrease in degradation as flow rate increased (p=0.002, cohen's f =

0.230) **c.** & **d.** DI_25 plotted by season (U.S) and by region (U.S.) the collection occurred. Midwest

collected samples showed a lower mean DI_25 compared to other regions (p = 0.023; cohen's f

= 0.127). Analysis was performed at the individual sequence level (n=1,135)

Figure 3: SNV detection in DNA from matched whole blood and menstrual effluence

samples

SNV summary statistics for the comparison of SNVs between 12 patients who were exome sequenced using DNA extracted from menstrual effluence and from a venous blood draw. **a.** Table summarizing the number of SNVs recorded in the venous blood and menstrual effluence, and the rate of venous blood SNVs that were confirmed also in menstrual effluence. **b.** Four patients with known pathogenic mutations in CFTR, GJB2, CAPN3 and HEXA were confirmed present in both menstrual effluence and venous blood. Charts show mutation plotted against the frequency of the SNV detected. Red points are menstrual and blue points are venous. 24 samples from 12 participants were used in this analysis.

Figure 4: Sample type comparisons in menstrual effluence compared to other tissue types

Menstrual effluence gene expression global differences to other tissue types. **a.** Spearman

correlation matrix of global RNA transcriptional activity in menstrual effluence (MB: tampon; MC:

menstrual cup) compared to public datasets for small intestine, ovary, vagina, fallopian tube,

cervix, uterus and blood. Red indicates a high level of transcriptional correlation between tissue

types, while white and blue indicate a low level of correlation to other tissue types. 4,235 samples

from GTeX and 218 samples from the NGJ dataset were used for this analysis. **b.** Top 25

differentially expressed genes between menstrual effluence and venous blood. Median gene

values for 20 sequences from each GTeX tissue type were taken for comparison. Shared genes

are grouped between reproductive tissue and menstrual (peach) vs menstrual samples with little

to now expression in GTeX tissues (magenta). Expression differences in KRT13 between

menstrual effluence collected in a menstrual cup or a tampon (p <0.0001; cohen's f = 2.072).

Analysis was done at the sequencing level (ESG) for 70 menstrual cup samples and 68 tampon

samples.

Figure 5: Flow rate and gene expression differences by sample collection characteristics

Flow rate and hemoglobin content by birth control and heavy menstrual bleeding and the

differential expression of endometrial genes by cycle day. **a.** Flow rate calculated as grams of

fluid collected per hour plotted against patient use of hormonal birth control. Patients on any form

of hormonal birth control show lower flow rates (p <0.0001; cohen's f = 0.204). Analysis was

performed at the kit level for 241 tampons with recorded flow rate. **b.** Flow rate multiplied by the

rfu (relative fluorescence units) spectrophotometer reading as a metric for hemoglobin within the

sample charted by whether patients self-report as heavy menstrual bleeders or normal menstrual

bleeders. Heavy bleeders show an increase in lost rate of hemoglobin per hour (p <0.0001;

cohen's f = 0.352). Analysis was performed at the kit level for 241 tampons with recorded flow

rate. **c.** GNLY expression, an endometrium-enriched gene, by the day in the cycle (first cycle day

represents first day of bleeding), shows significant expression changes by cycle day (p <0.0001;

cohen's f = 0.394). **d.** PAEP, another endometrium enriched gene charted by the day specific flow

type during the tampon collection (p <0.0001; cohen's f = 0.469). Mean and Median values are

represented by a red "+" and "——", respectively. C and d analyses were performed at the sequence

level (ESG) for 1,135 samples.

Figure 6: RNA transcriptional activity of key bacteria in menstrual effluence

a. *L. crispatus*, which is one beneficial lactobacillus species in the vaginal cavity alongside BV

causing bacteria at the genus level. CPM counts per genus were summed before log2

transformation. Mean and Median values are represented by a "+" and "—", respectively. **b.**

Genus level *Lactobacillus* shows a significant decrease in transcriptional activity on day 2 and

day 3 of menstruation, compared to day 1 (p <0.0001, cohen's f = 0.171). Analyses were

performed at the sequence level (ESG) for 1,135 samples.

Supplementary Table 1: Sample filtering parameters

Menstrual samples collected on cycle days 1-3 that had complete metadata for collection date,

bleeding phenotype, flow type, birth control, collection time and reproductive stage were included

in the analysis. Samples were further filtered by technical sequencing QC metrics including proper

strandedness of reads, ribosomal rRNA contamination and uniquely mapped reads.

All Samples	**1867**
Cycle day and Menstrual	**1600**
Cycle days 1-3	
Exclude vaginal only samples	
Missing Data (0, Not Recorded, or NA)	**1346**
Collection Date	
Bleeding Phenotype	
Flow type	
Birth Control	
Collection time	
Reproductive Stage	
Technical Filters	**1135**
Strandedness >0.8	
rRNA < 3%	
Uniquely mapped reads >8M	

Supplemental Figure 1: Supplemental analyses on menstrual cups

a. Expression differences in KRT13 between menstrual effluence collected in a menstrual cup or

a tampon (p <0.0001; cohen's f = 2.072). Analysis was done at the sequencing level (ESG) for 70

menstrual cup samples and 68 tampon samples. **b.** Menstrual effluence collected in menstrual

cups was either directly put into preservation buffer after collection or incubated at 37oC for 30

minutes before preservation (p=0.003; cohen's f = 0.606). A total of 10 diva cups from 5

participants were used in this analysis.