Hierarchical neurobiological changes in dorsolateral prefrontal cortex and subgenual anterior cingulate cortex connectivity induced by theta-burst stimulation

Armin Toghi[1,2], Ghazaleh Ghaffaripour Jahromi[3], Shayan Zarei[2,4], Babak Aliyari[3], Hamed Moqtaderi[5], Ali Motie Nasrabadi[6*]

Affiliations:

[1] Institute for Cognitive and Brain Sciences, Shahid Beheshti University, Tehran, Iran.

[2] Interdisciplinary Schools, Tehran, Iran.

[3] Neuroscience Research Center, Iran University of Medical Sciences, Tehran, Iran

[4] Tehran University of Medical Sciences, Tehran, Iran

[5] Department of Mechanical Engineering, Faculty of Engineering, Alzahra University, Tehran, Iran

[6*] Department of Biomedical Engineering, Shahed University, Tehran, Iran

* **Corresponding author:** Ali Motie Nasrabadi

Abstract

Recent studies show that left dorsolateral prefrontal cortex (DLPFC) stimulation induces acute and persistent changes in the subgenual anterior cingulate cortex (sgACC), and the degree of sgACC modulation predicts treatment response in patients. To understand how theta-burst stimulation (TBS) over the left DLPFC influences sgACC microcircuitry, we applied spectral dynamic causal modeling (DCM) with a conductance-based canonical microcircuit model (CMM-NMDA) to resting-state EEG data acquired at 2, 15, and 30 minutes following continuous TBS (cTBS), intermittent TBS (iTBS), and Sham stimulation. Using a multilevel parametric empirical Bayes framework, we observed that cTBS induce wider network effect 2 min after stimulation while iTBS made more lasting changes up to 30 min following stimulation. Unlike sham, both TBS protocols led to a transient reduction in NMDA-dependent effective connectivity from right DLPFC to left sgACC. Notably, TBS caused sustained inhibition within left sgACC microcircuits, characterized by increased self-inhibition of spiny-satellite cells. Together, these findings demonstrate that spectral DCM can resolve the dynamic, hierarchical neuromodulatory effects of TBS in healthy brains and highlight a potential mechanism underlying the antidepressant action of left DLPFC TBS.

Introduction

Repetitive transcranial magnetic stimulation (rTMS) has emerged as a powerful tool in both neuroscientific research and clinical applications[1]. This technique uses focal electromagnetic fields to induce neuronal action potentials in the brain[1]. A more recent advancement in rTMS is theta burst stimulation (TBS), which delivers magnetic pulses in a pattern that mimics natural brain oscillations [2]. Specifically, it consists of rapid gamma-frequency bursts (50 Hz) nested within slower theta-frequency oscillations (5 Hz) [2–4]. Previous works showed that intermittent TBS (iTBS) enhance cortical excitability, while continuous TBS (cTBS) have an inhibitory effect on cortical activity [3,5].

The effectiveness of TBS lies in its ability to induce lasting neural modulation beyond the stimulation period[2]. While its neuroplastic effects have been most extensively studied in the human primary motor cortex (M1), the underlying mechanisms by which TBS modify activity in nonmotor regions remain unclear[4,6,7]. This is particularly true for the left dorsolateral prefrontal cortex (DLPFC), a crucial region for therapeutic interventions. Despite the precise mechanisms of TBS still being under investigation, the technique has received FDA approval for MDD treatment [8]. However, a significant proportion of MDD patients do not respond to TBS, potentially due to poor understanding of TBS neural effects and optimal stimulation parameters[9].

Recent studies consistently highlight the relationship between the remote regulatory effects of left DLPFC rTMS on the subgenual anterior cingulate cortex (sgACC) and its role in the treatment improvement [1,10–12]. Aberrant connectivity between the DLPFC and sgACC has been implicated in the pathophysiology of depression across various neuroimaging modalities[13], and strong evidence suggests that this connectivity is altered following rTMS [1,9–11,14–16]. Notably, effective stimulation sites tend to have higher anticorrelation[10–12,17], and denser polysynaptic structural connection with the sgACC[18]. However, several challenges remain for understanding the left DLPFC TBS effect on sgACC activity and connectivity.

First, using functional magnetic resonance imaging (fMRI) to track prefrontal TBS effects yields mixed findings[19], commonly attributed to high intra- and inter-subject variability[3,19]. Moreover, applying concurrent TMS–fMRI to a full TBS protocol to track its network effect is challenging

due to insufficient temporal signal-to-noise ratio (tSNR) in subcortical regions[20]. Second,
electrophysiological studies showed that TBS does not produce a uniform inhibitory or excitatory
response[6], highly state-dependent[21], induces variable and complex changes in spectral power[4],
and a triggers widespread network connectivity effects[9]. Third, TMS creates a complex chain of
causation, spanning synaptic changes to broader network and behavioral outcomes[22], that
current methodological techniques cannot easily capture.

Instead, computational models with biologically plausible assumptions offer a bridge between
synaptic changes and electrophysiological observations[23–25]. In particular, dynamic causal
modeling (DCM) using conductance-based neural mass models of cortical microcircuitry has
become a popular approach for examining treatment-associated changes in extrinsic
interregional coupling, intrinsic connectivity, and cellular properties in each neural
population[24,26]. This approach enables tracking hierarchical neurobiological effects of
neuromodulation at multiple scales—macroscale (extrinsic interregional connections),
mesoscale (intrinsic microcircuit interactions), and microscale (cellular properties) [23,24] **(Figure**
1).

DCM inverts a generative neural mass model alongside an observation model to infer intrinsic
and extrinsic connectivity parameters that best explain the empirical data[27]. Spectral DCM
adopted for resting state data to estimate how endogenous neuronal fluctuations produce the
observed CSD[27], and has demonstrated strong reliability in resting-state electrophysiology[28]. In
this study, we used spectral DCM under a conductance-based canonical microcircuit model
(CMM_NMDA) to investigate TBS-induced effective connectivity changes between the bilateral
DLPFC and sgACC. Resting-state EEG data were recorded in a placebo-controlled, single-
blinded, crossover design involving 24 healthy adult participants, each completing three
sessions (iTBS, cTBS, sham). EEG recordings were obtained at baseline and at 2, 15, and 30
minutes post-stimulation **(Figure 2).**

After model inversion for each session, we applied a two-level hierarchical Bayesian
approach[24,29] to assess the network effects of iTBS, cTBS, and sham at 2 minutes and to model
subsequent dynamic changes up to 30 minutes. This method is a multilevel Bayesian estimation
of a general linear model, previously used in longitudinal DCM studies[24,29], which have two level
optimizations over posterior densities. At the first level, empirical priors from individual level
optimize posterior densities over parameters at the session level. Then, group-level means
serve as empirical priors that constrain parameter estimates at the individual level (**Figure 3**).
Crucially, this multilevel framework is particularly well-suited to account for the substantial intra-
and inter-subject variability inherent to TMS interventions, and finesses the local minima
problem inherent in the inversion of neural mass models (for the detailed description of this
approach compared to classical statistics such as ANOVA see **Method** section).

Results
In group-level analyses we only considered connections with posterior probability > 0.95,
reflecting strong evidence. We focus here on the difference connectivity matrices, where
positive values denote increased connectivity and negative values denote decreased
connectivity. Mean connectivity results are provided in the **Supplementary Material**.

TBS effect 2 min after stimulation
Using the PEB-of-PEB framework (**Figure 3A**), we compared Pre versus Post1 (2 min)
connectivity changes for iTBS, cTBS, and sham. Sham produced no significant extrinsic

changes, as expected. In contrast, cTBS induced the largest extrinsic connectivity modulations:
a significant increase in NMDA-mediated forward connectivity from left DLPFC to left sgACC.
For iTBS, we observed the opposite pattern—a decrease in forward NMDA connectivity from left
DLPFC to left sgACC and reduced backward AMPA coupling from left DLPFC to right sgACC.
Across extrinsic connections, cTBS also yielded bidirectional reductions between left sgACC
and right DLPFC, whereas iTBS did not significantly alter this pathway.

Intrinsic changes in the stimulation target further differentiated protocols: cTBS decreased SP →
DP excitatory connectivity in left DLPFC, indicating local hypoactivation, while iTBS reduced
DP→ II, reflecting local disinhibition. To quantify subject-level consistency, we extracted the sign
of these two intrinsic parameters from each subject's first-level PEB. Eighty-two percent of
participants showed the inhibitory effect of cTBS, and 77% exhibited the disinhibitory effect of
iTBS on left DLPFC microcircuit.

Additional intrinsic effects were found in sgACC and right DLPFC for cTBS only: left sgACC
displayed increased SP→ DP, and right sgACC showed reduced II→DP inhibition. The sole
significant change after sham was reduced SS self-inhibition in left sgACC. These immediate
connectivity alterations highlight protocol-specific modulation of prefrontal–subgenual circuits
and underscore the utility of DCM in resolving neuromodulatory effects (**Figure 4**).

132 Neural parameters that best explain TBS effect over time
To identify which neural parameters best explain TBS-induced changes across time, we
combined each subject's DCM estimates from all sessions into a single PEB and compared a
series of reduced models. Each reduced model varied the inclusion of intrinsic properties and
extrinsic connections, spanning Pre, Post1 (2 min), Post2 (15 min), and Post3 (30 min).

Bayesian model comparison revealed the full model, which simultaneously incorporates intrinsic
microcircuit parameters, intrinsic self-connections, and extrinsic interregional coupling, as
having the highest PEB free energy. This finding suggests that TBS induces hierarchical
changes from interregional connectivity down to cellular dynamics (**Figure 5**).

Based on prior rTMS literature demonstrating both sustained and transient neural effects[1,6,30],
we specified two temporal regressors: a sustained effect regressor ([-1 1 1 1]) to capture
changes persisting up to 30 minutes post-stimulation, and a transient effect regressor ([-1 1 1 -
1]) to detect changes that return to baseline after 30 minutes. Separate estimations for each
protocol (iTBS, cTBS, sham) showed evidence for both sustained and transient modulation
across protocols. To isolate parameters specific to each TBS condition, we performed between-
protocol contrasts against sham using a design matrix of [-1 1]. This analysis highlighted
protocol-specific intrinsic and extrinsic parameter changes, distinguishing neuromodulatory
profiles of iTBS and cTBS beyond shared effects observed in sham (**Figure 3.B**).

150 Sustained and transient changes following cTBS vs Sham
Figure 6A depicts the sustained and transient cTBS-induced changes relative to sham. We
observed a transient reduction in NMDA and AMPA-mediated forward connectivity from right
DLPFC to left sgACC, and in AMPA-mediated forward coupling from left sgACC to right DLPFC.
Interestingly, iTBS produced similar transient decreases, indicating a common short-lived
modulation for both TBS protocols beyond sham **(Table 2).**

The inhibitory effect of cTBS in the left DLPFC was transient: we found increase II self-inhibition
that returned to baseline by 30 minutes. No sustained intrinsic changes were observed in left

DLPFC. However, right DLPFC exhibited a sustained reduction in II→DP, reflecting prolonged
disinhibition. Extrinsic connectivity alterations following cTBS largely reverted to baseline by
30 minutes, with no enduring changes.

In the bilateral sgACC, microcircuit dynamics exhibited sustained hypoactivity after cTBS: left
sgACC showed increased self-inhibition of SS cells, while right sgACC demonstrated decreased
SS→SP and SS→II.

Sustained and transient changes following iTBS vs Sham
Figure 6B illustrates iTBS-specific effects compared to sham. Most extrinsic connectivity
modulations returned to baseline by 30 minutes, except for a sustained increase in AMPA-
mediated backward connectivity from right sgACC to left sgACC. Intrinsic parameters revealed
sustained changes in all four regions. In left DLPFC, II→SP increased, whereas right DLPFC
showed increase excitatory connection from SS→SP. The left sgACC additionally showed local
inhibition following iTBS, evidenced by increased self-inhibition in SS cells and enhanced
excitatory connection from DP→II **(Table 3).** .

Cellular property changes following TBS
Our model's conductance-based architecture also permits estimation of receptor time constants
for AMPA, NMDA, and GABA(A) channels. Compared to sham, iTBS induced sustained
increases in NMDA time constants and decreases in AMPA time constants in left DLPFC,
alongside prolonged increases in AMPA time constants in left sgACC. Conversely, cTBS
produced sustained decreases in NMDA time constants in right DLPFC and reductions in
GABA(A) time constants in right sgACC. Although channel dynamics interact nonlinearly with
intrinsic connectivity[24], these lasting receptor time constant changes in sgACC align with the
observed microcircuit hypoactivity—namely, increased AMPA time constants after iTBS and
reduced GABA(A) time constants after cTBS, both reflecting enduring inhibitory effects over
sgACC that is absent in sham.

Discussion
Spectral DCM with a canonical microcircuit model incorporating NMDA has recently gained
increasing attention in computational neuropharmacology for assessing drug effects across
hierarchical biological levels[24,31,32]. In this study, for the first time, we aimed to investigate TBS-
induced changes in the DLPFC-sgACC microcircuit network using this approach in healthy
adults. These regions have been repeatedly implicated in the pathophysiology of depression
and are known to play a crucial role in mediating the antidepressant effects of rTMS[11,16,33–35].

Recent TMS/fMRI studies consistently report that DLPFC stimulation elicits both acute and
lasting changes in sgACC activity[14,15,36]. Trapp et al. provided the first direct electrophysiological
evidence showing that sgACC activity is modulated by DLPFC stimulation, using TMS combined
with intracranial EEG[16]. Given the role of sgACC hyperactivity in depression, the therapeutic
effects of rTMS are thought to involve indirect regulatory modulation of this region [18].
Furthermore, fMRI studies have reported that sgACC–DLPFC connectivity is enhanced in MDD
patients[33,34] and that successful rTMS treatment downregulates this connectivity[37]. Notably, we
found that both TBS protocols, but not Sham, reduced right DLPFC to left sgACC effective
connectivity that is NMDA-dependent. Although these extrinsic connectivity changes were
transient, returning to baseline within 30 minutes post-stimulation, they induced lasting changes
within the sgACC microcircuit.

Unlike Sham, both TBS protocols reduced left sgACC activity by increasing self-inhibition on
spiny satellite cells and modulating receptor dynamics. Specifically, we found a decrease in the
GABA-A time constant in the right sgACC following cTBS and an increase in the AMPA time
constant in the left sgACC following iTBS. This suggests that dynamic changes in GABA-A and
AMPA receptors also contribute to sgACC inhibition. We ensured that these changes were not
driven by fatigue or other confounding factors by identifying sustained and transient changes
that differed from Sham in the between-subject PEB design. This finding aligns with a large
randomized clinical trial that analyzed source-localized TMS-evoked potentials in MDD patients
following rTMS treatment over the left DLPFC[37]. In particular, they found that higher subgenual
cingulate cortex (SGC) source activation and DLPFC-SGC connectivity, as measured by
significant current scattering, were attenuated after active rTMS and significantly correlated with
symptom improvement in MDD patients. Furthermore, they argued that this reduction in
connectivity is mediated by GABAergic neurotransmission, based on the observation of an
evoked potential at 100 milliseconds post-TMS pulse, associated with neuronal inhibitory
processes[37]. Our results, using source localization of resting-state EEG data and a model-based
effective connectivity approach, similarly found this effect in healthy individuals.

Additionally, we found evidence for an immediate inhibitory effect of cTBS and a disinhibitory
effect of iTBS on the left DLPFC at the group level. However, within-subject analysis showed
that these changes were not present in all participants, with some individuals even showing
opposite effects. Initially, we hypothesized that these changes would persist for up to 30
minutes after stimulation. However, we found that these effects were transient, and the DLPFC
microcircuit connections that changed and lasted over time did not follow this trend. In
particular, iTBS induce lasting inhibition over stimulation target with increasing inhibitory
connection to superficial pyramidal cells, while cTBS induce no sustain change over left DLPFC
that differed from Sham. This finding speaks against the notion of a sustained inhibitory or
excitatory effect of cTBS and iTBS over prefrontal regions. The local effects of TBS over the
DLPFC stimulation target have also been mixed in neuroimaging studies[4,19]. For example,
Shang et al. found no significant cerebral blood flow changes in the DLPFC following cTBS[38],
whereas Chang et al.[20] reported an increased BOLD response in the stimulation target following
iTBS. In a comprehensive review of concurrent TMS-fMRI studies, the authors argued that TMS
does not consistently increase BOLD activity at the site of stimulation, perhaps due to
alternating periods of increased and decreased neuronal firing that cancel each other out[39]. At
the individual neuronal level, Romero et al.[6] found non-uniform responses following cTBS over
parietal neurons in awake, behaving rhesus monkeys. While they observed an overall inhibitory
effect at the population level, peaking 30–40 minutes post-stimulation, transient phases of
hyperexcitability were also noted[6]. Together, our findings challenge the assumed sustained
effect of TBS protocols at the stimulation target and further demonstrate that the TBS effect is
variable between subjects and dynamically changes over time.

Direct comparisons between cTBS and iTBS effects over time revealed both shared and distinct
patterns. cTBS induced broader network-level effects at 2 minutes post-stimulation, consistent
with previous neuroimaging findings[2]. In contrast, iTBS produced more sustained changes in
intrinsic connectivity. Both protocols increased inhibition within the left sgACC and enhanced
excitation in the right DLPFC; however, only iTBS led to lasting modulation at the stimulation
target. A recent computational modeling study of longitudinal TMS-EEG data in 90 MDD
patients undergoing iTBS treatment showed that iTBS suppresses low-frequency TMS-evoked
potentials (TEPs) oscillatory power by modulating inhibition, with treatment responders

exhibiting stronger inhibitory interneuron-to-pyramidal cell connections using the Jansen-Rit
neural mass model[40]. In line with this, our results also revealed increased inhibition from
inhibitory interneurons to superficial pyramidal cell (II→SP) in the left DLPFC following iTBS, but
not after sham or cTBS. Furthermore, a meta-analysis of 23 randomized controlled trials
comparing TBS protocols in depression found that iTBS over the left DLPFC was associated
with greater symptom improvement and higher remission rates than cTBS. Together, these
findings suggest that the superior clinical efficacy of iTBS may be driven by its ability to induce
more durable intrinsic microcircuit changes, rather than simply acting as an excitatory protocol.

A recent DCM study by Pantazatos et al.,[41] which modeled fMRI time-varying connectivity
between the DLPFC and sgACC following TMS, found that TMS modulates both intrinsic
connectivity as well as bottom-up and top-down connectivity between these regions. Aligning
with this study, we found that our full model, incorporating a combination of intrinsic and
extrinsic connections, could best explain the TBS effect. This clearly highlights the complex
hierarchical neurobiological effects of TMS, creating a "chain of causation."[22] We further found
that these neuromodulatory effects dynamically change over time. For example, a few
parameters showed a negative change in the transient phase, followed by a positive shift in the
sustained regressor (see **Figure 6**), suggesting a potential compensatory neural mechanism
emerging during the later post-stimulation period (Post 3). A promising direction for future work
would be to apply data-driven clustering methods to the estimated neural parameters to better
capture possible connectivity changes over time.

This cascade of effects may be confounded by multiple factors, and we argue that such neural
changes should not be attributed solely to TBS without high-temporal-resolution measurements,
carefully controlled sessions, and robust statistical inference to capture dynamic processes over
time. In our modeling approach, we could clearly observe these confounding factors reflected in
the neural parameters. For example, previous studies have suggested that sgACC-induced
activity following DLPFC stimulation might be explained by emotional (e.g., pain) or
somatosensory experiences, due to the dense sgACC connections to limbic regions, including
the amygdala and its role in affective processing[16]. Yet, the question of whether sgACC-induced
activity is confounded by emotional or somatosensory experiences, or solely driven by DLPFC
stimulation, cannot be easily captured by the slower and indirect nature of fMRI BOLD signals.
Here, we found evidence for both. Immediately after stimulation, the only parameter change
following the Sham protocol was observed in the left sgACC, where we found reduced self-
inhibition in spiny satellite cells, indicating sgACC disinhibition. In the cTBS session, we also
found bilateral sgACC hyperactivity after Post1. We believe these early changes, particularly
following Sham and cTBS, could be better interpreted as an initial emotional or sensory reaction
rather than direct neuromodulatory effects of stimulation. However, including parameters from
post 2 (-15 minutes after stimulation) and post 3 (-30 minutes after stimulation) in our PEB
analysis, and comparing them with Sham, revealed sustained sgACC inhibition following TBS
that was not present in Sham. This finding could support the previously proposed
antidepressant effect of TBS on reducing sgACC hyperactivity[10,17,37]. Future studies are
necessary to replicate these results in MDD patients and, more importantly, to evaluate how
these parameters could predict clinical improvements.

The current dataset also includes TEPs recorded following the TBS protocols. Analysis of TEP
component amplitudes within a predefined left DLPFC ROI (i.e., average of F3, FC1, FC3) has
been reported elsewhere, showing no significant differences between active and sham

conditions[42]. Our present study, by contrast, focuses on resting-state EEG using a fundamentally different analytic pipeline. Although TEP components have been proposed as sensitive markers of cortical excitability, their underlying electrophysiological mechanisms remain poorly understood, and they exhibit substantial inter-individual variability[4]. In contrast, the biological plausibility of spDCM and neural mass models offers a more nuanced and mechanistic account of cortical excitability at the microcircuit level[26]. Additionally, spDCM approach using the CMM-NMDA model has recently demonstrated strong reliability and reproducibility in resting-state electrophysiology[28], and the multilevel Bayesian approach (PEB of PEB) incorporates group-level informative priors at both the within-subject and between-subject levels to address heterogeneity across individuals[29]. While we argue that this technique is especially promising for tracking dynamic neurophysiological responses to TMS, several methodological challenges remain. First, the choice of window length in resting state EEG for spDCM analysis has not been well established[43]; we used a 2-second window here following[24], although prior studies have also applied 1-second[28,43] 5-second[31] or even 10-second[44] window. Additionally, the current neuronal model cannot explain how on-level changes affect other neural parameters (e.g., how changes in time constants affect intrinsic connectivity, and vice versa), making the model blind to temporal orders[24]. Addressing this issue would require a biologically grounded model for the trans-hierarchical progression of TBS effects rather than the current all-in-one CMM-NMDA model[24]. Nonetheless, given the non-invasive nature, low cost, and portability of resting-state EEG compared to TMS-fMRI or TMS evoked potentials, this method holds considerable promise for tracking time-dependent changes in brain connectivity in future neuromodulation research.

There are important limitations to consider when interpreting the results. First, our finding only examines DLPFC-sgACC connectivity in healthy individuals, which may not be replicable in MDD patients. Although previous studies have shown that DLPFC-sgACC functional connectivity from normative connectomes can predict clinical response comparably to disease-matched connectomes[10,11], differences in connectivity have been reported in patients with depression[33,34,37,41]. For example, Pantazatos et al. showed that TMS modulates DLPFC-sgACC effective connectivity differently in healthy individuals compared to patients[41]. Future studies are needed to directly compare the current results in clinical populations. Second, TBS was applied at 75% of the RMT, following Chung et al.[45], who reported the strongest iTBS effects in the prefrontal cortex at this intensity. However, it is important to note that this relatively low stimulation intensity may limit comparability with clinical protocols, which typically use 100% or higher.

With these limitations in mind, we have uncovered protocol-specific and receptor-mediated dynamics of TBS between the DLPFC and sgACC by using spectral DCM with biophysically realistic microcircuit models. These results provide mechanistic insight into how TBS may achieve therapeutic modulation of the sgACC and offer a powerful computational toolkit for future precision neuromodulation studies.

Method

Dataset

The EEG recordings analyzed in this study were obtained from a peer-reviewed open-source dataset[46] (Available: https://doi.org/10.25452/figshare.plus.c.5910329). This dataset employed a single-blinded, sham-controlled crossover design. Twenty four healthy right-handed participants (11 females with mean age of 25.2 ± 9.9 years) attended five sessions scheduled at the same

time of day, each separated by at least one week to minimize carryover effects. Session order
was pseudo-randomized via computer-generated sequences: three initial sessions, each
comprising one of iTBS, cTBS, or sham (randomized), followed by two additional sessions (one
each of iTBS and cTBS). Participants were excluded if they met any of the following criteria: (1)
a history of psychiatric or neurological disorders, including seizures or stroke; (2) recent head
injury; (3) current use of medications that affect cognitive function; (4) drug or alcohol abuse
within the past three months; (5) smoking; (6) pregnancy; or (7) any contraindications to EEG or
MRI.

Resting motor thresholds (RMTs) were determined individually by identifying the minimum
stimulus intensity required to evoke at least 3 out of 6 motor-evoked potentials (MEPs) with a
peak-to-peak amplitude of ≥50 µV in the contralateral right first dorsal interosseous muscle.
Electromyographic (EMG) recordings were obtained using a 1401 laboratory interface and 1902
amplifier (Cambridge Electronic Design, Cambridge, UK), with data acquired via the Signal V4
software (Cambridge Electronic Design).

TBS and single-pulse TMS were administered using a MagPro® X100 stimulator (MagVenture,
Lucernemarken, Denmark) equipped with a 65 mm Cool-B65 figure-8 coil. The coil was
positioned tangentially to the scalp over the F3 electrode location to target the left DLPFC, and
TBS was delivered at 75% of each individual's RMT[46]. For the iTBS protocol, stimulation
consisted of two-second trains delivered every 10 seconds over 192 seconds, totaling 600
pulses[5]. The cTBS protocol involved 600 uninterrupted pulses delivered over 40 seconds[5]. The
sham condition used an inactive coil placed directly on the scalp, while a second active coil,
positioned 20 cm posterior and oriented away from the head, was used to simulate the TMS
clicking sound. This posterior coil operated at 20% higher intensity to account for the increased
distance from the ear and delivered either an iTBS or cTBS pattern (randomly assigned across
participants). To further enhance blinding, participants wore earphones playing white noise at a
volume sufficient to mask the auditory artifact of the stimulation, adjusted to the highest level
tolerable for each individual[46].

Four minutes of eyes-open resting-state EEG were recorded at the session's start and after
completing all four blocks of single-pulse TMS, as illustrated in **Figure 2**. EEG data were
recorded using a Refa amplifier (2048 Hz sampling rate; TMSi, Oldenzaal, The Netherlands)
and a 64-channel 10–20 EEG cap (EasyCap, GmbH, Herrsching, Germany), with cap size
chosen per participant. Electrodes were grounded at Fpz and captured with a common average
reference. Recordings took place in a dimly lit, sound-attenuated room, with participants fixating
on a cross displayed on a 58.4 cm (23-inch) monitor.

EEG preprocessing
The data were processed and analyzed offline.   This was done using a combination of open-
source toolboxes, including Fieldtrip[47], EEGLAB (version 2022b)[48], and custom scripts, all
implemented on the MATLAB platform (R2022b, The MathWorks, USA).

Eyes-open resting-state EEG data were preprocessed by down-sampling to 256 Hz, baseline
correction (demeaning), and detrending to remove any linear trends. A second-order bandpass
filter (1–50 Hz) and a 50 Hz notch filter were applied to eliminate electrical line noise. The data
were segmented into 2-second epochs, with an automated algorithm first rejecting epochs
exhibiting data ranges exceeding three standard deviations or absolute maxima surpassing 12
standard deviations of other epochs. Remaining noisy epochs were excluded through visual

inspection. Independent Component Analysis (ICA) was performed once to identify and remove
components associated with eye blinks, eye movement and muscle artifacts. The data were
then re-referenced to the common average reference. One of the subjects exclude at this stage
due to poor quality in Sham session.

Dynamic causal modelling for resting state EEG

Preprocessed data were imported into SPM25[49] (revision 25.01.rc3) to perform DCM for cross-
spectral densities (CSD) (https://www.fil.ion.ucl.ac.uk/spm/). This steady-state variant uses a
conductance-based canonical microcircuit model (cmm_NMDA) to estimate how endogenous
neuronal fluctuations produce the observed CSD[27]. DCM inverts a generative neural mass
model alongside an observation model to infer intrinsic and extrinsic connectivity parameters
that best explain the empirical data[27].

Conductance-Based Neural Mass Model

The Conductance-Based Neural Mass Model with NMDA (CMM-NMDA) is a variant of
electrophysiological DCM. This model describes the dynamic behavior of neural populations by
incorporating the conductance of three primary ionotropic receptors: AMPA, GABA-A, and
NMDA receptors. The conductance of these channels is determined by the cumulative synaptic
interactions within and between cortical regions. In the CMM-NMDA framework, neural activity
emerges from four key neural populations: excitatory spiny stellate cells (SS), superficial
pyramidal cells (SP), inhibitory interneurons (II), and deep pyramidal cells (DP). These
populations are interconnected via intrinsic connections within the cortical regions. Moreover,
the model captures the extrinsic connectivity between cortical regions[27]. We can describe the
CMM-NMDA model in 3 levels: intrinsic properties, intrinsic connections, and extrinsic
connections (**Figure 1**).

Neural population properties

405 Each population in a cortical region is modelled by a Morris–Lecar model[50] and contains four
406 hidden states: the membrane voltage V and the conductance gk for all three receptors: AMPA,
407 GABA-A, and NMDA[51,52]. The mean membrane voltage of population j is obtained by this
408 equation:

$$\dot{\mu}_V^{(j)} = \frac{1}{C^{(j)}} \left(\sum_k \mu_{g_k}^{(j)} \left(V_k - \mu_V^{(j)} \right) \right) + \mu_{g_{NMDA}}^{(j)} m\left(\mu_V^{(j)} \right) \left(V_{NMDA} - \mu_V^{(j)} \right)$$

$$k \in L, AMPA, GABA$$

Where:
j is the target neural population;
$\dot{\mu}_V$ is the time derivative of mean voltage;
μ_V is the mean voltage;
C is the membrane capacitance;
μ_{g_k} is the mean conductance of channel k;
$\mu_{g_{NMDA}}$ is the mean conductance of the NMDA channel;
V_k is the reversal potential of channel k;
V_{NMDA} is the mean reversal potential of the NMDA channel;
L is a passive leak channel, which means its conductance is a fixed number (1 in SPM25). Each
channel's conductance is voltage-time-dependent. The mean conductance of channel k of
population j is obtained by this equation:
422

$$\dot{\mu}_{g_k}^{(j)} = \kappa_k^{(j)}(\zeta_k^{(j)} - \mu_{g_k}^{(j)})$$

$$k \in L, AMPA, GABA, NMDA$$

424 Where:
425 $\dot{\mu}_{g_k}$ is the time derivative of the mean conductance of channel k;
426 μ_{g_k} is the mean conductance of channel k;
427 κ_k is the inverse time constant of channel k;
428 ζ_k is the input to the target neural population;
429
430 NMDA has a voltage-dependent magnesium block, too:
431

$$m(V) = \frac{1}{1 + 0.2 \cdot \exp(-\alpha_{NMDA} \cdot V)}$$

432
433 Where α_{NMDA} is the magnesium block parameter, which is a fixed parameter (Table 1).
434

Intrinsic connection
436 The populations are interconnected via intrinsic connections within the cortical regions. The
437 input to the target population is the cumulative output of activated origin populations:

$$\zeta_k^{(j)} = \sum_i \gamma_k^{(j,i)} \, \sigma(\mu_k^{(i)} - V_R, \Sigma^i)$$

Where:

j is the target neural population;
i is the origin neural population;
ζ_k is the input to the target neural population via channel k;
γ_k is the intrinsic coupling parameter from origin to target neural population via channel k;
α is the function for proportion of activated neurons in the origin population and is formulated as
a cumulative distribution of the univariate normal distribution $\mathcal{N}(\mu_k^{(i)} - V_R, \Sigma^i)$ where V_R is the
threshold potential (-40 in SPM25) and Σ^i is a free parameter.
448

Extrinsic connection
450 Extrinsic connections are divided into 2 types: Forward connections, which run from superficial
451 pyramidal cells of lower cortical regions to spiny stellate cells of higher regions. Backward
452 connections, which run from deep pyramidal cells of higher to superficial pyramidal and
453 inhibitory interneurons of lower brain regions.

Model specification and inversion
Anatomical coordinates of network nodes were defined a priori. Individual T1-weighted MRI
scans were utilized to construct subject-specific head models and lead fields, ensuring accurate
source localization. We specified a fully connected network comprising bilateral DLPFC and
sgACC, modeling each region as a cortical patch using the "IMG" option in SPM25. Nodes were
placed at the following MNI coordinates: left DLPFC[10] (–38, 44, 26), right DLPFC (38, 44, 26),
left sgACC (–10, 20, –15) and right sgACC (5, 15, –15)[53]. For each subject, the individuals
high-resolution T1-weighted MRI was first reoriented and co-registered to EEG electrode

coordinates of that session using fiducial landmarks (nasion and preauricular points) and head-
shape points, ensuring precise alignment of sensor and anatomical space. The MRI was then
segmented into scalp, skull and brain compartments to construct an individual three-layer
Boundary Element Method (BEM) head model, and a fine-resolution cortical mesh (mesh size =
3) was generated for accurate forward modeling and source localization[54]. Before inversion,
data registration and mesh quality were visually inspected and saved for each session.

DCM Model inversion was performed using variational Bayes under the Laplace approximation,
allowing for the estimation of parameters governing both neural dynamics and the observation
model[51,55]. Parameter estimation involved maximizing the variational free energy defined as:

$$F(q, y) = \langle \log p\,(y|\theta) \rangle_q - D_{KL}[q(\theta)|p(\theta)]$$

471   where $p(y \mid \theta)$ represents the likelihood function, and D_{KL} is the Kullback–Leibler divergence
472   between the recognition density $q(\theta)$ and the prior distribution $p(\theta)$. Upon convergence, the
473   recognition density $q(\theta)$ provides an approximate posterior distribution $p(\theta \mid y)$. Model fit was
474   assessed by explained variance; ; Which compute as:

$$EV = 100 \cdot \frac{\sum_{e,f} CSD(e,f)^2}{\sum_{e,f} CSD(e,f)^2 + \sum_{e,f}\left(CSD(e,f) - \widetilde{CSD}(e,f)\right)^2}$$

475   Where $CSD(e,f)$ denotes the observed cross-spectral density (CSD) at frequency f for edge e,
476   and $\widetilde{CSD}(e,f)$ is the model-estimated CSD at the same frequency and edge. Participants with
477   explained variance below 50% were excluded, resulting in removal of one subject prior to group-
478   level analysis **(Supplementary Material, Figure S1).**

Group level analysis using multilevel parametric empirical bayes
Following DCM parameter estimation, we applied a parametric empirical Bayesian (PEB)
framework with SPM25 to examine systematic variations in model parameters induced by TBS
and Sham over 22 subjects.

While an alternative approach would involve applying classical statistics (e.g., repeated-
measures ANOVA) to the expected posterior means (Ep), this method overlooks the underlying
uncertainty associated with DCM parameters[56]. Since DCM estimates are multivariate normal
distributions, relying solely on point estimates disregards the posterior covariance (Cp), which
reflects the confidence in parameter estimation[56]. In contrast, the PEB framework incorporates
both Ep and Cp in group-level inference. This means that participants with more precise
parameter estimates contribute more strongly to the group result, whereas those with noisier
data are down-weighted[56]. This property is particularly advantageous in clinical or high-
variability datasets, where reliable individual-level inference is critical.

A further challenge in working with nonlinear and ill-posed models—such as neural mass
models applied to resting-state EEG—is the risk of converging on different local optima when
fitting multiple full DCMs across sessions and subjects[29]. This issue emphasizes the importance
of hierarchical modeling frameworks like PEB that can stabilize inferences despite local
variations in model fits. To address the potential for session-level variability and local optima in
DCM inversion, we used a multilevel PEB framework specifically adapted for longitudinal DCM
analysis[29]. In this hierarchical model, within-subject longitudinal effects across sessions were
modeled at the second level, and between-subject effects were modeled at the third level.

Empirical priors derived at each level were used to iteratively optimize posterior densities over
parameters in the level below. At the first level, subject- and session-specific spDCMs were
estimated. At the second and third levels, Bayesian general linear models were used to capture
within- and between-subject effects, respectively[29]. The model hierarchy can be expressed as:

$$y_{ij} = \Gamma\big(\theta_{ij}^{(1)}\big) + \varepsilon_{ij}^{(1)}, \quad \varepsilon_{ij}^{(1)} \sim N\big(0, \Sigma^{(1)}\big)$$

$$\theta_{ij}^{(1)} = X_i \beta_j^{(2)} + \varepsilon_j^{(2)}, \quad \varepsilon_j^{(2)} \sim N\big(0, \Sigma^{(2)}\big)$$

$$\beta_j^{(2)} = X_2 \beta^{(3)} + \varepsilon^{(3)}, \quad \varepsilon^{(3)} \sim N\big(0, \Sigma^{(3)}\big)$$

In the first equation, the observed time series data yij for the ith session and jth subject are
modeled as the output of a nonlinear DCM function Γ plus observation noise $\varepsilon_{ij}^{(1)}$. The model
parameters $\theta_{ij}^{(1)}$ are sampled from the subject-specific second-level model jth , which
incorporates session-level effects (pre, post1, post2, post3) with design matrix $X1$ and
parameters $\beta_j^{(2)}$, with added noise $\varepsilon_j^{(2)}$.

The within-subject design matrix X_1 used in our study was defined as: X_1 = [1 1 1 1; -1 1 1 1; -1
1 1 -1]. This allowed us to separate baseline, sustained, and transient post-TBS effects for each
protocol. At the third level, subject-specific effects $\beta_j^{(2)}$ were modeled using the between-subject
design matrix X_2 and group-level parameters $\beta^{(3)}$. To compare TBS with Sham conditions
across all subjects, we used: X_2 = [1 1; -1 1]. This design tested for common and differential
effects of TBS relative to Sham (see **Figure 3**).

In this approach, extrinsic, intrinsic, and cellular parameters $\beta_j^{(2)}$ can be modeled as either fixed
or random effects by specifying appropriate priors on between-subject variability in the
hierarchical model[56]. To determine which parameters should be treated as random effects, we
performed Bayesian model comparison by evaluating the free energy of reduced models using
the "softmax" function[25]. This approach offers two key advantages: (1) it quantifies the
contribution of each neural parameter in explaining TBS effects, and (2) it identifies the simplest
model that best explains group differences.

We explored three classes of hypotheses regarding the parameters: (1) TBS effects are best
explained through changes in extrinsic (between-region) connections; (2) TBS effects are better
accounted for by modulations in intrinsic (within-region) microcircuit parameters; and (3) TBS
affects both intrinsic and extrinsic connectivity. The model yielding the highest free energy was
selected and treated as a random effect ("field" in the spm_dcm_peb function) for subsequent
analyses.

Following the estimation of group-level effects, we employed automatic model search using
Bayesian model reduction (BMR) and Bayesian model averaging (BMA) via the
spm_dcm_peb_bmc function[56]. This greedy search algorithm iteratively removes parameters
that do not contribute to model evidence. BMA was then performed across the 256 reduced
models from the final iteration. Comparing full and reduced models in this way is conceptually
similar to conducting a post-hoc F-test in classical statistics and is referred to as 'post-hoc' DCM
analysis[56]. Lastly, it is important to note that group-level Bayesian analysis does not rely on the
concept of significance in classical statistics. Instead, it provides posterior probabilities for each

parameter. However, it is a common practice in DCM studies to report the most probable effects
with posterior probability > 95, labelled as strong evidence based on Kass and Raftery[57].

Data and code availability
539 Raw EEG data available here: https://doi.org/10.25452/figshare.plus.c.5910329. Complete
540 MATLAB and Python code used to estimate DCM and plotting parameters are available in the
541 Github repository: https://github.com/ArminTi/Hierarchical-neurobiological-changes-in-DLPFC-
542 sgACC-connectivity-induced-by-Theta-Burst-Stimulation/tree/main

Authors contribution
544 A.T. conceptualized the study, analyzed the data, and wrote the manuscript. G.G. analyzed the
545 data, created graphical illustrations, and assisted with manuscript review and editing. S.Z.
546 participated in data analysis and manuscript editing. B.A. and H.M. contributed to data analysis.
547 A.N. supervised the project.

Competing interests.
549 The authors declare no competing interests.

References

1. Tik, M. *et al.* Towards understanding rTMS mechanism of action: Stimulation of the
DLPFC causes network-specific increase in functional connectivity. *Neuroimage* **162**,
289–296 (2017).

2. Hermiller, M. S. Effects of continuous versus intermittent theta-burst TMS on fMRI
connectivity. *Front Hum Neurosci* **18**, (2024).

3. Suppa, A. *et al.* Ten Years of Theta Burst Stimulation in Humans: Established
Knowledge, Unknowns and Prospects. *Brain Stimul* **9**, 323–335 (2016).

4. Speranza, B. E. *et al.* The Neurophysiological Effects of Theta Burst Stimulation as
Measured by Electroencephalography: A Systematic Review. *Biol Psychiatry Cogn*
Neurosci Neuroimaging **9**, 1083–1120 (2024).

5. Huang, Y.-Z., Edwards, M. J., Rounis, E., Bhatia, K. P. & Rothwell, J. C. Theta Burst
Stimulation of the Human Motor Cortex. *Neuron* **45**, 201–206 (2005).

6. Romero, M. C., Merken, L., Janssen, P. & Davare, M. Neural effects of continuous theta-
burst stimulation in macaque parietal neurons. *Elife* **11**, (2022).

7. Toghi, A., Chizari, M. & Khosrowabadi, R. A causal role of the right dorsolateral prefrontal
cortex in random exploration. *Sci Rep* **14**, 24796 (2024).

8. Blumberger, D. M. *et al.* Effectiveness of theta burst versus high-frequency repetitive
transcranial magnetic stimulation in patients with depression (THREE-D): a randomised
non-inferiority trial. *The Lancet* **391**, 1683–1692 (2018).

9. Huang, Y. *et al.* Theta-burst direct electrical stimulation remodels human brain networks.
Nat Commun **15**, 6982 (2024).

10.   Fox, M. D., Buckner, R. L., White, M. P., Greicius, M. D. & Pascual-Leone, A. Efficacy of
Transcranial Magnetic Stimulation Targets for Depression Is Related to Intrinsic
Functional Connectivity with the Subgenual Cingulate. *Biol Psychiatry* **72**, 595–603
(2012).

11.   Weigand, A. *et al.* Prospective Validation That Subgenual Connectivity Predicts
Antidepressant Efficacy of Transcranial Magnetic Stimulation Sites. *Biol Psychiatry* **84**,
28–37 (2018).

12.   Boes, A. D. *et al.* Rostral anterior cingulate cortex is a structural correlate of repetitive
TMS treatment response in depression. *Brain Stimul* **11**, 575–581 (2018).

13.   Benschop, L. *et al.* Reduced subgenual cingulate–dorsolateral prefrontal connectivity as
an electrophysiological marker for depression. *Sci Rep* **12**, 16903 (2022).

14.   Oathes, D. J. *et al.* Resting fMRI-guided TMS results in subcortical and brain network
modulation indexed by interleaved TMS/fMRI. *Exp Brain Res* **239**, 1165–1178 (2021).

15.   Duprat, R. J. *et al.* Resting fMRI-guided TMS evokes subgenual anterior cingulate
response in depression. *Neuroimage* **305**, 120963 (2025).

16.   Trapp, N. T. *et al.* Dorsolateral prefrontal cortex TMS evokes responses in the subgenual
anterior cingulate cortex: Evidence from human intracranial EEG. Preprint at
https://doi.org/10.1101/2024.12.20.629857 (2024).

17.   Cappon, D. *et al.* Transcranial magnetic stimulation (TMS) for geriatric depression.
Ageing Res Rev **74**, 101531 (2022).

18.   Seguin, C. *et al.* Neuroanatomical pathways of TMS therapy for depression. Preprint at
https://doi.org/10.1101/2025.02.10.25322034 (2025).

19.   Kirkovski, M. *et al.* A systematic review of the neurobiological effects of theta-burst
stimulation (TBS) as measured using functional magnetic resonance imaging (fMRI).
Brain Struct Funct **228**, 717–749 (2023).

20.   Chang, K.-Y. *et al.* Neural response during prefrontal theta burst stimulation: Interleaved
TMS-fMRI of full iTBS protocols. *Neuroimage* **291**, 120596 (2024).

21.   Grosshagauer, S. *et al.* Chronometric TMS-fMRI of personalized left dorsolateral
prefrontal target reveals state-dependency of subgenual anterior cingulate cortex effects.
Mol Psychiatry **29**, 2678–2688 (2024).

22.   Bergmann, T. O. & Hartwigsen, G. Inferring Causality from Noninvasive Brain Stimulation
in Cognitive Neuroscience. *J Cogn Neurosci* **33**, 195–225 (2021).

23.   Banaraki, A. K., Toghi, A. & Mohammadzadeh, A. RDoC Framework Through the Lens of
Predictive Processing: Focusing on Cognitive Systems Domain. *Computational*
Psychiatry **8**, 178–201 (2024).

24.   Eo, J., Kang, J., Youn, T. & Park, H.-J. Neuropharmacological computational analysis of
longitudinal electroencephalograms in clozapine-treated patients with schizophrenia
using hierarchical dynamic causal modeling. *Neuroimage* **275**, 120161 (2023).

25. Rosch, R. E., Auksztulewicz, R., Leung, P. D., Friston, K. J. & Baldeweg, T. Selective
Prefrontal Disinhibition in a Roving Auditory Oddball Paradigm Under N-Methyl-D-
Aspartate Receptor Blockade. *Biol Psychiatry Cogn Neurosci Neuroimaging* **4**, 140–150
(2019).

26. Bastos, A. M. *et al.* Canonical Microcircuits for Predictive Coding. *Neuron* **76**, 695–711
(2012).

27. Pereira, I. *et al.* Conductance-based dynamic causal modeling: A mathematical review of
its application to cross-power spectral densities. *Neuroimage* **245**, 118662 (2021).

28. Jafarian, A. *et al.* Reliability of dynamic causal modelling of resting state
magnetoencephalography. *Hum Brain Mapp* **45**, (2024).

29. Park, H.-J. *et al.* Hierarchical Dynamic Causal Modeling of Resting-State fMRI Reveals
Longitudinal Changes in Effective Connectivity in the Motor System after Thalamotomy
for Essential Tremor. *Front Neurol* **8**, (2017).

30. Romero, M. C., Davare, M., Armendariz, M. & Janssen, P. Neural effects of transcranial
magnetic stimulation at the single-cell level. *Nat Commun* **10**, 2642 (2019).

31. Symmonds, M. *et al.* Ion channels in EEG: isolating channel dysfunction in NMDA
receptor antibody encephalitis. *Brain* **141**, 1691–1702 (2018).

32. Shine, J. M. *et al.* Computational models link cellular mechanisms of neuromodulation to
large-scale neural dynamics. *Nat Neurosci* **24**, 765–776 (2021).

33. Philippi, C. L., Motzkin, J. C., Pujara, M. S. & Koenigs, M. Subclinical depression severity
is associated with distinct patterns of functional connectivity for subregions of anterior
cingulate cortex. *J Psychiatr Res* **71**, 103–111 (2015).

34. Davey, C. G., Harrison, B. J., Yücel, M. & Allen, N. B. Regionally specific alterations in
functional connectivity of the anterior cingulate cortex in major depressive disorder.
Psychol Med **42**, 2071–2081 (2012).

35. Siddiqi, S. H. *et al.* Brain stimulation and brain lesions converge on common causal
circuits in neuropsychiatric disease. *Nat Hum Behav* **5**, 1707–1716 (2021).

36. Tik, M. *et al.* Acute TMS/fMRI response explains offline TMS network effects – An
interleaved TMS-fMRI study. *Neuroimage* **267**, 119833 (2023).

37. Hadas, I. *et al.* Association of Repetitive Transcranial Magnetic Stimulation Treatment
With Subgenual Cingulate Hyperactivity in Patients With Major Depressive Disorder.
JAMA Netw Open **2**, e195578 (2019).

38. Shang, Y. *et al.* Theta-burst transcranial magnetic stimulation induced functional
connectivity changes between dorsolateral prefrontal cortex and default-mode-network.
Brain Imaging Behav **14**, 1955–1963 (2020).

39. Rafiei, F. & Rahnev, D. TMS Does Not Increase BOLD Activity at the Site of Stimulation:
A Review of All Concurrent TMS-fMRI Studies. *eNeuro* **9**, ENEURO.0163-22.2022
(2022).

40. Momi, D. *et al.* Excitation-Inhibition Balance and Fronto-Limbic Connectivity Drive TMS
Treatment Outcomes in Refractory Depression. Preprint at
https://doi.org/10.1101/2025.04.23.648963 (2025).

41. Pantazatos, S. P. *et al.* The timing of transcranial magnetic stimulation relative to the
phase of prefrontal alpha EEG modulates downstream target engagement. *Brain Stimul*
16, 830–839 (2023).

42. Moffa, A. H., Nikolin, S., Martin, D., Loo, C. & Boonstra, T. W. Assessing
neuromodulation effects of theta burst stimulation to the prefrontal cortex using TMS-
evoked potentials. Preprint at https://doi.org/10.1101/2021.06.21.449219 (2021).

43. Van de Steen, F., Almgren, H., Razi, A., Friston, K. & Marinazzo, D. Dynamic causal
modelling of fluctuating connectivity in resting-state EEG. *Neuroimage* **189**, 476–484
(2019).

44. Van de Steen, F. *et al.* Dynamic causal modelling shows a prominent role of local
inhibition in alpha power modulation in higher visual cortex. *PLoS Comput Biol* **18**,
e1009988 (2022).

45. Chung, S. W. *et al.* Impact of different intensities of intermittent theta burst stimulation on
the cortical properties during TMS□EEG and working memory performance. *Hum Brain*
Mapp **39**, 783–802 (2018).

46. Moffa, A. H. *et al.* Neuromodulatory effects of theta burst stimulation to the prefrontal
cortex. *Sci Data* **9**, 717 (2022).

47. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software
for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput*
Intell Neurosci **2011**, 1–9 (2011).

48. Fayaz, M. The bibliometric analysis of EEGLAB software in the Web of Science indexed
articles. *Neuroscience Informatics* **4**, 100154 (2024).

49. Tierney, T. M. *et al.* SPM 25: open source neuroimaging analysis software. (2025).

50. Morris, C. & Lecar, H. Voltage oscillations in the barnacle giant muscle fiber. *Biophys J*
35, 193–213 (1981).

51. Marreiros, A. C., Kiebel, S. J., Daunizeau, J., Harrison, L. M. & Friston, K. J. Population
dynamics under the Laplace assumption. *Neuroimage* **44**, 701–714 (2009).

52. Moran, R. J., Stephan, K. E., Dolan, R. J. & Friston, K. J. Consistent spectral predictors
for dynamic causal models of steady-state responses. *Neuroimage* **55**, 1694–1708
(2011).

53. Imperatori, C. *et al.* Exposure to nature is associated with decreased functional
connectivity within the distress network: A resting state EEG study. *Front Psychol* **14**,
(2023).

54. Litvak, V. *et al.* EEG and MEG Data Analysis in SPM8. *Comput Intell Neurosci* **2011**, 1–
32 (2011).

55. Zeidman, P. *et al.* A guide to group effective connectivity analysis, part 1: First level
analysis with DCM for fMRI. *Neuroimage* **200**, 174–190 (2019).

56. Zeidman, P. *et al.* A guide to group effective connectivity analysis, part 2: Second level
analysis with PEB. *Neuroimage* **200**, 12–25 (2019).

57. Kass, R. E. & Raftery, A. E. Bayes Factors. *J Am Stat Assoc* **90**, 773–795 (1995).

695
Figure 1. Hierarchical schematic of the CMM_NMDA neural model across three nested
scales. (i) macroscale extrinsic forward (A1) and backward (A2) AMPA-mediated connection
(same for NMDA) between four sources; (ii) mesoscale intrinsic laminar circuitry within each
source—superficial pyramidal, spiny satellite, inhibitory interneurons and deep pyramidal cells
across supra-granular, granular and infragranular layers 2 with excitatory (green) and inhibitory
(red) interactions; and (iii) microscale conductance-based receptor and ion-channel dynamics
(AMPA, NMDA, GABA☐) that are defined at the source level—rather than per individual
neuron—to capture aggregate postsynaptic currents.

Figure 2*. Overview of experimental design and analysis pipeline.* Each participant completed three single-blinded TMS-EEG sessions (iTBS, cTBS, sham; randomized crossover, ≥1 week apart). In each session, 4 min of eyes-open resting-state EEG was recorded before TBS and after each of four single-pulse TMS blocks (pre, post[1], post[2], post[3]). EEG data were preprocessed and cleaned. Subject-specific T1-weighted MRIs were used to build three-layer BEM head models and cortical meshes. Source-level cross-spectral densities were computed and inverted in SPM25 using the conductance-based CMM_NMDA microcircuit model for each time block. Hierarchical parametric empirical Bayes used for group level analysis and Bayesian model reduction and averaging were used as an efficient form of model selection.

Figure 3. Group-level analysis using PEB of PEB. (A) Within-subject PEB uses a 2×2 matrix [1 −1; 1 1] to estimate each subject's Pre→Post[1] effect (β_1), followed by a second-level PEB with a single-regressor [1] to compute the group mean differences (β_0). (B) Within-subject PEB fits a 4×3 matrix across the four EEG blocks (pre, post[1], post[2], post[3]) with regressors for baseline [1 1 1 1], sustained change [−1 1 1 1] and transient change [−1 1 1 −1] to yield β_0 (baseline), β_1 (sustained) and β_2 (transient). A second-level PEB then contrasts TBS (iTBS or cTBS) against sham using [1 −1; 1 1] to isolate TBS-specific modulations.

Figure 4. Connectivity changes two minutes after cTBS, iTBS, and sham (pre vs post1). Parameters with 95% or higher probability, indicating strong evidence, displayed for each protocol. Red and blue indicate positive (increasing connectivity) and negative (decreasing connectivity) effects. Receptor time constant (T) illustrate beside each source, where blue corresponds to AMPA, red to $GABA_A$, and green to NMDA. A1: forward AMPA extrinsic connection, A2: backward AMPA extrinsic connection, AN1: forward NMDA extrinsic connection, AN2: backward NMDA extrinsic connection.

Figure 5. Bayesian model comparison showing the full model could best explain TBS differences. Bar plots show PEB free energy (top) and posterior model probability (bottom) for three families of reduced models—intrinsic properties only, extrinsic connections only, and combined of both. The full model, which integrates all the intrinsic and extrinsic features, achieved the highest evidence for both protocols, indicating that TBS effect could best explain with hierarchical changes from synaptic to network scales.

Figure 6. Sustain and transient changes following TBS compared to sham. (A) cTBS vs Sham (B) iTBS vs Sham. Sustained effects reflect lasting changes up to 30 min post-TBS, while transient effects capture the parameters altered post-TBS but returning to baseline by 30 min. Red and blue indicate positive (increasing connectivity) and negative (decreasing connectivity) effects. Receptor time constant (T) illustrate beside each source, where blue corresponds to AMPA, red to GABA$_A$, and green to NMDA. A1: forward AMPA extrinsic connection, A2: backward AMPA extrinsic connection, AN1: forward NMDA extrinsic connection, AN2: backward NMDA extrinsic connection.

758   **Table 1**. Description of parameters

	Parameter	Description	Reference value	Prior
Intrinsic Property	$\alpha NMDA$	Magnesium block parameter	0.06	-
	$C\ (j)$	Membrane capacitance, $j \in \{ss, sp, ii, dp\}$	0.128,0.128,0.256,0.032	$N\ (0,\ 1/16)$
	S	Population variance	32	$N\ (0,\ 1/64)$
	$T_{ch}\ (\sim 1/\kappa ch)$	Channel time constant [ms], $ch \in \{GABA, AMPA, NMDA\}$	4,16,100	$N\ (0,\ 1/64)$
	E	Background activity	0.8	$N\ (0,\ 1/64)$
Intrinsic Connectivity	$\gamma\ k(i \rightarrow j)^*$	$(i, j)\ \in \{(ss, sp), (ss, ii), (sp, dp)\}$	4	$N\ (0,\ 1/32)$
	$\gamma\ k(i \rightarrow j)$	$(i, j)\ \in \{(dp, ii), (ii, ss), (ii, dp)\}$	2	$N\ (0,\ 1/32)$
	$\gamma\ k(i \rightarrow j)$	$(i, j)\ \in \{(ii, dp), (ss, ss), (sp, sp)\}$	2	$N\ (0,\ 1/32)$
	$\gamma\ k(i \rightarrow j)$	$(i, j)\ \in \{(ii, ii)\}$	32	$N\ (0,\ 1/32)$
	$\gamma\ k(i \rightarrow j)$	$(i, j)\ \in \{(dp, dp)\}$	128	$N\ (0,\ 1/32)$
Extrinsic Connectivity	A NMDA f/b $(A \rightarrow B)$	NMDA extrinsic forward (f)/ backward (b) connectivity from region A to B	0.11	$N\ (0,\ 1/8)$
	A AMPA f/b $(A \rightarrow B)$	AMPA extrinsic forward (f)/ backward (b) connectivity from region A to B	1	$N\ (0,\ 1/8)$

759   *Note: * Intrinsic connectivity from a neural population i to j via channel k. dp: excitatory deep*
760   *pyramidal cells, ss: excitatory spiny stellate cells, sp: excitatory superficial pyramidal cells, ii:*
761   *inhibitory interneuron;*

762   **Table 2.** PEB of PEB results; *Sustain and Transient changes following iTBS compared to*
763   *Sham.*

iTBS>Sham			
Sustain		*Transient*	
Parameters	*B_1*	*Parameters*	*B_1*
T_{AMPA} (L.DLPFC)	-0.0944	T_{AMPA} (L.sgACC)	-0.0535
T_{AMPA} (L.sgACC)	0.0952	$A_{f\ AMPA}$ (L. DLPFC→R.DLPFC)	0.1860
T_{NMDA} (L.DLPFC)	0.0644	$A_{f\ AMPA}$ (L.sgACC→R.DLPFC)	-0.1960
$A_{b\ AMPA}$ (R.sgACC → L.sgACC)	0.1929	$A_{f\ AMPA}$ (LsgACC → R.sgACC)	0.1838
$H_{L.DLPFC}$ (II → SP)	0.0882	$A_{b\ AMPA}$ (R.sgACC→LsgACC)	-0.1945
$H_{R.DLPFC}$ (SS → SP)	0.1517	$A_{f\ NMDA}$ (R.DLPFC→LsgACC)	-0.1647
$H_{R.sgAAC}$ (SS → II)	-0.0914	$H_{R.DLPFC}$ (SS → SP)	-0.1039
$H_{R.sgAAC}$ (SP → DP)	0.0908	$H_{L.sgAAC}$ (SS → II)	-0.0830
H_{LsgACC} (DP → II)	0.1036	$H_{L.sgAAC}$ (II → SS)	-0.0840
$H_{L.sgAAC}$ (SS → SS)	0.100	$H_{R.sgAAC}$ (SP → DP)	-0.1032
		$H_{L.DLPFC}$ (SS → II)	-0.0822

767   **Table 3.** PEB of PEB results; *Sustain and Transient changes following cTBS compared to*
768   *Sham.*

cTBS>Sham			
Sustain		*Transient*	
Parameters	*B_1*	*Parameters*	*B_1*
T_{GABA}(R.sgACC)	-0.0558	T_{AMPA}(R.DLPFC)	-0.0913
T_{NMDA}(R.DLPFC)	-0.0723	T_{AMPA}(R.sgACC)	-0.0699
$H_{R.DLPFC}$ (II → DP)	-0.1012	T_{NMDA}(R.DLPFC)	0.0776
$H_{L.sgAAC}$ (SS → SS)	0.1004	T_{NMDA}(R.sgACC)	-0.0730
$H_{R.sgAAC}$ (SS → SP)	-0.0842	CV(L.DLPFC)	-0.1323
$H_{R.sgAAC}$ (SS → II)	-0.1111	CV(R.DLPFC)	0.1151
		$A_{f\,AMPA}$ (R.DLPFC→LsgACC)	-0.1715
		$A_{f\,AMPA}$ (L.sgACC→R.DLPFC)	-0.3436
		$A_{f\,AMPA}$ (RsgACC → L.sgACC)	0.1720
		$A_{f\,NMDA}$ (R.DLPFC→LsgACC)	-0.2200
		$H_{L.DLPFC}$ (II → II)	0.0799
		$H_{R.DLPFC}$ (II → SP)	0.0867
		$H_{R.sgAAC}$ (DP → II)	0.0901

770