Community -Based Intervention to reach the last mile of childhood

vaccination coverage in Rwanda: Case study of Burera district

Name, affiliation and contact of the author(s):

1. Mr. Michee Nshimayesu, Center for Leadership in Global Health Equity, University of

Global Health Equity, Kigali, Rwanda. mnshimayesu@ughe.org ORCID: 0000-0001-

8127-0421

2. Dr. Barnabas Tobi Alayande, Center for Equity in Global Surgery, University of Global

Health Equity, Kigali, Rwanda balayande@ughe.org ORCID: 0000-0002-1326-6452

3. Dr. Sage M. C. Ishimwe, Institute of Global Health Equity Research, University of

Global Health Equity, Kigali, Rwanda. sishimwe@ughe.org ORCID: 0000-0001-9928-

5891

4. Mr. Aimable Uwimana, Department of Community Health and Social Medicine,

University of Global Health Equity, Kigali, Rwanda. auwimana@ughe.org ORCID:0000-

0001-6453-3760

5. Mr. Charif Niyiragira, Department of Clinical Medicine in School of Medicine,

University of Global Health Equity, Rwanda. cniyiragira@ughe.org

6. Dr. Ernest Tambo, Department of Monitoring, Evaluation, Research, and Learning,

University of Global Health Equity, Rwanda. etambo@ughe.org

Name and contact of the corresponding author:

Name: Michee Nshimayesu

Contact: mnshimayesu@ughe.org/Tel. +250788335534

Address: University of Global Health Equity (UGHE), Kigali Rwanda

Abstract

Background:

Vaccination remains a cornerstone of global child health, offering one of the most cost-effective interventions for reducing childhood morbidity and mortality caused by vaccine-preventable diseases (VPDs). Despite this, global vaccination coverage has recently declined, particularly in low- and middle-income countries. In Rwanda, although the national childhood immunization rate has reached 95%, gaps remain in reaching the last mile. Community Health Workers (CHWs) have played an instrumental role in promoting and delivering immunization services across hard-to-reach communities. Yet, little is known about the specific contributions of CHWs to these gains, particularly in rural settings such as Burera District, Northern Rwanda. The study aimed to (1) investigate the factors influencing CHWs' intervention in promoting childhood vaccination, (2) examine the impact of CHWs on vaccination coverage among children under five, and (3) explore the perspectives of caregivers on CHWs' intervention in childhood vaccination coverage.

Methods:

A mixed-methods approach was adopted, combining quantitative surveys with 345 CHWs and qualitative key informant interviews (KIIs) with selected caregivers of children under five benefiting from CHWs vaccination intervention. Quantitative data were collected through structured questionnaires and analyzed using descriptive statistics, and multivariable logistic regression in STATA 18. Thematic analysis was used to analyze qualitative data obtained from five KIIs across Burera's five major health center catchment areas.

Results:

CHWs in Burera were predominantly female (67.5%), married (98.8%), with primary-level education (86.1%), and most engaged in farming (98.6%). A large proportion (94.5%) had received training on vaccination services, and 98.3% regarded under-five vaccination as "very important."

Over 50% reported facilitating community vaccination events twice per month. CHWs were actively involved in community mobilization and advocacy, with 93.9% participating in campaign planning. CHW activities included referring children to health facilities (58.0%) and conducting home-based vaccination outreach (42.0%). From caregivers' perspectives, CHWs played critical roles in health education, timely reminders for appointments, and home-based vaccination services. Thematic analysis of interviews revealed strong trust in CHWs, who were seen as vital liaisons between the community and the formal health system. Barriers noted included logistical challenges and the need for more consistent training and communication support for CHWs. Over 55% of CHWs recommended more structured community awareness campaigns to further enhance immunization uptake.

Conclusion:

CHWs are central to Rwanda's success in achieving high childhood vaccination coverage, especially in rural areas like Burera District. Their interventions in education, mobilization, and service delivery have significantly contributed to bridging gaps in access and awareness. However, sustaining and enhancing these gains requires continued investment in CHW training, supervision, and logistical support. Findings from this study provide critical insights for policymakers and health planners to strengthen community-based vaccination strategies and reach the last mile by 2030 in alignment with Sustainable Development Goal (SDG) 3.b.

Keywords: Community, Intervention, Last Mile, Childhood, Vaccination, Coverage

BACKGROUND

Vaccination is one of the most effective and cost-efficient public health interventions for reducing child morbidity and mortality, particularly from vaccine-preventable diseases (VPDs)[1]. However, vaccination coverage remains a concern globally. In 2023, global immunization coverage showed mixed progress: 14.5 million children remained completely unvaccinated (zero-dose), diphtheria, tetanus, and pertussis (DTP3) and first-dose measles coverage plateaued at 84% and 83% respectively (below pre-pandemic levels), HPV first-dose coverage in girls improved from 20% to 27%, and yellow fever vaccination in at-risk countries lagged at 50%, far short of the recommended 80% threshold[2]. In 2019, an estimated 5.3 million children under the age of five died, nearly half (49.2%) of them from infectious causes, primarily preterm birth complications, lower respiratory infections, intrapartum-related events, and diarrhoea. Notably, 21.7% of these deaths were caused by vaccine-preventable diseases such as measles, meningitis, and respiratory infections, while many more could have been prevented with low-cost interventions. Since 2000, under-5 mortality has declined significantly, owing primarily to reductions in deaths from these preventable and treatable conditions[3].

Community Health Workers (CHWs) play a vital role in improving child health outcomes, especially in underserved communities. Their involvement in delivering curative care for malaria, pneumonia, and diarrhea has been instrumental in reducing childhood illness and mortality. In Sub-Saharan Africa, CHW-led interventions which decrease in child mortality through direct care, health education, and underscoring their critical contribution to closing healthcare gaps. CHWs identified and referred 42 probable measles cases for immunization, while in Kenya, CHW programs increased infant vaccination rates from 88.7% to 98.8%[4, 5, 6].

Rwanda implemented its CHW program in 1995 to enhance healthcare access and address medical professional shortages[7]. Since 2015, the number of CHWs has increased to four per village, with

two-thirds being women, significantly contributing to health interventions, including childhood vaccinations[8]. CHWs have been instrumental in improving vaccination rates, contributing to Rwanda's 95% childhood immunization coverage in alignment with Sustainable Development Goal (SDG) 3.b [9, 10, 11].

Despite Rwanda's success, little is known about the specific contributions of CHWs to national vaccination achievements. We sought 1) to investigate the factors influencing the intervention of CHWs in promoting childhood vaccination, 2) to examine the effects of CHWs on the childhood vaccination coverage, and 3) to explore the perspectives of mothers/caregivers on CHWs' intervention in childhood vaccination coverage in Burera district, Rwanda.

METHODS AND MATERIALS

Study design, setting and study population

This research employed a mixed-methods approach, combining quantitative and qualitative data collection and analysis techniques. This approach enabled a comprehensive understanding of the contribution of CHWs' intervention in the under-5 children's vaccine program in Burera district communities. This study was conducted in Burera District which is divided into 17 sectors with a total of 69 cells and 571 villages distributed in all sectors, in the Northern Province of Rwanda. Burera district occupies 645 km^2 and accounts for 397,754 habitants (199,706 males and 176,187 females), which represents 18% of the total population of the Northern province[12].

This study focused on CHWs and under-5 mothers in the Burera District of Rwanda's Northern Province. Burera district has a total of 2284 CHWs. Each village, which has 100-250 households, has four CHWs, one female "Agent de Sante Maternelle (ASM)" and a male-female pair known as "Binômes." ASMs provide maternal and newborn healthcare through home visits and follow-up, whereas "Binômes" provide a broader range of services, including childhood illness diagnosis

and treatment, antimalaria services for people of all ages, malnutrition screening, and contraceptive provision.[11,13]. Additional to the quantitative feedback from CHWs, the study focused on the mothers or caregivers' perceptions who have children benefiting from both vaccination services and CHWs intervention to collect the qualitative insights of services beneficiaries in Burera community.

Inclusion and Exclusion Criteria

The inclusion criteria for quantitative participants, included being a CHW living and working in Burera district, or having been a CHW in the same region for at least two years, and voluntary consenting to participate in the study. CHWs in charge of health services other than the vaccination excluded in the study, and the CHWs who are not active in the services during the study period were excluded For qualitative participants, we included mothers or primary caregivers of children under-five years who had , lived in the specific village for at least 2 years Mothers who did not use in community based-vaccination services or CHW services, and those unwilling to participate in the study were excluded a.

Sample size calculation and Sampling technique

 To calculate the quantitative participants from 2284 CHWS living and working in Burera, we calculated the sample size using a 95% confidence level with a 5% margin of error. Burera now has 5% of Rwanda's CHWs (2,284/45,516 total). Using Slovin's formula, we arrived at a minimum sample size of 340 CHWs[14]. A sampling frame technique used to select participants out of 2284 CHWs present in Burera district. Lists of CHWs were obtained from health centers, and a simple random sampling technique was applied every fourth name on the list was selected in each health center. Additionally, we targeted the health centers with the highest number of CHWs to ensure broader representation, inviting equal numbers of participants from each selected center. For the

qualitative component, five caregivers were selected one from each of the five health centers in

Burera district. Using a list provided by each health center, we randomly selected every fourth

individual who met the eligibility criteria within the respective catchment area. To ensure

participants met the study selection criteria, a purposive sampling approach was also applied to

identify mothers or caregivers of under-five children who were actively participating in

community-based vaccination services. The lists obtained from health centers served as the basis

for randomly selecting eligible participants.

Data collection

A data collection process took place between May 06th 2024 to June 16th, 2024. The trained data

collector was hired to facilitate the data collection process. Initially, participants' information,

including contact details, was extracted from health center records. The data collector then

contacted the participants to obtain a written consent before they decided whether or not to

participate. For quantitative data collection, a structured questionnaire was developed and

administered to CHWs. This questionnaire aimed to gather information on the factors influencing

their interventions, as well as their roles and impact on the vaccination program. It also collected

demographic data about the CHWs. The quantitative data collection involved face-to-face

administration of close-ended questionnaires, which consist of a series of questions designed to

elicit specific responses from participants[15]. In addition, KIIs were conducted with caregivers to

explore their experiences and perceptions related to the childhood vaccination program and the

interventions of CHWs. The KIIs were audio-recorded and subsequently transcribed for thematic

analysis.

Study Variables and measures

The study examined the perceived effectiveness of CHWs' interventions on childhood vaccination

coverage, considering various independent variables. These included socio-demographic factors,

infrastructure and logistics, training, knowledge, community awareness and perception,

community mobilization, services available at the CHW level, availability of health information

and communication for under-5 mothers, CHWs' availability for home visits, and equitable service

delivery for under-5 children and their mothers. The study measures focused on three key aspects

of CHW's interventions in childhood vaccination services: availability, accessibility, and quality.

These aspects were examined through two main approaches. The first approach assessed

community awareness and perceptions of CHWs' interventions. The second approach evaluated

changes in the likelihood of under-5 children being fully vaccinated before and after CHWs'

interventions. Accessibility was analyzed based on location, communication, appointment

availability, service costs, frequency of access, access to medical records, and age-related access.

Quality was assessed through confidentiality and privacy during visits, service provision,

accessibility, environmental factors, adherence to written guidelines, CHW characteristics and

competencies, and the involvement of under-5 mothers and caregivers.

Data management

All participants were required to sign a consent form approved by the UGHE IRB Committee

before the start and those signed the consent form assigned a unique study ID code to be used

throughout the study period. During data collection, the information was cross-checked and further

verified by the principal investigator to ensure data quality and integrity. Quantitative data was

collected using REDCap (Research Electronic Data Capture) and subsequently transferred to

STATA (Stata Corp Inc.) version 18 for analysis (Ref). Key Informant Interviews (KIIs) were

audio-recorded and subjected to thematic analysis to identify common themes and patterns related

to the interventions of community health workers (CHWs) on vaccination coverage. All data files

associated with the study were securely stored on the researchers' computers, protected by strong

passwords.

Data analysis

Quantitative data were checked for completeness, edited, coded, and exported for analysis into

STATA (Stata corp inc.) version 18. Descriptive statistics were performed for prevalence

estimation and participant demographic characteristics after cleaning the data for inconsistencies.

To determine the relationship between different variables, the Chi-square test, and the t-test were

used. Also, multivariable analyses used to investigate the relationship between under-5 vaccination

coverage and CHW intervention. A statistically significant p-value of 0.05 was considered during

the analysis. For qualitative data (KIIs), thematic analysis was employed to identify common

themes and patterns in the data. Transcripts were coded accordingly, and codes were then grouped

into themes. The findings are presented using narrative descriptions, illustrative quotes, and

thematic matrices.

RESULTS

Socio-demographic characteristics of the community health workers

202 The highest percentage (38.8%) of the community health workers (CHWs) were aged between 40

203 and 49 years whereas only a small proportion (7.5%) were aged less than 30 years. Females

204 constituted the majority (67.5%) and about half were identified as Catholic (48.4%). Al most all

205 (98.8%) were married and 86.1% of them attained primary level of education. A large percentage

206 (98.6%) were engaged in farming (Table 1).

Table 1. Socio-demographic characteristics of community health workers

Variables	N = 345	%

Age

<30 years	26	7.5
30 to 39 years	81	23.5
40 to 49 years	134	38.8
50 years and above	104	30.1

Sex

Female	233	67.5
Male	112	32.5

Religion

Protestant	81	23.5
Catholic	167	48.4
Adventist	47	13.6
Others	50	14.5

Marital status

Married	341	98.8
Widowed	4	1.2

Education

Primary	297	86.1
Secondary	46	13.3
None	2	0.6

Occupation

No occupation	2	0.6
Farmer	340	98.6
Others	3	0.9

Knowledge and training on childhood vaccination program

Most of the CHWs (50.4%) reported that vaccination services are offered twice a month. The

majority (94.5%) of CHWs had received education or training on vaccination services with 64.1%

having participated in training not more than twice. Significant percentage (98.3%) indicated that

vaccination under 5 years is very important and 94.2% reported that missing vaccination could

cause diseases (Table 2).

This was also supported in the qualitative data as one pregnant woman expressed "*Community*

health workers explain it well each time we come for vaccination, starting from when you get

tested for pregnancy, they explain how you have to get your child vaccinated on time" (KII-3).

Another woman also endorsed this as "*They inform and educate us on how to vaccinate our*

children, starting from the beginning of pregnancy. As soon as you find out you are pregnant, they

tell you that you will give birth at a health facility and follow up with the first vaccination. They

ensure vaccinations at one and a half months, two and a half months, and continue at nine and

fifteen months" (KII-4).

Table 2. Knowledge and training on childhood vaccination program

Variables	N = 345	%
Frequency of offering vaccination services in one month		
0 time	15	4.3
1 time	69	20.0
2 times	174	50.4
3 times	44	12.8

4 times	27	7.8
5+ times	16	4.6
Ever received any education or training about vaccination services		
Yes	326	94.5
No	19	5.5
Frequency of training received about vaccination services in year (n = 326)		
1-2 times	209	64.1
3 - 4 times	92	28.2
5 times and above	25	7.7
How important is the under-5 vaccination program in the community		
Not important	1	0.3
Important	5	1.4
Very important	339	98.3
Main effects of missing vaccine for under-5		
Cause disability	23	6.7
Cause death	1	0.3
Cause of disease	325	94.2
Don't know	1	0.3

225 **Community mobilization and participation towards childhood vaccination program**

226 As indicated in Table 3, the majority of CHWs (93.9%) were fully engaged in conducting

227 vaccination program campaign organized by community leaders during planning phase (Table 3).

228 This is reinforced by the qualitative data where one pregnant woman indicated:

229   *"They encourage us to be there, they pay us a visit to let us know that we will be meeting*

230   *on the vaccination site and do what is required from us* (KII-3).

231   Another woman also stated, *"Community health workers manage it efficiently. When it*

232   *comes to pills or drops, they are available at the local service centers. They maintain a list*

233   *of children and ensure each child receives the appropriate vaccines according to their area*

234   *(KII-2).*

235 Majority (61.4%) reported that community members were fully involved when vaccination event

236 was planned. The main responsibilities of CHWs included referring children to healthcare facilities

237 (58.0%) and administering vaccines during home visits (42.0%). Similarly, these findings are

238 reflected in the qualitative data where the pregnant women interviewed indicated that they get

239 advocacy and support through providing vaccinations and follow-ups as well as reminding them

240 for vaccination appointment. Some of the pregnant women's quotations were as follows:

241   *"What is great is that they advocate in taking care of my child and reminding me to have*

242   *my child vaccinated according to the given appointment* (KII 1)".

243   *"A community health worker visits our home, informs us about new vaccines, and advises*

244   *us to take our child for vaccination* (KII-4)"; and another woman indicated "*It makes me*

245   *happy when they bring vaccinations promptly or provide necessary information directly to*

246   *my home* (KII-5)".

247 Over half (55.4%) of the CHWs believed that mobilization or awareness to be conducted in order

248 to enhance vaccination coverage. About 7.0% of the CHWs indicated that community members

249 complained about vaccination. It was also reported that there are still ignorance and poor

250 understanding according to the pregnant women participated in the qualitative study. These

251 expressions include:

"It is due to poor understanding and lack of education. Families who are better informed

understand the importance of vaccinations and ensure their children receive them". KII-4

"Some parents do not understand how vital vaccinations are because they lack proper

education on the topic. Those who are educated take it seriously and follow through with

the schedule." KII-1

"I cannot understand why a child wouldn't be vaccinated if they are present. Perhaps some

parents refuse vaccinations, but I cannot comprehend their reasoning". KII-5

Table 3. Community mobilization and participation towards childhood vaccination

program

Variables	N = 345	%
Involved to conduct a vaccination program campaign when there is planning by community leaders		
Somehow involved	21	6.1
Fully involved	324	93.9
Village members involvement during on your plan an event or campaign on vaccination program		
Not involved	1	0.3
Somehow involved	6	1.7
Involved	125	36.2
Full involved	212	61.4
At which level as CHW you intervene on vaccination program.		
Referring a child to healthcare facility	200	58.0
Administering vaccines by home visit	145	42.0

On your opinion what do you think, and community needs to enhance vaccination coverage

Mobilization or awareness	191	55.4
Increase home visit	44	12.8
Increase vaccination services at CHW level	58	16.8
Improved equitable health services	1	0.3
Support from government	31	9.0
Information and communication	67	19.4
Don't know	8	2.3
Complaining of community members on your intervention on vaccination program		
Yes	24	7.0
No	297	86.1
Don't know	24	7.0

Infrastructure and logistics on childhood vaccination program

The primary methods of communication and transportation for follow-ups on child vaccination

appointments were physical visits (89.6%), predominantly conducted on foot (99.7%). All CHWs

reported covering their own communication expenses. All vaccines were stored and distributed

exclusively from health centers (Table 4).

However, limited vaccination sites and long distance emerged as the key infrastructure and

logistics barriers on implementing of childhood vaccination program. They indicated that in a cell

there is only one vaccination site, which is limited vaccination site. The distance is identified as a

burden for those coming from afar and is not easy for them to travel long distances just for a

vaccination. This is how the key informants expressed their views:

"The problem is that there is just one vaccination site per cell. It makes it hard for everyone

to reach it, especially those who live far away (KII-3)."

"*For families living far away, getting to the vaccination site is a real struggle. The distance*

makes it difficult for them to attend appointments on time (KII-1).

" To overcome these challenges the pregnant women suggested that all the vaccines should

be administered by CHWs "*Maybe some CHWs should be in charge and administrate the*

vaccines so that the issue get resolved and people with those issues be included." KII-3

Table 4. Infrastructure and logistics on childhood vaccination program

Variables	N = 345	%
Communication with caregivers during follow up on child vaccination appointment		
Call on phone	5	1.4
Physical visit	309	89.6
Others*	31	9.0
Means of transport during home visit		
Bicycle	1	0.3
Foot	344	99.7
Cover communication expenses/money		
Own	345	100.0
Vaccines storage before distributed		
Health center	345	100.0

*Others: General village meeting, community services (umuganda), and during saving group

meeting

Suggestions achieving remaining 5% vaccination rate

When asked about recommendations for achieving full vaccination rate, CHWs prioritized monetary incentives or monthly salary along with additional trainings (51.9%) as the top suggestions. This was followed by increased mobilization about the importance of vaccination programs (13.3%) and provision of some materials such as digital tools (7.2%) as shown in Figure 1. These findings are further reinforced by the qualitative data from pregnant women, as demonstrated by the following quotes:

> *"I think CHWs are always complaining that they do not get any form of incentives or support when they visit pregnant women. It discourages them from doing their work effectively." KII-5*

> *"A message should be sent as a reminder, not like in the past when it was just papers with dates. With technology, a mobile text message would be much better. It would help a mother to remember, instead of relying on papers." KII-4*

> *We need more mobilization to help people understand how important vaccinations are for our children. KII-3*

Figure 1. Suggests achieving remaining 5% vaccination rate

DISCUSSION

Given the persistent challenges of equitable access, vaccine hesitancy and health worker shortages,

leveraging CHWs as vaccinators is an important opportunity [16]. The deployment of Community

Health Workers (CHWs) is widely promoted as a strategy for reducing health inequities in low-

and middle-income countries (LMIC) [17].

The study found that the majority of CHWs were female (67.5%) and aged between 40 and 49

years (38.8%), with primary education being the highest level attained by most (86.1%). These

demographic characteristics align with previous studies in sub-Saharan Africa, where CHWs are

predominantly middle-aged women with limited formal education but extensive community

engagement [18]. The dominance of women among CHWs is critical, as they are more likely to

communicate effectively with mothers and caregivers about childhood vaccination. However, the

limited educational background of CHWs may affect their ability to effectively deliver health

messages, a challenge that has been noted in other studies [19]. Our study highlights that 94.5% of

CHWs received training on vaccination services, with most (64.1%) attending training at least

once or twice a year, and this supporting the argument of WHO that adequate training is essential

for effective immunization programs. These findings are consistent with studies in in Uganda and

Kenya, which emphasize the importance of continuous CHW training to improve knowledge and

vaccination uptake [20]. Qualitative responses from caregivers also emphasized the effectiveness of

CHWs in disseminating vaccination information, which echoes findings from a study that

highlighted CHWs' role in bridging knowledge gaps among mothers [21].

The findings indicated that CHWs significantly contribute to increasing immunization rates

through education, household visits, and follow-up which align with previous research that

highlights the crucial role of CHWs in increasing childhood vaccination coverage. Similar to a

study conducted in Ethiopia, which found that CHWs played a pivotal role in immunization

services, our results confirm that CHWs actively engage in community mobilization and follow-

up visits [22]. The predominance of female CHWs in our study (67.5%) also mirrors global trends,

where women often take the lead in community health interventions due to their trusted roles

within communities [17]. The qualitative findings further reinforce the effectiveness of CHWs in

ensuring timely vaccination, as echoed by caregivers who acknowledged the proactive role of

CHWs in reminding them about vaccination schedules. This aligns with research from Nigeria,

where CHWs' home visits significantly increased vaccine uptake [20].

The study found that 93.9% of CHWs were fully involved in vaccination campaigns, a crucial

factor in vaccine acceptance and coverage. This is in line with findings from Ghana, where strong

community engagement through CHWs improved childhood immunization rates. Additionally,

over half (55.4%) of CHWs emphasized the need for continued mobilization efforts to enhance

vaccination coverage. Similar findings were reported in Kenya, where CHWs' proactive

community engagement led to increased routine immunization uptake [22]. However, 7.0% of CHWs

reported resistance from community members, often due to misinformation and lack of awareness,

which is a common barrier in many low-resource settings [19].

Despite their critical role, CHWs faced logistical challenges, with 99.7% relying on foot transport

and covering communication expenses themselves. Vaccine storage was centralized at health

centers, and only one vaccination site existed per cell, making access difficult for remote

populations. Studies in Uganda and Nigeria have similarly highlighted transportation and cold

chain constraints as key barriers to effective vaccine delivery [23]. The limited vaccination sites, as

indicated by the qualitative data, reinforce the need for decentralizing vaccination services to

improve accessibility [24]. The perspectives of mothers and caregivers in this study further reinforce

the importance of CHWs in vaccination programs. The qualitative findings revealed that CHWs

are trusted sources of information and play a key role in appointment reminders, follow-ups, and

vaccine advocacy. These findings are consistent with previous studies highlighting the trust placed

in CHWs as frontline healthcare providers in rural settings [25]. However, caregivers also identified

gaps in community understanding, with some parents still showing reluctance due to

misinformation or lack of education. Similar hesitancy has been reported in other studies,

emphasizing the need for intensified awareness campaigns [26].

Challenges and recommendations for improving vaccination coverage

Infrastructure and logistical barriers emerged as major constraints, particularly the limited number

of vaccination sites and the long distances some families must travel. This is consistent with

findings from other studies where inadequate access to vaccination services, community awareness

and misinformation was identified as a leading factor in incomplete immunization [26]. Additionally,

CHWs reported a lack of incentives, which negatively affects their motivation. Studies have shown

that providing CHWs with financial incentives or non-monetary support, such as mobile

technology for reminders, significantly enhances their effectiveness [27]. Further, the frequency of training remains relatively low, with 64.1% of CHWs receiving training not more than twice in year, suggesting the need for enhanced refresher courses to maintain competency. To address these challenges, CHWs recommended increased awareness campaigns, government support, and financial incentives. These recommendations align with global best practices, where strengthening CHW training, providing stipends, and integrating digital health tools have proven to enhance vaccination outcomes [28]. Mobile technology, such as SMS reminders for vaccination appointments, was also suggested and has been successfully implemented in other low-resource settings to improve compliance rates [29].

CONCLUSION

The findings of this study highlight the significant role of CHWs in promoting childhood vaccination and the barriers they face. While CHWs have positively influenced vaccination coverage through mobilization and home visits, infrastructural and logistical challenges remain. Strengthening CHW training, increasing incentives, and improving access to vaccination services could enhance childhood immunization rates. Future research should explore the long-term impact of CHW-led interventions and innovative solutions such as digital health tools to optimize vaccination efforts in rural settings. Findings provide insights for policymakers and communities, helping to bridge the last mile of remaining 5% vaccination coverage by 2030.

Ethical approval

This study was approved by the University of Global Health Equity (UGHE) Institutional Review Board on the 9th of April 2024 (reference number UGHE-IRB/2024/240). Prior to participation, all participants provided written informed consent.

380 **Funding:** The study funded by the Dean's Office, School of Medicine of the University of

381 Global Health Equity Rwanda

REFERENCES

1. Rodrigues CMC, Plotkin SA. Impact of vaccines: Health, economic and social perspectives. *Front Microbiol.* 2020;11:1526. doi:10.3389/fmicb.2020.01526

2. World Health Organization (WHO). Immunization coverage. 2024 Jul 15 [cited 2025 May 18]. Available from: https://www.who.int/news-room/fact-sheets/detail/immunization-coverage

3. Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong K, et al. Global, regional, and national causes of under-5 mortality in 2000–19: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet Child &Adolesc Health.* 2021;6:106–15. doi:10.1016/S2352-4642(21)00311-4

4. Bagonza J, Kibira S, Rutebemberwa E. Performance of community health workers managing malaria, pneumonia and diarrhea under the community case management programme in central Uganda: A cross-sectional study. *Malar J.* 2014;13:367. doi:10.1186/1475-2875-13-367

5. Vouking MZ, Binde T, Tadenfok CN, Ekani JME, Ekra D. Contribution of community health workers to surveillance of vaccine-preventable diseases in the Obala health district. *Pan Afr Med J.* 2017;28:207. doi:10.11604/pamj.2017.28.207.11537

6. Nzioki JM, Onyango RO, Ombaka JH. Efficiency and factors influencing efficiency of Community Health Strategy in providing Maternal and Child Health services in Mwingi District, Kenya: An expert opinion perspective. *Pan Afr Med J.* 2015;20:88. doi:10.11604/pamj.2015.20.88.4711

7. Rwanda Governance Board (RGB). Rwanda Community Health Workers Programme: 1995–2015. Kigali: RGB; 2017 [cited 2025 May 18]. Available from: https://www.rgb.rw/fileadmin/user_upload/RGB/Publications/HOME_GROWN_SOLUTIONS/Rwanda_Community_Healt_Workers_Programme_2017.pdf

8. Ministry of Health (Rwanda). Rwanda elects fourth community health worker. 2018 [cited 2025 May 18]. Available from: https://www.moh.gov.rw/news-detail/rwanda-elects-fourth-community-health-worker

9. Schurer JM, Fowler K, Rafferty E, Masimbi O, Muhire J, Rozanski O, et al. Equity for health delivery: Opportunity costs and benefits among Community Health Workers in Rwanda. *PLoS One.* 2020;15(9):e0236255. doi:10.1371/journal.pone.0236255

10. Bao J, McAlister H, Robson J, Wang A, Koswin K, Sayinzoga F, et al. Near universal childhood vaccination rates in Rwanda: how was this achieved, and can it be duplicated? *Lancet Glob Health.* 2018;6:S47. doi:10.1016/S2214-109X(18)30176-1

11. Wightman P, McCue K, Sabo S, et al. Community health worker intervention improves early childhood vaccination rates: results from a propensity-score matching evaluation. *BMC Public Health.* 2022;22:1854. doi:10.1186/s12889-022-14239-w

12. Burera District. Menya Akarere. 2022 [cited 2025 May 18]. Available from: https://www.burera.gov.rw/about/menya-akarere

13. Niyigena A, Girukubonye I, Barnhart DA, Cubaka VK, Niyigena PC, Nshunguyabahizi M, et al. Rwanda's community health workers at the front line: A mixed-method study on perceived needs and challenges for community-based healthcare delivery during COVID-19 pandemic. *BMJ Open.* 2022;12(4):e055119. doi:10.1136/bmjopen-2021-055119

14. Ellen S. Slovin's formula sampling techniques. *Sciencing.* 2022 Mar 24 [cited 2025 May 18]. Available from: https://www.sciencing.com/slovins-formula-sampling-techniques-5475547/

15. Kabir S. Methods of data collection. 2016 [cited 2025 May 18]. Available from: https://www.researchgate.net/publication/325528948_Methods_of_Data_Collection

16. Gavi. Community health workers as vaccinators: A pathway to achieving global immunization goals. 2022 [cited 2025 May 18]. Available from: https://www.gavi.org/vaccineswork/community-health-workers-vaccinators-pathway-achieving-global-immunisation-goals

17. Ahmed S, Chase LE, Wagnild J, Akhter N, Sturridge S, Clarke A, et al. Community health workers and health equity in low- and middle-income countries: systematic review and recommendations for policy and practice. *Int J Equity Health.* 2022;21(1):49. doi:10.1186/s12939-021-01615-y

18. Mbugua R. Effect of socio-demographic characteristics on performance of Community Health Workers: A cross sectional study in Makueni County, Kenya. *Nurs Health Sci.* 2022;6:14–18. doi:10.9790/1959-0606061418

19. Ogutu EA, Ellis AS, Hester KA, Rodriguez K, Sakas Z, Jaishwal C, et al. Success in vaccination programming through community health workers: a qualitative analysis of interviews and focus group discussions from Nepal, Senegal and Zambia. *BMJ Open.* 2024;14(4):e079358. doi:10.1136/bmjopen-2023-079358

20. Alban R, Gibson E, Payne J, et al. Leveraging community health workers as vaccinators: a case study exploring the role of Malawi's Health Surveillance Assistants in delivering routine immunization services. *Hum Resour Health.* 2023;21:42. doi:10.1186/s12960-023-00827-3

21. Gibson E, Zameer M, Alban R, Kouwanou LM. Community Health Workers as Vaccinators: A Rapid Review of the Global Landscape, 2000–2021. *Glob Health Sci Pract.* 2023;11(1):e2200307. doi:10.9745/GHSP-D-22-00307

22. Bakkabulindi P, Ampeire I, Ayebale L, Mubiri P, Feletto M, Muhumuza S. Engagement of community health workers to improve immunization coverage through addressing inequities and enhancing data quality and use is a feasible and effective approach: An implementation study in Uganda. *PLoS One.* 2023;18(10):e0292053. doi:10.1371/journal.pone.0292053

23. Malande OO, Munube D, Afaayo RN, Annet K, Bodo B, Bakainaga A, et al. Barriers to effective uptake and provision of immunization in a rural district in Uganda. *PLoS One.* 2019;14(2):e0212270. doi:10.1371/journal.pone.0212270

24. Gavi. Effective ways to increase vaccination rates. 2018 [cited 2025 May 18]. Available from: https://cms.wellcome.org/sites/default/files/2020-12/effective-ways-to-increase-vaccination-rates.pdf

25. Kowalczyk M, Yao N, Gregory L, Cheatham J, DeClemente T, Fox K, et al. Community health worker perspectives: examining current responsibilities and strategies for success. *Arch Public Health.* 2024;82(1):26. doi:10.1186/s13690-024-01313-5

26. Sinuraya RK, Nuwarda RF, Postma MJ, Suwantika AA. Vaccine hesitancy and equity: Lessons learned from the past and how they affect the COVID-19 countermeasure in Indonesia. *Glob Health.* 2024;20(1):11. doi:10.1186/s12992-023-00987-w

27. Bangura JB, Xiao S, Qiu D, Ouyang F, Chen L. Barriers to childhood immunization in sub-Saharan Africa: A systematic review. *BMC Public Health.* 2020;20:1108. doi:10.1186/s12889-020-09169-4

28. World Health Organization (WHO). Guideline on health policy and system support to optimize community health worker programmes. Geneva: WHO; 2018. Licence: CC BY-NC-SA 3.0 IGO. Available from: https://iris.who.int/bitstream/handle/10665/275474/9789241550369-eng.pdf

29. Eze P, Lawani LO, Acharya Y. Short message service (SMS) reminders for childhood immunisation in low-income and middle-income countries: a systematic review and meta-analysis. *BMJ Glob Health.* 2021;6(7):e005035. doi:10.1136/bmjgh-2021-005035