Comparison of low eGFR prevalence and prediction for mortality using 2009 and

2021 CKD-EPI equations in Mexican adults

Daniel Ramírez-García[1,2*], Carlos A. Fermín-Martínez[1,3*], Paulina Sánchez Castro[1,2*],

Brenda Guadalupe Cortez-Flores[4], Jerónimo Perezalonso Espinosa[1,3], Juan Pablo Díaz-

Sánchez[1,3], Karime Berenice Carrillo-Herrera[1,3], Leslie Alitzel Cabrera-Quintana[1,3], Jaime

Berumen-Campos[5], Pablo Kuri-Morales[6], Roberto Tapia-Conyer[5], Jacqueline A. Seiglie[7,8],

Jesus Alegre-Díaz[5], Neftali Eduardo Antonio-Villa[9], Omar Yaxmehen Bello-Chavolla[1]

These authors contributed equally to the drafting of this work

[1]Research Division, Instituto Nacional de Geriatría, Mexico City, Mexico. [2]Facultad de

Medicina, Universidad Nacional Autónoma de México, Mexico City, Mexico. [3]MD/PhD

(PECEM) Program, Facultad de Medicina, Universidad Nacional Autónoma de México,

Mexico City, Mexico. [4]Department of Nephrology, National Institute of Cardiology Ignacio

Chavez, Mexico City, Mexico. [5]Experimental Research Unit, Faculty of Medicine, National

Autonomous University of Mexico, Mexico City, Mexico. [6]Instituto Tecnológico y de

Estudios Superiores de Monterrey, Monterrey, Mexico. [7]Diabetes Unit, Massachusetts

General Hospital, Harvard Medical School, Boston, Massachusetts, USA. [8]Department of

Medicine, Harvard Medical School, Boston, Massachusetts, USA. [9]Department of

Endocrinology, Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico.

Correspondence: Omar Yaxmehen Bello-Chavolla. Division of Research. Instituto

Nacional de Geriatría. Anillo Perif. 2767, San Jerónimo Lídice, La Magdalena Contreras,

10200, Mexico City, Mexico. Phone: +52 (55) 5548486885. E-mail: oybello@inger.gob.mx

Disclosures: The authors declare that they have no conflict of interests.

Funding: This research was supported by Instituto Nacional de Geriatría in Mexico.

Word Count: 2,977**. References:** 33 **references. Figures:** 4**. Tables:** 3.

Short Title: Comparison of CKD equations in Mexican adults

ABSTRACT (300 WORDS)

Accurate estimation of glomerular filtration rate (eGFR) is essential for identifying and managing chronic kidney disease (CKD). The CKD-EPI 2021 equation removed the race coefficient from the 2009 version, but its impact in Mexican populations remains unclear. Here, we compared eGFR category prevalence, and predictive performance between the CKD-EPI 2009 and 2021 creatinine-based eGFR equations, as well as the prognostic relevance of reclassification in eGFR categories using the 2021 equation in Mexicans. We evaluated 25,110 adults ≥20 years from the 2016-2023 cycles of the Mexican National Health and Nutrition Survey (ENSANUT) to estimate national low eGFR and eGFR category prevalence using both equations. We also assessed 5-year and 10-year risk of all-cause, cardiovascular, and kidney-related mortality in 142,884 adults from the Mexico City Prospective Study (MCPS) using Cox proportional hazards and Fine & Gray regression models. In ENSANUT 2023, prevalence of eGFR <60mL/min/1.73m2 was lower with CKD-EPI 2021 (2.9%, 95%CI 1.56–4.24%) compared to the 2009 equation (3.6%, 95%CI 1.99–5.21). Use of the 2021 equation resulted in upward eGFR reclassification in 6.52% (95%CI 4.07-8.97) of adults ≥20 years, particularly among older adults and those with hypertension or diabetes, yielding a reduction in 486,532 adults identified with eGFR <60mL/min/1.73m^2 compared to the 2009 equation. In MCPS, despite both equations showing similar C-statistics, the 2021 equation showed slightly improved predictive performance for 5-year and 10-year mortality outcomes. The 2021 equation reclassified 8.3% of participants to higher eGFR categories, and reclassification was associated with decreased risk of all-cause, cardiovascular, and kidney-related mortality, particularly for participants reclassified upward from G3a-G5 categories. The CKD-EPI 2021 equation yields lower prevalence of low eGFR but leads to prognostically relevant eGFR category reclassification compared to the 2009 equation. Our findings support the implementation of the 2021 equation for population health monitoring in Mexico without compromising prognostic utility.

Keywords: *CKD-EPI, Chronic kidney disease, eGFR, Mexico, mortality.*

INTRODUCTION

Chronic kidney disease (CKD) has become a significant health problem worldwide, especially in low- and middle-income countries, where inequities in access to preventive services, renal replacement therapy (RRT), and the increasing prevalence of cardio-metabolic diseases have contributed to growing prevalence and mortality rates[1]. CKD diagnosis and stratification typically involves estimation of the glomerular filtration rate (eGFR) and measuring albuminuria[2]. Previous clinical guidelines recommended using the 2009 CKD Epidemiology Collaboration (CKD-EPI) equation which accounts for age, sex, and race-related variability[3,4]. However, using race in clinical algorithms has been recently questioned, considering it as a social and not a biological construct[5]. Consequently, a new CKD-EPI equation without a race coefficient was developed in 2021[6]; which, compared to directly measured GFR, leads to higher CKD prevalence estimates amongst black individuals, and lower amongst non-black individuals[6].

Mexicans have an admixed ancestry, comprising of ~50% of Amerindian native-American, 45% European and <5% African and East Asian contributions[7]. This admixture profile contrasts with the predominantly European ancestry of non-Hispanic white individuals, and African ancestry of non-Hispanic Black individuals in the US[8]. Despite representing a large share of the US population, Mexicans and broader Latino populations remain underrepresented in kidney disease studies, despite showing significant inter-population variability in CKD markers[9,10]. Furthermore, eGFR equations commonly used in clinical settings frequently classify Mexican individuals as white, disregarding emerging evidence of population-specific differences in serum creatinine levels[10], and the influence of genetic ancestry on the accuracy of creatinine-based eGFR estimations[11]. To date, no direct estimations of CKD prevalence have been conducted in Mexico, with prior studies focusing on forecasting estimates based on statistical models[1]. Moreover, the impact of replacing the 2009 with the 2021 equation on CKD prevalence, and the utility of these equations for

prediction of relevant outcomes, including all-cause and cause-specific mortality in

Mexican populations, has not been previously validated. In this study, we aimed to

estimate the prevalence of eGFR categories and low eGFR using the 2009 and 2021

CKD-EPI equations in adults using nationally representative surveys from 2016 to 2023 in

Mexico, and to assess the performance of both equations for prediction of all-cause,

cardiovascular, and kidney-related mortality in prospective cohort of adults living in Mexico

City.

METHODS

Study populations

Mexican National Health and Nutrition Survey

We used data from the cross-sectional Mexican National Health and Nutrition Survey

(ENSANUT) cycles 2016, 2018, 2020, 2021, 2022, and 2023. ENSANUT is a probabilistic

survey representative at a regional and national level which uses a multi-stage

probabilistic cluster stratified sampling design[12]. Sample weights are corrected using the

response rate observed from the immediate previous cycle and calibrated to account for

representation of specific subgroups. Participants completed a questionnaire including

demographic, socioeconomic, and health-related information, along with a physical

examination. Additionally, serum samples were collected from a random subsample per

cycle to measure fasting glucose, serum creatinine, lipid profile, and glycated

hemoglobin[12]. Further details on ENSANUT's methodology are reported elsewhere[12–15],

and in **Supplementary Methods**. For this study, we included individuals ≥20 years with

measured serum creatinine to calculate eGFR.

Mexico City Prospective Study

The Mexico City Prospective Study (MCPS) is a population-based, prospective cohort

study with a baseline survey conducted from 1998 to 2004 on two urban districts of Mexico

City[16]. In total, 159,755 individuals aged ≥35 years were recruited and underwent a

comprehensive questionnaire collecting sociodemographic and health-related information,

clinical measurements (anthropometry, and blood pressure), and a non-fasting 10-mL

blood sample. We included all participants with complete mortality, covariate, and serum

creatinine data. Mortality follow-up of participants was done through electronic probabilistic

linkage to death registries, with deaths tracked up to September 30th, 2022, and registered

according to the International Classification of Diseases 10th Revision (ICD-10)[17]. Mortality

was codified in accordance with the underlying causes of death determined by study

clinicians as all-cause, cardiovascular disease (CVD) (myocardial infarction for codes I20-

I25, stroke for codes I60-I69), and kidney-related mortality (codes N17-N19, E10-E14, I12-

I13, N28, Q61, N00-N16, Y84). Details on approval, mortality follow-up, comorbidities, and

creatinine measurements are reported in **Supplementary Materials**.

Estimated glomerular filtration rate and chronic kidney disease

eGFR was calculated using the CKD-EPI 2009 and 2021 creatinine-based equations[6], and

classified into six categories as recommended by KDIGO guidelines, specifically: G1

normal or high (≥90 mL/min/1.73m^2), G2 mildly decreased (60–89 mL/min/1.73m^2), G3a

mildly to moderately decreased (45–59 mL/min/1.73m^2), G3b moderately-to-severely

decreased (30–44 mL/min/1.73 m^2), G4 severely decreased (15-29 mL/min/1.73m^2), and

G5 kidney failure (<15 mL/min/1.73m^2)[3]. Given that albuminuria was not measured in

either study, low eGFR was used as a proxy for CKD, defined as eGFR <60

mL/min/1.73m^2. We also explored the prevalence of CKD defined as the combination of

previous self-reported medical diagnosis of CKD or low eGFR.

Statistical analyses

We obtained weighted prevalence estimates of eGFR categories and low eGFR using the

2021 CKD-EPI creatinine-based equation, considering complex survey design and sample

weights from the ENSANUT venous blood subsample using the *survey* R package.

Additionally, we estimated low eGFR prevalence stratified by age categories (<60 or ≥60

years old), sex, diabetes, hypertension, and obesity status. Next, we compared these

estimations against the 2009 CKD-EPI creatinine-based equation, exploring the

proportion, and total number of individuals that would be reclassified to different eGFR

categories. Additional details on ENSANUT survey design are available in **Supplementary**

Methods.

To evaluate the predictive performance of both CKD-EPI equations for all-cause, CVD, and

kidney-related mortality, we fitted a baseline Cox proportional hazards model stratified by

baseline age category and adjusted for sex, place of residence, education, smoking,

diabetes, history of CVD, systolic blood pressure, and total cholesterol, which is similar to

the CKD Prognosis Consortium model for all-cause and CVD mortality risk[18]. eGFR

estimated with either the 2009 or 2021 equations, was added separately to this baseline

model to assess discrimination and performance metrics[19,20]. For CVD and kidney-related

mortality, we used Fine & Gray competing risk models, adjusted for the same covariates

using the *survival* R package. Administrative censoring was applied at both 5 and 10 years

to assess all outcomes. The non-linear relationship between eGFR and mortality risk was

investigated with penalized smoothing splines[21], and eGFR was modeled using restricted

cubic splines with 5 knots[22]. Fixed-time point, and time-range discrimination was assessed

using Harrell's c-statistic, and overall performance was determined evaluating changes in

the model's chi-square statistic, the Brier score, and the scaled Brier score. While relative

hazards might not capture a marker's added predictive value[19], we report the hazard ratios

(HR) and sub-distribution HRs (sHR)s for eGFR 2009 or 2021 from extended models. To

assess the prognostic significance of reclassification from eGFR categories based on the

2021 equation to those estimated with the 2009 equation, we generated an indicator

variable identifying participants who were 1) reclassified upward from G2 to G1 or 2)

reclassified upward from G3a-G5. We then estimated the relative risks for all-cause,

cardiovascular, and kidney-related mortality by comparing these participants with those

who remained in the same categories (i.e., not reclassified). Outcomes were visualized

using Kaplan-Meier curves adjusted using inverse probability of treatment weighting

(IPTW) groups with similar covariates as previous models using the *adjustedCurves* R

package. Additional details on modeling, missing data, and underlying assumptions are

reported in **Supplementary Methods**. All statistical analyses were conducted using R

software version 4.4.2.

RESULTS

eGFR categories and CKD prevalence in Mexican population

We analyzed data from 25,110 adults ≥20 years from six ENSANUT cycles during the

2016-2023 period (**Supplementary Table 1**). Using the 2021 CKD-EPI equation, the

prevalence of eGFR categories remained stable over time. The proportion of individuals on

the G1 category reached 87.08% (95%CI 84.31–89.84%) in 2023, with a slight decrease

over time (**Figure 1A**). G2 prevalence was much lower, reaching 9.98% (95%CI 7.76–

12.20%) by 2023 (**Figure 1B**). G3a and G3b categories had a prevalence of 1.76%

(95%CI 0.77–2.76%) and 0.90% (95%CI 0.22–1.57%) in 2023, respectively (**Figure 1C**),

and G4 and G5 prevalence estimations from 2016 to 2023 were below 0.6% and 0.35%,

respectively. (**Table 1**). Categories G3a-G5 were stable over time but presented an

increase during the 2020-2023 period. In 2023, low eGFR prevalence was 2.94% (95%CI

1.61–4.28%), remaining stable from 2016 to 2022, and showing a slight increase in 2023

(**Figure 1D, Table 1**). Prevalence was similar between men and women, but higher among

adults ≥60 years (**Figure 1E, F**), in individuals with diabetes or hypertension (**Figure 1G,**

H), and no differences were observed in individuals with obesity (**Figure 1I**). The estimated

number of individuals with low eGFR in Mexico ranged from 1,264,500 (95%CI 822,068–

1,706,931) in 2016 to 2,251,125 (95%CI 1,240,552–3,261,697) in 2023 (**Supplementary**

Table 2).

Comparison of low eGFR prevalence using different CKD-EPI equations in ENSANUT

eGFR was consistently higher using the 2021 CKD-EPI equation, with a median of 111.27

(IQR 99.1–121.1) compared to a median of 107.9 (IQR 95.1–119.4) using the 2009 CKD-

EPI equation (**Figure 2A**). Consequently, the prevalence of G3a-G5 categories was also

higher when using the 2009 compared with the 2021 equation (**Figure 2B,**

Supplementary Table 3 and 4). This trend persisted regardless of survey cycle, sex, age,

diabetes, or hypertension status (**Figure 2C–F**). The difference between equations was

much lower among individuals with higher eGFR (i.e. participants <60 years, without

diabetes, or without hypertension). Prevalence estimates of a CKD proxy comprising

previous self-reported medical diagnosis of CKD or low eGFR for both equations can be

found in **Supplementary Table 5,** and **Supplementary Figure 1.**

Reclassification in eGFR categories using the 2021 CKD-EPI equation

Using the 2021 CKD-EPI equation resulted in reclassification of 6.52% (95%CI 4.07-8.97)

of adults to higher eGFR categories in 2023, with similar estimates over the 2016-2023

period (**Table 1**). Reclassified individuals were predominately men, older, had a higher

proportion of diabetes, and hypertension compared to those not reclassified

(**Supplementary Table 6**). Adoption of 2021 CKD-EPI equation would result in 247,962

less individuals with low eGFR in 2016 and 486,532 less in 2023 (**Supplementary Table**

2); moreover, a total of 4,986,795 individuals (95%CI 3,009,252 to 6,964,338) would be

reclassified upwards when using the 2021 CKD-EPI equation in 2023.

Predictive performance of 2009 and 2021 CKD-EPI equations for cause-specific mortality

Among 159,517 participants recruited from 1998 to 2004 in MCPS, 142,884 had complete

covariate and mortality data, and were included in the analyses (**Supplementary Figure**

2, **Supplementary Table 7**). At the 10-year administrative censoring, there were 12,823

all-cause, 4,277 cardiovascular, and 1,562 kidney-related deaths (**Supplementary Figure**

3). Similar to our findings in ENSANUT, eGFR distribution was higher using the 2021

compared to the 2009 equation (**Figure 3A**), and the prevalence of each eGFR category

and low eGFR were lower with the 2021 equation (**Supplementary Tables 8-9**). Risk for

all-cause mortality was similar between equations in both extended models, with slightly

lower HRs when using the 2021 equation, particularly at higher eGFR values (**Figure 3B**).

When evaluating discrimination and overall performance for all-cause, cardiovascular, and

kidney-related mortality at 5 and 10-years, we found similar discrimination and

performance metrics when adding eGFR calculated with either the 2009 or the 2021

equation to the baseline model. Albeit small in absolute terms, relative larger

improvements were observed in Brier scores, Chi-squared statistics and log-likelihood

when adding the 2021 equation to the baseline model for all 5-year outcomes and for 10-

year cardiovascular mortality. In contrast, adding the 2009 equation to the baseline model

improved prediction for 10-year all-cause, and kidney-related mortality **(Figures 3C-D,**

Table 3 and **Supplementary Table 10**). Full report of all fitted Cox and Fine & Gray

models are reported in **Supplementary Tables 11-22**.

Prognostic significance of reclassification in eGFR categories

Using the 2021 equation, 11,838/142,884 participants (8.29%) were reclassified to higher

eGFR categories compared to categories classified using the 2009 equation. Amongst

them, 10,414/138,256 participants (7.53%) were reclassified from G2 to G1, and

1,061/4,628 (22.92%) were reclassified from G3a-G5 upward. Participants who were

reclassified from categories G3a-G5 upward using the 2021 equation had a lower risk of

all-cause, cardiovascular, and kidney-related mortality at 5 and 10 years compared to

those not reclassified in the same categories. Similarly, participants classified upwards

from G2 to G1 had a lower risk of 5- and 10-year all-cause, and 5-year cardiovascular

mortality. No association was observed for cardiovascular mortality for those reclassified

upward from G2 to G1 (**Figure 4, Table 3**).

DISCUSSION

To our knowledge, this is the first study estimating the prevalence of eGFR categories and

low eGFR using a nationally representative sample in Mexico. Our results using the 2021

CKD-EPI creatinine-based equation, show stable prevalence estimates of eGFR

categories from 2016 to 2023. eGFR estimates were consistently lower using the 2009

CKD-EPI equation and, consequently, there was a pattern of upward reclassification

adopting the newest 2021 CKD-EPI creatinine-based equation, particularly for older adults

with comorbidities. Accordingly, CKD prevalence based only on low eGFR with the 2009

equation was higher compared to the 2021 equation. The predictive performance was

similar for both equations; however, individuals reclassified to higher eGFR categories

using the 2021 equation showed a lower risk of mortality outcomes compared to similar

categories using the 2009 equation. Our results support the use of the 2021 equation to

estimate GFR in Mexican population as it shows prognostically relevant reclassification of

eGFR categories; however, we argue that the choice of equation should be made based

on individual countries' population structures, and on a wider array of kidney-related

outcomes.

Previous reports have estimated CKD prevalence in Mexico relying on heterogeneous

data sources and statistical methods[23,24], or in specific geographical locations[25–28]. An

analysis for the Global Burden of Disease estimated around 14.5 million prevalent cases of

CKD in Mexico on 2017[1], which contrast significantly with our estimations. Based solely on

eGFR, we estimate that ~2.25 million individuals are living with CKD in 2023, which

increases to ~2.4 million when considering prior CKD self-reported diagnosis in ENSANUT.

These estimates of low eGFR prevalence in Mexico are lower compared to estimates for

white adults in the US (2.9% vs. 6.3%), and similar to estimates from Hispanic adults

(2.2%) derived from NHANES 2017-2020. This indicates either ethnic-specific differences

in kidney function or CKD burden, or potential eGFR overestimation in adults of Hispanic

descent. Further studies are warranted to provide more accurate estimates of CKD in

Mexico, considering chronicity, albuminuria, cystatin-C measures, or the number of RRT

users.

Recent clinical guidelines in the US have advocated for the implementation of eGFR

equations without a race coefficient[2]. This recommendations has, however, raised

concerns given that the 2021 CKD-EPI equation may lead to higher eGFR, and lower

prevalence estimates in non-black individuals[6]. Since no specific equations have been

developed for Hispanic of specifically for Mexican population, the use of eGFR equations

without a race coefficient may lead to more precise estimates over prior alternatives.

Similar to our estimations, previous studies in diverse populations have reported higher

eGFR distribution, and lower CKD prevalence using the 2021 CKD-EPI equation, with

upward reclassification especially among men, and older individuals[29–31]. Notably, Mexican

clinical guidelines on CKD which were last updated in 2019 recommend GFR estimation

with either Cockroft-Gault equation, the MDRD equation, or the 2009 CKD-EPI creatinine-

based equation[32]. It should be noted that Cockroft-Gault is a creatinine clearance

equation, and MDRD systematically underestimates higher GFR values[33]. We observed

that the 2021 CKD-EPI equation yielded a similar but slightly improved prediction for all-

cause, cardiovascular and kidney-related mortality compared to the 2009 equation, and

reclassifies individuals at lower risk of these outcomes to higher eGFR categories. Our

results support that the 2021 CKD-EPI creatinine-based equation should a reasonable

choice for populations like Mexico, which is relatively homogeneous and comprised mainly

of individuals with Hispanic descent. Additionally, creatinine measurements are widely

used across the country to evaluate kidney function in clinical practice, and due to cost

constraints, widespread implementation of cystatin C testing in Mexico remains a distant

prospect.

Our study has several strengths. To our knowledge, this is the first study estimating GFR

and CKD prevalence directly from creatinine measurements in Mexico, rather than relying

on statistical methods to infer estimates. Additionally, given the data availability since 2016,

we were able to obtain trends of eGFR and CKD estimates over the last six nationally

representative surveys. Furthermore, access to data from a prospective study with

mortality follow-up, allowed us to evaluate the predictive performance of eGFR equations

for cause-specific outcomes in a predominately Hispanic population, broadening similar

results in mostly white and European countries. However, some limitations should be

considered to adequately interpret our results. First, CKD prevalence was estimated

considering only low eGFR, and we were unable to include albuminuria to classify

individuals as recommended by clinical guidelines[2]. Second, given the cross-sectional

nature of the study, our definition of CKD relied on a single creatinine measurement, which

lacks the three-month chronicity criterion, leading to a potential overestimation of CKD

prevalence. Notably, our estimates did not account for the use of RRT. Moreover, due to

the lack of data, we were unable to evaluate eGFR and CKD prevalence using the

creatinine and cystatin C equation, which has been reported as the most accurate[6]. Finally,

despite there being Afro-Mexican communities, its representation in ENSANUT and MCPS

is uncertain. Given the admixture of Mexican population, most participants self-recognize

as Mestizos[7], and therefore neither ENSANUT nor MCPS routinely collect ethnicity data,

which precluded us from being able to assess differences in prevalence and predictive

performance for black Mexican adults.

In conclusion, eGFR and CKD prevalence based on low eGFR in Mexico have been stable

over time, with some recent increases in 2023. As reported in predominately white

Europeans, the new 2021 CKD-EPI equation consistently estimates higher GFR levels,

which translates to lower CKD prevalence estimates and upward reclassification

predominately in men and older individuals. Additionally, the 2021 equation yields a better

predictive performance and adequately reclassified participants according to their risk for

all-cause, cardiovascular and kidney-related morality compared to the 2009 equation.

Therefore, we recommend implementing the creatinine-based 2021 CKD-EPI equation as

a reliable choice for estimating eGFR and tracking CKD prevalence in Mexico.

ACKNOWLEDGMENTS

This project was registered and approved by the Research Committee at Instituto Nacional

de Geriatría, project number DI-PI-009-2024. JPE and CAFM are enrolled at the PECEM

Program of the Faculty of Medicine at UNAM and are supported by CONACyT. JAS was

supported by Grant Number K23DK135798 from the NIH/NIDDK and by the

Massachusetts General Hospital Executive Committee and Center for Diversity and

Inclusion Physician-Scientist Development Award. This research was conducted using

Mexico City Prospective Study (MCPS) data obtained through an open-access data

request (application number 2022-012). MCPS is a long-standing scientific collaboration

between researchers at the National Autonomous University of Mexico and the University

of Oxford, and has received funding from the Mexican Health Ministry, the National Council

of Science and Technology for Mexico, Wellcome, Cancer Research UK, the British Heart

Foundation, Kidney Research UK and the UK Medical Research Council. The authors

thank the participants for their willingness to take part in this prospective study 20 years

ago.

AUTHOR CONTRIBUTIONS

Research idea and study design: DRG, CAFM, PSC, OYBC; data acquisition: CAFM,

DRG, PSC, JAD, PKM, JBC, RTC, OYBC; analysis/interpretation: DRG, CAFM, OYBC;

statistical analysis: DRG, CAFM, OYBC; manuscript drafting: DRG, CAFM, PSC, BGCF,

JPE, JPDS, KBCH, LACQ, JCB, PKM, RTC, JAD, NEAV, JAS, OYBC; supervision or

mentorship: OYBC. Each author contributed important intellectual content during

manuscript drafting or revision and accepts accountability for the overall work by ensuring

that questions pertaining to the accuracy or integrity of any portion of the work are

appropriately investigated and resolved.

DATA AVAILABILITY: All code and materials regarding ENSANUT are available for

reproducibility of results http://github.com/oyaxbell/ckdepi_mcps_ensanut/. Data from the

Mexico City Prospective Study are available to bona fide researchers. For more details,

the study's Data and Sample Sharing policy may be downloaded (in English or Spanish)

from https://www.ctsu.ox.ac.uk/research/mcps. Available study data can be examined in

detail through the study's Data Showcase, available at

https://datashare.ndph.ox.ac.uk/mexico/.

CONFLICT OF INTEREST/FINANCIAL DISCLOSURE: Nothing to disclose.

FUNDING: This research was supported by Instituto Nacional de Geriatría in Mexico.

ETHICAL DISCLOSURES: This project was registered and approved by the Research

Committee at Instituto Nacional de Geriatría, project number DI-PI-009-2024. MCPS was

approved by Ethics Committees at the Mexican Ministry of Health, the Mexican National

Council for Science and Technology, and the University of Oxford, UK. All MCPS

participants provided written informed consent.

REFERENCES
1.   Bikbov B, Purcell CA, Levey AS, et al. Global, regional, and national burden of
chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of
Disease Study 2017. *The Lancet*. 2020;395(10225):709-733. doi:10.1016/S0140-
6736(20)30045-3
2.   Stevens PE, Ahmed SB, Carrero JJ, et al. KDIGO 2024 Clinical Practice Guideline
for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int*.
2024;105(4):S117-S314. doi:10.1016/j.kint.2023.10.018
3.   Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO
2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney
Disease. *Kidney Inter*. 2013;Suppl.(3):1-150.
4.   Levey AS, Stevens LA, Schmid CH, et al. A New Equation to Estimate Glomerular
Filtration Rate. *Ann Intern Med*. 2009;150(9):604. doi:10.7326/0003-4819-150-9-
200905050-00006
5.   Delgado C, Baweja M, Crews DC, et al. A Unifying Approach for GFR Estimation:
Recommendations of the NKF-ASN Task Force on Reassessing the Inclusion of Race in
Diagnosing Kidney Disease. *J Am Soc Nephrol*. 2021;32(12):2994-3015.
doi:10.1681/ASN.2021070988
6.   Inker LA, Eneanya ND, Coresh J, et al. New Creatinine- and Cystatin C–Based

Equations to Estimate GFR without Race. *N Engl J Med.* 2021;385(19):1737-1749.
doi:10.1056/NEJMoa2102953

7.   Sohail M, Palma-Martínez MJ, Chong AY, et al. Mexican Biobank advances
population and medical genomics of diverse ancestries. *Nature.* 2023;622(7984):775-783.
doi:10.1038/s41586-023-06560-0

8.   Sharma S, Nagar SD, Pemu P, et al. Genetic ancestry and population structure in
the All of Us Research Program cohort. *Nat Commun.* 2025;16(1):4123.
doi:10.1038/s41467-025-59351-8

9.   Ricardo AC, Flessner MF, Eckfeldt JH, et al. Prevalence and Correlates of CKD in
Hispanics/Latinos in the United States. *Clin J Am Soc Nephrol CJASN.* 2015;10(10):1757-
1766. doi:10.2215/CJN.02020215

10.   Rodriguez RA, Hernandez GT, O'Hare AM, Glidden DV, Pérez-Stable EJ.
Creatinine levels among Mexican Americans, Puerto Ricans, and Cuban Americans in the
Hispanic Health and Nutrition Examination Survey. *Kidney Int.* 2004;66(6):2368-2373.
doi:10.1111/j.1523-1755.2004.66025.x

11.   Hsu CY, Yang W, Parikh RV, et al. Race, Genetic Ancestry, and Estimating Kidney
Function in CKD. *N Engl J Med.* 2021;385(19):1750-1760. doi:10.1056/NEJMoa2103753

12.   Romero-Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, et al. Metodología de
la Encuesta Nacional de Salud y Nutrición 2022 y Planeación y diseño de la Ensanut
Continua 2020-2024. *Salud Pública México.* 2022;64(5, sept-oct):522-529.
doi:10.21149/14186

13.   Romero-Martínez M, Shamah-Levy T, Cuevas-Nasu L, et al. Diseño metodológico
de la Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. *Salud Pública*
México. 2017;59(3, may-jun):299. doi:10.21149/8593

14.   Romero-Martínez M, Shamah-Levy T, Vielma-Orozco E, et al. Encuesta Nacional
de Salud y Nutrición 2018-19: metodología y perspectivas. *Salud Pública México.*
2019;61(6, nov-dic):917-923. doi:10.21149/11095

15.   Romero Martínez M, Barrientos-Gutiérrez T, Cuevas-Nasu L, et al. Metodología de
la Encuesta Nacional de Salud y Nutrición 2021. *Salud Pública México.* 2021;63(6, Nov-
Dic):813-818. doi:10.21149/13348

16.   Tapia-Conyer R, Kuri-Morales P, Alegre-Díaz J, et al. Cohort Profile: The Mexico
City Prospective Study. *Int J Epidemiol.* 2006;35(2):243-249. doi:10.1093/ije/dyl042

17.   Alegre-Díaz J, Herrington W, López-Cervantes M, et al. Diabetes and Cause-
Specific Mortality in Mexico City. *N Engl J Med.* 2016;375(20):1961-1971.
doi:10.1056/NEJMoa1605368

18.   Chronic Kidney Disease Prognosis Consortium. Association of estimated
glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in
general population cohorts: a collaborative meta-analysis. *The Lancet.*
2010;375(9731):2073-2081. doi:10.1016/S0140-6736(10)60674-5

19.   Steyerberg EW, Pencina MJ, Lingsma HF, Kattan MW, Vickers AJ, Van Calster B.
Assessing the incremental value of diagnostic and prognostic markers: a review and
illustration. *Eur J Clin Invest.* 2012;42(2):216-228. doi:10.1111/j.1365-2362.2011.02562.x

20.   McLernon DJ, Giardiello D, Van Calster B, et al. Assessing Performance and
Clinical Usefulness in Prediction Models With Survival Outcomes: Practical Guidance for
Cox Proportional Hazards Models. *Ann Intern Med.* 2023;176(1):105-114.
doi:10.7326/M22-0844

21.   Therneau TM, Grambsch PM. *Modeling Survival Data: Extending the Cox Model.*
Springer; 2000.

22.   Harrell FE. *Regression Modeling Strategies: With Applications to Linear Models,*
Logistic and Ordinal Regression, and Survival Analysis. Second edition. Springer; 2015.

23.   Argaiz ER, Morales-Juárez L, Razo C, et al. La carga de enfermedad renal crónica

en México. Análisis de datos basado en el estudio *Global Burden of Disease* 2021. *Gac*
Médica México. 2023;159(6):12575. doi:10.24875/GMM.23000393

24.   Agudelo-Botero M, Valdez-Ortiz R, Giraldo-Rodríguez L, et al. Overview of the
burden of chronic kidney disease in Mexico: secondary data analysis based on the Global
Burden of Disease Study 2017. *BMJ Open*. 2020;10(3):e035285. doi:10.1136/bmjopen-
2019-035285

25.   Gutierrez-Peña M, Zuñiga-Macias L, Marin-Garcia R, et al. High prevalence of end-
stage renal disease of unknown origin in Aguascalientes Mexico: role of the registry of
chronic kidney disease and renal biopsy in its approach and future directions. *Clin Kidney*
J. 2021;14(4):1197-1206. doi:10.1093/ckj/sfaa229

26.   Aguilar-Ramirez D, Raña-Custodio A, Villa A, et al. Decreased kidney function and
agricultural work: a cross-sectional study in middle-aged adults from Tierra Blanca,
Mexico. *Nephrol Dial Transplant*. 2021;36(6):1030-1038. doi:10.1093/ndt/gfaa041

27.   Amato D, Alvarez-Aguilar C, Castaneda-Limones R, et al. Prevalence of chronic
kidney disease in an urban Mexican population. *Kidney Int*. 2005;68:S11-S17.
doi:10.1111/j.1523-1755.2005.09702.x

28.   Obrador GT, García-García G, Villa AR, et al. Prevalence of chronic kidney disease
in the Kidney Early Evaluation Program (KEEP) México and comparison with KEEP US.
Kidney Int. 2010;77:S2-S8. doi:10.1038/ki.2009.540

29.   Fu EL, Coresh J, Grams ME, et al. Removing race from the CKD-EPI equation and
its impact on prognosis in a predominantly White European population. *Nephrol Dial*
Transplant. 2023;38(1):119-128. doi:10.1093/ndt/gfac197

30.   Vestergaard SV, Heide-Jørgensen U, Birn H, Christiansen CF. Effect of the
Refitted Race-Free eGFR Formula on the CKD Prevalence and Mortality in the Danish
Population. *Clin J Am Soc Nephrol*. 2022;17(3):426-428. doi:10.2215/CJN.14491121

31.   Gregg LP, Richardson PA, Akeroyd J, Matheny ME, Virani SS, Navaneethan SD.
Effects of the 2021 CKD-EPI Creatinine eGFR Equation among a National US Veteran
Cohort. *Clin J Am Soc Nephrol*. 2022;17(2):283-285. doi:10.2215/CJN.10000721

32.   CENETEC. Prevención, Diagnóstico y Tratamiento de la Enfermedad Renal
Crónica. Guía de Evidencias y Recomendaciones: Guía de Práctica Clínica. Published
online 2019.

33.   Stevens LA, Coresh J, Feldman HI, et al. Evaluation of the modification of diet in
renal disease study equation in a large diverse population. *J Am Soc Nephrol JASN*.
2007;18(10):2749-2757. doi:10.1681/ASN.2007020199

TABLES

eGFR categories						
Category	**2016**	**2018**	**2020**	**2021**	**2022**	**2023**
G1	88.90% (87.24-90.57)	87.39% (86.36-88.43)	89.88% (88.47-91.30)	85.82% (83.57-88.07)	89.02% (86.96-91.07)	87.08% (84.31-89.84)
G2	9.30% (7.74-10.85)	10.63% (9.67-11.60)	8.23% (6.96-9.50)	12.06% (9.98-14.14)	9.13% (7.21-11.06)	9.98% (7.76-12.20)
G3a	0.93% (0.53-1.34)	1.06% (0.81-1.31)	0.97% (0.54-1.40)	1.09% (0.49-1.70)	0.73% (0.36-1.10)	1.76% (0.77-2.76)
G3b	0.37% (0.15-0.60)	0.52% (0.29-0.74)	0.48% (0.13-0.82)	0.32% (0.03-0.61)	0.40% (0.12-0.69)	0.90% (0.22-1.57)
G4	0.14% (0.01-0.27)	0.26% (0.14-0.38)	0.22% (0.03-0.41)	0.57% (0.00-1.38)	0.53% (0.00-1.24)	0.00% (0.00-0.01)
G5	0.35% (0.00-0.71)	0.14% (0.06-0.22)	0.22% (0.05-0.39)	0.14% (0.00-0.37)	0.18% (0.01-0.36)	0.28% (0.00-0.59)
Low eGFR						
eGFR <60 mL/min 1.73 m^2	1.80% (1.17–2.43)	1.97% (1.59–2.35)	1.89% (1.28–2.50)	2.12% (1.05–3.19)	1.85% (0.95–2.73)	2.94% (1.61–4.28)
eGFR category reclassification with CKD-EPI 2021						
Upward reclassification	4.72% (3.56-5.88)	5.16% (4.50-5.82)	5.27% (4.15-6.39)	4.67% (3.33-6.02)	4.62% (3.38-5.86)	6.52% (4.07-8.97)

Table 1. Nationally representative prevalence with 95% confidence intervals from 2016 to 2023 of eGFR categories, low eGFR (<60 mL/min 1.73 m^2) and eGFR category reclassification using ENSANUT data. eGFR was estimated using the CKD-EPI 2021 creatinine-based equation. G1 normal or high (≥90 mL/min per 1.73 m^2), G2 mildly decreased (60–89 mL/min/1.73 m^2), G3a mildly to moderately decreased (45–59 mL/min/1.73 m^2), G3b moderately to severely decreased (30–44 mL/min/1.73 m^2), G4 severely decreased (15-29 mL/min/1.73 m^2), and G5 kidney failure (<15 mL/min/1.73 m^2). The table also shows the prevalence of eGFR category reclassification using the 2021 CKD-EPI equation from categories classified originally using the 2009 CKD-EPI equation. **Abbreviations:** eGFR: Estimated glomerular filtration rate, CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration.

Outcome	Model	Discrimination	5-year overall performance		10-year overall performance	
		Harrell C-statistic (95%CI)	Brier score (95%CI)	Scaled Brier score	Brier score (95%CI)	Scaled Brier score
All-cause mortality	**Baseline**	0.674 (0.669-0.68)	0.036 (0.035–0.037)	9.31%	0.055 (0.054-0.056)	26.62%
	+ eGFR 2009	0.690 (0.684-0.695)	0.035 (0.034–0.036)	12.77%	0.053 (0.052-0.054)	28.65%
	+ eGFR 2021	0.689 (0.684-0.695)	0.035 (0.034–0.036)	12.79%	0.053 (0.052-0.054)	28.66%
Cardiovascular mortality	**Baseline**	0.672 (0.662-0.681)	0.013 (0.013-0.014)	5.41%	0.027 (0.026-0.028)	9.78%
	+ eGFR 2009	0.681 (0.671-0.690)	0.013 (0.013-0.014)	6.44%	0.027 (0.026-0.028)	10.46%
	+ eGFR 2021	0.680 (0.671-0.690)	0.013 (0.013-0.014)	6.44%	0.027 (0.026-0.028)	10.46%
Kidney-related mortality	**Baseline**	0.834 (0.824-0.844)	0.014 (0.013-0.014)	1.61%	0.029 (0.028-0.03)	2.58%
	+ eGFR 2009	0.878 (0.869-0.887)	0.014 (0.013-0.014)	0.94%	0.029 (0.029-0.03)	1.83%
	+ eGFR 2021	0.878 (0.869-0.887)	0.014 (0.013-0.014)	0.92%	0.029 (0.029-0.03)	1.82%

471 **Table 2.** Discrimination assessment and overall performance assessment of all-cause, cardiovascular and kidney-related mortality for the

472 baseline model and both extended models in participants of the Mexico City Prospective Study. The baseline model was stratified by age

473 category and adjusted for sex, place of residence (Coyoacán or Iztapalapa), educational level (elementary, high school, university, other),

smoking, diabetes, history of cardiovascular disease, systolic blood pressure, and total cholesterol. eGFR: Estimated glomerular filtration rate.

Outcome	Reclassification with CKD-EPI 2021	5-year			10-year		
		Event rate at 5 years (R/NR)	HR*	95% CI[1]	Event rate at 10 years (R/NR)	HR*	95% CI[1]
All-cause mortality	G2 to G1	13.05 / 15.12	0.90	0.82, 0.97	15.89 / 17.47	0.93	0.88, 0.99
	Upward from G3a-G5	29.51 / 75.97	0.50	0.42, 0.59	29.76 / 63.43	0.60	0.53, 0.68
Cardiovascular mortality	G2 to G1	5.02 / 5.77	0.84	0.73, 0.96	5.82 / 6.46	0.93	0.85, 1.02
	Upward from G3a-G5	11.96 / 23.70	0.58	0.44, 0.76	12.46 /18.19	0.69	0.56, 0.84
Kidney-related mortality	G2 to G1	0.86 / 0.95	0.88	0.63, 1.23	1.07 / 1.31	0.89	0.73, 1.10
	Upward from G3a-G5	6.54 / 32.06	0.33	0.22, 0.48	5.81 / 24.32	0.44	0.33, 0.58

Table 3. Five- and ten-year risk of all-cause, cardiovascular, and kidney-related mortality, as well as age and sex-standardized outcome rates per 1,000 person-years related to reclassification of CKD-EPI 2009 eGFR categories with the CKD-EPI 2021 equation in participants of the Mexico City Prospective Study. All models were stratified by age category and adjusted for sex, place of residence, educational level (elementary, high school, university, other), smoking, diabetes, history of cardiovascular disease, systolic blood pressure, and total cholesterol. All-cause mortality was evaluated using Cox proportional hazard regression models, and cause-specific mortality using Fine & Gray sub distribution hazard models; reference group is No reclassification within the same category (G1-G2 and G3a-G5, respectively). *For all-cause mortality adjusted hazard ratios are presented, for cardiovascular and kidney-related mortality adjusted subdistribution hazard

484   ratios are presented. eGFR: Estimated glomerular filtration rate. Abbreviations: HR, Hazard Ratio; CI, Confidence Interval; R,

485   Reclassification;   NR,   No   reclassification.

FIGURES LEGENDS

Figure 1. Prevalence of eGFR categories and CKD based on low eGFR in Mexico using the 2021 CKD-EPI creatinine equation in ENSANUT 2016-2023 data. (A) G1, (B) G2, (C) G3a, G3b, G4, and G5, (D) Overall CKD prevalence and CKD prevalence stratified by (E) sex, (F) age category, (G) diabetes, (H) hypertension, and (I) obesity status. G1 normal or high (≥90 mL/min per 1.73 m^2), G2 mildly decreased (60–89 mL/min/1.73 m^2), G3a mildly

493   to moderately decreased (45–59 mL/min/1.73 m^2), G3b moderately to severely decreased

494   (30–44 mL/min/1.73 m^2), G4 severely decreased (15-29 mL/min/1.73 m^2), and G5 kidney

495   failure (<15 mL/min/1.73 m^2). eGFR: Estimated glomerular filtration rate, CKD: Chronic

496   Kidney Disease, CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration.

Figure 2. eGFR distribution and comparison of CKD prevalence using either 2021 or 2009 CKD-EPI creatinine equations in ENSANUT 2016-2023 data. (A) eGFR distribution, (B) Comparison of overall CKD prevalence using both equations, (C) Comparison of CKD prevalence in men, (D) age category, (E) diabetes status, and (F) hypertension status. eGFR: Estimated glomerular filtration rate, CKD: Chronic Kidney Disease, CKD-EPI: Chronic Kidney Disease Epidemiology Collaboration.

Figure 3. eGFR distribution and all-cause mortality risk prediction performance of eGFR

estimated with either 2021 or 2009 CKD-EPI creatinine equation in the Mexico City

Prospective Study. (A) Comparison of eGFR distribution in MCPS, (B) Comparison of HRs

estimations across continuous eGFR levels, (C) 5-year discrimination assessment of

eGFR estimated with either 2009 or 2021 CKD-EPI creatinine equation, (D) 10-year

discrimination assessment of eGFR estimated with either 2009 or 2021 CKD-EPI

creatinine equation. In both C and D, the gray line represents the baseline model without

the eGFR covariate. All models were stratified by age category and adjusted for sex, place

of residence, educational level, smoking, diabetes, history of cardiovascular disease,

systolic blood pressure, and total cholesterol. To this baseline model, GFR estimated with

either the 2021 or 2009 CKD-EPI creatinine equation was added. eGFR: Estimated

glomerular filtration rate, CKD: Chronic Kidney Disease, CKD-EPI: Chronic Kidney

Disease Epidemiology Collaboration.

523 **Figure 4.** Kaplan-Meier cumulative incidence curves adjusted using inverse probability of treatment weighting (IPTW) by age, sex,

524 community of origin, educational level, smoking, diabetes, prior cardiovascular disease, systolic blood pressure and total cholesterol

525 for all-cause, cardiovascular and kidney-related mortality in MCPS participants. Panels A-C compare participants reclassified from G2

526   to G1 and Panels D-F participants reclassified from G3a-G5 upward using the 2021 eGFR creatinine-based CKD-EPI equation

527   compared to non-reclassified participants in the same categories for risk of these outcomes.