A natural experiment in Kenya reveals durable immunosuppressive effects of early

childhood malaria: a longitudinal cohort study

Mercy S. Safari[1,2,6], Timothy O. Makori[1,2], Elijah T. Gicheru[1,2], Maureen W. Mburu[1,2], Omar

Nyawa[1,2], Faiz Shee[6], James Nyagwange[1], Eunice W. Kagucia[1], Francis Ndungu[1,3], Timothy

Chege[,5], James O. Tuju[1#], Charles J. Sande[1,2,3,4*#]

1. Bioscience Department, KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya

2.Epidemiology and Demography Department, KEMRI-Wellcome Trust Research

Programme, Kilifi, Kenya

3. Centre for Tropical Medicine & Global Health, University of Oxford, Oxford, UK

4. Kenya Medical Research Institute, Centre for Geographic Medicine Research, Coast,

Kilifi, Kenya

5. Department of Microbiology and Immunology, Columbia University Irving Medical Center,

New York, NY 100322 USA

6. Pwani University, Kilifi, Kenya

* Correspondence to:

Charles J. Sande, KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya P.O. Box 230

- 80108 (csande@kemri-wellcome.org)

# Contributed equally

Keywords: malaria, antibody immunosuppression, microarray

Abstract

Background:

Chronic malaria exposure has been proposed to modulate immune function, but its long-term effects on antibody-mediated responses to unrelated pathogens remain poorly defined. Whether these effects persist beyond periods of active infection, and how early-life exposure shapes humoral immunity over time, is not well understood.

Methods:

We leveraged a natural experiment in coastal Kenya - where two regions (Junju and Ngerenya) diverged sharply in malaria transmission from around 2004 - to evaluate the long-term immunological consequences of malaria exposure in childhood. Using a protein microarray platform, we measured IgG responses to vaccine and pathogen antigens in 123 children sampled longitudinally over a 15-year period. Active weekly malaria surveillance enabled precise reconstruction of individual exposure histories.

Results:

IgG responses to *Plasmodium falciparum* apical membrane antigen 1 (AMA1) tracked closely with clinical malaria episodes, confirming the ability of the microarray platform to detect biologically meaningful variation in antigen-specific immunity. Despite comparable vaccination histories, children from the high malaria transmission setting (Junju) exhibited persistently lower measles-specific IgG levels than children from the low-transmission setting (Ngerenya), a pattern validated by ELISA. At 10 years of age, Junju children showed significantly reduced antibody levels to a wide range of unrelated pathogens, including *Bordetella pertussis*, CMV, rubella, and measles. Within the Ngerenya cohort, children with documented early-life malaria had broadly lower IgG responses at age 10 compared to malaria-naïve peers, despite identical geography, vaccines, and follow-up duration.

Conclusions:

These findings suggest that malaria exposure during early childhood is linked with durable suppression of antibody responses to unrelated pathogens and vaccines. This effect persists long after infection and may partially explain the overall diminished long-term vaccine effectiveness in malaria-endemic settings.

Introduction

Malaria remains a major cause of childhood morbidity and mortality in sub-Saharan Africa. In 2023, Africa accounted for 95% of global malaria deaths, with over three-quarters occurring in children under five years old[1]. Alongside malaria, children in these settings face high exposure to a wide array of pathogens - including respiratory, enteric, and helminth infections [2], making immune competence critical for survival and long-term health. Despite widespread vaccination efforts, several studies report reduced vaccine efficacy in malaria-endemic regions compared to malaria-naïve populations [3–5]. Although multiple factors may contribute[6], growing evidence suggests that malaria itself can impair host immunity[5,7,8]. Repeated P. falciparum infection has been associated with immunomodulatory changes, including expansion of regulatory T cells [9,10], regulatory B cells[11], and atypical memory B cells with limited effector function[12,13]. These alterations promote host tolerance for persistent parasitaemia, resulting from recurrent exposure in endemic settings, but they also suppress effector immune responses[14], potentially to unrelated antigens, including vaccines. While malaria-induced immunosuppression has been described in both experimental and observational studies[7,8,15,16], its duration and broader impact on the developing immune system remain poorly understood. Most prior work has focused on short-term or antigen-specific outcomes, leaving open the question of whether early-life malaria exposure durably attenuates antibody responses over the long term. Conflicting findings across settings highlight the need for context-specific, and longitudinal data studies[16,17].

Here, we address this gap using two longitudinal cohorts from coastal Kenya with contrasting malaria transmission histories. One region (Junju) has experienced a sustained malaria burden, while the other (Ngerenya) underwent a rapid decline in transmission after 2004[18]. Intensive weekly malaria surveillance was conducted over more than a decade, enabling precise classification of individual exposure histories. We combined these data with serial serological measurements using a high-throughput in-house protein microarray. This design enabled us to investigate whether clinical malaria in early life leaves a lasting immunological imprint that compromises antibody responses to common childhood pathogens and vaccines. Our findings reveal a durable suppression of humoral immunity

linked to malaria exposure in early childhood, with implications for vaccine effectiveness,
serosurveillance interpretation, and immune recovery in endemic regions.

Materials and Methods

Study setting and cohort design - This study was conducted in Kilifi County, a rural region on
the northern coast of Kenya, within the catchment area of the Kilifi Health and Demographic
Surveillance System (KHDSS), a long-term population-based platform maintained by the
KEMRI-Wellcome Trust Research Programme[19]. Serum samples were obtained
retrospectively from two well-characterized paediatric cohorts - Ngerenya and Junju -
enrolled between 1998 and 2017 as part of annual malaria cross-sectional surveys[20].
Ngerenya and Junju were selected for their contrasting malaria transmission intensities;
Ngerenya experienced a sharp decline in malaria transmission beginning in the early
2000s[18], whereas Junju maintained moderate malaria endemicity throughout the study
period, with *P.falciparum* prevalence approximating 30% during the rainy seasons[21]. All
children were visited weekly at home for the detection of febrile episodes, and any child
with an axillary temperature ≥37.5°C was tested for P. falciparum parasitaemia using a rapid
diagnostic test, with confirmation by microscopic examination of Giemsa-stained thick and
thin blood smears. A clinical malaria episode was defined as fever in the presence of ≥2,500
parasites/µL. In addition to active malaria surveillance, serum samples were collected
annually from each child and for future serological analysis. The vaccination status of each
child for routine childhood vaccines was assessed using digitized immunization records
stored at the KEMRI-Wellcome Trust Research Programme.

Protein microarray antibody profiling - Antibody responses were measured using an in-
house protein microarray platform. Recombinant and whole-virus lysate antigens
(Supplementary table 1) were reconstituted in a glycerol-based buffer containing 1% Triton
X-100 and printed in quadruplicate onto epoxy-coated glass slides using a non-contact
microarrayer (Marathon Argus, Arrayjet, Scotland). Slides were fitted into hybridization
cassettes, washed with PBST (0.05% Tween-20 in PBS), and blocked for 1 hour at 37°C using
PBST containing 5% BSA. Serum samples were diluted 1:30 in PBST with 5% BSA and
incubated on the slides for 3 hours at room temperature. Slides were washed and probed
with secondary antibodies: goat anti-human IgG conjugated to Alexa Fluor 647

(SouthernBiotech, cat. no. 2040-03) and goat anti-human IgA conjugated to Alexa Fluor 555

(SouthernBiotech, cat. no. 2050-32), each incubated for 1.5 hours at 25°C. Slides were

rinsed, disassembled, and scanned using a GenePix 4300A scanner (635 nm at PMT 600, 532

nm at PMT 700). Mean fluorescence intensities (MFIs) were extracted from duplicate mini-

arrays per slide and averaged across four replicate spots per antigen. Data processing and

quality control of microarray data were performed in R (version 4.4.2).

Enzyme-linked immunosorbent assay (ELISA) - Measles-specific IgG was quantified using a

conventional direct ELISA. Plates were coated overnight at 4°C with 2.30 μg/mL measles

antigen diluted in PBS. After blocking with 5% skimmed milk for 1 hour at 37°C, serum

samples (1:100 dilution) were added and incubated for 1.5 hours at 37°C. Plates were

washed with PBST and incubated with HRP-conjugated secondary antibody (1:100 dilution)

for 1 hour. Following additional washes, 100 μL of OPD substrate solution (30 mg OPD in 30

mL PBS with 30 μL H_2O_2) was added and incubated in the dark for 10 minutes. Reactions

were stopped with 50 μL of 2.5 M sulfuric acid, and absorbance was measured at 490 nm.

Statistical analysis - All statistical analyses were performed using R (version 4.4.2). Antibody

responses between cohorts were compared using the Wilcoxon rank-sum test. P-values

<0.05 were considered statistically significant. Data were anonymized and delinked from all

personally identifiable information.

Ethics statement - The study was approved by the Scientific and Ethics Review Unit (SERU)

of the Kenya Medical Research Institute. All procedures were conducted in accordance with

the principles of Good Clinical Laboratory Practice (GCLP).

Results

A natural experiment in coastal Kenya reveals sharply divergent malaria exposure

trajectories in early childhood - Between 1998 and 2017, two rural communities in coastal

Kenya - Junju and Ngerenya - underwent markedly different transitions in malaria

transmission. Both regions experienced a high malaria burden in the 1990s and early 2000s.

However, beginning in 2004, transmission in Ngerenya declined sharply and remained near

zero for more than a decade, while Junju continued to experience sustained endemicity well

into the mid-2010s. To quantify the scale and timing of this transition, we analyzed

surveillance data collected from August 1998 to April 2017. A total of 1,243 children were followed in Ngerenya and 659 in Junju. In Ngerenya, the proportion of febrile surveillance visits with confirmed P. falciparum parasitaemia declined from 22.4% before 2004 (1,378 of 6,148 visits) to just 1.1% after 2004 (48 of 4,389 visits). In contrast, Junju children continued to experience high rates of febrile malaria, with 17% of visits testing positive between 2007 and 2017 (869 of 5,130 visits) - **Fig. 1**.

To assess whether the protein microarray platform accurately captured longitudinal P. falciparum-specific immune responses, we examined IgG levels against the plasmodium apical membrane antigen 1 (AMA1) in individual children with known malaria exposure histories. Among children with multiple confirmed episodes of febrile malaria detected via weekly active surveillance, AMA1-specific IgG levels increased sharply over time, tracking closely with the timing of clinical infections (example shown in **Fig. 2a**), while children from Ngerenya who remained malaria-free throughout follow-up exhibited consistently low IgG levels against AMA1 over more than a decade of surveillance (example shown in **Fig. 2b**). These patterns mirrored expected immunological trajectories of repeated exposure versus non-exposure, confirming that the microarray platform is capable of detecting meaningful variation in P. falciparum-specific humoral responses. We then extended this analysis to assess longitudinal IgG responses to AMA1 in a subset of 123 children drawn from the two surveillance cohorts. These children contributed a total of 1,194 serum samples, with a mean of 10 samples per child, collected between 2002 and 2017 (**Supplementary Fig. 1**). This subset was selected on the basis of the relative completeness of their longitudinal follow-up serum sampling and included 58 children from Junju (505 samples) and 65 from Ngerenya (689 samples). AMA1-specific IgG levels diverged sharply between cohorts early in life and remained distinct throughout follow-up. In Ngerenya, levels declined rapidly after 2003, stabilizing at lower levels by mid-childhood. In contrast, Junju children maintained elevated levels across all time points (**Fig. 2c**).

Malaria-exposed children exhibit lower antibody levels to non-malarial antigens - To examine the impact of differential malaria endemicity on the antibody response to non-malarial antigens, we first compared IgG responses to the measles virus among Junju and Ngerenya children. We started by validating the ability of the in-house protein microarray

platform to detect biologically meaningful measles-specific IgG responses by examining
longitudinal measles levels in individual children with known vaccination histories. Among
children with a documented history of receiving all three recommended doses of measles
vaccine, we observed a sharp rise in IgG levels following immunization, followed by
sustained levels into later childhood (example shown in **Fig. 3a**). In contrast, unvaccinated
children exhibited consistently low IgG trajectories across all time points (example shown in
Fig. 3b). These patterns were reproducible across the cohort and recapitulated expected
vaccine-induced versus naïve antibody dynamics, supporting the use of this platform for
population-level serological inference. Using this platform, we then compared IgG responses
to measles virus among children from Junju and Ngerenya. This analysis was restricted only
to children with a documented record of measles vaccination. Despite matched vaccination
histories, children from Junju - where malaria transmission remained high - consistently
exhibited lower measles-specific IgG titers than children from Ngerenya, where malaria
transmission had declined. This difference was evident from the earliest timepoints and
persisted throughout childhood. Annual antibody measurements showed that mean
measles-specific IgG levels were higher in Ngerenya than in Junju in every sampling year (**Fig.**
3c). To validate these findings, we measured measles-specific IgG levels by ELISA in a subset
of 3-year-old children who had completed measles vaccination at least one year prior. In this
assay, Junju children again displayed significantly lower IgG levels than their Ngerenya
counterparts (**Fig. 3d**).

Early-life malaria exposure is associated with long-term suppression of antibody responses
To assess whether the attenuation of antibody responses extended beyond measles, we
compared IgG levels to a broader panel of antigens at 10 years of age. This analysis included
vaccine-preventable pathogens (*Bordetella pertussis*, H1N1 influenza virus, rubella virus, and
measles virus) and common childhood infections (cytomegalovirus (CMV), Epstein-Barr virus
(EBV), herpes simplex virus 1 (HSV-1), coxsackievirus B1). Children from Junju exhibited
significantly lower IgG levels for most antigens compared to their Ngerenya counterparts
(**Fig. 4**). This pattern was observed for both viral and bacterial pathogens and was
particularly marked for coxsackievirus, EBV, HSV-1, and measles. Antibody levels against P.
falciparum AMA1 remained elevated in Junju. Antibody levels to the 2009 pandemic strain

of H1N1 were the least differentiated between cohorts, although Ngerenya children still

exhibited significantly higher levels than Junju participants (**Fig. 4**).

To determine whether the attenuation of antibody responses could still be attributed to

early-life malaria exposure independent of geographic or environmental differences, we

conducted a stratified analysis within the Ngerenya cohort, which experienced a sharp

decline in malaria transmission beginning in 2004. Due to stochastic differences in malaria

infection around the time of this inflection, children in Ngerenya had different malaria

exposure histories despite living in the same area. At the 10-year-of-age time point, sera

were available for 62 out of the 65 children that were originally selected in the Ngerenya

cohort subset for serological analysis. Of these, 20 experienced one or more episodes of

febrile malaria during early childhood prior to the decline in malaria transmission, while 42

remained entirely malaria-free throughout follow-up (**Fig. 5a**). At 10 years of age, Ngerenya

children who had experienced early-life malaria exhibited significantly lower IgG levels to a

wide range of antigens compared to their malaria-naïve peers (**Fig. 5b**). These included

responses to Coxsackievirus B1, CMV, H1N1 influenza, HSV-1 and rubella. Differences in

antibody level to *B. pertussis* and EBV did not reach statistical significance.

Discussion

This study demonstrates that early-life exposure to malaria is associated with broad and

durable impairments in antibody-mediated immunity to unrelated pathogens and vaccines.

By leveraging a natural experiment in coastal Kenya - where malaria transmission diverged

sharply between adjacent communities during the early 2000s - we were able to disentangle

the immunological effects of malaria exposure from confounding geographic and temporal

factors. Our findings reveal that children exposed to malaria in early childhood not only

generate lower antibody titres to non-malarial antigens but maintain these attenuated

responses well into adolescence, long after malaria transmission has ceased. The

attenuating effect of malaria on antibody responses has long been suspected [15] but difficult

to quantify. Prior studies have shown reduced vaccine efficacy in malaria-endemic settings

[3,16], and experimental models have implicated regulatory T cells, atypical memory B cells,

and B cell exhaustion as potential mediators of malaria-induced immune suppression[9,12,13,22].

However, most human studies to date have focused on short-term immune responses or

outcomes in the setting of concurrent parasitaemia. Our data extend this work by

demonstrating that transient exposure to malaria in early childhood is sufficient to imprint

long-lasting changes on the humoral immune repertoire.

The strength of this study lies in its integration of long-term active malaria surveillance with

serial antibody profiling. Children were visited weekly for febrile illness surveillance, allowing

for precise documentation of clinical malaria episodes. In parallel, our use of a validated

protein microarray platform enabled us to track IgG responses to a wide panel of pathogen

and vaccine antigens at multiple timepoints. We first confirmed that the microarray platform

captured biologically relevant responses by comparing antibody trajectories in individual

children with known measles vaccination and malaria exposure histories. The platform

reliably detected both vaccine-induced and infection-associated rises in IgG, providing

confidence in the longitudinal patterns observed. We found that children from Junju, a

region of persistent transmission, had significantly lower antibody levels to multiple

pathogens compared to children from Ngerenya, where malaria transmission declined in the

mid-2000s. These differences were most pronounced for antigens such as measles virus,

Bordetella pertussis, CMV, rubella virus, and HSV-1. Importantly, all children had comparable

vaccination histories and were followed through the same longitudinal infrastructure,

minimizing the likelihood of differential healthcare access or vaccine uptake.

To test whether early-life malaria exposure specifically contributed to long-term immune

suppression, we examined antibody responses within the Ngerenya cohort. Because the

transmission decline occurred rapidly, children born just before and after the inflection point

experienced markedly different levels of malaria exposure while living in the same

geographic area. Among children followed longitudinally, those with even limited early-life

exposure to malaria had significantly lower antibody titers at 10 years of age than their

malaria-naïve peers. This within-cohort contrast strongly implicates early-life infection as the

critical window for immune programming. Taken together, these findings support a model in

which malaria exposure during critical developmental windows modulates long-term

maintenance of immune memory. This may occur through direct effects on B cell

maturation, altered antigen presentation[23], or long-lived changes in lymphoid

microenvironments[24–26]. While the mechanistic basis of this phenomenon remains to be fully

explained, the consistency of the attenuation on multiple antigens and timepoints suggests a
generalizable effect rather than antigen-specific phenomenon.
276
This work has important implications for vaccine policy and infection risk in malaria-endemic
regions. Reduced antibody titres to vaccine-preventable diseases may translate into
diminished long-term protection, even when vaccine coverage is high. Our findings raise the
possibility that children in high-transmission settings may require different immunization
strategies, such as delayed dosing and boosting, to improve vaccine-induced immunity.
Moreover, as malaria control efforts continue to shift disease epidemiology, these data
highlight the value of longitudinal serological surveillance for understanding the broader
immunological legacy of malaria exposure. Our study has several limitations. Although the
microarray platform was validated internally and against ELISA, quantitative comparisons
across platforms are inherently constrained. Additionally, while the sample size for antibody
profiling was modest, the inclusion of dense longitudinal sampling and detailed clinical
surveillance adds depth to the dataset. Finally, we cannot fully exclude the influence of other
infections or environmental exposures that may have differed subtly between subgroups,
although the within-Ngerenya analysis provides strong evidence for malaria as a primary
driver of long-term immune attenuation.
292
In summary, our findings reveal that early-life malaria exposure is associated with long-term
suppression of antibody responses to unrelated pathogens and vaccines. This effect is
detectable many years after infection, and appears to persist even in the absence of ongoing
transmission. As global malaria control efforts continue, understanding the immunological
legacy of childhood malaria may be critical for improving vaccination strategies and
mitigating susceptibility to other infections.

Figures

Figure 1

Figure 2

Figure　3

Figure 4

Figure   5

 Figure captions

Figure 1

Divergent malaria exposure histories in coastal Kenya - Monthly malaria case counts from

active surveillance in two adjacent regions of Kilifi County, Kenya, between 1998 and 2017.

Junju (red) maintained sustained malaria transmission throughout the study period, while

Ngerenya (blue) experienced a rapid collapse in transmission beginning in 2004. Points

represent total cases per month; lines show LOESS-smoothed trends.

Figure 2

AMA1-specific IgG trajectories mirror individual and regional malaria exposure - (A–B)

Longitudinal AMA1 IgG levels in individual children measured by protein microarray. Vertical

red lines denote confirmed febrile malaria episodes. Panel A shows a Junju child with

multiple documented infections; Panel B shows a Ngerenya child who remained malaria-free

throughout follow-up. Each blue spot is a single antibody measurement. Each time point was

measured in quadruplicate **(C)** Mean AMA1-specific IgG levels with 95% confidence intervals

for all children in the microarray subset plotted by year of sampling. Junju children showed

persistently elevated antibodies, while AMA1 antibody levels in Ngerenya declined sharply

after 2004.

Figure 3

Malaria exposure is associated with reduced measles-specific antibody levels. Example

plots of temporal changes in measles-specific antibody in vaccinated and unvaccinated

children are shown **(A)** Longitudinal IgG levels in an individual child measured by microarray.

The dashed vertical red line indicates the timing of the last dose of the routine measles

vaccine. **(B)** Shows a similar temporal trend for a child with no history of measles

vaccination. **(C)** Longitudinal IgG responses to measles virus by cohort, measured by protein

microarray in vaccinated children. Junju children exhibited consistently lower levels of

measles-specific antibody than Ngerenya counterparts. The circles indicate means, and the

whiskers denote 95% confidence intervals. **(C)** Measles-specific IgG levels in 3-year-old

children from Junju and Ngerenya, measured by ELISA in children with a documented history

of measles vaccination. Each dot represents an individual participant.

Figure 4

Broad suppression of antibody responses to diverse pathogens in malaria-exposed children - cross-sectional IgG titres at 10 years of age for a panel of pathogen and vaccine antigens, measured by protein microarray. Each dot represents an individual child; boxes indicate interquartile ranges, with the midline denoting the median. Junju children show significantly lower antibody levels across most antigens compared to Ngerenya children, including *Bordetella pertussis*, H1N1 influenza virus, rubella virus, measles virus, cytomegalovirus (CMV), Epstein-Barr virus (EBV), herpes simplex virus 1 (HSV-1), and coxsackievirus B1.

Figure 5

Early-life malaria exposure predicts long-term suppression of antibody responses within the same geographic region - (A) Active malaria surveillance records for children in the Ngerenya cohort. Each row represents an individual child, and each column represents a surveillance timepoint. Dark red boxes indicate one or more confirmed febrile malaria episodes; light grey boxes indicate surveillance visits without malaria detection. Children are grouped by early-life exposure status (top: malaria-naïve; bottom: previously exposed). **(B)** IgG levels at 10 years of age among Ngerenya children, stratified by early-life malaria exposure. Children with ≥1 confirmed febrile malaria episode during early childhood (n = 20) show significantly lower titres to multiple unrelated pathogens compared to malaria-naïve peers (n = 42). All children lived in the same geographic area and received identical vaccines and follow-up. The black dots are means and error bars are 95% confidence intervals.

Supplementary Figure 1

A summary of the sampling frame for the study cohorts (Junju and Ngerenya). Each vertical line represents the longitudinal time series for a single individual, and each dot represents a timepoint where a serum sample was collected.

Supplementary Table 1

378   Summary of pathogens, strains or subtypes, and corresponding antigens included in the

379   immunoassay panel.

Data and Code Availability

The datasets and R scripts used for the analysis in this study are available from the corresponding author upon reasonable request. Due to local laws and regulations, individual-level data from the Kilifi Health and Demographic Surveillance System (KHDSS), cannot be made publicly available. All data processing and visualization were performed using open-source R packages; package versions and analysis pipelines are available upon request.

Funding statement

This study was supported by fellowship funding to C.J.S. from the Wellcome Trust (WT105882MA). The funder played no role in the conceptualization, design, data collection, analysis, decision to publish, or preparation of the manuscript. M.S.S was funded in whole by Science for Africa Foundation to the Developing Excellence in Leadership, Training, and Science in Africa (DELTAS Africa) program [DEL-22-012] with support from Wellcome Trust and the UK Foreign, Commonwealth & Development Office and is part of the EDCPT2 programme supported by the European Union. For purposes of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

References

1 World Health Organization. World malaria report 2024. Geneva, 2024
https://www.who.int/publications/i/item/9789240086173.

2 Reiner RC, LBD Triple Burden Collaborators, Welgan CA, *et al.* The overlapping burden of
the three leading causes of disability and death in sub-Saharan African children. *Nat*
Commun 2022; **13**: 7457.

3 Bejon P, Mwacharo J, Kai O, *et al.* The induction and persistence of T cell IFN-gamma
responses after vaccination or natural exposure is suppressed by Plasmodium falciparum.
J Immunol 2007; **179**: 4193–201.

4 RTS,S Clinical Trials Partnership, Agnandji ST, Lell B, *et al.* A phase 3 trial of RTS,S/AS01
malaria vaccine in African infants. *N Engl J Med* 2012; **367**: 2284–95.

5 Cunnington AJ, Riley EM. Suppression of vaccine responses by malaria: insignificant or
overlooked? *Expert Rev Vaccines* 2010; **9**: 409–29.

6 P Z, N C. Factors That Influence the Immune Response to Vaccination. *PubMed* 2019.
https://pubmed.ncbi.nlm.nih.gov/30867162/ (accessed April 25, 2025).

7 Bediako Y, Adams R, Reid AJ, *et al.* Repeated clinical malaria episodes are associated with
modification of the immune system in children. *BMC Med* 2019; **17**: 60.

8 Calle CL, Mordmüller B, Singh A. Immunosuppression in Malaria: Do Plasmodium
falciparum Parasites Hijack the Host? *Pathogens* 2021; **10**: 1277.

9 Kurup SP, Obeng-Adjei N, Anthony SM, *et al.* Regulatory T cells impede acute and long-
term immunity to blood-stage malaria through CTLA-4. *Nat Med* 2017; **23**: 1220–5.

10 Walther M, Tongren JE, Andrews L, *et al.* Upregulation of TGF-beta, FOXP3, and
CD4+CD25+ regulatory T cells correlates with more rapid parasite growth in human
malaria infection. *Immunity* 2005; **23**: 287–96.

11 Han X, Yang J, Zhang Y, *et al.* Potential Role for Regulatory B Cells as a Major Source of
Interleukin-10 in Spleen from Plasmodium chabaudi-Infected Mice. *Infect Immun* 2018;
86: e00016-18.

12 Illingworth J, Butler NS, Roetynck S, *et al.* Chronic exposure to Plasmodium falciparum is
associated with phenotypic evidence of B and T cell exhaustion. *J Immunol* 2013; **190**:
1038–47.

13 Weiss GE, Crompton PD, Li S, *et al.* Atypical Memory B Cells Are Greatly Expanded in
Individuals Living in a Malaria-Endemic Area. *The Journal of Immunology* 2009; **183**:
2176–82.

14 Portugal S, Moebius J, Skinner J, *et al.* Exposure-dependent control of malaria-induced
inflammation in children. *PLoS Pathog* 2014; **10**: e1004079.

15 Greenwood BM, Bradley-Moore AM, Bryceson AD, Palit A. Immunosuppression in
children with malaria. *Lancet* 1972; **1**: 169–72.

16 Zirimenya L, Natukunda A, Nassuuna J, *et al.* The Effect of Malaria on Responses to
Unrelated Vaccines in Animals and Humans: A Systematic Review and Meta-Analysis.
Parasite Immunology 2024; **46**: e13067.

17 Crawley J, Sismanidis C, Goodman T, Milligan P. Effect of intermittent preventive
treatment for malaria during infancy on serological responses to measles and other
vaccines used in the Expanded Programme on Immunization: results from five
randomised controlled trials. *The Lancet* 2012; **380**: 1001–10.

18 O'Meara WP, Mwangi TW, Williams TN, McKenzie FE, Snow RW, Marsh K. Relationship
between exposure, clinical malaria, and age in an area of changing transmission intensity.
Am J Trop Med Hyg 2008; **79**: 185–91.

19 Scott JAG, Bauni E, Moisi JC, *et al.* Profile: The Kilifi Health and Demographic Surveillance
System (KHDSS). *Int J Epidemiol* 2012; **41**: 650–7.

20 Mwangi TW, Ross A, Snow RW, Marsh K. Case definitions of clinical malaria under
different transmission conditions in Kilifi District, Kenya. *J Infect Dis* 2005; **191**: 1932–9.

21 Bejon P, Williams TN, Liljander A, *et al.* Stable and unstable malaria hotspots in
longitudinal cohort studies in Kenya. *PLoS Med* 2010; **7**: e1000304.

22 Portugal S, Obeng-Adjei N, Moir S, Crompton PD, Pierce SK. Atypical memory B cells in
human chronic infectious diseases: An interim report. *Cell Immunol* 2017; **321**: 18–25.

23 Yap XZ, Lundie RJ, Beeson JG, O'Keeffe M. Dendritic Cell Responses and Function in
Malaria. *Front Immunol* 2019; **10**: 357.

24 Cadman ET, Abdallah AY, Voisine C, *et al.* Alterations of splenic architecture in malaria are
induced independently of Toll-like receptors 2, 4, and 9 or MyD88 and may affect
antibody affinity. *Infect Immun* 2008; **76**: 3924–31.

25 Obeng-Adjei N, Portugal S, Holla P, *et al.* Malaria-induced interferon-γ drives the
expansion of Tbethi atypical memory B cells. *PLoS Pathog* 2017; **13**: e1006576.

26 Weiss GE, Traore B, Kayentao K, *et al.* The Plasmodium falciparum-specific human
memory B cell compartment expands gradually with repeated malaria infections. *PLoS*
Pathog 2010; **6**: e1000912.

Supplementary Materials

470   Supplementary figure 1

Pathogen	Strain/subtype	Antigen
Measles virus	Edmonston	Whole virus antigen
Coxsackie B virus	B1	VP1
Cytomegalovirus	TB40E	PP150
Epstein-Barr virus	Type 1	Nuclear antigen 1
Herpes simplex virus	Type 1	Extract
H1N1 Influenza A	A/California/07/2009	Haemagluttinin
Plasmodium falciparum	3D7	AMA-1
Rubella virus	HPV-77	Whole virus antigen

476   Supplementary table 1