1

Dharma: A novel machine learning framework for pediatric appendicitis—diagnosis, severity assessment and evidence-based clinical decision support.

Anup Thapa[1*], Subash Pahari[2], Shashank Timilsina[1], Binay Chapagain[3]

[1]Department of Emergency Medicine and General Practice, Matri-Sishu Miteri Provincial

Hospital, Pokhara, Gandaki, Nepal.

[2]Department of Computer Engineering, Faculty of Science and Technology, Pokhara

University, Pokhara, Gandaki, Nepal.

[3]Department of Critical Care Medicine, Tribhuvan University Teaching Hospital,

Kathmandu, Bagmati, Nepal.

Corresponding author:

E-mail: ajungthapa11@gmail.com (AT)

Abstract

Background: Acute appendicitis remains a challenging diagnosis in pediatric populations, with high rates of misdiagnosis and negative appendectomies despite advances in imaging modalities. Current diagnostic tools, including clinical scoring systems like Alvarado and Pediatric Appendicitis Score (PAS), lack sufficient sensitivity and specificity, while reliance on CT scans raises concerns about radiation exposure, contrast hazards and sedation in children. Moreover, no established tool effectively predicts progression from uncomplicated to complicated appendicitis, creating a critical gap in clinical decision-making.

Objective: To develop and evaluate a machine learning model that integrates clinical, laboratory, and radiological findings for accurate diagnosis and complication prediction in pediatric appendicitis and to deploy this model as an interpretable web-based tool for clinical decision support.

Methods: We analyzed data from 780 pediatric patients (ages 0-18) with suspected appendicitis admitted to Children's Hospital St. Hedwig, Regensburg, between 2016 and 2021. For severity prediction, our dataset was augmented with 430 additional cases from published literature and only the confirmed cases of acute appendicitis(n=602) were used. After feature selection using statistical methods and recursive feature elimination, we developed a Random Forest model named

Dharma, optimized through hyperparameter tuning and cross-validation. Model performance was

evaluated on independent test sets and compared with conventional diagnostic tools.

Results: Dharma demonstrated superior diagnostic performance with an AUC-ROC of 0.96 (±0.02

SD) in cross-validation and 0.97-0.98 on independent test sets. At an optimal threshold of 64%,

the model achieved specificity of 88%-98%, sensitivity of 89%-95%, and positive predictive value

of 93%-99%. For complication prediction, Dharma attained a sensitivity of 93% (±0.05 SD) in

cross-validation and 96% on the test set, with a negative predictive value of 98%. The model

maintained strong performance even in cases where the appendix could not be visualized on

ultrasonography (AUC-ROC 0.95, sensitivity 89%, specificity 87% at the threshold of 30%).

Conclusion: Dharma is a novel, interpretable machine learning based clinical decision support tool

designed to address the diagnostic challenges of pediatric appendicitis by integrating easily

obtainable clinical, laboratory, and radiological data into a unified, real-time predictive framework.

Unlike traditional scoring systems and imaging modalities, which may lack specificity or raise

safety concerns in children, Dharma demonstrates high accuracy in diagnosing appendicitis and

predicting progression from uncomplicated to complicated cases, potentially reducing unnecessary

surgeries and CT scans. Its robust performance, even with incomplete imaging data, underscores

its utility in resource-limited settings. Delivered through an intuitive, transparent, and interpretable

web application, Dharma supports frontline providers—particularly in low- and middle-income

settings—in making timely, evidence-based decisions, streamlining patient referrals, and

improving clinical outcomes. By bridging critical gaps in current diagnostic and prognostic tools,

Dharma offers a practical and accessible 21st-century solution tailored to real-world pediatric

surgical care across diverse healthcare contexts. Furthermore, the underlying framework and

concepts of Dharma may be adaptable to other clinical challenges beyond pediatric appendicitis,

providing a foundation for broader applications of machine learning in healthcare.

Keywords: pediatric appendicitis, machine learning, clinical decision support system, evidence-

based medicine, artificial intelligence, Dharma, Novel

Author Summary

Accurate diagnosis of pediatric appendicitis remains challenging, with current clinical scores and

imaging tests limited by sensitivity, specificity, predictive values, and safety concerns. We

developed Dharma, an interpretable machine learning model that integrates clinical, laboratory,

and radiological data to assist in diagnosing appendicitis and predicting its severity in children.

Evaluated on a large dataset supplemented by published cases, Dharma demonstrated strong

diagnostic and prognostic performance, including in cases with incomplete imaging—making it

potentially especially useful in resource-limited settings for early decision-making and streamlined

referrals. Available as a web-based tool, it provides real-time support to healthcare providers in

making evidence-based decisions that could reduce negative appendectomies while avoiding

hazards associated with advanced imaging modalities such as sedation, contrast, or radiation

exposure.

Furthermore, the open-access concepts and framework underlying Dharma have the potential to

address diverse healthcare challenges beyond pediatric appendicitis.

Background

Acute abdominal pain (AAP) is a common complaint of Pediatric Patients which accounts for up

to 10 % of total visits to the Emergency Department while acute appendicitis is the major cause of

AAP in children over the age of 1 [1,2]. Despite being a routinely encountered clinical dilemma,

diagnosis of acute appendicitis still poses a major challenge for the attending healthcare

professionals who use patterns of signs, symptoms, laboratory values and radiological findings to

make an informed decision, probably, due to the variability in clinical presentation and the frequent

absence of classic symptoms in pediatric population. Different scoring systems, with Alvarado

being the oldest and the most common, have been incorporated in the workflow to help the

physicians with quantifying the risk of child having acute appendicitis. However, none of the

scoring systems provide sufficient sensitivity, specificity or predictive values to be used as an

exclusive standard in setting the diagnosis. [3-5].

The increasing reliance on CT and USG for evaluating acute abdominal pain has not been

accompanied by a corresponding improvement in appendicitis detection rates [19]. Misdiagnosis

rates remain alarmingly high, ranging from 70% to 100% in children aged three years or younger,

with a gradual decrease as age increases [21]. This diagnostic challenge is often attributed to the

atypical presentation of the disease in younger children and the anatomical variability of the

diameter of the appendix. Consequently, the lack of a reliable diagnostic tool may contribute to

the persistently high incidence of Negative Appendectomy (NA) worldwide, reported between 5%

and 20% in various centers [7-10]. NA is a dreaded outcome for any operating surgeon due to the

inherent risks associated with general anesthesia and surgery, particularly in the pediatric

population.

Moreover, no traditional inflammatory biomarkers, clinical signs and symptoms, or imaging

modalities can reliably predict the progression of uncomplicated to complicated appendicitis

[12,42]. This is particularly concerning in pediatric patients, where the incidence of perforated

appendix is alarmingly high and increases as the age decreases, reaching over 80% in children

younger than three years old [22]. This highlights the urgent need to refine diagnostic and

prognostic strategies for pediatric appendicitis.

Machine Learning (ML), a subfield of Artificial Intelligence (AI), involves analyzing data to

identify patterns and applying that information to make predictions in similar future scenarios [38],

thus, offers a promising avenue for improving diagnostic and prognostic precision, particularly in

pediatric patients, by integrating clinical, laboratory, and imaging data into a comprehensive

predictive framework.

ML has been applied to assist in the diagnosis of acute appendicitis since the 1990s [27], with

models specifically targeting pediatric patients emerging in the early 2010s [28]. These models

aim to predict pediatric appendicitis, its complications, and postoperative recovery outcomes

[11,24-31]. Despite the development of numerous ML-based tools, none have undergone

independent validation, highlighting a critical gap in their clinical applicability.

ML-based tools have the potential to enhance the accuracy of appendicitis diagnosis and severity

prediction [39], thereby reducing unnecessary radiation exposure from CT scans, minimizing

negative appendectomy rates, and optimizing referral patterns from primary care settings. An

effective ML tool should provide reliable predictions even in cases with missing data, ensuring

applicability across varied clinical environments, from well-resourced urban hospitals to resource-

limited rural settings. Consequently, there is a pressing need for a transparent, interpretable,

adaptable and easy-to-use diagnostic tool that can reliably predict or rule out appendicitis and its

complications in pediatric patients.

Thus, we defined two primary objectives for this study:

● To develop and evaluate an optimized, generalizable machine learning model for the

136   diagnosis and complication prediction of pediatric appendicitis, and to compare its

137   performance with conventional diagnostic tools, including the Alvarado Score (AS),

138   Pediatric Appendicitis Score (PAS), and Ultrasonography (USG).

140   ● To deploy the model as an interpretable, web-based diagnostic and prognostic tool to

141   support evidence-based clinical decision-making in the management of suspected

142   appendicitis cases.

Materials And Methods

Data Curation

146 This study utilizes data from a cohort of pediatric patients with suspected appendicitis who were

147 admitted with abdominal pain to Children's Hospital St. Hedwig in Regensburg, Germany,

148 between 2016 and 2021. The original data collection was approved by the Ethics Committee of

149 the University of Regensburg (approval numbers: 18-1063-101, 18-1063_1-101, and 18-1063_2-

150 101). The dataset has since been anonymized and is now publicly available for secondary use [44].

Data Description And Preprocessing

This study utilizes a dataset comprising 780 suspected pediatric appendicitis cases with 55

independent variables and three dependent variables: Diagnosis, Treatment Modality and Severity.

● Diagnosis: appendicitis(n=463) and no appendicitis(n=317)

● Severity: complications(n=119) and no complications or no appendicitis(n=661)

● Treatment Modality: Conservative(n=483) and Surgical(n=297)

The dependent variables with more than 30% missing data were excluded from the study, except

for appendix diameter. Despite a substantial amount of missing data, this variable was retained

owing to the known challenges associated with appendix visualization through ultrasonography

(USG). Moreover, the diameter of the appendix holds well-established clinical significance in the

pathophysiology and diagnosis of acute appendicitis.

Binary categorical variables were mapped as 0 and 1 while the ordinal categorical variables were

mapped as 0,1 ,2 and 3. The missing values were imputed as 0 to simulate the real-world difficulty

in obtaining complete medical data, especially in resource- limited settings.

For the primary classification task (diagnosis), the full dataset comprising 780 cases was used. For

the complication prediction task, the dataset was augmented by integrating 430 suspected

appendicitis cases from the dataset published by Marcinkevics et al. [11]. After removing the

duplicates, the combined dataset consisted of 1,015 unique patient records. Only cases with a

confirmed diagnosis of appendicitis (n = 602) were included for training and testing the severity

assessment model. Treatment modality was not analyzed in the study directly, as we believe, it

should be determined by the operating surgeon based on the diagnosis and the anticipated risk of

complications in the diagnosed cases.

Feature Selection

The Chi-Square test was performed to assess the association between categorical variables and
disease state, while Cramér's V measured the strength of these associations. Variables with a
significant association ($p < 0.001$) and a moderate to strong effect size (Cramér's V > 0.10) were
selected for the development of the model.

For continuous variables, the Mann-Whitney U test was used to compare distributions between
positive and negative classes, and rank-biserial correlation measured the strength of association.
Features with a significant difference ($p < 0.001$) and a correlation coefficient > 0.40 were included
in the model.

All statistical analyses of features with the targets were conducted using Python, employing
libraries: Pandas, NumPy, and SciPy, on the original dataset comprising 780 cases of suspected
appendicitis.

RFECV was used to identify the parameters with the greatest influence over the decision-making
process. Different subsets of selected parameters were then tested independently to select the
optimum combination and arrangement of features for the model.

Data mining was done independently for both the targets.

Model Development

The Random Forest model was selected for its suitability in handling non-parametric medical data.

For diagnosis prediction, the dataset was split into training, validation, and test sets in a 3:1:1 ratio.

The model was optimized through hyperparameter tuning using grid search, followed by

randomized search and stratified cross-validation, ensuring robust performance. To mitigate

overfitting and ensure consistent performance across diverse clinical presentations, 10-fold cross-

validation was performed on the training set.

The optimal classification threshold was determined using Youden's Index and the Euclidean

Distance method on the Receiver Operating Characteristic (ROC) curve applied to the validation

set. We aimed to strike a balance between sensitivity and specificity to minimize both false

positives and false negatives in appendicitis diagnosis.

We named the final random forest model **Dharma** after selecting the optimal hyperparameters and

decision threshold. The model's performance was subsequently evaluated on the independent test

set using sensitivity, specificity, AUC-ROC, and other relevant metrics to ensure an unbiased

assessment.

Additionally, we curated an independent test set of 218 cases by removing duplicates from the

dataset of 430 cases used by Marcinkevics et al [11]. Dharma's performance was further evaluated

on this fresh set. The performance of commonly used pediatric appendicitis diagnostic tools was

also evaluated on this dataset.

For severity prediction, we combined our dataset with a cohort of 430 suspected appendicitis cases

from Marcinkevics et al [11] and removed the duplicates, resulting in a combined dataset (n =

1015). Only the confirmed appendicitis cases (n = 602) were included for model development and

testing.

The complication prediction dataset was highly imbalanced, with a positive-to-negative class ratio

of 148:454. To address this, we applied a combination of oversampling and undersampling

techniques—using SMOTE to increase the positive instances and NearMiss (n=1) to reduce the

negative class—after splitting the dataset into a temporary set (adjusted from 122:359 to 359:179)

and a test set (26:95). The temporary set was subsequently divided into a training set (262:141)

and a validation set (97:38). The test set remained unaltered to preserve the original class

distribution and ensure an unbiased performance evaluation. We prioritized optimizing Dharma's

sensitivity at the expense of specificity to quantify the risk of possible complications. This trade-

off was considered appropriate, as missing true positives could result in severe, life-threatening

outcomes like gangrenous or perforated appendicitis, whereas false positives are more acceptable

given the patient is already a diagnosed case of acute appendicitis and can be managed surgically.

Model development was conducted using Scikit-learn in Python.

Explainability Analysis

To understand the decision-making process of Dharma, we analyzed the model using SHAP's

Tree Explainer (SHapley Additive exPlanations Tree Explainer), a game theory-based approach

that assigns contribution values (SHAP values) to each feature by comparing them to a baseline

prediction before any data is introduced (base SHAP value). Each feature's SHAP value

quantifies its influence on shifting the probability of a positive outcome either towards or away

from this baseline.

We used SHAP's summary plot method to visualize these contributions and provide a global

interpretation of how Dharma assessed suspected appendicitis cases for both diagnosis and

severity prediction tasks in our test set. Additionally, we integrated SHAP values to explain

individual predictions in our web app, along with the 95% confidence interval of predictions

from each decision tree in our random forest providing an additional layer of uncertainty

quantification. This integration is intended to assist attending physicians in exploring the

reasoning behind the model's predictions thus enhancing transparency, interpretability and trust.

Results

Data mining

A total of 780 pediatric patients (ages 0–18 years; mean age: 11.35 years, 95% CI: 11.10–11.59) with suspected appendicitis were included in the diagnostic prediction cohort, with a male-to-female ratio of 1.07 (403:377). For complication prediction, 602 confirmed appendicitis cases from the combined dataset, as described earlier, were included.

A Chi-square test was performed to assess the significance of associations, and Cramér's V was used to evaluate the strength of association of the categorical features. Seven features showed significant association with appendicitis diagnosis and six features were significantly associated with complication prediction ($p < 0.001$). Among the diagnostic features, peritonitis demonstrated the strongest association (Cramér's V = 0.37), followed by neutrophilia (Cramér's V = 0.23). For complication prediction, peritonitis again showed the strongest association (Cramér's V = 0.30), followed by urinary ketones (Cramér's V = 0.29), Table 1 and 2..

Table 1. Diagnosis and Categorical Variables (n=780)				
Features	Chi-Square	P-value	Cramer's V	Samples
Peritonitis	108.24	<0.001	0.37	773

Neutrophilia	67.17	<0.001	0.30	732
Free Fluids	32.48	<0.001	0.21	719
Ketones in Urine	24.34	<0.001	0.20	582
Contralateral Rebound Tenderness	25.08	<0.001	0.18	767
Nausea/Vomiting	23.19	<0.001	0.17	774
Loss of Appetite	19.24	<0.001	0.16	772
Ipsilateral Rebound Tenderness	8.64	<0.01	0.12	619
Sex	10.78	<0.01	0.12	779
RIF Tenderness	8.18	<0.01	0.10	774
Migratory Pain	6.70	0.01	0.09	773
Coughing Pain	6.30	0.01	0.09	766
Psoas Sign	4.00	0.05	0.07	745
Stool	3.04	0.22	0.06	765
RBC in Urine	2.02	0.57	0.06	576
WBC in Urine	1.25	0.74	0.05	583
Dysuria	1.61	0.20	0.05	753

Table 2. Complications and Categorical variables(n=463)				
Features	**Chi-Square**	**P-value**	**Cramer's V**	**Samples**
Peritonitis	41.29	<0.001	0.3	457

Ketones in Urine	26.45	<0.001	0.29	318
Neutrophilia	23.75	<0.001	0.23	434
Nausea/Vomiting	23.17	<0.001	0.22	458
Loss of Appetite	20.56	<0.001	0.21	456
Free Fluids	16.72	<0.001	0.2	430
RBC in Urine	10.65	0.01	0.18	317
Ipsilateral Rebound Tenderness	5.62	<0.05	0.14	305
WBC In Urine	4.9	0.18	0.12	319
Stool	5.12	0.08	0.11	451
RIF Tenderness	2.66	0.1	0.08	458
Psoas Sign	1.58	0.21	0.06	433
Contralateral Rebound Tenderness	1.5	0.22	0.06	451
Migratory Pain	0.59	0.44	0.04	457
Dysuria	0.54	0.46	0.04	437
Sex	0.54	0.46	0.03	462
Coughing Pain	0.01	0.94	0.004	450

277 A Mann-Whitney U test was conducted to assess the statistical significance of differences in the

278 distributions of continuous variables between positive and negative cases, and rank-biserial

279 correlation was used to evaluate the effect size of these differences Table 3 and 4. Six variables

were significantly associated with appendicitis diagnosis, and five variables were significantly

associated with complication prediction (p < 0.001). Among the diagnostic variables, appendix

diameter demonstrated the strongest effect (rank-biserial correlation = 0.9), followed by C-reactive

protein (CRP) level (rank-biserial correlation = 0.46). For complication prediction, CRP level

exhibited the strongest effect (rank-biserial correlation = 0.64), followed by white blood cell

(WBC) count (rank-biserial correlation = 0.45).

Table 3. Diagnosis and Continuous variables (n=780)						
Feature	U_statistic	P_value	Rank-Biserial correlation	Mean (Positive Class)	Mean(Negative Class)	Samples
Appendix Diameter	2341	<0.001	0.9	8.7	5.04	498
CRP	38686	<0.001	0.46	44.9	11.72	771
WBC Count	39910	<0.001	0.45	14.28	10.33	776
Neutrophil Percentage	32518.5	<0.001	0.42	76.03	65.6	679
Body Temperature	57410	<0.001	0.21	37.52	37.24	775
BMI	79280	<0.001	-0.15	18.45	19.56	753
Weight	82617	<0.01	-0.13	41.72	45.25	778
Age	80879	0.02	-0.1	11.08	11.72	780
Height	74276.5	0.08	-0.07	146.93	149.51	755
RDW	65536.5	0.21	0.05	13.4	12.87	756
Hemoglobin	72577.5	0.52	-0.03	13.38	13.38	764

RBC Count	70347	0.92	0.004	4.79	4.82	764
Thrombocyte Count	70459.5	0.95	0.003	285.79	284.48	764

Table 4. Complications and Continuous variables(n=463)						
Feature	**U_statistic**	**P_value**	**Rank-Biserial Correlation**	**Mean (Positive Class)**	**Mean(Negative Class)**	**Samples**
CRP	6952.5	<0.001	0.64	107.68	24.52	457
WBC Count	10960	<0.001	0.45	17.43	13.23	460
Body Temperature	12010.5	<0.001	0.4	37.93	37.38	459
Neutrophil Percentage	8803.5	<0.001	0.38	82.44	74.14	403
Appendix Diameter	7296	<0.001	0.28	9.53	8.51	371
Weight	24201.5	<0.01	-0.2	38.73	42.75	461
Age	24327	<0.01	-0.2	10.08	11.42	463
Height	21988.5	<0.01	-0.17	141.56	148.79	443
BMI	21284.5	0.01	-0.16	17.92	18.64	441
Hemoglobin	21018.5	0.06	-0.12	13.36	13.39	450
Thrombocyte Count	17298	0.2	0.08	296.65	282.24	450

| RBC Count | 20250.5 | 0.23 | -0.08 | 4.75 | 4.8 | 450 |
| RDW | 17530 | 0.44 | 0.05 | 13.56 | 13.35 | 445 |

Performance metrics

The Alvarado Score (AS) and Pediatric Appendicitis Score (PAS) were significantly associated

with appendicitis diagnosis (p < 0.001), demonstrating AUC-ROC values of 0.78 and 0.72,

respectively. The median scores for positive and negative cases were 7 vs. 5 for AS, and 6 vs. 4

for PAS, indicating higher scoring trends among the confirmed cases. Ultrasonography (USG)

exhibited the highest diagnostic performance among the available tools, with an AUC-ROC of

0.93. However, appendix diameter measurements were unavailable in 34% of suspected cases,

likely due to the inherent challenges in appendix visualization. A summary of diagnostic

associations and performance metrics of these conventional tools are presented in Table 5 and 6.

Table 5. Association of Diagnostic Tools					
Scoring Systems	U Statistic	P-value	Rank-Biserial Correlation	Median(Positive Class)	Median(Negative Class)
Alvarado Score	33163.5	<0.001	0.48	7	5

Pediatric Appendicitis Score	38445.5	<0.001	0.4	6	4

Table 6. Performance of Diagnostic tools					
Tools	**AUC-ROC**	**Sensitivity**	**Specificity**	**PPV**	**NPV**	**Accuracy**
Alvarado Score	0.78	91%	45%	71%	77%	72%
Pediatric Appendicitis Score	0.72	48%	76%	74%	50%	59%
Ultrasonography	0.93	93%	94%	98%	79%	93%

Dharma demonstrated high diagnostic and prognostic performance in predicting appendicitis and its complications. The model achieved a diagnostic AUC-ROC of 0.96 (±0.02 SD) in 10-fold cross-validation. On the split and curated test sets, the AUC-ROC, specificity, positive predictive value (PPV), sensitivity, and negative predictive value (NPV) of |0.97-0.98|88% -98%|93%- 99% |89%- 95%| and |86%-91%| was achieved.

For complication prediction, Dharma achieved a sensitivity of 0.93 (±0.05 SD) in 10-fold cross-validation. On the unaltered test set, the sensitivity and NPV were 96% and 98%, respectively.

The optimal hyperparameters and decision thresholds for both diagnosis and complication prediction tasks are detailed in Table 7 and 8, respectively.

Table 7. Hyperparameters and Threshold for Diagnosis	
n_estimators	555
min_samples_split	12
min_samples_leaf	1
max_depth	35
class_weight	"balanced"
random_state	17
Threshold	0.64
Threshold(appendix not visualized)	0.3

Table 8. Hyperparameters and Threshold for Complication Prediction	
n_estimators	555
min_samples_split	3
min_samples_leaf	1
max_depth	33
random_state	17
Threshold	0.52

320 Performance metrics and comparative evaluations for diagnostic and severity prediction are

321 summarized in Table 9 and 10. The list of features and their arrangement is listed in Table 11.

Table 9. Dharma's Diagnostic Performance						
Dharma	**AUC-ROC**	**Specificity**	**PPV**	**Sensitivity**	**NPV**	**Accuracy**
10-fold cross- validation	0.96±0.02	-	-	92%±0.02	-	91%±0.03
Split Test Set(n=156)	0.98	88%	93%	95%	91%	93%
Curated Test Set(n=216)	0.97	98%	99%	89%	86%	93%
Appendix not visualized(n=127)	0.95	87%	74%	90%	95%	87%

Table 10. Dharma's Performance for Severity Prediction					
Dharma	**Sensitivity**	**NPV**	**AUC-ROC**	**Specificity**	**PPV**
10-fold cross-validation	93±0.05	-	-	-	-
Unaltered Test Set	96%	98%	0.88	64%	42%

Table 11.List of features and their arrangement	
Diagnosis	**Complications**
Nausea	Nausea
Appetite Loss	Appetite Loss
Peritonitis	Peritonitis
WBC Count	Urinary Ketones
Neutrophil_Percentage	Free Fluids
CRP	CRP
Urinary Ketones	WBC Count
Appendix Diameter	Body Temperature
Free Fluids	Appendix Diameter

SHAP Analysis

329 The SHAP (Shapley Additive Explanations) analysis revealed the following insights about the

330 Dharma model's predictions:

Baseline (Base) Values

332 For the diagnosis task, the base value (i.e., the average model output before any feature input) was

333 0.499 and for the severity assessment task, the base value was 0.649.

Feature Importance

In the diagnostic predictions, the appendix diameter had the largest impact, followed by the white blood cell (WBC) count, presence of peritonitis, and C-reactive protein (CRP) levels.

In the complications prediction task, CRP exhibited the highest influence, followed by WBC count, appendix diameter, body temperature, and appetite loss. Ineterestingly, high values of appendix diameter showed both positive and negative effects on the model's complications risk assessment, indicating a complex and context-dependent relationship. This observation warrants further exploration in future studies.

Absence of Appendix diameter measurement

Notably, the absence of appendix diameter data decreased the model's probability of a positive diagnosis prediction from the base value. Subsequent inputs of other features then pushed the prediction upward to arrive at the final output.

Visual Summary

The SHAP summary dot plots illustrating the global feature importance for the diagnosis and severity prediction tasks are presented in S1 and S2 Figs. for the diagnosis task and in S3 Fig. for the severity assessment task.

S1 Fig. SHAP dot plot for diagnosis prediction

353 **S2 Fig. SHAP dot plot for diagnosis prediction(cases with no appendix diameter data)**

354 **S3 Fig. SHAP dot plot for severity assessment**

Discussion

In our study, the Alvarado Score (AS) and Pediatric Appendicitis Score (PAS) demonstrated

extremely weak abilities to differentiate between positive and negative cases, with AUC-ROC

values of 0.75 and 0.71 respectively. As expected, the median AS and PAS scores were

significantly higher among confirmed positive cases compared to negative ones (7 vs 5 for AS and

6 vs 4 for PAS) [p<0.001]. The extremely low specificity of these systems, 45% and 76% for AS

and PAS respectively, as also reported by Sefa Sağ et al. [6] (14.3%-57.1%), contributes to a high

rate of false-positive diagnoses, ultimately leading to unnecessary negative appendectomies (NA).

Ultrasonography (USG) demonstrated strong diagnostic performance in , with an AUC-ROC of

0.94 and using a 6 mm appendix diameter cutoff, sensitivity and specificity of 93%, and 94%

respectively. However, the appendix was not visualized in 35% of the cases. This limitation aligns

with findings by Duncan et al. [40] and Jenny et al. [41], who reported an inability to visualize the

appendix in 71%-76% of the suspected appendicitis cases. The failure to visualize the appendix

may be attributed to variable anatomical positioning, patient-specific anthropometric differences,

or the operator-dependent nature of USG [13]. Moreover, advanced findings, other than the

diameter of the appendix like peri-appendiceal fluid collection (PALC), peri-appendiceal fat

stranding, target sign, hyperemia on color doppler, free fluids can only be reliably estimated by

an experienced radiologist, which may not be easily available in resource-limited primary care

settings. Additionally, USG findings should also be correlated with clinical and laboratory findings

to formulate a complete diagnosis [43].

Although advanced imaging modalities like CT and MRI provide high diagnostic accuracy for

appendicitis [14-16,18], the risks associated with sedation, contrast use, and high-dose ionizing

radiation [17] make these modalities less favorable for pediatric populations. Moreover, these

modern imaging tools are often inaccessible in the primary and secondary healthcare settings,

particularly in resource-limited regions like ours. Furthermore, like other diagnostic tools, they do

not effectively distinguish between uncomplicated and complicated appendicitis, which is crucial

for guiding optimal management [20].

Significant progress has been made in applying machine learning to the diagnosis, severity

stratification, and management of pediatric appendicitis [11,24–27]. However, many of these

studies are technically complex and AI-centric, making their outputs difficult for clinicians to

interpret and integrate into practice. Unlike traditional scoring systems, these machine learning

models also require a dedicated, easily accessible interface to be of use in real-life clinical

scenarios. For instance, the image-based approach by Byun et al. [26] is computationally intensive

and difficult to implement in real-time clinical settings. While Marcinkevics et al. [11] attempted

to improve accessibility through an online tool, their model relies heavily on complex

ultrasonographic features that require expert radiological interpretation. Moreover, it includes

variables that are not directly involved in the model's decision-making process, limiting its utility

in primary care or resource-constrained emergency departments. Consequently, this tool remains

better suited for academic research than for frontline clinical use.

To address the diagnostic challenges associated with pediatric appendicitis, we present a novel

Machine Learning framework, Dharma, designed to guide attending physicians in making optimal

management decisions based on diagnosis and anticipated risk of complications. Dharma assesses

suspected cases of acute appendicitis by analyzing a comprehensive pattern of clinical features—

mirroring the approach of an attending pediatric surgeon. These include symptoms such as nausea

or vomiting and loss of appetite; signs like peritonitis; physiological parameters such as body

temperature; laboratory findings including white blood cell (WBC) count, percentage of

neutrophils, C-reactive protein (CRP), and urinary ketones; as well as radiological findings such

as appendiceal diameter and the presence of free fluid on ultrasonography. These features are

already well-established in existing literature for the diagnosis and risk stratification of acute

appendicitis [32-37].

In our study, Dharma achieved consistently high diagnostic performance, with AUC-ROC values

of 0.96 ± 0.02, 0.98, and 0.97 across 10-fold cross-validation and two independent test sets. This

demonstrates a robust discriminatory ability to differentiate between positive and negative

appendicitis cases, regardless of threshold. Furthermore, this consistent performance of Dharma

across varied clinical presentations underscores its generalizability and strong pattern recognition

capabilities.

At a threshold of 64%, Dharma achieved a high specificity and positive predictive value (PPV) of

88%-98% and 93%-99%, respectively, alongside an equally strong sensitivity of 89%-95%. This

balance, also supported by the base SHAP value of 0.499, makes Dharma an ideal diagnostic tool,

minimizing false positives and reducing the rate of negative appendectomies (NA) while

maintaining an extremely low risk of false negatives. Such performance characteristics allow

Dharma to provide reliable objective evidence to support clinical suspicion of acute appendicitis,

ideally following ultrasonography and before exposing pediatric patients to the potential hazards

of CT imaging and sedation. This model offers a viable strategy for reducing CT utilization,

particularly in settings where its use is limited by accessibility or concerns about radiation sedation,

contrast or radiation hazards.

Even in the test cases where the appendix measurements were unavailable, Dharma demonstrated

strong diagnostic performance, achieving an AUC-ROC of 0.95, misclassifying only 16 out of 127

cases (12.5%). High sensitivity and specificity of 89% and 87% respectively, at a threshold of

30%, makes it a reliable tool for affirming clinical suspicion of a positive diseased state even with

missing appendix diameter value. Additionally, the excellent negative predictive value (NPV) of

95% underscores the model's ability in ruling-out appendicitis and prompting consideration of

alternative diagnoses. In this context, Dharma may be particularly beneficial in resource-limited

rural and semi-urban primary-to-secondary care settings of low- and middle-income

countries(LMICs), supporting early decision-making and guiding referrals. It could also prove

valuable in oversaturated tertiary-care emergency departments (EDs) of LMICs, offering timely

support to clinical assessments even when USG is unavailable or yields inconclusive results.

Additionally, even in well-equipped centers, Dharma could assist in evaluating probable cases with

ambiguous imaging findings. This usability of Dharma in diverse clinical scenarios makes it a

reliable clinical decision-support tool, in both resource-limited and resource-sufficient settings.

In addition to diagnosis, Dharma is trained to estimate the risk of progression from uncomplicated

to complicated appendicitis. With a sensitivity of 0.93 (±0.05 SD) and 96% in cross-validation and

test sets, along with a high negative predictive value (NPV) of 98%, Dharma can reliably exclude

complications in confirmed appendicitis cases. This feature equips attending surgeons with

evidence-based guidance for deciding between surgical and conservative management, optimizing

patient outcomes and potentially reducing unnecessary interventions in uncomplicated cases [45].

Similarly, in centers with established surgical management guidelines, Dharma's risk stratification

ability through severity assessment may prove useful in scheduling emergency operation theatres

as per the urgency of need of surgical care.

The severity assessment dataset (n=602) exhibited significant class imbalance, with a heavy

predominance of uncomplicated cases (75%). Training the model on this asymmetry introduced a

bias towards negative predictions, increasing the risk of false negatives—i.e., missing possible

complications. To address this, we applied a combination of oversampling using the Synthetic

Minority Oversampling Technique (SMOTE) and undersampling via the NearMiss (n=1)

algorithm, adjusting the class distribution of the combined training and validation sets from

122:359 to 359:179 in favor of the positive (complicated) class. Training Dharma on this

synthetically generated imbalance towards the positive class, reversed its bias, improving the

model's ability to identify almost all the possible complicated cases. The base SHAP value of 0.65

for positive-class confirms this bias adjustment. By prioritizing sensitivity, the model minimizes

false negative instances and effectively flags almost all of the potentially complicated cases,

making it an appropriate screening tool for severity grading in confirmed appendicitis patients.

While the assessment of signs and symptoms can be particularly challenging in younger children,

certain features remain comparatively more accessible. History-based symptoms such as

nausea/vomiting and poor feeding are generally easier to elicit, even in preverbal children through

caregiver observations. Similarly, signs of peritonitis can often be assessed using surrogate

physical findings, including tenderness, rebound tenderness, guarding, and rigidity—either

localized to the right iliac fossa (RIF) or generalized across the abdomen. Radiological parameters

such as appendiceal diameter and the presence of peritoneal free fluid are also reliably assessable

in most pediatric patients, even with screening ultrasonography or Point-Of-Care Ultrasound

(POCUS), making them particularly valuable in settings where advanced imaging modalities or

experienced radiologists may be unavailable or limited.

Dharma addresses the limitations of similar machine learning models in existing literature by

providing not only superior diagnostic and prognostic performance [11,24–31], but also a practical,

interpretable, and easily accessible web-based decision support tool tailored for real-world clinical

environments. Our web app takes patient features, analyzes the case, and provides a Dharma

Score—calculated as the mean probability of a positive disease state across all 555 decision trees

in Dharma's diagnostic random forest. For severity assessment, the risk of complications is

similarly estimated by the trained 555 prognostic trees. In addition, the app provides the probability

range within a 95% confidence interval, along with SHAP values for each feature to explain the

prediction confidence and the reasoning behind the model's decisions. This prevents the models

from being mere "black boxes" and ensures that their results are explained in a way that is easy to

understand and interpret by any attending physician.

Nevertheless, our study had certain limitations that should be acknowledged. A major limitation

was the lack of access to reliable, high-quality medical datasets, which significantly hampers

progress in leveraging statistical and machine-learning tools—tools we view as critical for

advancing 21st-century medicine [23]. We strongly advocate for the open sharing of anonymized

medical datasets to support broader, more impactful research. Additionally, as our model was

developed using single-center data, its adaptability to other populations is unknown. The absence

of exclusively histologically confirmed appendicitis cases may have constrained the model's

ability to optimize pattern recognition for both diagnostic and prognostic tasks. Future work will

prioritize multicenter validation—both prospective and retrospective—including studies

conducted in low- and middle-income countries (LMICs) to rigorously evaluate the real-world

clinical utility and generalizability of Dharma across diverse healthcare settings. We also

encourage fellow researchers and med-tech enthusiasts to conduct independent validation studies

on Dharma.

Conclusion

In this study, we developed Dharma, a novel machine learning–based clinical decision support tool

that integrates readily available clinical, laboratory, and radiological findings into a unified

predictive framework. By offering real-time, evidence-based diagnostic and prognostic assistance

through an easily accessible and interpretable web-based tool, Dharma aims to reduce diagnostic

uncertainty and facilitate timely surgical decision-making in pediatric appendicitis. With its high

diagnostic specificity and positive predictive value demonstrated during testing, Dharma may also

help reduce the rate of negative appendectomies—an important clinical and economic

consideration in pediatric surgical care—without the hazards associated with CT or sedation.

Furthermore, Dharma provides a practical, data-driven solution to this common and critical

pediatric surgical challenge, particularly in low-resource settings where imaging and subspecialty

access may be limited or unavailable. By leveraging readily available clinical and laboratory data,

the model can support frontline providers in making early and accurate decisions. In doing so,

Dharma may improve pediatric health outcomes by reducing diagnostic errors and guiding timely

referrals.

Thus, Dharma offers a reliable, transparent, interpretable, and evidence-based clinical decision

support system for the common yet challenging clinical scenario of pediatric appendicitis. We also

encourage others to build upon the concepts and framework we have developed, which we believe

are adaptable to a wide range of healthcare challenges beyond pediatric appendicitis.

In alignment with open-access principles, we have made the source code publicly available at

GitHub Repository Link [46] and the web-based tool accessible at [dharma-ai.org], promoting

transparency, reproducibility, and continuous advancement in clinical decision support and

medicine as a whole through knowledge-sharing.

Acknowledgement

529   We thank Marcinkevics R and colleagues [11] for their inspiring work and for publicly providing

530   anonymized data that facilitated our study. We also appreciate the contributions of our colleague

531   Amrit Neupane for designing the user interface and user experience of our web tool.

Author Contributions

Conceptualization and study design: Anup Thapa, Subash Pahari

Data curation: Anup Thapa, Subash Pahari

Literature review: Anup Thapa

Formal analysis: Anup Thapa

Methodology: Anup Thapa

Software: Anup Thapa (model development), Subash Pahari (web-app development)

Writing – original draft: Anup Thapa (except introduction), Sashank Timilsina, Binay Chapagain
(introduction)

Writing – review & editing: Anup Thapa

Writing – Manuscript Formating: Subash Pahari

References

[1] Hijaz NM, Friesen CA. Managing acute abdominal pain in pediatric patients: current
perspectives. Pediatric Health Med Ther. 2017 June 29;8:83-91. doi: 10.2147/PHMT.S120156.
PMID: 29388612; PMCID: PMC5774593.

[2] Tseng YC, Lee MS, Chang YJ, Wu HP. Acute abdomen in pediatric patients admitted to the
pediatric emergency department. Pediatr Neonatol. 2008 Aug;49(4):126-34. doi: 10.1016/S1875-
9572(08)60027-3. PMID: 19054918.

[3] Schneider C, Kharbanda A, Bachur R. Evaluating appendicitis scoring systems using a
prospective pediatric cohort. Ann Emerg Med. 2007 Jun;49(6):778-84, 784.e1. doi:
10.1016/j.annemergmed.2006.12.016. Epub 2007 Mar 26. PMID: 17383771.

[4] Sencan A, Aksoy N, Yıldız M, Okur Ö, Demircan Y, Karaca I. The evaluation of the validity

of Alvarado, Eskelinen, Lintula and Ohmann scoring systems in diagnosing acute appendicitis in

children. Pediatr Surg Int. 2014 Mar;30(3):317-21. doi: 10.1007/s00383-014-3467-0. PMID:

24448910.

[5] Pogorelić Z, Rak S, Mrklić I, Jurić I. Prospective validation of Alvarado score and Pediatric

Appendicitis Score for the diagnosis of acute appendicitis in children. Pediatr Emerg Care. 2015

Mar;31(3):164-8. doi: 10.1097/PEC.0000000000000375. PMID: 25706925.

[6] Sağ S, Basar D, Yurdadoğan F, Pehlivan Y, Elemen L. Comparison of Appendicitis Scoring

Systems in Childhood Appendicitis. Turk Arch Pediatr. 2022 Sep;57(5):532-537. doi:

10.5152/TurkArchPediatr.2022.22076. PMID: 36062441; PMCID: PMC9524470.

[7] Jukić M, Nizeteo P, Matas J, Pogorelić Z. Trends and Predictors of Pediatric Negative

Appendectomy Rates: A Single-Centre Retrospective Study. Children (Basel). 2023 May

15;10(5):887. doi: 10.3390/children10050887. PMID: 37238435; PMCID: PMC10217643.

[8] Kundiona I, Chihaka OB, Muguti GI. Negative appendicectomy: evaluation of ultrasonography

and Alvarado score. Cent Afr J Med. 2015 Sep-Dec;61(9-12):66-73. PMID: 29144064.

[9] Oyetunji TA, Ong'uti SK, Bolorunduro OB, Cornwell EE 3rd, Nwomeh BC. Pediatric negative

appendectomy rate: trend, predictors, and differentials. J Surg Res. 2012 Mar;173(1):16-20. doi:

10.1016/j.jss.2011.04.046. Epub 2011 May 19. PMID: 21696768.

[10] Patel H, Kamel M, Cooper E, Bowen C, Jester I. The Variable Definition of "Negative

Appendicitis" Remains a Surgical Challenge. Pediatr Dev Pathol. 2024 Nov-Dec;27(6):552-558.

doi: 10.1177/10935266241255281. Epub 2024 June 6. PMID: 38845117.

[11] Marcinkevics R, Reis Wolfertstetter P, Wellmann S, Knorr C, Vogt JE. Using Machine

Learning to Predict the Diagnosis, Management and Severity of Pediatric Appendicitis. Front

Pediatr. 2021 Apr 29;9:662183. doi: 10.3389/fped.2021.662183. PMID: 33996697; PMCID:

PMC8116489.

[12] Bălănescu L, Băetu AE, Cardoneanu AM, Moga AA, Bălănescu RN. Predictors of

Complicated Appendicitis with Evolution to Appendicular Peritonitis in Pediatric Patients.

Medicina (Kaunas). 2022 Dec 22;59(1):21. doi: 10.3390/medicina59010021. PMID: 36676645;

PMCID: PMC9866196.

[13] Callahan MJ, Rodriguez DP, Taylor GA. CT of appendicitis in children. Radiology. 2002

Aug;224(2):325-32. doi: 10.1148/radiol.2242010998. PMID: 12147823.

[14] Kim DW, Yoon HM, Lee JY, Kim JH, Jung AY, Lee JS, Cho YA. Diagnostic performance

of CT for pediatric patients with suspected appendicitis in various clinical settings: a systematic

review and meta-analysis. Emerg Radiol. 2018 Dec;25(6):627-637. doi: 10.1007/s10140-018-

1624-9. Epub 2018 Jul 12. PMID: 30003463.

[15] Sivit CJ, Applegate KE, Stallion A, Dudgeon DL, Salvator A, Schluchter M, Berlin SC, Myers

MT, Borisa VJ, Weinert DM, Morrison SC, Grisoni ER. Imaging evaluation of suspected

appendicitis in a pediatric population: effectiveness of sonography versus CT. AJR Am J

Roentgenol. 2000 Oct;175(4):977-80. doi: 10.2214/ajr.175.4.1750977. PMID: 11000147.

[16] Stephen AE, Segev DL, Ryan DP, Mullins ME, Kim SH, Schnitzer JJ, Doody DP. The

diagnosis of acute appendicitis in a pediatric population: to CT or not to CT. J Pediatr Surg. 2003

Mar;38(3):367-71; discsussion 367-71. doi: 10.1053/jpsu.2003.50110. PMID: 12632351.

[17] Brennan GD. Pediatric appendicitis: pathophysiology and appropriate use of diagnostic

imaging. CJEM. 2006 Nov;8(6):425-32. doi: 10.1017/s1481803500014238. PMID: 17209492.

[18] Mittal MK. Appendicitis: Role of MRI. Pediatr Emerg Care. 2019 Jan;35(1):63-66. doi:

10.1097/PEC.0000000000001710. PMID: 30608328.

[19] Pines JM. Trends in the rates of radiography use and important diagnoses in emergency

department patients with abdominal pain. Med Care. 2009 Jul;47(7):782-6. doi:

10.1097/MLR.0b013e31819748e9. PMID: 19536032.

[20]Lastunen K, Leppäniemi A, Mentula P. Perforation rate after a diagnosis of uncomplicated

appendicitis on CT. BJS Open. 2021 Jan 8;5(1):zraa034. doi: 10.1093/bjsopen/zraa034. PMID:

33609386; PMCID: PMC7893470.

[21] Almaramhy HH. Acute appendicitis in young children less than 5 years: review article. Ital J

Pediatr. 2017 Jan 26;43(1):15. doi: 10.1186/s13052-017-0335-2. PMID: 28257658; PMCID:

PMC5347837.

[22] Pogorelić Z, Domjanović J, Jukić M, Poklepović Peričić T. Acute Appendicitis in Children

Younger than Five Years of Age: Diagnostic Challenge for Pediatric Surgeons. Surg Infect

(Larchmt). 2020 Apr;21(3):239-245. doi: 10.1089/sur.2019.175. Epub 2019 Oct 16. PMID:

31618143.

[23] Handelman GS, Kok HK, Chandra RV, Razavi AH, Lee MJ, Asadi H. eDoctor: machine

learning and the future of medicine. J Intern Med. 2018 Dec;284(6):603-619. doi:

10.1111/joim.12822. Epub 2018 Sep 3. PMID: 30102808.

[24] Harmantepe AT, Dikicier E, Gönüllü E, Ozdemir K, Kamburoğlu MB, Yigit M. A different

way to diagnosis acute appendicitis: machine learning. Pol Przegl Chir. 2023 Oct 13;96(2):38-43.

doi: 10.5604/01.3001.0053.5994. PMID: 38629278.

[25] Reismann J, Romualdi A, Kiss N, Minderjahn MI, Kallarackal J, Schad M, Reismann M.

Diagnosis and classification of pediatric acute appendicitis by artificial intelligence methods: An

investigator-independent approach. PLoS One. 2019 Sep 25;14(9):e0222030. doi:

10.1371/journal.pone.0222030. PMID: 31553729; PMCID: PMC6760759.

[26] Byun J, Park S, Hwang SM. Diagnostic Algorithm Based on Machine Learning to Predict

Complicated Appendicitis in Children Using CT, Laboratory, and Clinical Features. Diagnostics

(Basel). 2023 Mar 1;13(5):923. doi: 10.3390/diagnostics13050923. PMID: 36900066; PMCID:

PMC100010Forsström JJ, Irjala K, Selén G, Nyström M, Eklund P. Using data preprocessing and

single layer perceptron to analyze laboratory data. Scand J Clin Lab Invest Suppl. 1995;222:75-

81. doi: 10.3109/00365519509088453. PMID: 7569750.49.

[27] Forsström JJ, Irjala K, Selén G, Nyström M, Eklund P. Using data preprocessing and single

layer perceptron to analyze laboratory data. Scand J Clin Lab Invest Suppl. 1995;222:75-81. doi:

10.3109/00365519509088453. PMID: 7569750.

[28] Grigull L, Lechner WM. Supporting diagnostic decisions using hybrid and complementary

data mining applications: a pilot study in the pediatric emergency department. Pediatr Res. 2012

Jun;71(6):725-31. doi: 10.1038/pr.2012.34. Epub 2012 Mar 22. PMID: 22441377.

[29] Chadaga K, Khanna V, Prabhu S, Sampathila N, Chadaga R, Umakanth S, Bhat D, Swathi

KS, Kamath R. An interpretable and transparent machine learning framework for appendicitis

detection in pediatric patients. Sci Rep. 2024 Oct 18;14(1):24454. doi: 10.1038/s41598-024-

75896-y. Erratum in: Sci Rep. 2025 Jan 22;15(1):2841. doi: 10.1038/s41598-025-86494-x. PMID:

39424647; PMCID: PMC11489819.

[30] Hua R, O'Brien MK, Carter M, Pitt JB, Kwon S, Ghomrawi HMK, Jayaraman A, Abdullah

F. Improving Early Prediction of Abnormal Recovery after Appendectomy in Children using Real-

world Data from Wearables. Annu Int Conf IEEE Eng Med Biol Soc. 2024 Jul;2024:1-4. doi:

10.1109/EMBC53108.2024.10782031. PMID: 40039430.

[31] Ghomrawi HMK, O'Brien MK, Carter M, Macaluso R, Khazanchi R, Fanton M, DeBoer C,

Linton SC, Zeineddin S, Pitt JB, Bouchard M, Figueroa A, Kwon S, Holl JL, Jayaraman A,

Abdullah F. Applying machine learning to consumer wearable data for the early detection of

complications after pediatric appendectomy. NPJ Digit Med. 2023 Aug 16;6(1):148. doi:

10.1038/s41746-023-00890-z. PMID: 37587211; PMCID: PMC10432429.

[32] Pop GN, Costea FO, Lungeanu D, Iacob ER, Popoiu CM. Ultrasonographic findings of child

acute appendicitis incorporated into a scoring system. Singapore Med J. 2022 Jan;63(1):35-41.

doi: 10.11622/smedj.2020102. Epub 2020 Jul 15. PMID: 32668829; PMCID: PMC9251212.

[33] Adir A, Braester A, Natalia P, Najib D, Akria L, Suriu C, Masad B, Igor W. The role of blood

inflammatory markers in the preoperative diagnosis of acute appendicitis. Int J Lab Hematol. 2024

Feb;46(1):58-62. doi: 10.1111/ijlh.14163. Epub 2023 Aug 29. PMID: 37644670.

[34] Moris D, Paulson EK, Pappas TN. Diagnosis and Management of Acute Appendicitis in

Adults: A Review. JAMA. 2021 Dec 14;326(22):2299-2311. doi: 10.1001/jama.2021.20502.

PMID: 34905026.

[35] Yajima H, Fujita T, Yanaga K. Profile of signs and symptoms in mild and advanced acute

appendicitis. Int Surg. 2010 Jan-Mar;95(1):63-6. PMID: 20480844.

[36] Hajibandeh S, Hajibandeh S, Hobbs N, Mansour M. Neutrophil-to-lymphocyte ratio predicts

acute appendicitis and distinguishes between complicated and uncomplicated appendicitis: A

systematic review and meta-analysis. Am J Surg. 2020 Jan;219(1):154-163. doi:

10.1016/j.amjsurg.2019.04.018. Epub 2019 Apr 27. PMID: 31056211.

[37] Biehl CM, Elliver M, Gudjonsdottir J, Salö M. Utility of Urine Dipstick Testing in Pediatric

Appendicitis: Assessing its Role in Identifying Complicated Cases and Retrocecal Appendicitis.

Eur J Pediatr Surg. 2024 Dec 19. doi: 10.1055/a-2490-1156. Epub ahead of print. PMID:

39701137.

[38] Singh, C. Machine Learning in Pattern Recognition. European Journal of Engineering and

Technology Research. 2023 Apr;8(2):63–68. doi:10.24018/ejeng.2023.8.2.3025.

[39] Issaiy M, Zarei D, Saghazadeh A. Artificial Intelligence and Acute Appendicitis: A

Systematic Review of Diagnostic and Prognostic Models. World J Emerg Surg. 2023 Dec

19;18(1):59. doi: 10.1186/s13017-023-00527-2. PMID: 38114983; PMCID: PMC10729387.

[40] Scrimgeour DS, Driver CP, Stoner RS, King SK, Beasley SW. When does ultrasonography

influence management in suspected appendicitis? ANZ J Surg. 2014 May;84(5):331-4. doi:

10.1111/ans.12415. Epub 2014 Jan 9. PMID: 24405944.

[41] Held JM, McEvoy CS, Auten JD, Foster SL, Ricca RL. The non-visualized appendix and

secondary signs on ultrasound for pediatric appendicitis in the community hospital setting. Pediatr

Surg Int. 2018 Dec;34(12):1287-1292. doi: 10.1007/s00383-018-4350-1. Epub 2018 Oct 6. PMID:

30293146.

[42] Nijssen DJ, van Amstel P, van Schuppen J, Eeftinck Schattenkerk LD, Gorter RR, Bakx R.

Accuracy of ultrasonography for differentiating between simple and complex appendicitis in

children. Pediatr Surg Int. 2021 Jul;37(7):843-849. doi: 10.1007/s00383-021-04872-8. Epub 2021

Mar 7. PMID: 33677613; PMCID: PMC8172400.

[43] Mills LD, Mills T, Foster B. Association of clinical and laboratory variables with ultrasound

findings in right upper quadrant abdominal pain. South Med J. 2005 Feb;98(2):155-61. doi:

10.1097/01.SMJ.0000129927.88863.65. PMID: 15759944.

[44] Marcinkevičs R, Reis Wolfertstetter P, Klimiene U, Ozkan E, Chin-Cheong K, Paschke A,

et al. Regensburg Pediatric Appendicitis Dataset. Zenodo; 2023.

[45] Steiner Z, Buklan G, Stackievicz R, Gutermacher M, Litmanovitz I, Golani G, Arnon S.

Conservative treatment in uncomplicated acute appendicitis: reassessment of practice safety. Eur

J Pediatr. 2017 Apr;176(4):521-527. doi: 10.1007/s00431-017-2867-2. Epub 2017 Feb 16. PMID:

28210834

[46] Thapa A. Dharma-AppendicitisModel [Internet]. 2025 [cited 2025 May 26]. Available from:

https://github.com/ajung17/Dharma-AppendicitisModel

Figure

Figure

Figure