1

Dharma: A novel, clinically grounded machine learning framework for pediatric appendicitis—diagnosis, severity assessment and evidence-based clinical decision support.

Anup Thapa Kshetri[1*], Subash Pahari[2], Shashank Timilsina[1], Binay Chapagain[3]

[1]Department of Emergency Medicine and General Practice, Matri-Sishu Miteri Provincial Hospital, Pokhara, Gandaki, Nepal.

[2]Department of Computer Engineering, Faculty of Science and Technology, Pokhara University, Pokhara, Gandaki, Nepal.

[3]Department of Critical Care Medicine, Tribhuvan University Teaching Hospital, Kathmandu, Bagmati, Nepal.

Corresponding author: ajungthapa11@gmail.com (ATK)

Abstract

Acute appendicitis is a common but diagnostically challenging surgical emergency in children.

Existing linear scoring systems lack sufficient accuracy for standalone use, while advanced

imaging is constrained by risks of sedation, contrast, and radiation. Furthermore, no available

tools provide prognostic guidance.

We introduce *Dharma*, a machine learning framework consisting of a clinically grounded

imputer and two random forest classifiers for diagnosis and severity assessment. Designed for

real-world bedside use, Dharma is open-sourced and accessible through a web application.

Dharma achieved excellent diagnostic performance, with an AUC-ROC of 0.98 [0.97–0.99] and

accuracy of 93% [91–95]. For prognostic classification, it identified complicated appendicitis

with high sensitivity (96% [93–99]) and negative predictive value (97% [94–99]). In cases

without appendix visualization—a frequent limitation in resource-constrained settings—Dharma

maintained strong performance (AUC-ROC 0.96 [0.93–0.99]), with specificity of 97% [93–100]

and PPV of 93% [84–100] at a 44% threshold, and sensitivity of 92% [84–98] with NPV of 95%

[91–99] at a 25% threshold. These threshold-dependent trade-offs enable Dharma to support both

ruling in and ruling out appendicitis within diverse clinical workflows.

Beyond pediatric appendicitis, Dharma's open-source framework and clinically grounded design

provide a generalizable foundation for developing equitable and practical decision-support tools

in healthcare.

Keywords: Dharma, appendicitis, pediatric appendicitis, machine learning, clinical decision

support , evidence-based medicine, artificial intelligence, Clinical AI, Hybrid AI, AI

Author Summary

Accurate diagnosis and risk stratification of pediatric appendicitis remain challenging due to

heterogeneous clinical presentations, the absence of definitive biomarkers, and limited access to

reliable imaging, particularly in resource-constrained settings. Existing clinical scoring systems,

while simple and widely used, rely on fixed linear assumptions, lack interpretability at the

individual patient level, and are restricted to diagnostic decision-making without addressing

disease severity or prognosis.

To address these limitations, we developed **Dharma**, a clinically grounded, interpretable

machine learning-based framework designed to support both diagnosis and severity assessment

of pediatric appendicitis using routinely available clinical, laboratory, and radiological data.

Rather than replacing clinical judgment, Dharma is intentionally designed to mirror real-world

clinical reasoning, explicitly model non-linear interactions among features, and provide

transparent explanations for its predictions.

Implemented as an open-access, web-based clinical decision support tool, Dharma delivers real-

time, evidence-based risk estimates that are adaptable to varying resource settings. Beyond

pediatric appendicitis, this work demonstrates a generalizable, medicine-first approach to

developing equitable and interpretable AI systems that complement clinician decision-making

and align with the evolving needs of 21st-century healthcare.

Background

Acute abdominal pain (AAP) is a common complaint of Pediatric Patients which accounts for up

to 10 % of total visits to the Emergency Department while acute appendicitis is the major cause

of AAP in children over the age of 1 [1,2]. Despite being a routinely encountered clinical

dilemma, acute appendicitis remains a challenging diagnosis for healthcare professionals. This is

likely due to variability in clinical presentation and the frequent absence of classic symptoms in

children. Various scoring systems, with Alvarado being the oldest and most widely used, help

physicians quantify the risk of pediatric appendicitis. However, none of the scoring systems

provide sufficient sensitivity, specificity or predictive values to be used as an exclusive standard

in setting the diagnosis. [3-5].

The increasing reliance on CT and USG for evaluating acute abdominal pain has not been

accompanied by a corresponding improvement in appendicitis detection rates [19]. Misdiagnosis

rates remain alarmingly high, ranging from 70% to 100% in children aged three years or

younger, with a gradual decrease as age increases [21]. This diagnostic challenge is often

attributed to the atypical presentation of the disease in younger children and the anatomical

variability of the diameter of the appendix. Consequently, the lack of a reliable diagnostic tool

may contribute to the persistently high incidence of Negative Appendectomy (NA) worldwide,

reported between 5% and 20% in various centers [7-10]. NA is a dreaded outcome for any

operating surgeon due to the inherent risks associated with general anesthesia and surgery,

particularly in the pediatric population.

Moreover, no traditional inflammatory biomarkers, clinical signs and symptoms, or imaging

modalities can reliably predict the progression of uncomplicated to complicated appendicitis

[12,42]. This is particularly concerning in pediatric patients, where the incidence of perforated

appendix is alarmingly high and increases as the age decreases, reaching over 80% in children

younger than three years old [22]. This highlights the urgent need to refine diagnostic and

prognostic strategies for pediatric appendicitis.

Machine Learning (ML), a subfield of Artificial Intelligence (AI), involves learning patterns

from data and applying this knowledge to make predictions in future scenarios [38]. In medicine,

ML offers the potential to enhance diagnostic, prognostic, and management precision by

integrating heterogeneous clinical, laboratory, and imaging data into predictive frameworks.

Accordingly, AI and ML have seen growing application across medical domains, with recent

emphasis on Large Language Models (LLMs) and other data-driven architectures.

However, in high-stakes clinical settings, predictive performance alone is insufficient. Clinical

decision-making occurs under uncertainty, incomplete information, and asymmetric risk, where

the consequences of false negatives and false positives differ substantially. As a result, AI

systems are most effective when they are designed around clinical reasoning, workflow

constraints, and patient safety considerations, with ML serving as a supporting component rather

than the primary driver. Hybrid, human-centered approaches—where model behavior,

interpretability, and optimization objectives are explicitly aligned with clinical priorities—are

more likely to generalize, earn clinician trust, and operate safely in real-world environments [49–

51].

Placing human understanding of disease progression, decision pathways, and clinical constraints

at the core of system design, while leveraging ML as an enabling tool, may better realize AI's

potential to deliver accurate, efficient, and accessible healthcare globally [51,52].

ML has been applied to appendicitis diagnosis since the 1990s, with pediatric-specific models

emerging in the early 2010s [28]. These models have primarily focused on predicting pediatric

appendicitis, associated complications, and postoperative recovery outcomes [11,24–31]. Despite

the development of numerous ML-based tools reporting promising retrospective performance,

none have been widely adopted as real-world clinical decision support systems.

ML-based approaches have the potential to enhance both diagnostic accuracy and severity

prediction in pediatric appendicitis [39], thereby reducing unnecessary radiation exposure from

computed tomography, minimizing sedation-related risks, lowering negative appendectomy

rates, and optimizing referral pathways from primary care. However, achieving these benefits

requires models that go beyond retrospective performance metrics and demonstrate real-time

clinical utility [51]. Such systems must perform reliably even with missing or incomplete data

and remain applicable across diverse healthcare environments, from well-resourced urban

hospitals to resource-limited rural settings. Therefore, there is a pressing need for a transparent,

interpretable, adaptable, and user-friendly clinical decision support system that can reliably assist

in diagnosis, rule out disease, stratify complication risk, and guide management decisions in

pediatric appendicitis.

Thus, we defined two primary objectives for this study:

⬜ To develop and evaluate a clinically informed, optimized, and generalizable ML

128   framework for diagnosing pediatric appendicitis and predicting complications, and to

129   compare its performance with conventional diagnostic tools (AS, PAS, AIR, Tzanaki,

130   USG) as well as state-of-the-art ML architectures (XGBoost, LightGBM).

132 ⬚ Deploy the ML framework as an interpretable, web-based diagnostic and prognostic

133 tool that facilitates evidence-based clinical decision-making for pediatric appendicitis,

134 with particular applicability in resource-constrained settings.

Materials And Methods

Ethics Statement

140 The original dataset was collected in accordance with the ethical guidelines and approval

141 procedures of the University of Regensburg(approval numbers: 18-1063-101, 18-1063_1-101,

142 and 18-1063_2-101). For the present study, only anonymized secondary data were analyzed. In

143 line with these institutional regulations, no identifiable patient information was included, and

144 therefore additional ethical approval and individual informed consent were not required.

Data Source

This study analyzed anonymized secondary data from pediatric patients with suspected

appendicitis. In the primary cohort, diagnosis data were available for 780 of 782 patients

admitted with abdominal pain to Children's Hospital St. Hedwig in Regensburg, Germany,

between 2016 and 2021 [44]. For testing the diagnostic model, 289 unique records from 430

suspected appendicitis cases collected at the same hospital between January 1, 2016, and

December 31, 2018, were used [11]. To develop the severity assessment model, the training

dataset was augmented with 49 complicated cases from 301 records collected between 2015 and

2022 at the Department of Pediatric Surgery and Pediatric Traumatology, Florence-Nightingale

Hospital, Düsseldorf, Germany [47]. All datasets were already anonymized, with patient

identifiers removed, and are publicly accessible from the respective sources.

Dataset Description

The primary dataset consisted of 782 suspected pediatric appendicitis cases collected over a

three-year period at a tertiary children's hospital in Germany [44]. Each record contained 55

demographic, clinical, laboratory, and radiological variables, along with labels for diagnosis,

treatment modality, and severity. The cohort included children and adolescents aged 0–18 years

who presented with abdominal pain and were evaluated for suspected appendicitis. Patients with

a prior appendectomy, concurrent abdominal pathologies, or pre-evaluation antibiotic use for

unrelated infections were not included. After applying these criteria and excluding two records

lacking diagnostic information, 780 cases with confirmed diagnostic labels were available for

analysis.

☐   Diagnosis: appendicitis (n = 463) and no appendicitis (n = 317)

☐   Severity: complications (n = 119) and no complications or no appendicitis (n = 661)

□   Treatment Modality: conservative (n = 483) and surgical (n = 297)

For severity assessment modeling, the primary dataset was augmented with 289 additional

unique records from the cohort used in literature [11], yielding a combined dataset of 1,069

patients after deduplication. Only patients with confirmed appendicitis were retained for

prognostic modeling, resulting in a final sample of 650 cases, of which 490 had uncomplicated

disease and 160 presented with complications. This subset of the dataset allowed us to train and

evaluate a clinically meaningful model focused specifically on prognostic discrimination

following diagnostic classification.

Given the scarcity of publicly available pediatric appendicitis datasets, we used all curated cases

from the three high-quality cohorts [11, 44, 47], applying task-specific inclusion criteria to derive

the diagnostic and prognostic subsets used in this study, as described in detail in the Model

Development section.

Data Preprocessing

Data preprocessing involved standardizing categorical encodings: binary variables were mapped

to 0/1, urinary ketones were coded as 0 (absence/trace), 1 (1+), 2 (2+), and 3 (3+), peritonitis was

coded as 0 (absent), 1 (local), and 2 (generalized), and stool changes were coded as 0 (normal)

and 1 (any alteration, including diarrhea or constipation).

Missing values were present in almost all variables (see Supplementary Table 1). The appendix

diameter variable had 36% missingness, yet it was retained because of both the inherent

challenges of appendix visualization on ultrasonography (USG) and its well-established

diagnostic and pathophysiological relevance in appendicitis.

S1 Table: Percentage of missing values in each feature.

For initial exploratory analyses, statistical tests were conducted on the available data without

imputation. For model development, missing values in selected predictors were handled using a

custom imputation pipeline, the Dharma_Imputer. This approach incorporated domain

knowledge by grouping inflammatory markers (WBC count, neutrophil percentage, CRP, and

temperature) and imputing them through iterative imputation with scikit-learn's decision tree

regressor. Categorical variables were then imputed separately, with each feature modeled by a

dedicated decision tree classifier. The appendix diameter feature was treated distinctly: missing

values were flagged by an indicator column and imputed with a clinically impossible placeholder

value of -1, reflecting that non-visualization of the appendix on ultrasound is common and

diagnostically meaningful. This strategy preserved the clinical nuance that decision-making

differs depending on whether the appendix is visualized or not.

To preserve data integrity, imputations for training, validation, and test sets were performed

independently. The trained Dharma_Imputer was also integrated into the web-based clinical

decision support tool, enabling clinically informed, real-time handling of missing values.

For the primary classification task, we used the full dataset of 780 cases. Additionally, 239

unique cases from the 430 suspected appendicitis cases [11] were used exclusively for external

evaluation of our models and existing linear scoring systems.

For the severity assessment task, the dataset was augmented with the full set of 430 suspected

cases [11]. After duplicate removal, the combined dataset comprised 1,069 unique patient

records. Among these, only patients with a confirmed diagnosis of appendicitis (n = 650) were

included for training and testing the severity assessment model.

Treatment modality was not analyzed in this study, as we consider it a clinical decision to be

determined by the operating surgeon based on the diagnosis and anticipated risk of

complications.

221   Feature Selection

Advanced radiological features such as fecal impaction, bowel wall thickening, or pathological

lymph nodes were excluded from the study, as these findings can be challenging to assess even

for experienced radiologists. Our aim was to develop a model that could augment pediatric

surgical decision-making, particularly in resource-constrained settings where access to advanced

radiology is almost always limited.

The Chi-Square test was performed to assess the association between categorical variables and

disease state, while Cramér's V measured the strength of these associations. Variables with a

significant association ($p < 0.001$) and a moderate to strong effect size (Cramér's $V > 0.10$) were

selected for the development of the model. Contralateral rebound tenderness, despite meeting the

initial inclusion criteria, was excluded from the model's feature pool as it is not a classically

defined or standardized clinical sign in the diagnostic framework of acute appendicitis.

For continuous variables, the Mann–Whitney U test was applied to compare distributions

between positive and negative classes, and rank-biserial correlation was calculated to quantify

effect size. Features with significant differences ($p < 0.001$) and correlation coefficients > 0.40

were retained for the feature pool.

All statistical analyses were conducted in Python using Pandas, NumPy, and SciPy on the

original dataset of 780 suspected appendicitis cases.

To refine feature selection, Recursive Feature Elimination with Cross-Validation (RFECV) was

employed to identify the most influential features. Different subsets of selected features were

then evaluated independently to determine the optimal combination and arrangement for model

development.

Data mining was performed separately for diagnostic and prognostic targets. The final feature

sets used in each model are summarized in Table 1.

Table 1. List of Dharma's Features	
Diagnosis	**Complications**
Nausea	Nausea
Appetite Loss	Appetite Loss
Peritonitis	Peritonitis
WBC Count	Urinary Ketones
Neutrophil Percentage	Free Fluids
CRP	CRP
Urinary Ketones	WBC Count
Appendix Diameter	Body Temperature
Free Fluids	Appendix Diameter
	Neutrophil Percentage

Model Development

The Random Forest model was selected for its suitability in handling non-parametric medical

data. For diagnosis prediction, the dataset was split into training, validation, and test sets in a

3:1:1 ratio. The model was optimized through hyperparameter tuning using grid search, followed

by randomized search and stratified cross-validation, ensuring robust performance. To mitigate

overfitting and ensure consistent performance across diverse clinical presentations, 10-fold

stratified cross-validation was performed on the 555 bootstrap samples of the training set.

The optimal classification threshold was determined using Youden's Index and the Euclidean

Distance method on the Receiver Operating Characteristic (ROC) curve applied to the validation

set. We aimed to strike a balance between sensitivity and specificity to minimize both false

positives and false negatives in appendicitis diagnosis.

The model's performance was evaluated on independent test sets using AUC-ROC, sensitivity,

specificity, and predictive values to ensure an unbiased clinical assessment. AUC-ROC was

chosen as the primary evaluation metric for the diagnostic task because it captures threshold-

independent discriminative performance, allowing decision thresholds to be adjusted post-hoc to

suit different clinical subgroups or population cohorts without retraining the entire model.

The 650 diagnosed appendicitis cases, curated as described in the data preprocessing section,

were split into training, validation, and test sets in a 3:1:1 ratio. The severity classes were highly

imbalanced, with negative-to-positive ratios of 286:104 in the training set, 98:32 in the validation

set, and 106:24 in the test set. The higher prevalence of the negative class in the training set

biased the model towards predicting uncomplicated cases. To address this, we augmented the

training set with 49 complicated cases from the literature [47] and applied NearMiss

undersampling to reduce the abundance of easily identifiable uncomplicated cases. This resulted

in a more balanced dataset with a ratio of 235:153 (1.54:1).

We further adjusted the class weights by assigning the positive class a weight of 5:1, thereby

penalizing false negatives more heavily than false positives. This deliberate design choice biased

the prognostic model toward identifying complicated appendicitis and was considered clinically

appropriate, as missing complications carries substantially greater risk—such as perforation and

gangrene—than incorrectly flagging non-complicated cases. Importantly, all patients in this

cohort had a confirmed diagnosis of appendicitis; therefore, a false-positive prediction would

typically result in an appendectomy, which is an accepted and established management even for

non-complicated appendicitis. In contrast, a false-negative prediction could delay timely surgical

intervention in patients with complicated disease, potentially leading to life-threatening

outcomes.

The primary evaluation metric for the complication model was recall (sensitivity) for the positive

class, reflecting the clinical priority of minimizing missed complications.

We named the final machine-learning framework **Dharma**, which comprises the clinically

grounded *Dharma_Imputer* and two Random Forest classifier models: one for diagnosis and the

other for prognosis among positively diagnosed cases.

The hyperparameters used for Dharma's Diagnostic and Severity Assessment models are in

Table 2.

Table 2. Hyperparameters for Dharma

Hyperparameters	Diagnostic Model	Prognostic Model
n_estimators	555	888
min_samples_split	12	5
min_samples_leaf	1	5
max_depth	35	3
class_weight	"balanced"	{0: 1, 1: 5}
max_features	sqrt	sqrt
random_state	17	17

Model Evaluation

We applied a multi-layer evaluation strategy for Dharma across both diagnostic and prognostic

tasks to assess robustness and generalization. First, model performance was estimated using 10-

fold stratified cross-validation across 555 bootstrap samples of the training set. Within each

bootstrap sample, Dharma was benchmarked against state-of-the-art models appropriate for our

dataset size, namely XGBoost and LightGBM.

The diagnostic model was further evaluated on two independent test sets: (i) 5,555 bootstrap

samples of the unaltered test split from the primary dataset, and (ii) 5,555 bootstrap samples of

239 unique cases extracted from [11]. For the prognostic (complications) model, the test set

contained only 18% complicated cases (106:24). Dharma's severity feature model was

benchmarked against XGBoost under two settings: one tuned with simple hyperparameters and

another with complex hyperparameters highly regularized for high recall.

Model performance was reported in terms of AUC-ROC, sensitivity, specificity, predictive

values, and corresponding 95th-percentile confidence intervals. Hyperparameter tuning for all

benchmark models was performed using grid search with stratified 10-fold cross-validation to

ensure a fair comparison.

The hyperparameter settings of all benchmark models for diagnostic and prognostic tasks are

reported in Tables 3 and 4 respectively.

Table 3. Hyperparameters of SOTA models for diagnosis			
XGBoost		**LightGBM**	
n_estimators	555	**n_estimators**	333
max_depth	3	**max_depth**	3
min_child_weight	6	**min_child_samples**	3
learning_rate	0.01	**learning_rate**	0.01
gamma	0	**min_gain_to_split**	0
subsample	0.6	**bagging_fraction**	0.6
colsample_bytree	0.6	**feature_fraction**	0.6

reg_alpha	0.1	lambda_l1	0
reg_lamda	1	lambda_l2	1
random_state	17	random_state	17

Table 4. Hyperparameters of SOTA models for complications		
Hyperparameters	**XGB(Simple)**	**XGB(Complex)**
n_estimators	111	111
max_depth	3	3
min_child_weight	6	6
scale_pos_weight	7	7
learning_rate	0.01	0.01
gamma	-	1
subsample	-	0.8
colsample_bytree	-	0.6
reg_alpha	-	0.1
reg_lamda	-	10

We further evaluated the robustness of Dharma against different imputation strategies for

missing values. For the markers of inflammation (WBC, Neutrophil Percentage, Body

temperature, CRP), we applied Iterative Imputer from the Scikit-learn library with decision tree

regressor and linear regressor estimators, as well as a K-nearest neighbors (KNN) imputer. For

categorical variables, we paired these regression imputation strategies with a decision tree

classifier custom-trained for each feature. In addition, we tested simple strategies using mean

imputation for continuous variables and mode imputation for categorical variables. Across all

methods, no statistically significant difference in model performance was observed, indicating

that Dharma's predictive ability was stable irrespective of the imputation approach

(Supplementary Table 2).

For the final Dharma_Imputer, we selected a combination of decision tree regressor imputation

for continuous variables and decision tree classifier imputation for categorical variables. This

approach enables real-time, clinically informed imputation, which is particularly important in

real-world settings where missing data are frequent, especially in resource-constrained

environments.

S1 Table. Performance stability of Dharma with different imputation strategies.

Explainability Analysis

To examine Dharma's decision-making process, we applied SHAP's Tree Explainer (SHapley Additive exPlanations), a game theory–based approach that quantifies the contribution of each feature relative to a baseline prediction (base SHAP value). A feature's SHAP value indicates its influence in shifting the predicted probability toward or away from this baseline. Global feature contributions were visualized using SHAP summary plots for both diagnostic and prognostic tasks. Additionally, we employed the Random Forest's native *feature_importance* method to assess the relative importance assigned to each variable by the models.

At the individual level, SHAP values were integrated into our web application alongside 95th-percentile confidence intervals derived from the 555 decision trees in Dharma's random forest, providing real-time uncertainty estimates. This integration enables attending physicians to explore the reasoning behind model predictions, thereby enhancing transparency, interpretability, and clinical trust.

Results

Data Mining

A total of 780 pediatric patients (ages 0–18 years; mean age: 11.35 years, 95% CI: 11.10–11.59) with suspected appendicitis were included in the diagnostic prediction cohort, with a male-to-

female ratio of 1.07 (403:377). For complication prediction, 650 confirmed appendicitis cases from the combined dataset, as described earlier, were included.

A Chi-square test was performed to assess the significance of associations, and Cramér's V was used to evaluate the strength of association of the categorical features. Seven features showed significant association with appendicitis diagnosis and six features were significantly associated with complication prediction ($p < 0.001$). Among the diagnostic features, peritonitis demonstrated the strongest association (Cramér's V = 0.37), followed by neutrophilia (Cramér's V = 0.23). For complication prediction, peritonitis again showed the strongest association (Cramér's V = 0.30), followed by urinary ketones (Cramér's V = 0.29), Table 5 and 6.

Table 5. Diagnosis and Categorical Variables (n=780)				
Features	**Chi-Square**	**P-value**	**Cramer's V**	**Samples**
Peritonitis	108.24	<0.001	0.37	773
Neutrophilia	67.17	<0.001	0.30	732
Free Fluids	32.48	<0.001	0.21	719
Ketones in Urine	24.34	<0.001	0.20	582
Contralateral Rebound Tenderness	25.08	<0.001	0.18	767
Nausea/Vomiting	23.19	<0.001	0.17	774

Loss of Appetite	19.24	<0.001	0.16	772
Ipsilateral Rebound Tenderness	8.64	<0.01	0.12	619
Sex	10.78	<0.01	0.12	779
RIF Tenderness	8.18	<0.01	0.10	774
Migratory Pain	6.70	0.01	0.09	773
Coughing Pain	6.30	0.01	0.09	766
Psoas Sign	4.00	0.05	0.07	745
Stool	3.04	0.22	0.06	765
RBC in Urine	2.02	0.57	0.06	576
WBC in Urine	1.25	0.74	0.05	583
Dysuria	1.61	0.20	0.05	753

Table 6. Complications and Categorical variables(n=463)				
Features	Chi-Square	P-value	Cramer's V	Samples
Peritonitis	41.29	<0.001	0.3	457
Ketones in Urine	26.45	<0.001	0.29	318
Neutrophilia	23.75	<0.001	0.23	434
Nausea/Vomiting	23.17	<0.001	0.22	458
Loss of Appetite	20.56	<0.001	0.21	456
Free Fluids	16.72	<0.001	0.2	430

RBC in Urine	10.65	0.01	0.18	317
Ipsilateral Rebound Tenderness	5.62	<0.05	0.14	305
WBC In Urine	4.9	0.18	0.12	319
Stool	5.12	0.08	0.11	451
RIF Tenderness	2.66	0.1	0.08	458
Psoas Sign	1.58	0.21	0.06	433
Contralateral Rebound Tenderness	1.5	0.22	0.06	451
Migratory Pain	0.59	0.44	0.04	457
Dysuria	0.54	0.46	0.04	437
Sex	0.54	0.46	0.03	462
Coughing Pain	0.01	0.94	0.004	450

A Mann-Whitney U test was conducted to assess the statistical significance of differences in the distributions of continuous variables between positive and negative cases, and rank-biserial correlation was used to evaluate the effect size of these differences, Table 7 and 8. Six variables were significantly associated with appendicitis diagnosis, and five variables were significantly associated with complication prediction ($p < 0.001$). Among the diagnostic variables, appendix diameter demonstrated the strongest effect (rank-biserial correlation = 0.9), followed by C-reactive protein (CRP) level (rank-biserial correlation = 0.46). For complication prediction, CRP level exhibited the strongest effect (rank-biserial correlation = 0.64), followed by white blood cell (WBC) count (rank-biserial correlation = 0.45).

Table 7. Diagnosis and Continuous variables (n=780)						
Feature	U_statistic	P_value	Rank-Biserial correlation	Mean (Positive Class)	Mean(Negative Class)	Samples
Appendix Diameter	2341	<0.001	0.9	8.7	5.04	498
CRP	38686	<0.001	0.46	44.9	11.72	771
WBC Count	39910	<0.001	0.45	14.28	10.33	776
Neutrophil Percentage	32518.5	<0.001	0.42	76.03	65.6	679
Body Temperature	57410	<0.001	0.21	37.52	37.24	775
BMI	79280	<0.001	-0.15	18.45	19.56	753
Weight	82617	<0.01	-0.13	41.72	45.25	778
Age	80879	0.02	-0.1	11.08	11.72	780
Height	74276.5	0.08	-0.07	146.93	149.51	755
RDW	65536.5	0.21	0.05	13.4	12.87	756
Hemoglobin	72577.5	0.52	-0.03	13.38	13.38	764
RBC Count	70347	0.92	0.004	4.79	4.82	764
Thrombocyte Count	70459.5	0.95	0.003	285.79	284.48	764

Table 8. Complications and Continuous variables(n=463)						
Feature	U_statistic	P_value	Rank-Biserial Correlation	Mean (Positive Class)	Mean(Negative Class)	Samples
CRP	6952.5	<0.001	0.64	107.68	24.52	457
WBC Count	10960	<0.001	0.45	17.43	13.23	460
Body Temperature	12010.5	<0.001	0.4	37.93	37.38	459
Neutrophil Percentage	8803.5	<0.001	0.38	82.44	74.14	403
Appendix Diameter	7296	<0.001	0.28	9.53	8.51	371
Weight	24201.5	<0.01	-0.2	38.73	42.75	461
Age	24327	<0.01	-0.2	10.08	11.42	463
Height	21988.5	<0.01	-0.17	141.56	148.79	443
BMI	21284.5	0.01	-0.16	17.92	18.64	441
Hemoglobin	21018.5	0.06	-0.12	13.36	13.39	450
Thrombocyte Count	17298	0.2	0.08	296.65	282.24	450
RBC Count	20250.5	0.23	-0.08	4.75	4.8	450
RDW	17530	0.44	0.05	13.56	13.35	445

Diagnostic Performance

Conventional Tools

All conventional scoring systems—Alvarado Score (AS), Pediatric Appendicitis Score (PAS), Appendicitis Inflammatory Response (AIR) Score, and Tzanaki Score—were significantly associated with appendicitis diagnosis (p < 0.001). The discriminatory performance, expressed as

395 AUC-ROC with 95% confidence intervals, was 0.76 [0.71–0.82] for AS, 0.70 [0.64–0.76] for

396 PAS, 0.74 [0.68–0.80] for AIR, and 0.91 [0.87–0.94] for the Tzanaki Score. Median values were

397 consistently higher among confirmed cases compared to negatives (AS: 7 vs. 5; PAS: 5 vs. 4;

398 AIR: 6 vs. 4; Tzanaki: 12 vs. 6), indicating higher scoring trends among positive cases, Table 9.

Table 9. Association of scoring systems with diagnosis				
Scoring Systems	P-value	Rank-Biserial Correlation	Median(Positive Class)	Median(Negative Class)
Alvarado Score	<0.001	0.52	7	5
Pediatric Appendicitis Score	<0.001	0.4	5	4
AIR Score	<0.001	0.48	6	4
Tzanaki Score	<0.001	0.82	12	6

The Pediatric Appendicitis Score (PAS) with a low cut-off of 4 demonstrated the highest

sensitivity, 89 [84–93], making it most effective for ruling out the disease. Conversely, the

Appendicitis Inflammatory Response (AIR) Score with a cut-off of 9 achieved very high

specificity, 99 [97–100], supporting its role in ruling in the disease. A stepwise approach—using

PAS ($\geq$ 4) for initial screening and AIR ($\geq$ 9) for final diagnostic confirmation may provide a

promising framework for future studies. However, none of these linear scoring systems, when

used in isolation, demonstrated sufficient performance to serve as a reliable standalone

diagnostic method, Table 10.

Table 10. Performance of Diagnostic tools on curated test set (n=289)						
Tools	AUC-ROC	Sensitivity %	Specificity %	PPV %	NPV %	Accuracy %
Alvarado Score(≥5)	0.76 [0.71-0.82]	86 [81-91]	45 [35-55]	68 [74-80]	64 [53-75]	72 [66-78]
Alvarado Score(≥7)	0.76 [0.71-0.82]	55 [48-62]	83[76-90]	86 [79-92]	50 [43-58]	65 [60-70]
PAS Score(≥4)	0.70 [0.64-0.76]	89 [84-93]	38 [29-48]	72 [67-78]	65 [53-76]	71 [66-76]
PAS Score(≥6)	0.70 [0.64-0.76]	48 [40-55]	76 [68-85]	79 [71-86]	44 [3]	58 [52-63]
AIR Score(≥5)	0.74 [68-80]	67 [60-73]	67 [57-76]	79 [72-85]	52 [44-61]	67 [61-72]
AIR Score(≥9)	0.74 [68-80]	16 [11-21]	99 [97-100]	97 [89-100]	39 [33-45]	45 [40-51]
Tzanaki Score(≥8)	0.91 [87-94]	85 [80-90]	88 [81-94]	93 [89-97]	76 [68-84]	86 [82-90]
USG(≥6 mm)	0.94 [0.89-0.99]	94 [91-97]	95 [87-100]	99 [97-100]	80 [67-91]	94 [91-97]

Ultrasonography (USG) demonstrated the highest diagnostic performance among the available

tools, with an AUC-ROC of 0.94 [0.89–0.99]. However, appendix diameter measurements were

unavailable in 34% of suspected cases, reflecting the inherent challenges in appendix

visualization. A similar limitation was observed with the Tzanaki Score, which relies heavily on

USG findings for its calculation despite its strong discriminatory ability.

Dharma

Dharma demonstrated consistently high diagnostic performance across cross-validation and independent test sets.

In 10-fold cross-validation, the model achieved an AUC-ROC of 0.98 [0.97–0.99] with an overall accuracy of 93% [91–95]. The performance was balanced, with sensitivity 93% [91–95], specificity 93% [90–96], PPV 95% [93–97], and NPV 90% [87–94].

In the first independent test set, the model maintained robust discriminative ability with an AUC-ROC of 0.98 [0.96–0.99] and accuracy of 91% [87–95]. Sensitivity remained high (93% [88–98]) with slightly lower specificity (87% [78–95]), while predictive values were balanced (PPV 93% [88–97], NPV 88% [78–96]).

In the second curated test set, performance was further reinforced, achieving an AUC-ROC of 0.98 [0.96–0.99] and accuracy of 94% [91–97]. The model demonstrated high specificity (98% [95–100]) and PPV (99% [97–100]) while maintaining sensitivity (92% [88–96]) and NPV (87% [80–93]).

Even in the cohort with an unvisualized appendix, Dharma maintained strong discriminatory ability with an AUC-ROC of 0.96 [0.93–0.99]. At a threshold of 44%, it achieved a very high specificity of 97% [93–100] with a PPV of 93% [84–100]. At a threshold of 25%, it achieved a high NPV of 95% [91–99] and sensitivity of 92% [84–98].

435 Overall, Dharma showed excellent and stable diagnostic accuracy with consistently high AUC

436 (>0.96) across all evaluations, demonstrating both generalizability and clinical robustness.

437 Detailed diagnostic results are summarized in Table 11 and Table 12.

Table 11. Dharma's Diagnostic Performance						
Dharma	AUC-ROC	Specificity	PPV	Sensitivity	NPV	Accuracy
10-fold cross- validation	0.98 [0.97-0.99]	93% [90-96]	95% [93-97]	93% [90-95]	90% [87-94]	93% [91-95]
Split Test Set(n=156)	0.98 [0.96-0.99]	87% [78-95]	93% [88-97]	93% [88-98]	88% [78-96]	91% [87-95]
Curated Test Set(n=289)	0.98 [0.96-0.99]	98% [95-100]	99% [97-100]	92% [88-96]	87% [80-93]	94% [91-97]

Table 12. Diagnostic Performance on Unvisualized appendix cohort (n = 144)						
Threshold	AUC-ROC	Specificity	PPV	Sensitivity	NPV	Accuracy
44%	0.96 [0.93-0.99]	97% [93-100]	93% [84-100]	75% [62-86]	87% [80-93]	89% [83-94]
25%	0.96 [0.93-0.99]	89% [83-95]	82% [72-92]	92% [84-98]	95% [91-99]	90%[85-95]

Prognostic Performance

For complication prediction, Dharma achieved a sensitivity of 96% [93–99] and NPV of 97% [94–99] in 10-fold cross-validation, with corresponding specificity 65% [53–75], PPV 65% [59–72], and AUC-ROC 0.92 [0.90–0.95]. This performance profile, characterized by high sensitivity and NPV, reflects the intended design of Dharma's severity assessment module as a screening tool to safely rule out complications and to guide decisions regarding surgical versus conservative management, as well as the urgency of surgery.

In the unaltered independent test set, the model again demonstrated strong recall for the positive class, with sensitivity 88% [68–97] and NPV 95% [85–99]. Specificity was lower (49% [39–59]) with a corresponding PPV of 28% [18–40]. Confidence intervals for this evaluation were calculated using Clopper–Pearson exact binomial estimation, which is appropriate for small sample sizes and rare events. This was necessary given the pronounced class imbalance (106 negative vs. 24 positive cases). The wide confidence interval observed for sensitivity likely reflects the limited number of positive cases (24, 18% of the test set).

Overall, Dharma demonstrated strong prognostic performance for complication prediction, with high sensitivity and NPV supporting its role as a reliable screening tool. Detailed prognostic results are summarized in Table 13.

Table 13. Dharma's Performance for Severity Assessment				
Dharma	**Sensitivity**	**NPV**	**Specificity**	**PPV**
10-fold cross-validation	96% [93-99]	97% [94-99]	65% [53-75]	65% [59-72]
Unaltered Test Set (n=130)	88% [68-97]	95% [85-99]	49% [39-59]	28% [18-40]

Benchmarking Analyses

Diagnostic Value

We compared Dharma, the Imputer–Diagnostic Classifier, against existing clinical scoring systems, ultrasonography (USG), and two state-of-the-art models suited for medium-sized tabular data (XGBoost and LightGBM). Evaluation was performed using 5,555 bootstrap samples of the curated test set (n = 289) and the split test set.

Dharma consistently outperformed all existing scoring systems across every key metric (Table 14). Against SOTA models, Dharma was slightly inferior in the split test set but showed a slight edge in the curated test set (Table 15). Taken together, these findings suggest Dharma performs

474 on par with contemporary machine-learning benchmarks while clearly surpassing conventional

475 diagnostic tools.

Table 14. Dharma minus Conventional tools on curated test set (n=289)						
Tool	AUC-ROC	Accuracy	Sensitivity	Specificity	PPV	NPV
AS (≥5)	0.22 [0.16–0.27]	0.23 [0.17–0.28]	0.06 [0.01–0.11]	0.53 [0.42–0.63]	0.25 [0.19–0.31]	0.23 [0.12–0.34]
AS (≥7)	0.22 [0.16–0.27]	0.29 [0.24–0.35]	0.37 [0.30–0.44]	0.15 [0.07–0.23]	0.13 [0.07–0.20]	0.37 [0.30–0.44]
PAS (≥4)	0.28 [0.22–0.34]	0.23 [0.17–0.29]	0.03 [–0.03–0.09]	0.60 [0.49–0.70]	0.26 [0.21–0.32]	0.22 [0.10–0.34]
PAS (≥6)	0.28 [0.22–0.34]	0.36 [0.30–0.42]	0.44 [0.36–0.52]	0.22 [0.13–0.31]	0.20 [0.13–0.28]	0.43 [0.35–0.50]
AIR (≥5)	0.24 [0.18–0.30]	0.27 [0.21–0.33]	0.25 [0.18–0.32]	0.31 [0.22–0.41]	0.20 [0.14–0.27]	0.35 [0.27–0.43]
AIR (≥9)	0.24 [0.18–0.30]	0.49 [0.43–0.55]	0.76 [0.69–0.82]	–0.01 [–0.04–0.02]	0.02 [–0.02–0.10]	0.48 [0.41–0.54]
Tzanaki	0.07 [0.04–0.10]	0.08 [0.04–0.12]	0.07 [0.03–0.11]	0.10 [0.04–0.16]	0.06 [0.03–0.10]	0.11 [0.05–0.17]
USG	0.04 [0.00–0.09]	0.01 [0.00–0.03]	0.01 [0.00–0.03]	0.00 [0.00–0.00]	0.00 [0.00–0.00]	0.04 [0.00–0.10]

Table 15. Dharma minus SOTA models for diagnostic task in two test sets				
	Split test set (n=156)		Curated test set (n=289)	
Metrics	XGBoost	LightGBM	XGBoost	LightGBM
AUC-ROC	0.00[-0.01-0.00]	0.00[-0.01-0.00]	0.01[0.00-0.02]	0.01[0.00-0.01]
Accuracy	-0.02[-0.04-0]	-0.01[-0.03-0.01]	0.00[-0.01-0.01]	0.00[-0.01-0.01]
Sensitivity	-0.03[-0.07-0]	-0.01[-0.04-0.02]	-0.01[-0.02-0.01]	
Specificity	0.00[0.00-0.00]	0.00[0.00-0.00]	0.01[0.00-0.03]	0.01[0-0.03]
PPV	0.00[-0.01-0]	0.00[0.00-0.00]	0.01[0.00-0.02]	0.01[0-0.02]
NPV	-0.05[-0.11-0]	-0.02[-0.07-0.04]	-0.01[-0.03-0.02]	-0.01[-0.03-0.02]

Prognostic Performance

485 Dharma was marginally inferior to the XGBoost models in key prognostic metrics, namely

486 sensitivity and NPV, but outperformed them markedly in secondary metrics, including accuracy,

487 specificity, and PPV. A detailed comparison is provided in Table 16.

Table 16. Dharma minus SOTA models for prognostic task		
Metric	XGBoost (Simple Hyperparameters)	XGBoost (Complex Hyperparameters)
Sensitivity	–0.01 [–0.05–0.01]	–0.03 [–0.07–0.01]
NPV	–0.01 [–0.04–0.02]	–0.01 [–0.05–0.10]
Specificity	0.18 [0.08–0.27]	0.40 [0.29–0.51]
AUC-ROC	0.01 [–0.01–0.02]	0.00 [–0.01–0.01]
Accuracy	0.10 [0.04–0.15]	0.23 [0.17–0.29]
PPV	0.10 [0.05–0.14]	0.18 [0.14–0.23]

SHAP Analysis

SHAP (Shapley Additive Explanations) analysis was used to interpret the Dharma model's predictions. For the diagnostic model, the SHAP base value—representing the expected model output prior to incorporating feature-specific contributions—was 0.50. In contrast, the severity assessment model exhibited a higher base value of 0.73. This difference reflects an intentional and clinically motivated design choice: the diagnostic model was optimized for balanced discrimination, whereas the severity model, applied exclusively to confirmed appendicitis cases, was deliberately biased toward the positive (complicated) class to minimize the risk of missing potentially complicated disease (false negatives), which carries substantial clinical consequences if surgical intervention is delayed.

In the diagnostic task, appendix diameter exerted the greatest influence on predictions, followed by white blood cell (WBC) count, C-reactive protein (CRP), and neutrophil percentage. For the complications model, CRP had the strongest impact, followed by appendix diameter and peritonitis. Interestingly, larger appendix diameters were associated with lower predicted complication risk, suggesting an inverse relationship between diameter and severity that warrants further exploration.

The absence of appendix diameter data reduced the model's diagnostic probability below the base value; subsequent contributions from other features then shifted the prediction upward to reach the final output. Under these circumstances, peritonitis had the strongest influence, followed by CRP, WBC count, and neutrophil percentage. The global feature importance

patterns for both diagnostic and prognostic tasks are illustrated in the SHAP summary dot plots (S1–S3 Figs).

S1 Fig. SHAP summary plot for Dharma's diagnostic model.

S2 Fig. SHAP summary plot for Dharma's diagnostic model in cases with missing appendix diameter data.

S3 Fig. SHAP summary plot for Dharma's prognostic model.

Discussion

In this study, we present *Dharma*, a novel and clinically grounded machine learning framework specifically designed for pediatric appendicitis. Dharma integrates its native imputer (*Dharma_Imputer*) with a stacked architecture comprising two Random Forest models.

The first model, optimized for unbiased diagnostic performance, achieved an AUC–ROC of 0.98 [0.97–0.99] and an accuracy of 93% [91–95] in identifying acute appendicitis during 10-fold cross-validation on 555 bootstrap samples of the training set. Model performance was well

balanced, with a sensitivity of 93% [91–95], specificity of 93% [90–96], positive predictive

value (PPV) of 95% [93–97], negative predictive value (NPV) of 90% [87–94], and a base

SHAP value of 0.50. This balanced performance was achieved through the use of a relatively

balanced training dataset and hyperparameter tuning with AUC–ROC as the primary

optimization metric, and using class weight as balanced. These design choices reflect the clinical

implications of misclassification in appendicitis diagnosis: false negatives may increase

morbidity by delaying treatment and predisposing patients to complications, whereas false

positives may lead to unnecessary interventions, including negative appendectomy or advanced

but complex imaging in pediatric cohorts.

For patients classified as having acute appendicitis by the diagnostic model, a second Random

Forest model was deployed for prognostic assessment. This severity model was intentionally

optimized to prioritize detection of complicated appendicitis, achieving a sensitivity of 96% [93–

99] and a negative predictive value (NPV) of 97% [94–99], with a specificity of 65% [53–75],

positive predictive value (PPV) of 65% [59–72], and an AUC–ROC of 0.92 [0.90–0.95] based on

cross-validation. The base SHAP value of 0.73 reflects this deliberate operating point favoring

the positive (complicated) class. For the severity assessment task, Dharma was designed to

minimize false negatives, as delayed recognition of complicated appendicitis carries substantial

clinical risk. In contrast, false-positive predictions in this setting primarily lead to appendectomy,

which remains the standard management even for uncomplicated appendicitis. This clinically

motivated optimization was achieved through augmentation of complicated cases from the

existing literature, selection of recall (sensitivity) for the positive class as the primary

hyperparameter optimization metric, and assignment of a 5:1 class weight favoring the positive

class.

Through this approach, Dharma provides a highly reliable and well-calibrated diagnostic tool for

a common yet challenging pediatric emergency, while also functioning as a risk-aware screening

tool for potential complications, supporting safe exclusion and early, evidence-based

management decisions.

To assess its performance, Dharma was compared with established linear scoring systems,

including both older models (Alvarado Score, Pediatric Appendicitis Score [PAS]) and newer

models (Appendicitis Inflammatory Response [AIR] Score, Tzanaki Score), on a common,

previously unseen test cohort (n=289). Each scoring system showed strengths and weaknesses:

for instance, AIR with a high cut-off of 9 demonstrated excellent specificity for appendicitis

diagnosis (99 [97–100]), whereas PAS with a low cut-off of 4 was highly sensitive (89 [84–93]).

Using PAS (≤4) for ruling out and AIR (≥9) for ruling in may provide a simple yet pragmatic

framework for clinical decision-making, which warrants investigation in future studies.

However, none of these scoring systems alone is sufficiently robust for diagnosing acute

appendicitis. Their commonly used intermediate cut-offs have relatively poor specificity (38%–

67%), leading to a high rate of false positives and subsequent negative appendectomies. Similar

patterns were noted in [6], who reported specificities as low as (14.3%–57.1%). These

limitations underscore the need for more accurate, balanced and data-driven non-linear

approaches such as Dharma.

As expected, ultrasonography (USG) demonstrated excellent diagnostic performance in our test

cohort (n=289), with an AUC-ROC of 0.94 [0.89–0.99]. Using a 6 mm appendix diameter cutoff,

sensitivity and specificity reached 95% [87–100] and 95% [87–100], respectively. However, in

34% of cases the appendix could not be visualized. This limitation aligns with prior reports [40-

41], that described non-visualization rates as high as 71–76% in suspected appendicitis. Failure

to identify the appendix may be attributed to anatomical variation, patient habitus, or the

operator-dependent nature of USG [13]. Moreover, advanced features beyond diameter—such as

peri-appendiceal fluid collection (PALC), fat stranding, target sign, hyperemia on Color Doppler,

or free fluid—are typically recognized only by experienced radiologists, who are usually not

available in primary care or resource-limited contexts. Importantly, USG findings must be

integrated with clinical and laboratory parameters for accurate diagnosis [43], and they provide

little prognostic information about disease severity. In LMICs and overcrowded emergency

departments, even timely USG access may be limited, further constraining its practical utility.

Advanced imaging modalities such as CT and MRI provide high diagnostic accuracy for

appendicitis [14–16,18]. However, their use in children is limited by the risks of sedation,

contrast exposure, and ionizing radiation [17]. In addition, these modalities remain largely

inaccessible in many primary and secondary healthcare settings, particularly in resource-limited

regions like ours. Crucially, while they improve diagnostic certainty, they do not reliably

differentiate between uncomplicated and complicated appendicitis—an essential distinction for

guiding timely and appropriate management [20].

Significant progress has been made in applying machine learning to the diagnosis, severity

stratification, and management of pediatric appendicitis [11,24–29]. Yet, many of these studies

are highly technical and AI-centric, which makes their predictions difficult for clinicians to

interpret and incorporate into day-to-day practice. Unlike traditional bedside scoring systems,

machine learning models also require dedicated, user-friendly interfaces to be clinically useful.

For example, the image-based approach [26], although innovative, is computationally intensive

and impractical for real-time decision-making in busy emergency departments. Similarly, the

online tool developed by [11] improved accessibility but remains dependent on complex

ultrasonographic variables requiring expert radiological input. Furthermore, several features

included in their dataset were not directly used in the model's decision process, which

diminishes transparency and clinical applicability. As a result, such tools are better suited to

academic exploration and dataset development rather than to frontline clinical deployment.

Thus, to address the common yet challenging clinical problem of diagnosing and grading acute

appendicitis, we present **Dharma**, a novel non-linear, multimodal framework deployed as a real-

time clinical decision-support web application (dharma-ai.org). Dharma first performs robust

imputation of features frequently missing in resource-constrained settings—such as C-reactive

protein (CRP), appendiceal diameter, free fluid, and urinary ketones—using the **Dharma**

Imputer, a model pre-trained on the appendicitis cohort. The complete feature set is then passed

to the downstream, class-balanced diagnostic model, which outputs a probability score (the

Dharma score) indicating the likelihood of acute appendicitis. For cases predicted as positive, a

second downstream prognostic model—optimized for high recall for the positive (complicated)

class—stratifies the case as *complicated* or *non-complicated*. Both model decisions are

accompanied by corresponding SHAP-based explanations to enhance interpretability and clinical

transparency.

Importantly, Dharma not only substantially surpasses traditional linear scoring systems—as

demonstrated in our benchmarking analyses—but also outperforms all previously published

machine-learning models for both diagnostic classification and severity prediction of

appendicitis [11, 24–29, 39, 48]. Beyond these performance gains, Dharma has been translated

into a fully deployed, real-time clinical decision support system accessible through a functional

web interface. We consider this practical implementation to be a more meaningful contribution

than incremental improvements in model metrics alone, as it directly addresses the translational

gap that limits most existing research models.

When benchmarked against contemporary state-of-the-art machine learning architectures such as

XGBoost and LightGBM, which are designed for handling incomplete non-parametric tabular

datasets, Dharma demonstrated broadly comparable performance across both diagnostic and

prognostic tasks on unseen test sets. Although some differences reached statistical significance,

they were clinically negligible, indicating that Dharma provides comparable discriminative

performance while offering distinct advantages in simplicity, accessibility, interpretability, and

clinical alignment.

Dharma assesses suspected cases of acute appendicitis by analyzing a comprehensive non-linear

pattern of clinical features. These include symptoms such as nausea or vomiting and loss of

appetite; signs like peritonitis; physiological parameters such as body temperature; laboratory

findings including white blood cell (WBC) count, percentage of neutrophils, C-reactive protein

(CRP), and urinary ketones; as well as radiological findings such as appendiceal diameter and

the presence of free fluid on ultrasonography. These features are already well-established in

existing literature for the diagnosis and risk stratification of acute appendicitis [32-37].

While the assessment of signs and symptoms can be particularly challenging in younger children,

certain features remain comparatively more accessible. History-based symptoms such as

nausea/vomiting and poor feeding are generally easier to elicit, even in preverbal children

through caregiver observations. Similarly, signs of peritonitis can often be assessed using

surrogate physical findings, including tenderness, rebound tenderness, guarding, and rigidity—

either localized to the right iliac fossa (RIF) or generalized across the abdomen. Radiological

parameters such as appendiceal diameter and the presence of peritoneal free fluid are also

reliably assessable in most pediatric patients, even with screening ultrasonography or Point-Of-

Care Ultrasound (POCUS), making them particularly valuable in settings where advanced

imaging modalities or experienced radiologists may be unavailable or limited.

Dharma is a medicine-first, hybrid machine learning framework designed by integrating clinical

reasoning into every aspect of its development. From clinically informed feature selection and

data imputation to an unbiased diagnostic model, a risk-averse severity assessment model, and

deployment through a fully functional web application, Dharma follows a structured, stepwise

approach. This workflow demonstrates that clinical reasoning can be directly encoded into

models and systems through careful, medicine-first architectural design, increasing the likelihood

of developing models that generalize effectively, provide explainable outputs, and operate within

real-world clinical constraints [49–50].

A key contribution of this work is the translation of Dharma into a practical, transparent, and

interpretable web-based tool. The platform provides Dharma Score, case-level probability

estimate with 95% confidence intervals derived from the 555 estimators in the diagnostic random

forest, as well as the predicted probability of chances of complications, thereby supporting

surgeons in selecting appropriate treatment modalities. To enhance interpretability, Dharma also

generates feature-level SHAP explanations for both diagnostic and prognostic tasks, allowing

clinicians to understand and trust the model's reasoning. For real-time deployment, the

Dharma_Imputer has been integrated into the web application pipeline to intelligently impute

variables such as CRP, appendix diameter, and free fluid, which are often difficult to assess in

resource-limited settings. Other variables have been made mandatory, as they are relatively

easier to obtain even under such constraints.

Thus, Dharma is not merely another high-performing algorithm but a purpose-built, clinically

grounded system designed for real-world, bedside use, Fig S4. More broadly, we suggest that

future progress in clinical AI may hinge less on incremental algorithmic optimization and more

on the development of systems that are clinically aligned, resource-sensitive, interpretable, and

usable.

S4 Fig. Dharma as a Clinical Decision Support system (CDSS)

We believe that knowledge should be freely available and openly shared. In alignment with this

principle, we have made our source code publicly accessible on GitHub [46], and the web

application is freely accessible at dharma-ai.org , promoting transparency, reproducibility, and

continuous improvement in medical science. Screenshots of the interface are provided in Figures

S4–S6.

S5 Fig. User interface of Dharma's web-based clinical decision support tool.

S6 Fig. Example of diagnostic and prognostic predictions generated by Dharma.

S7 Fig. Feature-level SHAP explanations for individual predictions.

Nevertheless, this study has certain limitations that warrant acknowledgment. Foremost, the lack

of access to large, high-quality medical datasets remains a significant barrier, limiting the full

potential of statistical and machine learning approaches that we believe are essential for

advancing 21st-century medicine [23]. We strongly advocate for the open sharing of anonymized

clinical datasets to accelerate broader and more impactful research. Another limitation is that our

model was developed on single-center data, raising questions about its adaptability across

diverse populations. Furthermore, the absence of exclusively histologically confirmed

appendicitis cases may have constrained the model's ability to fully optimize pattern recognition

in both diagnostic and prognostic tasks. Future work should prioritize multicenter validation—

both prospective and retrospective—including studies in low- and middle-income countries

(LMICs) to rigorously assess Dharma's real-world clinical utility and generalizability. We also

encourage independent researchers and med-tech innovators to validate Dharma further, and

where necessary, calibrate its thresholds to suit the epidemiological and clinical nuances of their

own populations.

Despite these limitations, the Dharma represents more than an incremental step in clinical

decision support. By integrating readily available clinical, laboratory, and radiological findings

into a unified predictive framework, it provides real-time, evidence-based diagnostic and

prognostic support for one of the most common pediatric surgical emergencies. With its high

specificity and positive predictive value, Dharma has the potential to meaningfully reduce

unnecessary negative appendectomies while maintaining patient safety through very low false-

negative rates. In doing so, the model could lessen reliance on advanced imaging such as CT or

MRI, thereby reducing the risks associated with sedation, contrast exposure, and ionizing

radiation in children—offering both clinical and economic benefits.

Importantly, Dharma is designed to function across diverse healthcare settings, from resource-

limited facilities to advanced tertiary centers. In primary and secondary care, where even basic

imaging and surgical expertise may be unavailable, Dharma leverages accessible clinical and

laboratory features to support early recognition, guide timely referrals, and reduce delays in

management. In high-volume tertiary emergency departments, it may help prioritize cases when

imaging is inconclusive or radiology services are saturated. At advanced surgical centers,

calibrated thresholds optimized for high specificity and PPV allow Dharma to be used as a

reliable rule-in tool. Combined with its severity assessment feature, it can also assist in triaging

patients for appendectomy and informing decisions between conservative versus surgical

management.

Taken together, Dharma illustrates how data-driven tools can complement, rather than replace,

clinical judgment. By bridging bedside intuition with machine learning–derived insights, it

provides a pathway toward equitable, AI-augmented care. While prospective multicenter validation remains essential, this work underscores the potential of interpretable, accessible, medicine-first, AI-powered, clinical decision support systems—not only for pediatric appendicitis but also as a clinically grounded framework adaptable to broader healthcare challenges.

Acknowledgement

We sincerely thank our dear friend, Amrit Neupane, for his continued support and invaluable contributions to the UI/UX and logo design of Dharma. We also extend our gratitude to Marcinkevics and team for making their anonymized datasets publicly available, which greatly facilitated this work.

Author Contributions

Conceptualization and study design: Anup Thapa Kshetri, Subash Pahari

Data curation: Anup Thapa Kshetri, Subash Pahari

Literature review: Anup Thapa Kshetri

Formal analysis: Anup Thapa Kshetri

Methodology: Anup Thapa Kshetri

Software: Anup Thapa Kshetri (Machine Learning Framework and Web-app Backend

Development), Subash Pahari (web-app development and deployment)

Writing – original draft: Anup Thapa Kshetri, Sashank Timilsina, Binay Chapagain

Writing – review & editing: Anup Thapa Kshetri, Subash Pahari

Writing – Manuscript Formating: Subash Pahari

References

[1] Hijaz NM, Friesen CA. Managing acute abdominal pain in pediatric patients: current

perspectives. Pediatric Health Med Ther. 2017 Jun 29;8:83-91. doi: 10.2147/PHMT.S120156.

PMID: 29388612; PMCID: PMC5774593.

[2] Tseng YC, Lee MS, Chang YJ, Wu HP. Acute abdomen in pediatric patients admitted to the

pediatric emergency department. Pediatr Neonatol. 2008 Aug;49(4):126-34. doi: 10.1016/S1875-

9572(08)60027-3. PMID: 19054918.

[3] Schneider C, Kharbanda A, Bachur R. Evaluating appendicitis scoring systems using a

prospective pediatric cohort. Ann Emerg Med. 2007 Jun;49(6):778-84, 784.e1. doi:

10.1016/j.annemergmed.2006.12.016. Epub 2007 Mar 26. PMID: 17383771.

[4] Sencan A, Aksoy N, Yıldız M, Okur Ö, Demircan Y, Karaca I. The evaluation of the validity

of Alvarado, Eskelinen, Lintula and Ohmann scoring systems in diagnosing acute appendicitis in

children. Pediatr Surg Int. 2014 Mar;30(3):317-21. doi: 10.1007/s00383-014-3467-0. PMID:

24448910.

[5] Pogorelić Z, Rak S, Mrklić I, Jurić I. Prospective validation of Alvarado score and Pediatric

Appendicitis Score for the diagnosis of acute appendicitis in children. Pediatr Emerg Care. 2015

Mar;31(3):164-8. doi: 10.1097/PEC.0000000000000375. PMID: 25706925.

[6] Sağ S, Basar D, Yurdadoğan F, Pehlivan Y, Elemen L. Comparison of Appendicitis Scoring

Systems in Childhood Appendicitis. Turk Arch Pediatr. 2022 Sep;57(5):532-537. doi:

10.5152/TurkArchPediatr.2022.22076. PMID: 36062441; PMCID: PMC9524470.

[7] Jukić M, Nizeteo P, Matas J, Pogorelić Z. Trends and Predictors of Pediatric Negative

Appendectomy Rates: A Single-Centre Retrospective Study. Children (Basel). 2023 May

15;10(5):887. doi: 10.3390/children10050887. PMID: 37238435; PMCID: PMC10217643.

[8] Kundiona I, Chihaka OB, Muguti GI. Negative appendicectomy: evaluation of

ultrasonography and Alvarado score. Cent Afr J Med. 2015 Sep-Dec;61(9-12):66-73. PMID:

29144064.

[9] Oyetunji TA, Ong'uti SK, Bolorunduro OB, Cornwell EE 3rd, Nwomeh BC. Pediatric

negative appendectomy rate: trend, predictors, and differentials. J Surg Res. 2012

Mar;173(1):16-20. doi: 10.1016/j.jss.2011.04.046. Epub 2011 May 19. PMID: 21696768.

[10] Patel H, Kamel M, Cooper E, Bowen C, Jester I. The Variable Definition of "Negative

Appendicitis" Remains a Surgical Challenge. Pediatr Dev Pathol. 2024 Nov-Dec;27(6):552-558.

doi: 10.1177/10935266241255281. Epub 2024 Jun 6. PMID: 38845117.

[11] Marcinkevics R, Reis Wolfertstetter P, Wellmann S, Knorr C, Vogt JE. Using Machine

Learning to Predict the Diagnosis, Management and Severity of Pediatric Appendicitis. Front

Pediatr. 2021 Apr 29;9:662183. doi: 10.3389/fped.2021.662183. PMID: 33996697; PMCID:

PMC8116489.

[12] Bălănescu L, Băetu AE, Cardoneanu AM, Moga AA, Bălănescu RN. Predictors of

Complicated Appendicitis with Evolution to Appendicular Peritonitis in Pediatric Patients.

Medicina (Kaunas). 2022 Dec 22;59(1):21. doi: 10.3390/medicina59010021. PMID: 36676645;

PMCID: PMC9866196.

[13] Callahan MJ, Rodriguez DP, Taylor GA. CT of appendicitis in children. Radiology. 2002

Aug;224(2):325-32. doi: 10.1148/radiol.2242010998. PMID: 12147823.

[14] Kim DW, Yoon HM, Lee JY, Kim JH, Jung AY, Lee JS, et al. Diagnostic performance of

CT for pediatric patients with suspected appendicitis in various clinical settings: a systematic

review and meta-analysis. Emerg Radiol. 2018 Dec;25(6):627-637. doi: 10.1007/s10140-018-

1624-9. Epub 2018 Jul 12. PMID: 30003463.

[15] Sivit CJ, Applegate KE, Stallion A, Dudgeon DL, Salvator A, Schluchter M, et al. Imaging

evaluation of suspected appendicitis in a pediatric population: effectiveness of sonography

versus CT. AJR Am J Roentgenol. 2000 Oct;175(4):977-80. doi: 10.2214/ajr.175.4.1750977.

PMID: 11000147.

[16] Stephen AE, Segev DL, Ryan DP, Mullins ME, Kim SH, Schnitzer JJ, et al. The diagnosis

of acute appendicitis in a pediatric population: to CT or not to CT. J Pediatr Surg. 2003

Mar;38(3):367-71; discussion 367-71. doi: 10.1053/jpsu.2003.50110. PMID: 12632351.

[17] Brennan GD. Pediatric appendicitis: pathophysiology and appropriate use of diagnostic

imaging. CJEM. 2006 Nov;8(6):425-32. doi: 10.1017/s1481803500014238. PMID: 17209492.

[18] Mittal MK. Appendicitis: Role of MRI. Pediatr Emerg Care. 2019 Jan;35(1):63-66. doi:

10.1097/PEC.0000000000001710. PMID: 30608328.

[19] Pines JM. Trends in the rates of radiography use and important diagnoses in emergency

department patients with abdominal pain. Med Care. 2009 Jul;47(7):782-6. doi:

10.1097/MLR.0b013e31819748e9. PMID: 19536032.

[20] Lastunen K, Leppäniemi A, Mentula P. Perforation rate after a diagnosis of uncomplicated

appendicitis on CT. BJS Open. 2021 Jan 8;5(1):zraa034. doi: 10.1093/bjsopen/zraa034. PMID:

33609386; PMCID: PMC7893470.

[21] Almaramhy HH. Acute appendicitis in young children less than 5 years: review article. Ital J

Pediatr. 2017 Jan 26;43(1):15. doi: 10.1186/s13052-017-0335-2. PMID: 28257658; PMCID:

PMC5347837.

[22] Pogorelić Z, Domjanović J, Jukić M, Poklepović Peričić T. Acute Appendicitis in Children

Younger than Five Years of Age: Diagnostic Challenge for Pediatric Surgeons. Surg Infect

(Larchmt). 2020 Apr;21(3):239-245. doi: 10.1089/sur.2019.175. Epub 2019 Oct 16. PMID:

31618143.

[23] Handelman GS, Kok HK, Chandra RV, Razavi AH, Lee MJ, Asadi H. eDoctor: machine

learning and the future of medicine. J Intern Med. 2018 Dec;284(6):603-619. doi:

10.1111/joim.12822. Epub 2018 Sep 3. PMID: 30102808.

[24] Harmantepe AT, Dikicier E, Gönüllü E, Ozdemir K, Kamburoğlu MB, Yigit M. A different

way to diagnosis acute appendicitis: machine learning. Pol Przegl Chir. 2023 Oct 13;96(2):38-

43. doi: 10.5604/01.3001.0053.5994. PMID: 38629278.

[25] Reismann J, Romualdi A, Kiss N, Minderjahn MI, Kallarackal J, Schad M, et al. Diagnosis

and classification of pediatric acute appendicitis by artificial intelligence methods: An

investigator-independent approach. PLoS One. 2019 Sep 25;14(9):e0222030. doi:

10.1371/journal.pone.0222030. PMID: 31553729; PMCID: PMC6760759.

[26] Byun J, Park S, Hwang SM. Diagnostic algorithm based on machine learning to predict

complicated appendicitis in children using CT, laboratory, and clinical features. Diagnostics

(Basel). 2023;13(5):923. doi: 10.3390/diagnostics13050923. PMID: 36900066; PMCID:

PMC10001049.

[27] Forsström JJ, Irjala K, Selén G, Nyström M, Eklund P. Using data preprocessing and single

layer perceptron to analyze laboratory data. Scand J Clin Lab Invest Suppl. 1995;222:75-81. doi:

10.3109/00365519509088453. PMID: 7569750.

[28] Grigull L, Lechner WM. Supporting diagnostic decisions using hybrid and complementary

data mining applications: a pilot study in the pediatric emergency department. Pediatr Res. 2012

Jun;71(6):725-31. doi: 10.1038/pr.2012.34. Epub 2012 Mar 22. PMID: 22441377.

[29] Chadaga K, Khanna V, Prabhu S, Sampathila N, Chadaga R, Umakanth S, et al. An

interpretable and transparent machine learning framework for appendicitis detection in pediatric

patients. Sci Rep. 2024 Oct 18;14(1):24454. doi: 10.1038/s41598-024-75896-y. Erratum in: Sci

Rep. 2025 Jan 22;15(1):2841. doi: 10.1038/s41598-025-86494-x. PMID: 39424647; PMCID:

PMC11489819.

[30] Hua R, O'Brien MK, Carter M, Pitt JB, Kwon S, Ghomrawi HMK, et al. Improving Early

Prediction of Abnormal Recovery after Appendectomy in Children using Real-world Data from

Wearables. Annu Int Conf IEEE Eng Med Biol Soc. 2024 Jul;2024:1-4. doi:

10.1109/EMBC53108.2024.10782031. PMID: 40039430.

[31] Ghomrawi HMK, O'Brien MK, Carter M, Macaluso R, Khazanchi R, Fanton M, et al.

Applying machine learning to consumer wearable data for the early detection of complications

after pediatric appendectomy. NPJ Digit Med. 2023 Aug 16;6(1):148. doi: 10.1038/s41746-023-

00890-z. PMID: 37587211; PMCID: PMC10432429.

[32] Pop GN, Costea FO, Lungeanu D, Iacob ER, Popoiu CM. Ultrasonographic findings of

child acute appendicitis incorporated into a scoring system. Singapore Med J. 2022 Jan;63(1):35-

41. doi: 10.11622/smedj.2020102. Epub 2020 Jul 15. PMID: 32668829; PMCID: PMC9251212.

[33] Adir A, Braester A, Natalia P, Najib D, Akria L, Suriu C, et al. The role of blood

inflammatory markers in the preoperative diagnosis of acute appendicitis. Int J Lab Hematol.

2024 Feb;46(1):58-62. doi: 10.1111/ijlh.14163. Epub 2023 Aug 29. PMID: 37644670.

[34] Moris D, Paulson EK, Pappas TN. Diagnosis and Management of Acute Appendicitis in

Adults: A Review. JAMA. 2021 Dec 14;326(22):2299-2311. doi: 10.1001/jama.2021.20502.

PMID: 34905026.

[35] Yajima H, Fujita T, Yanaga K. Profile of signs and symptoms in mild and advanced acute

appendicitis. Int Surg. 2010 Jan-Mar;95(1):63-6. PMID: 20480844.

[36] Hajibandeh S, Hajibandeh S, Hobbs N, Mansour M. Neutrophil-to-lymphocyte ratio

predicts acute appendicitis and distinguishes between complicated and uncomplicated

appendicitis: A systematic review and meta-analysis. Am J Surg. 2020 Jan;219(1):154-163. doi:

10.1016/j.amjsurg.2019.04.018. Epub 2019 Apr 27. PMID: 31056211.

[37] Biehl CM, Elliver M, Gudjonsdottir J, Salö M. Utility of Urine Dipstick Testing in Pediatric

Appendicitis: Assessing its Role in Identifying Complicated Cases and Retrocecal Appendicitis.

Eur J Pediatr Surg. 2024 Dec 19. doi: 10.1055/a-2490-1156. Epub ahead of print. PMID:

39701137.

[38] Singh C. Machine learning in pattern recognition. Eur J Eng Technol Res. 2023

Apr;8(2):63-68. doi:10.24018/ejeng.2023.8.2.3025.

[39] Issaiy M, Zarei D, Saghazadeh A. Artificial Intelligence and Acute Appendicitis: A

Systematic Review of Diagnostic and Prognostic Models. World J Emerg Surg. 2023 Dec

19;18(1):59. doi: 10.1186/s13017-023-00527-2. PMID: 38114983; PMCID: PMC10729387.

[40] Scrimgeour DS, Driver CP, Stoner RS, King SK, Beasley SW. When does ultrasonography

influence management in suspected appendicitis? ANZ J Surg. 2014 May;84(5):331-4. doi:

10.1111/ans.12415. Epub 2014 Jan 9. PMID: 24405944.

[41] Held JM, McEvoy CS, Auten JD, Foster SL, Ricca RL. The non-visualized appendix and

secondary signs on ultrasound for pediatric appendicitis in the community hospital setting.

Pediatr Surg Int. 2018 Dec;34(12):1287-1292. doi: 10.1007/s00383-018-4350-1. Epub 2018 Oct

6. PMID: 30293146.

[42] Nijssen DJ, van Amstel P, van Schuppen J, Eeftinck Schattenkerk LD, Gorter RR, Bakx R.

Accuracy of ultrasonography for differentiating between simple and complex appendicitis in

children. Pediatr Surg Int. 2021 Jul;37(7):843-849. doi: 10.1007/s00383-021-04872-8. Epub

2021 Mar 7. PMID: 33677613; PMCID: PMC8172400.

[43] Mills LD, Mills T, Foster B. Association of clinical and laboratory variables with ultrasound

findings in right upper quadrant abdominal pain. South Med J. 2005 Feb;98(2):155-61. doi:

10.1097/01.SMJ.0000129927.88863.65. PMID: 15759944.

[44] Marcinkevičs R, Reis Wolfertstetter P, Klimiene U, Ozkan E, Chin-Cheong K, Paschke A,

et al. Regensburg Pediatric Appendicitis Dataset [Internet]. Zenodo; 2023 [cited 2025 May 26].

Available from: https://zenodo.org/records/7669442

[45] Steiner Z, Buklan G, Stackievicz R, Gutermacher M, Litmanovitz I, Golani G, et al.

Conservative treatment in uncomplicated acute appendicitis: reassessment of practice safety. Eur

J Pediatr. 2017 Apr;176(4):521-527. doi: 10.1007/s00431-017-2867-2. Epub 2017 Feb 16.

PMID: 28210834.

[46] Thapa A. Dharma-AppendicitisModel [Internet]. 2025 [cited 2025 May 26]. Available from:

https://github.com/ajung17/Dharma-AppendicitisModel

[47] Marcinkevičs R, Sokol K, Paulraj A, Hilbert MA, Rimili V, Wellmann S, et al. External

Validation of Predictive Models for Diagnosis, Management and Severity of Pediatric

Appendicitis   [Preprint].   medRxiv.   2024   Oct   28;2024.10.28.24316300.

doi:10.1101/2024.10.28.24316300.

[48] Erman A, Ferreira J, Ashour WA, Guadagno E, St-Louis E, Emil S, et al. Machine-learning-

assisted Preoperative Prediction of Pediatric Appendicitis Severity. J Pediatr Surg. 2025

Jun;60(6):162151. doi: 10.1016/j.jpedsurg.2024.162151. Epub 2025 Jan 13. PMID: 39855986.

[49] Samadi ME, Guzman-Maldonado J, Nikulina K, Mirzaieazar H, Sharafutdinov K, Fritsch

SJ, et al. A hybrid modeling framework for generalizable and interpretable predictions of ICU

mortality across multiple hospitals. Sci Rep. 2024 Mar 8;14(1):5725. doi: 10.1038/s41598-024-

55577-6. PMID: 38459085; PMCID: PMC10923850.

[50] Alonge M, Isreal O. Hybrid AI Models: Combining Machine Learning with Domain

Knowledge [Internet]. 2025 [cited 2025 May 26]. Available from

https://www.researchgate.net/publication/392166442_Hybrid_AI_Models_Combining_Machine

_Learning_with_Domain_Knowledge

[51] Cutillo CM, Sharma KR, Foschini L, et al. Machine intelligence in healthcare—perspectives

on trustworthiness, explainability, usability, and transparency. NPJ Digit Med. 2020;3:47.

doi:10.1038/s41746-020-0254-2.

[52] Rajpurkar P, Chen E, Banerjee O, et al. AI in health and medicine. Nat Med. 2022;28:31-38.

doi: 10.1038/s41591-021-01614-0.

922 **Legends:**

Figure S1. SHAP summary plot for Dharma's diagnostic model.

Figure S2. SHAP summary plot for Dharma's diagnostic model in patients with missing appendix diameter data.

Figure S3. SHAP summary plot for Dharma's prognostic model.

Figure S4. Dharma as a Clinical Decision Support system (CDSS).

Figure S5. User interface of Dharma's web-based clinical decision support tool. Accessible through dharma-ai.org

Figure S6. Example of diagnostic and prognostic predictions generated by Dharma. Dharma score and severity risk estimation, combined into a decision framework.

Figure S7. Feature-level SHAP explanations for both the diagnostic and prognostic estimations.

Table S1. Percentage of missing values in each feature.

Table S2. Performance stability of Dharma with different imputation strategies.

NAUSEA

Select option ⌄

LOSS OF APPETITE

Select option ⌄

PERITONITIS

Select option ⌄

WBC COUNT × 10^3 (per µL)

10.0

BODY TEMPERATURE (°C)

38.0

NEUTROPHIL PERCENTAGE (%)

75.0

CRP (mg/L)

0.0

KETONES IN URINE

Select option ⌄

APPENDIX DIAMETER (MM)

CRP (mg/L)

KETONES IN URINE

No

APPENDIX DIAMETER (MM)

FREE FLUIDS

No

PREDICTION

EXPLAINATION

DHARMA SCORE: 97%

95% Confidence Interval: 96% - 98%
Result: Very high likelihood of acute appendicitis.
Note: Management in line with acute appendicitis is recommended.

Severity Assessment

Probability: 35%
95% Confidence Interval: 34% - 37%
Result: Low likelihood of developing complications.
Note: Conservative management can be considered.

Prediction Explanation

Diagnosis Explanation

Base Value: 0.50

Feature	SHAP Value
Appendix_Diameter	0.3058
Peritonitis	0.0900
CRP	0.0501
Neutrophil_Percentage	0.0396
Nausea	0.0117
Loss_of_Appetite	0.0016
Ketones_in_Urine	0.0010
Free_Fluids	-0.0074
WBC_Count	-0.0438
Final Prediction	0.95

Complication Explanation

Base Value: 0.65

Feature	SHAP Value
CRP	0.2197
Loss_of_Appetite	0.0509
Peritonitis	0.0457
Body_Temperature	0.0241
Nausea	0.0059
Free_Fluids	-0.0014
Ketones_in_Urine	-0.0047
WBC_Count	-0.0923
Appendix_Diameter	-0.0992
Final Prediction	0.80

Download Full Report