Associations of Lisinopril/Candesartan, Anti-CGRP Treatments vs. Topiramate on Cardiovascular Outcomes in Patients with Migraine

Kaicheng Wang, MD MPH [1,2], Brenda T. Fenton, PhD [3], John P. Ney, MD MPH [4, 5], Vinh X. Dao, PharmD [6], Alexander B. Guirguis, PharmD [4, 7], Emmanuelle A. Schindler, MD PhD [4, 5], Adam de Havenon, MD MS [5], Melissa Skanderson, MSW [1], Sarah E. Anthony, MS [1], Laura J. Burrone, BS [1], Jason J. Sico, MD MHS [1,5,8]

[1] Research, Education, Evaluation and Engagement Activities Center for Headache, VA Connecticut Healthcare System, Orange, Connecticut
[2] Yale Center for Analytical Sciences, Yale School of Public Health, New Haven, Connecticut
[3] Pain Research, Informatics, Multi-Morbidities, and Education Center, VA Connecticut Healthcare System, West Haven, Connecticut
[4] Headache Center of Excellence, VA Connecticut Healthcare System, West Haven, Connecticut
[5] Department of Neurology, Yale School of Medicine, New Haven, Connecticut
[6] Headache Center of Excellence, VA Minneapolis Health Care System, Minneapolis, Minnesota
[7] Pharmacy Benefits Management Services, VA Connecticut Healthcare System, West Haven, Connecticut
[8] Department of Internal Medicine, Yale School of Medicine, New Haven, Connecticut

Corresponding Author
Kaicheng Wang, MD MPH
689-VA Connecticut Healthcare System
VA Annex Orange Campus – National Program Center
200 Edison Road
Orange, CT 06477
kaicheng.wang@va.gov

ABSTRACT

Background and Objectives

Migraine is a prevalent neurological condition and is associated with an increased risk of myocardial infarction (MI) and ischemic stroke. However, the cardiovascular risks or benefits of specific migraine preventive medications remain unknow. The objective of this study is to compare the risk of major adverse cardiovascular events (MACE) among patients with migraine disorder initiating lisinopril/candesartan or anti-calcitonin gene-related peptide (aCGRP) treatments versus topiramate.

Methods

Two target trials were emulated comparing patients who were prescribed lisinopril/candesartan or aCGRP treatment to those receiving topiramate for migraine prevention within the U.S. Department of Veterans Affairs between June 1, 2018 and September 30, 2024. Outcome was MACE including MI, ischemic stroke, intracerebral and subarachnoid hemorrhage (ICH/SAH) and all-cause mortality. Five-year cumulative incidences, risk differences, risk ratios and overall hazard ratios (HRs) were estimated.

Results

Among 48,610 person-trials initiating lisinopril/candesartan (mean [SD] age, 52.3 [12.3] years; 18.9% women) and 25,635 initiating aCGRP treatment (mean [SD] age, 47.2 [12.2] years; 39.4% women), lisinopril/candesartan was associated with an increased risk of MACE (HR 1.21; 95%CI, 1.10-1.34), particularly MI (HR 1.28; 95%CI, 1.03-1.59). The MI risk was greater in patients without documented cardiovascular indications (HR 2.75; 95%CI, 2.01-3.77), patients aged <40 years (HR 3.18; 95%CI, 1.63-6.20), or with baseline systolic blood pressure (SBP) <130 mm Hg (HR 1.86; 95%CI, 1.44-2.39). aCGRP use was associated with a reduced risk of MI (HR 0.80; 95% CI, 0.62-1.02) and was not associated with ischemic stroke (HR 1.11; 95% CI, 0.84-1.49) or ICH/SAH (HR 0.95; 95% CI, 0.65-1.38).

Discussion

Lisinopril/candesartan may increase the risk of MACE or MI in patients with migraine disorder, particularly in those with minimal atherosclerotic risks. aCGRP treatment appears safe regarding cardiovascular events and potentially protective against MI. Future research is needed to confirm these findings before clinical recommendations can be made.

Introduction

Migraine is a prevalent and incapacitating neurological disorder that affects 15.9% of adults in the United States.[1] Migraine patients are predisposed to an increased risk of myocardial infraction (MI) and ischemic stroke,[2,3] a susceptibility that poses challenges in selecting effective migraine treatments while accounting for cardiovascular risks.[4] Conventional migraine abortive and preventive medications are often associated with considerable adverse effects, including ischemic events. For instance, triptan use is associated with at least a threefold risk of major adverse cardiovascular events (MACE)[5,6], and tricyclic antidepressants are associated with a 20% higher risk of MACE in patients with chronic pain[7]. Although trials of anti-calcitonin gene-related peptide (aCGRP) treatments have reported fewer adverse events, the long-term cardiovascular safety profile of these treatments remains uncertain. Preclinical studies suggest that CGRP may exhibit vasodilation and cardioprotective properties[8], but real-world studies on the impact of aCGRP treatments on blood pressure have been equivocal.[9]

Despite the poor understanding on the pathophysiology of migraine-associated cardiovascular risks, angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) may offer a dual benefit for patients with migraine and cardiovascular risks. ACEIs/ARBs have been shown to reduce the composite risk of all-cause and cardiovascular-specific mortality, MI and stroke by 11% in patients with or at risk for atherosclerotic disease regardless of systolic blood pressure (SBP)[10]. In particular, lisinopril (Class C[11]; weak[12]) and candesartan (Class C[11]; strong[12]) are recommended for migraine prevention by clinical guidelines, making them a potential option for primary prevention of MACE in patients with migraine.

In this study, we evaluated the impact of two classes of migraine preventive medications on the risk of MACE among patients with migraine. We hypothesized that initiating lisinopril/candesartan may reduce the risk of MACE, whereas initiating aCGRP treatments may increase the risk compared to topiramate.

Methods
Study design and participants

This retrospective cohort study utilized data from the U.S. Department of Veterans Affairs (VA) Headache Centers of Excellence Administrative Data Cohort.[13] The source population included incident migraine patients between Oct 1, 2007 and Sep 30, 2023. From this population, we emulated two target trials analyzing patients who were (i) prescribed lisinopril/candesartan or (ii) aCGRP treatment to those receiving topiramate for migraine prevention between June 1, 2018 and September 30, 2024. This study was approved by the VA Connecticut Healthcare System Research & Development Committee and granted a waiver of informed consent (IRBNet# 1784787). The study design followed the PRINCIPLED guideline[14], with reporting follows the STROBE guideline for cohort studies. A detailed statistical analysis plan (SAP) is provided in *eSAP*.

Key protocol components of the target trials were outlined in the *Supplement Table 1*. Migraine patients were identified using International Classification of Diseases 9[th] and 10[th] versions codes. Exclusion criteria included prior use of study medications before migraine diagnosis, use within the past 3 months, or no primary care visit in the past 12 months. For lisinopril/candesartan vs. topiramate target trial, patients with prior use of other ACEIs (e.g., enalapril) or ARBs (e.g., losartan), or with concurrent prescriptions of topiramate and any ACEI or ARB, were excluded. Similarly, for aCGRP vs. topiramate target trial, patients with concurrent abortive/preventive aCGRP and topiramate were excluded. All treatment episodes (person-trials) that met the eligibility criteria were included, with baseline defined as the date of the first prescription dispensed.

Data sources

The primary data source was the VA Corporate Data Warehouse (CDW) which contains structured electronic health records, including patient demographics, vital status, inpatient and outpatient encounters, laboratory, pharmacy dispensing data and provider information since October 1, 1999. To capture healthcare occurred outside the VA, we included Medicare and Medicaid claims from October 1, 2007, to December 31, 2022, and VA Community Care Data up to September 30, 2024. Cause of death data were not included, as they were only available through December 31, 2021.

Treatment strategies

Details of medications are provided in *Supplement Table 2*. Topiramate, a migraine preventive agent that has not demonstrated effects on cardiovascular events[15,16] was used as the comparator. Galcanezumab (100 mg/mL) for cluster headache, topiramate/phentermine for weight management, and lisinopril/candesartan combinations (e.g., lisinopril and hydrochlorothiazide indicated for hypertension) were not included. Rimegepant for migraine prevention was defined as ≥12 tablets dispensed per 30 days. We defined adherence as receiving a refill within 90 days after the end of supply. The medication possession ratio (MPR) was calculated by summing of days' supply of all prescriptions and divided by the number of days from baseline to the end of treatment and truncated at the maximum value of 1.0.

Outcomes

The primary outcome was MACE, defined as a composite of MI, ischemic stroke, intracerebral and subarachnoid hemorrhage (ICH/SAH), and all-cause mortality. Secondary outcomes were MI, ischemic stroke, ICH/SAH, respectively. Patients were followed from the baseline and ended at the occurrence of an event, 60 months after the baseline, or administrative end of the study by September 30, 2024, whichever happened first.

Covariates

We used a priori selection strategy[17] to identify potential confounders based on subject-matter knowledge, literature and a directed acyclic graph (*Supplement Figure 1a*). We assumed treatment strategies were exchangeable conditional on baseline covariates, including demographics, social determinants of health (SDOH), healthcare-seeking behavior, cardiovascular risk factors and related medications, migraine severity and treatments (*Supplement Table 3*). Missing data for continuous covariates were imputed with median values, and categorical variables with missing data were coded as "Unknown".

Statistical analysis

Baseline characteristics were summarized as frequencies and percentages, means and standardized deviations (SDs), or medians and interquartile ranges (IQRs) for non-normally distributed variables. Both intention-to-treat and per-protocol analyses were performed. Our

approach to interpreting data is based on an evaluation of the magnitude, direction, and precision of the effect estimates rather than binary significance testing.[18] Data were managed using SQL Server Management Studio 19.0, and statistical analyses were conducted using R version 4.3.1. All SQL queries and analytical code are available at https://github.com/kw375/aCGRP_CV/.

The intention-to-treat analyses examined the association between initiating lisinopril/candesartan or aCGRP treatments versus topiramate on the risk of MACE. Observational analogs were estimated by adjusting for baseline confounders using stabilized inverse probability of treatment initiating weights (IPTiW) via logistic regression models (details in SAP). Covariate balance was assessed using standardized mean difference (SMD), with a value less than 0.1 indicating good balance between comparisons. Next, cumulative incidence curves were estimated using IPTiW-weighted pooled logistic regressions, with an indicator for treatment strategy, a flexible time-varying intercept, and a treatment-by-time interaction term. Five-year risk differences (RDs) and risk ratios (RRs) were compared. Confidence intervals (CIs) for cumulative incidence rates, RDs, and RRs were computed nonparametrically using 500 bootstrapping samples. The overall hazard ratios (HRs) over five years were obtained using pooled logistic regressions without the interaction term, with robust sandwich estimator for standardized errors.

For per-protocol analyses, observations not adherent to their treatment strategies were censored. Stabilized inverse probability of treatment adherence weights (IPTaW) were estimated to account for potential selection bias associated with non-adherence, using logistical regression models additional adjusted for time-updated covariates, including body mass index, blood pressures, hospitalizations, diagnoses of chronic kidney disease (CKD), heart failure (HF), hypertension or chronic migraine, and medications (β-blockers, diuretics, antiplatelets, triptans, and non-steroid anti-inflammatory drugs [NSAIDs]). Final weights for each person-trial-month were calculated as the product of IPTiW and cumulative IPTaW up to that month. Due to computational constraints, only overall HRs were estimated via weighted pooled logistic regression.

To assess the robustness of intention-to-treat estimates, we conducted the following sensitivity analyses. (1) To address potential confounding by indication[19], patients having cardiovascular indications (CKD, HF, hypertension, or history of MI) were excluded in the

lisinopril/candesartan target trial, and (2) further stratified by baseline age and SBP in the overall sample and among those without documented cardiovascular indications. (3) We repeated the primary analyses using outcome regressions to estimate the overall HRs, adjusting for the exact set of baseline confounders used in the IPTiW models, (4) and additionally estimated the sub-distribution hazards to account for competing risks from death for secondary outcomes.

We also conducted three quantitative bias analyses. We (1) computed E-values[20] based on the estimation from outcome regressions to quantify the magnitude unmeasured confounding have to be to negate the observed associations, and then (2) conducted probabilistic bias analyses specified in the *Supplement Methods* to assess the impact of residual confounding by indication from an unknown baseline binary confounder. (3) To detect spurious associations arising from selection bias, misclassification, residual confounding, and statistical analysis, we repeated the analyses on both positive and negative outcome controls. Given the well-documented efficacy of topiramate for weight loss[21] and findings from the Food and Drug Administration Adverse Event Reporting System[22], we used weight change and primary malignancy as positive outcome controls. Influenza and non-pathological fracture, which have no known etiologic or pathophysiologic links to lisinopril/candesartan and aCGRP treatment, were selected as negative outcome controls[23,24] to assess residual confounding from healthcare-seeking behavior and frailty. Influenza diagnosis depends on vaccination uptake and symptom-driven testing, making it a proxy for healthcare-seeking behavior. Similarly, non-pathological fractures are often associated with frailty, which in turn is associated with cardiovascular events.

Results

The study included 48,610 person-trials that initiated lisinopril/candesartan vs. 115,047 person-trials initiated topiramate, 25,635 person-trials that initiated aCGRP treatment vs. 135,740 person-trials that initiated topiramate (*Table 1 & Figure 1*).

Lisinopril/Candesartan versus *Topiramate Target Trial*

Patients initiating lisinopril/candesartan were older (mean [SD], 52.3 [12.3] vs. 43.2 [11.2] years), with fewer women (18.9% vs. 40.0%), Black (23.3% vs. 35.1%) and Hispanic individuals (9.8%

vs. 12.8%). These patients had higher SBP (136.4 [16.6] vs. 123.8 [13.4] mm Hg) and cardiovascular risks at baseline, including hypertension (76.3% vs. 23.8%), diabetes (31.4% vs. 8.2%), dyslipidemia (65.0% vs. 37.1), and history of MI (4.1% vs. 0.6%) or ischemic stroke (4.3% vs. 1.5%). Use of antiplatelets (5.7% vs. 2.0%), β-blockers (11.6% vs. 7.0%), and lipid-modifying agents (21.4% vs. 10.3%) was also higher, while chronic migraine diagnosis (11.2% vs. 15.6%) and triptans use (7.3% vs. 16.8%) were less frequent. Notably, comorbid conditions commonly managed by primary care, such as hypertension and dyslipidemia, were well captured in the VA CDW data, while cardiovascular events were more accurately captured using VA Community Care Data (*Supplement Table 4*).

In the intention-to-treat analysis, the median [IQR] follow-up time was 35.0 [20.0-55.0] months for lisinopril/candesartan and 39.0 [23.0-57.0] months in the topiramate. After applying IPTiW, baseline covariates were well balanced (*Supplement Figure 2*). The 5-year cumulative incidence (*Figure 2*) of MACE was 7.08% (95% CI: 6.57-7.64%) in the lisinopril/candesartan group vs. 6.01% (5.54-6.54%) in the topiramate group, which results in a RD of 1.06% (0.35-1.85%), a RR of 1.18 (1.05-1.33), and an average HR of 1.21 (1.10-1.34) by 5 years (*Table 2*).

For secondary outcomes, the 5-year cumulative incidence after initiating lisinopril/candesartan was 2.49% (2.18-2.93%) for MI, 1.73% (1.49-2.07%) for ischemic stroke and 0.24% (0.18-0.31%) for ICH/SAH (*Figure 3*). A higher risk of MI was observed in the lisinopril/candesartan arm (RD 0.36% [-0.25-0.95%]; RR 1.17 [0.90-1.54]; and HR 1.28 [1.03-1.59]). The risk of ischemic stroke was similar in both groups (HR 1.04; 95% CI, 0.85-1.27), while the risk of ICH/SAH was lower (HR 0.83; 95% CI, 0.54-1.27) in the lisinopril/candesartan group.

In the per-protocol analysis, the median follow-up time was 8.0 [3.0-19.0] months for lisinopril/candesartan and 4.0 [2.0-11.0] months for the topiramate. The mean IPTaW was 1.000 and ranged from 0.028 to 26.747 (*Supplement Table 5*). The risk of MACE (HR 1.30; 95% CI, 1.09-1.58) and MI (HR 1.52; 95% CI, 1.07-2.17) remained higher in the lisinopril/candesartan group (*Table 3*).

aCGRP versus Topiramate Target Trial

The aCGRP and topiramate groups had similar age (47.2 [12.2] vs. 45.0 [11.9] years) and proportion of women (39.4% vs. 36.9%), but a lower proportion of black individuals (24.1% vs. 35.7%). Both groups had comparable SBP (125.3 [14.3] mm Hg), although the aCGRP group had slightly higher cardiovascular risks, including hypertension (40.9% vs. 35.3%), history of MI (2.1% vs. 1.2%) and ischemic stroke (3.8% vs. 2.2%). The aCGRP group also had more severe migraine characteristics, such as traumatic brain injury (31.1% vs. 24.9%), chronic migraine (50.7% vs. 15.3), ≥1 headache-related neurology visits (72.4% vs. 28.7%), and higher triptan (24.6% vs. 16.5%) and neurotoxins use (12.8% vs. 2.2%).

In the intention-to-treat analysis, median [IQR] follow-up was 28.0 [16.0-45.0] months for aCGRP and 38.0 [22.0-57.0] for topiramate. The 5-year cumulative incidence of MACE was 4.57% (3.96-5.27%) in the aCGRP group and 4.78% (3.96-5.27%) in the topiramate group. The risks of MACE (HR 0.99; 95% CI, 0.86-1.14), ischemic stroke (HR 1.11; 95% CI, 0.94, 1.49) and ICH/SAH (HR 0.95; 95% CI, 0.65-1.38) were similar between groups. However, the risk of MI appeared lower in the aCGRP group (cumulative incidence: 1.21% vs. 1.50%), with a 5-year RD of -0.29% (-0.61-0.07%), RR of 0.80 (0.60-1.05), and an average HR of 0.80 (0.62-1.02).

In the per-protocol analysis, median follow-up time was 7.0 [3.0-17.0] months for aCGRP and 4.0 [2.0-11.0] for topiramate. The mean IPTaW was 1.001 and ranged from 0.021 to 42.704. The risk of MACE remained similar (HR 0.92; 95% CI 0.70-1.21) between groups, and the risk of MI was also lower in the aCGRP arm (HR 0.71; 95% CI, 0.45-1.14).

Sensitivity and Quantitative Bias Analyses
Results of the sensitivity analyses are presented in *Supplement Tables 6-10*. Excluding patients with documented cardiovascular indications for lisinopril/candesartan yielded HRs of 1.70 (95% CI 1.44-2.03) for MACE and 2.75 (95% CI 2.01-3.77) for MI. Stratification of the overall sample (*Supplement Figure 3*) showed higher risks of MACE (HR 1.63; 95% CI 1.16-2.29) and MI (HR 3.18; 95% CI 1.63-6.20) in younger patients < 40 years old, or those with SBP < 130 (MACE 1.63, 95% CI 1.43-1.86; MI 1.86, 95% CI 1.44-2.39) and between 130-139 mm Hg (MACE 1.16, 95% CI 1.00-1.34; MI 1.46, 95% CI 1.11-1.92). The increased risks of MACE and MI among younger patients or those with SBP < 140 mm Hg were more pronounced among

those without cardiovascular indications for lisinopril/candesartan. In contrast, the risks of MACE (HR 0.79; 95% CI, 0.65-0.97) and ischemic stroke (HR 0.64; 95% CI, 0.47-0.88) were lower in the overall sample of patients with SBP ≥ 140 mm Hg. After accounting for competing risks from death, the lisinopril/candesartan group showed higher risks of MI (HR 1.48; 95% CI, 1.31-1.67). Conversely, the aCGRP group showed a lower risk of MI (HR 0.82; 95% CI, 0.70-0.97). E-values were 2.32 and 1.74, respectively.

In the probabilistic bias analyses, the association between aCGRP treatment and MI was attenuated after adjustment for potential unmeasured confounder: monthly migraine days (RR 0.88; 95% simulation interval [SI], 0.62 – 1.19), monthly headache days (RR 0.92; 95% SI, 0.71 – 1.18), or migraine with aura (RR 0.90; 95% SI, 0.69 – 1.16). For the association between lisinopril/candesartan on the risk of MI among individuals without documented cardiovascular indications, bias-adjusted RRs excluded null under scenarios with moderate to strong unmeasured confounding, and only when assuming an unknown cardiovascular risk as strong as a history of MI, did the 95% SI cross the null (RR, 1.45; 95% SI, 0.90 – 2.14). after adjustment for strong unknown confounding using a hypothetical cardiovascular risk factor, the bias-adjusted RRs excluded the null, only when an unknown cardiovascular risk factor as strong as history of MI, the 95% SI crossed 1 (RR 1.45; 95% SI 0.90 – 2.14). For outcome controls, both lisinopril/candesartan and aCGRP groups had higher risks of primary malignancy and weight gain compared to topiramate. The risk of non-pathological fracture was comparable in the lisinopril/candesartan group (HR 1.06; 95% CI, 0.98-1.14) but elevated in the aCGRP group (HR 1.13; 95% CI, 1.01-1.26). Risks of influenza were similar between comparisons.

Discussion

In this retrospective study, we emulated two target trials to evaluate the association of lisinopril/candesartan and aCGRP treatments, compared to topiramate, for migraine prevention with the risk of MACE. Contrary to our hypotheses, lisinopril/candesartan use was associated with an increased risk of MACE, primarily driven by MI, in the overall sample. This risk was greater in younger patients or those without documented cardiovascular indications or SBP <140 mmHg. However, the risk of ischemic stroke among individuals with SBP ≥ 140 mm Hg was 36% lower, which is consistent with the guideline recommendation to initiate ACEIs/ARBs to prevent

stroke in hypertensive adults.[25] Meanwhile, aCGRP treatment was association with a reduced risk of MI, and was not associated with ischemic stroke or ICH/SAH. Lastly, we did not observed associations of lisinopril/candesartan and aCGRP treatment on the risk of ICH/SAH, possibly due to lower incident rate of ICH/SAH and insufficient power.

To interpret these seemingly contradictory observations that lisinopril/candesartan may increase risk of cardiovascular events, it is important to note that migraine-associated cardiovascular risk is distinct to atherosclerotic risk. In fact, two population-level observational studies have reported greater risk of cardiovascular events in younger migraine patients. For instance, a Danish study[26] reported HRs for MI of 1.62 (95% CI, 1.33-1.98) in those aged 40 years or younger, 1.56 (1.38-1.77) in those aged 40-59 years, and 1.41 (1.21-1.64) in those 60 years or older. Similarly, a Korean study[27] found HRs for stroke were 1.74 (1.65-1.84) for individuals under 60 years old, and 1.22 (1.18-1.26) for those aged 60 and older. These findings support that migraine-associated cardiovascular risks differ from the atherosclerotic cardiovascular risks, which typically increase with age. Thus lisinopril/candesartan may not prevent MI driven by non-atherosclerotic mechanisms related to migraine.

Although the pathophysiology linking migraine-associated cardiovascular risks remains unclear, cortical spreading depression (CSD)[28] and trigeminal nerve activation may have contributed to the non-atherosclerotic risks in migraine patients and could explain our counterintuitive findings. The release of CGRP and substance P (SP) following trigeminal nerve activation can lead to vasodilation and neurogenic inflammation. Furthermore, SP may also promote platelet aggregation and clots formation.[29] Higher plasma SP levels have been associated with recurrent infraction or death by 2 years among post-MI patients.[30] In this context, blocking ACE, which degrades SP, may exacerbate neurogenic inflammation[31] and increase MI risk, especially if ACE expression may be down-regulated in migraine patients.[32] In our sample predominantly with lisinopril (98.3%), the increased MI risk could reflect accumulation of SP and neurogenic inflammation among individuals with minimal atherosclerotic risks (younger patients, SBP < 140 mmHg, and without documented CKD, HF, hypertension or history of MI). Additionally, this lisinopril-induced neurogenic inflammation may also affect cerebral vascular system, as the risk of ischemic stroke was higher in the lisinopril/candesartan group with SBP < 130 mm Hg

(eTable 7, HR 1.56; 95% CI, 1.15-2.21). Notably, delayed cerebral ischemia after SAH, which shares CSD as one of the underline causes, is often managed by raising mean arterial pressure rather than lowering it with lisinopril. However, due to small sample size of younger patients or those with low atherosclerotic risks and relatively low incident rate of ischemic stroke, this association should be further validated in a larger cohort.

Current evidence from *in vitro* and animal studies suggests that CGRP may mediate a cardioprotective effect following myocardial injuries through its vasodilation, inotropy and chronotropy effects.[33] However, the causal relationship of CGRP as the exposure has yet to be directly studied. To date, only two observational studies have compared the cardiovascular safety of aCGRP vs. neurotoxin[34]. One study, based on an underpowered Medicare sample, reported a HR of 0.86 (95% CI 0.30 - 2.48)[34], but relied on machine learning-estimated propensity scores, which may yield biased estimates in causal inference[35,36]. Another insurance claim-based study reported a relative risk of 0.87 (95% CI 0.19 – 1.55) for MI.[37] Despite their limitations, the direction of these point estimates aligns with our findings from the intention-to-treat analysis (HR 0.80; 95% CI 0.62-1.02) and the competing risk-adjusted estimate (HR 0.82; 95%CI 0.70-0.97). The smaller effect sizes observed in prior studies are likely attributable to differences in comparator selection. Neurotoxins are typically prescribed for patients with migraine patients (≥15 migraine days per month) who have failed multiple preventive treatments. As such, migraine frequency and severity may be more comparable between patients initiated with aCGRP and neurotoxin than between patients initiated with aCGRP and topiramate. In our probabilistic bias analysis adjusting for potential confounding by monthly migraine days, the bias-adjusted estimate (RR 0.88; 95% SI, 0.62 - 1.19) closely mirrored the estimates from neurotoxin comparisons. These findings are consistent with the hypothesis that migraine-associated cardiovascular risk may be mediated by neuroinflammatory process involving CGRP and substance P released during migraine attacks. By blocking CGRP or its receptor, aCGRP treatment could reduce the long-term neurogenic inflammation in patients with migraine and thus lower the risk of MI.

Limitations

This study has several limitations. The primary concern is confounding by indication, a type of confounding bias[38] arises when the indication for treatment (disease or disease severity) is unmeasured or misclassified. The lisinopril/candesartan group was older, had higher SBP and had greater cardiovascular comorbidities, consistent with clinical practice where these agents are more frequently prescribed to migraine patients with or at risk of atherosclerotic disease. While β-blockers could be a closer equipoise active comparator, particularly if the increased risk of MACE/MI is driven by SP-related neuroinflammation, the positivity assumption in our analysis holds, as both lisinopril/candesartan and topiramate are indicated for migraine prevention. Although the precise indication, whether for migraine prevention, hypertension, or both, could not be ascertained with EHR data, conditional exchangeability remains plausible given extensive adjustment for atherosclerotic risk factors and other confounders. Besides, baseline covariates balances (SMDs) were achieved after weighting, large E-values which exceeded the HRs of known cardiovascular risk factors included in the model (*Supplement* Table 8), and findings from probabilistic bias analysis adjusting for strong unknown cardiovascular risk factors (*Supplement Table 9*), suggesting that the increased risk of MACE with lisinopril/candesartan is unlikely biased by residual confounding or unmeasured cardiovascular indications. Additionally, negative outcome control analyses for influenza (confounding by healthcare-seeking behavior) and non-pathological fracture (frailty) did not find meaningful associations, supporting the robustness of our estimates. Future studies could consider instrumental variable analyses to further address potential selection bias and confounding by indication. Second, the administrative data did not capture information on migraine aura, frequency and severity, which could better quantify the migraine-associated cardiovascular risk (*Supplement Figure 1b*). Although the probabilistic bias analysis accounted for these unmeasured confounding, they were based on assumptions and parameters informed by the literature, and should be interpreted with caution. Third, although outpatient encounter diagnoses within VHA demonstrated high sensitivity and specificity to capture cardiovascular comorbidities[39], false-negative misclassification cannot be eliminated. By restricting the study sample among those with primary care visits and leveraging VA Community Care and CMS claims, the risk of misclassification bias is likely minimal. Lastly, gaps between prescription fills were defined "retrospectively" in this study.[40] If patients did not redeem a new prescriptions within the 90-day grace period, follow-up was censored at day 0 of the grace period. This approach may introduce a dependency to treatment status on future prescription refills.

However, conclusions of this study are not biased as they were drawn based on intention-to-treat analyses.

Conclusions

Findings from this retrospective study suggest that initiation of lisinopril/candesartan may be associated with a higher risk of MACE, particularly MI, with greater risk observed in migraine patients with minimal risk of atherosclerotic diseases. In contrast, aCGRP treatment appears safe regarding cardiovascular events and may reduce the risk of MI. While these associations are biologically plausible, given mechanisms involving neurogenic inflammation mediated by CGRP and SP, future research is needed to confirm dose-response relationships and the consistency of these findings on migraine-associated non-atherosclerotic cardiovascular risks before informing clinical practice.

Acknowledgement

This research was supported by the resources and computational environment of the VA Informatics and Computing Infrastructure (VINCI) (DART# 2024-02-057-D), as well as by the use of Centers for Medicare and Medicaid data provided by the VA Information Resource Center (VIReC) (DUI# Wang-05-1). This publication does not necessarily represent the views of the Department of Veterans Affairs or the United States Government.

This manuscript was previously submitted to *Circulation*, *Annals of Internal Medicine*, and *Neurology*, where it was rated as a low priority. Despite extensive sensitivity analyses, quantitative bias analyses, and a cautious interpretation of the findings as hypothesis-generating rather than practice-changing, reviewers remained concerned about residual confounding inherent to observational studies. To promote transparency and open discussion, the corresponding author has included all reviewer comments and corresponding responses in the supplementary materials, allowing readers to evaluate and interpret the study independently.

Reference

1. Burch R, Rizzoli P, Loder E. The prevalence and impact of migraine and severe headache in the United States: Updated age, sex, and socioeconomic□specific estimates from government health surveys. *Headache: The Journal of Head and Face Pain*. 2021;61(1):60-68.

2. Ng CYH, Tan BY, Teo YN, et al. Myocardial infarction, stroke and cardiovascular mortality among migraine patients: a systematic review and meta-analysis. *Journal of neurology*. 2022:1-13.

3. Hvitfeldt Fuglsang C, Pedersen L, Schmidt M, Vandenbroucke JP, Bøtker HE, Sørensen HT. Migraine and risk of premature myocardial infarction and stroke among men and women: A Danish population-based cohort study. *Plos Medicine*. 2023;20(6):e1004238.

4. Mohebi R, Chen C, Ibrahim NE, et al. Cardiovascular disease projections in the United States based on the 2020 census estimates. *Journal of the American College of Cardiology*. 2022;80(6):565-578.

5. Petersen CL, Hougaard A, Gaist D, Hallas J. Risk of stroke and myocardial infarction among initiators of triptans. *JAMA neurology*. 2024;81(3):248-254.

6. Wang Z, VanderPluym JH, Singh RBH, et al. Safety of Triptans in Patients Who Have or Are at High Risk for Cardiovascular Disease: A Target Trial Emulation. Elsevier; 2024:1722-1731.

7. Jang HY, Kim JH, Song Y-K, et al. Antidepressant use and the risk of major adverse cardiovascular events in patients without known cardiovascular disease: a retrospective cohort study. *Frontiers in Pharmacology*. 2020;11:594474.

8. Kee Z, Kodji X, Brain SD. The role of calcitonin gene related peptide (CGRP) in neurogenic vasodilation and its cardioprotective effects. *Frontiers in physiology*. 2018;9:1249.

9. Charles AC, Digre KB, Goadsby PJ, et al. Calcitonin gene□related peptide□targeting therapies are a first□line option for the prevention of migraine: An American Headache Society position statement update. *Headache: The Journal of Head and Face Pain*. 2024;64(4):333-341.

10. McAlister FA, Investigators RASMM-A. Angiotensin-converting enzyme inhibitors or angiotensin receptor blockers are beneficial in normotensive atherosclerotic patients: a collaborative meta-analysis of randomized trials. *European heart journal*. 2012;33(4):505-514.

11. Silberstein S, Holland S, Freitag F, Dodick D, Argoff C, Ashman E. Evidence-based guideline update: pharmacologic treatment for episodic migraine prevention in adults: report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. 2012;78(17):1337-1345.

12. Sico JJ, Antonovich NM, Ballard-Hernandez J, et al. 2023 US Department of Veterans Affairs and US Department of Defense Clinical Practice Guideline for the Management of Headache. *Annals of Internal Medicine*. 2024;177(12):1675-1694.

13. Sico JJ, Seng EK, Wang K, et al. Characteristics and gender differences of headache in the Veterans Health Administration: a national cohort study, fiscal year 2008–2019. *Neurology*. 2022;99(18):e1993-e2005.

14. Desai RJ, Wang SV, Sreedhara SK, et al. Process guide for inferential studies using healthcare data from routine clinical practice to evaluate causal effects of drugs (PRINCIPLED): considerations from the FDA Sentinel Innovation Center. *bmj*. 2024;384

15.   Wilding J, Van Gaal L, Rissanen A, Vercruysse F, Fitchet M. A randomized double-blind placebo-controlled study of the long-term efficacy and safety of topiramate in the treatment of obese subjects. *International journal of obesity*. 2004;28(11):1399-1410.

16.   Hoffman V, Xue F, Ezzy SM, et al. Risk of cardiovascular and cerebrovascular events and mortality in patients with migraine receiving prophylactic treatments: an observational cohort study. *Cephalalgia*. 2019;39(12):1544-1559.

17.   Staerk C, Byrd A, Mayr A. Recent methodological trends in Epidemiology: No need for data-driven variable selection? *American journal of epidemiology*. 2024;193(2):370-376.

18.   Savitz DA, Wise LA, Bond JC, et al. Responding to Reviewers and Editors About Statistical Significance Testing. American College of Physicians; 2024. p. 385-386.

19.   Sendor R, Stürmer T. Core concepts in pharmacoepidemiology: confounding by indication and the role of active comparators. *Pharmacoepidemiology and drug safety*. 2022;31(3):261-269.

20.   VanderWeele TJ, Ding P. Sensitivity analysis in observational research: introducing the E-value. *Annals of internal medicine*. 2017;167(4):268-274.

21.   Kramer C, Leitao C, Pinto L, Canani L, Azevedo M, Gross J. Efficacy and safety of topiramate on weight loss: a meta‐analysis of randomized controlled trials. *Obesity reviews*. 2011;12(5):e338-e347.

22.   Takada M, Fujimoto M, Motomura H, Hosomi K. Inverse association between sodium channel-blocking antiepileptic drug use and cancer: data mining of spontaneous reporting and claims databases. *International journal of medical sciences*. 2016;13(1):48.

23.   Sloot R, Breskin A, Colantonio LD, et al. Comparing PCSK9 Monoclonal Antibody Treatment Strategies Following Myocardial Infarction Using Negative Control Outcomes: A Target Trial Emulation Study. *Epidemiology*. 2024:10.1097.

24.   Zafari Z, Park J-e, Shah CH, et al. The state of use and utility of negative controls in pharmacoepidemiologic studies. *American journal of epidemiology*. 2024;193(3):426-453.

25.   Bushnell C, Kernan WN, Sharrief AZ, et al. 2024 guideline for the primary prevention of stroke: a guideline from the American Heart Association/American Stroke Association. *Stroke*. 2024;55(12):e344-e424.

26.   Bonnesen K, Fuglsang CH, Schmidt M. Impact of nonsteroidal anti-inflammatory drug use on the migraine-associated risk of myocardial infarction and ischemic stroke: a population-based cohort study. *Neurology*. 2024;103(12):e210034.

27.   Kwon MJ, Choi HG, Kim YH, et al. A higher probability of subsequent stroke and ischemic heart disease in migraine patients: A longitudinal follow-up study in Korea. *The journal of headache and pain*. 2023;24(1):98.

28.   Binder NF, Glück C, Middleham W, et al. Vascular response to spreading depolarization predicts stroke outcome. *Stroke*. 2022;53(4):1386-1395.

29.   Azma T, Sugimoto Y, Kinoshita H, et al. Detection of the full-length transcript variant for neurokinin-1 receptor in human whole blood associated with enhanced reinforcement of clot by substance-P. *Journal of thrombosis and thrombolysis*. 2012;33:329-337.

30.   Ng LL, Sandhu JK, Narayan H, et al. Pro-substance p for evaluation of risk in acute myocardial infarction. *Journal of the American College of Cardiology*. 2014;64(16):1698-1707.

31.   de Mos M, Huygen F, Stricker BC, Dieleman J, Sturkenboom M. The association between ACE inhibitors and the complex regional pain syndrome: Suggestions for a neuro-inflammatory pathogenesis of CRPS. *PAIN®*. 2009;142(3):218-224.

32.   Bhering Martins L, Silva de Miranda A, Rodrigues AMdS, et al. Altered serum levels of renin‐angiotensin system markers in migraine. *Headache: The Journal of Head and Face Pain*. 2020;60(9):1995-2002.

33.   Argunhan F, Brain SD. The vascular-dependent and-independent actions of calcitonin gene-related peptide in cardiovascular disease. *Frontiers in Physiology*. 2022;13:833645.

34.   Yang S, Orlova Y, Park H, et al. Cardiovascular Safety of Anti-CGRP Monoclonal Antibodies in Older Adults or Adults With Disability With Migraine. *JAMA neurology*.

35.   Zivich PN, Breskin A. Machine learning for causal inference: on the use of cross-fit estimators. *Epidemiology*. 2021;32(3):393-401.

36.   Naimi AI, Mishler AE, Kennedy EH. Challenges in obtaining valid causal effect estimates with machine learning algorithms. *American Journal of Epidemiology*. 2023;192(9):1536-1544.

37.   Dodick DW, Tepper SJ, Ailani J, et al. Effect of erenumab versus other migraine preventive medications on cardiovascular and cerebrovascular outcomes: A United States claims database‐based observational cohort study. *Headache: The Journal of Head and Face Pain*. 2025;

38.   Porta MS, Greenland S, Hernán M, dos Santos Silva I, Last JM. *A dictionary of epidemiology*. Oxford university press; 2014.

39.   Hoffman BB, Goldstein MK. Accuracy of computerized outpatient diagnoses in a Veterans Affairs general medicine clinic. *Am J Manag Care*. 2002;8(January):37-43.

40.   Nielsen LH, Løkkegaard E, Andreasen AH, Keiding N. Using prescription registries to define continuous drug use: how to fill gaps between prescriptions. *Pharmacoepidemiology and drug safety*. 2008;17(4):384-388.

Table 1. Baseline Characteristics of Person-Trials in the Study Sample

	Target trial (i)		Target trial (ii)	
	Lisinopril/Candesartan (N = 48,610)	Topiramate (N = 115,047)	aCGRP (N = 25,635)	Topiramate (N = 135,740)
Age, mean (SD), y	52.3 (12.3)	43.2 (11.2)	47.2 (12.2)	45.0 (11.9)
Gender, women	9,173 (18.9)	45,980 (40.0)	10,113 (39.4)	50,112 (36.9)
Race				
American Indian/Alaska Native	493 (1.0)	1,177 (1.0)	247 (1.0)	1,329 (1.0)
Asian	987 (2.0)	2,231 (1.9)	368 (1.4)	2,645 (1.9)
Black	11,308 (23.3)	40,382 (35.1)	6,176 (24.1)	48,468 (35.7)
Multi-Race	712 (1.5)	1,913 (1.7)	361 (1.4)	2,212 (1.6)
Native Hawaiian/Pacific Islander	744 (1.5)	1,369 (1.2)	256 (1.0)	1,707 (1.3)
White	30,886 (63.5)	58,165 (50.6)	16,244 (63.4)	68,325 (50.3)
Unknown	3,480 (7.2)	9,810 (8.5)	1,983 (7.7)	11,054 (8.1)
Ethnicity, Hispanics	4,778 (9.8)	14,769 (12.8)	2,885 (11.3)	16,666 (12.3)
Residence, non-urban	17,128 (35.2)	31,788 (27.6)	8,008 (31.2)	38,349 (28.3)
Area deprivation index, mean (SD)	54.0 (24.6)	51.8 (24.0)	51.6 (23.6)	52.4 (24.2)
N miss	4,579	10,131	4,125	12,469
Service-connected disability	42,807 (88.1)	110,277 (95.9)	24,544 (95.7)	129,215 (95.2)
Smoking status				
Never	20,260 (41.7)	61,955 (53.9)	13,372 (52.2)	71,799 (52.9)
Current	17,227 (35.4)	31,606 (27.5)	6,652 (25.9)	37,714 (27.8)
Former	10,939 (22.5)	20,967 (18.2)	5,554 (21.7)	25,658 (18.9)
Unknown	184 (0.4)	519 (0.5)	57 (0.2)	569 (0.4)
Body mass index, mean (SD)	32.1 (6.2)	31.2 (6.0)	31.0 (5.9)	31.5 (6.1)
Underweight/Normal	4,904 (10.1)	15,393 (13.4)	3,642 (14.2)	16,763 (12.3)
Overweight	13,957 (28.7)	36,593 (31.8)	8,236 (32.1)	41,743 (30.8)
Obese	29,161 (60.0)	60,753 (52.8)	13,434 (52.4)	74,800 (55.1)
Unknown	588 (1.2)	2,308 (2.0)	323 (1.3)	2,434 (1.8)
Blood pressure, mean (SD), **mmHg**				

	Target trial (i)		Target trial (ii)	
	Lisinopril/Candesartan (N = 48,610)	Topiramate (N = 115,047)	aCGRP (N = 25,635)	Topiramate (N = 135,740)
Systolic	136.4 (16.6)	123.8 (13.4)	125.3 (14.3)	125.3 (14.3)
Diastolic	84.0 (11.0)	78.1 (9.0)	78.9 (9.2)	78.7 (9.4)
N miss	334	1,512	141	1,566
Care Assessment Need score, median [IQR]	20.0 [10.0, 40.0]	10.0 [5.0, 25.0]	15.0 [5.0, 30.0]	15.0 [5.0, 25.0]
N miss	1,029	2,395	209	2,592
Primary care visits, median [IQR]	2.0 [1.0, 3.0]	2.0 [1.0, 3.0]	2.0 [1.0, 4.0]	2.0 [1.0, 3.0]
Influenza vaccination	23,265 (47.9)	50,656 (44.0)	12,875 (50.2)	62,084 (45.7)
Comorbidity				
Atrial fibrillation/flutter	2,143 (4.4)	1,586 (1.4)	905 (3.5)	2,892 (2.1)
Alcohol-related disorder	9,828 (20.2)	19,240 (16.7)	4,063 (15.8)	23,863 (17.6)
Cardiomyopathy	1,281 (2.6)	591 (0.5)	423 (1.7)	1,592 (1.2)
Chronic inflammatory conditions*	2,934 (6.0)	4,668 (4.1)	1,718 (6.7)	6,101 (4.5)
Chronic kidney disease	3,905 (8.0)	3,522 (3.1)	1,815 (7.1)	6,536 (4.8)
COPD	5,743 (11.8)	5,393 (4.7)	2,337 (9.1)	8,481 (6.2)
Diabetes	15,240 (31.4)	9,433 (8.2)	4,409 (17.2)	18,289 (13.5)
DVT/PE	906 (1.9)	1,164 (1.0)	516 (2.0)	1,632 (1.2)
Dyslipidemia	31,608 (65.0)	42,669 (37.1)	13,124 (51.2)	58,047 (42.8)
Heart failure	2,601 (5.4)	1,266 (1.1)	974 (3.8)	3,277 (2.4)
Hypertension	37,100 (76.3)	27,376 (23.8)	10,481 (40.9)	47,880 (35.3)
Depression	29,569 (60.8)	72,969 (63.4)	18,539 (72.3)	87,448 (64.4)
Obstructive sleep apnea	5,679 (11.7)	6,348 (5.5)	2,672 (10.4)	9,944 (7.3)
Peripheral artery disease	3,425 (7.0)	2,377 (2.1)	1,361 (5.3)	4,369 (3.2)
Substance-related disorder	7,411 (15.2)	12,418 (10.8)	3,265 (12.7)	15,562 (11.5)
Traumatic brain injury†	11,876 (24.4)	29,339 (25.5)	7,963 (31.1)	33,739 (24.9)
Valvular heart disease	2,553 (5.3)	2,077 (1.8)	1,082 (4.2)	3,547 (2.6)
History of MACE	3956 (8.1)	2382 (2.1)	1,485 (5.8)	4,550 (3.4)
Myocardial infraction	1991 (4.1)	683 (0.6)	529 (2.1)	1,608 (1.2)

	Target trial (i)		Target trial (ii)	
	Lisinopril/Candesartan (N = 48,610)	Topiramate (N = 115,047)	aCGRP (N = 25,635)	Topiramate (N = 135,740)
Ischemic stroke	2109 (4.3)	1699 (1.5)	975 (3.8)	3,047 (2.2)
ICH/SAH	262 (0.5)	206 (0.2)	141 (0.6)	357 (0.3)
Medication (%)				
Any ACEIs/ARBs	48,610 (100.0)	0 (0.0)	3,949 (15.4)	16,574 (12.2)
β-blockers	5,639 (11.6)	8,049 (7.0)	4,788 (18.7)	12,923 (9.5)
Calcium channel blockers	4,306 (8.9)	4,925 (4.3)	2,704 (10.5)	9,444 (7.0)
Diuretics	4,705 (9.7)	4,728 (4.1)	2,218 (8.7)	9,359 (6.9)
MRAs	480 (1.0)	1,002 (0.9)	542 (2.1)	1,765 (1.3)
Other antihypertensives ‡	4,811 (9.9)	10,235 (8.9)	4,019 (15.7)	14,177 (10.4)
Anti-coagulants	1,113 (2.3)	1,015 (0.9)	642 (2.5)	1,822 (1.3)
Anti-platelets	2,794 (5.7)	2,244 (2.0)	1,450 (5.7)	4,879 (3.6)
Lipid modifying agents	10,381 (21.4)	11,817 (10.3)	5,842 (22.8)	20,039 (14.8)
Hormone replacement therapy	414 (0.9)	1,905 (1.7)	667 (2.6)	2,101 (1.5)
GLP-1 agonists	624 (1.3)	519 (0.5)	536 (2.1)	1,129 (0.8)
SGLT2i	1,170 (2.4)	568 (0.5)	640 (2.5)	1,752 (1.3)
Anti-psychotics	2,246 (4.6)	6,433 (5.6)	2,293 (8.9)	7,730 (5.7)
Migraine duration, median [IQR], mo	69.0 [30.0, 111.0]	33.0 [7.0, 75.0]	56.0 [23.0, 99.0]	34.0 [8.0, 77.0]
Chronic migraine	5,462 (11.2)	17,939 (15.6)	13,006 (50.7)	20,749 (15.3)
Any headache-related ED visits	1,591 (3.3)	6,965 (6.1)	2,046 (8.0)	8,413 (6.2%)
Headache-related neurology visits				
None	43,657 (89.8)	81,862 (71.2)	7,085 (27.6)	96,728 (71.3)
Once	2,265 (4.7)	18,607 (16.2)	6,392 (24.9)	22,180 (16.3)
Twice	1,308 (2.7)	7,928 (6.9)	4,324 (16.9)	9,298 (6.8)
Three times and more	1,380 (2.8)	6,650 (5.8)	7,834 (30.6)	7,534 (5.6)
Any aCGRP	892 (1.8)	1,859 (1.6)	25,635 (100.0)	0 (0.0)
Ergotamines	NA§	25 (<0.1)	31 (0.1)	29 (<0.1)
Triptans	3,562 (7.3)	19,322 (16.8)	6,304 (24.6)	22,409 (16.5)
NSAIDs	7,022 (14.4)	16,081 (14.0)	5,079 (19.8)	21,017 (15.5)

	Target trial (i)		Target trial (ii)	
	Lisinopril/Candesartan (N = 48,610)	Topiramate (N = 115,047)	aCGRP (N = 25,635)	Topiramate (N = 135,740)
Other anticonvulsants (%)	1,480 (3.0)	3,009 (2.6)	1,848 (7.2)	3,788 (2.8)
SNRI	1,622 (3.3)	3,974 (3.5)	1,451 (5.7)	4,777 (3.5)
TCAs	1,423 (2.9)	3,428 (3.0)	1,708 (6.7)	4,240 (3.1)
Neurotoxins (%)	877 (1.8)	2,713 (2.4)	3,273 (12.8)	3,047 (2.2)
MPR, median [IQR]	1.00 [0.92, 1.00]	1.00 [0.86, 1.00]	0.97 [0.79, 1.00]	1.00 [0.86, 1.00]
Treatment length, median [IQR], d	270.0 [90.0, 629.0]	111.0 [60.0, 327.0]	256.0 [90.0, 605.0]	120.0 [60.0, 340.0]

Abbreviations: ACEIs/ARBs, angiotensin-converting enzyme inhibitors/angiotensin II receptor blockers; aCGRP, anti-calcitonin gene-related peptide; COPD, chronic obstructive pulmonary disease; DVT/PE, deep vein thrombosis/pulmonary embolism; ED, emergency department; GLP-1, glucagon-like peptide-1; ICH/SAH, intracerebral/subarachnoid hemorrhage; IQR, interquartile range; MACE, major adverse cardiovascular events; MPR, medication possession ratio; MRAs, mineralocorticoid receptor antagonists; NSAIDs, non-steroidal anti-inflammatory drugs; SD, standard deviation; SGLT2i, sodium-glucose cotransporter-2 inhibitors; SNRI, serotonin and norepinephrine reuptake inhibitor; TCAs, tricyclic antidepressants.

* Chronic inflammatory conditions included ankylosing spondylitis, HIV, inflammatory bowel disease, psoriasis or psoriatic arthritis, rheumatoid arthritis, systemic lupus erythematosus, and systemic sclerosis.
† Combined with ICD diagnoses and TBI screening from health factor domain.
‡ Other antihypertensives included α-blockers, centrally acting sympathetic agonists, nitrates, direct renin inhibitor, and vasodilators.
§ Not available because of the CMS cell size suppression policy.

Table 2. Intention-to-treat estimates of cumulative incidence, risk differences (RD), risk ratio (RR) and overall hazard ratio (HR) for major adverse cardiovascular events and secondary outcomes over 5 years.

	5-year cumulative incidence		Risk differences (95% CI)	Risk ratio (95% CI)	Overall HR (95% CI)*
	Intervention, % (95% CI)	Topiramate, % (95% CI)			
Lisinopril/Candesartan					
MACE	7.08 (6.57, 7.64)	6.01 (5.54, 6.54)	1.06 (0.35, 1.85)	1.18 (1.05, 1.33)	1.21 (1.10, 1.34)
MI	2.49 (2.18, 2.93)	2.13 (1.70, 2.63)	0.36 (-0.25, 0.95)	1.17 (0.90, 1.54)	1.28 (1.03, 1.59)
Ischemic stroke	1.73 (1.49, 2.07)	1.79 (1.54, 2.10)	-0.06 (-0.40, 0.34)	0.97 (0.79, 1.20)	1.04 (0.85, 1.27)
ICH/SAH	0.24 (0.18, 0.31)	0.29 (0.20, 0.41)	-0.05 (-0.18, 0.07)	0.82 (0.54, 1.34)	0.83 (0.54, 1.27)
aCGRP					
MACE	4.57 (3.96, 5.27)	4.78 (4.62, 4.95)	-0.21 (-0.87, 0.53)	0.96 (0.82, 1.11)	0.99 (0.86, 1.14)
MI	1.21 (0.89, 1.57)	1.50 (1.41, 1.60)	-0.29 (-0.61, 0.07)	0.80 (0.60, 1.05)	0.80 (0.62, 1.02)
Ischemic stroke	1.55 (1.11, 1.97)	1.43 (1.34, 1.51)	0.12 (-0.34, 0.55)	1.08 (0.77, 1.39)	1.11 (0.84, 1.49)
ICH/SAH	0.20 (0.13, 0.27)	0.22 (0.19, 0.25)	-0.02 (-0.10, 0.05)	0.91 (0.58, 1.28)	0.95 (0.65, 1.38)

Abbreviations: aCGRP, anti-calcitonin gene-related peptide; CI, confidence interval; HR, hazard ratio; ICH/SAH, intracerebral or subarachnoid hemorrhage; MACE, major adverse cardiovascular events; MI, myocardial infraction.

* Robust sandwich estimator

Table 3. Per-protocol estimates of overall hazard ratio (HR) of major adverse cardiovascular events and secondary outcomes over 5 years, after censoring individuals who did not adhere to the treatment.

	Lisinopril/Candesartan HR (95% CI)*	aCGRP HR (95% CI)*
MACE	1.30 (1.09, 1.58)	0.92 (0.70, 1.21)
MI	1.52 (1.07, 2.17)	0.71 (0.45, 1.14)
Ischemic stroke	1.16 (0.87, 1.55)	1.00 (0.58, 1.73)
ICH/SAH	1.71 (0.90, 3.23)	0.73 (0.37, 1.46)

Abbreviations: aCGRP, anti-calcitonin gene-related peptide; CI, confidence interval; HR, hazard ratio; ICH/SAH, intracerebral/subarachnoid hemorrhage; MACE, major adverse cardiovascular events; MI, myocardial infraction.

*Robust sandwich estimator

Figure Legends

Figure 1. Study Flowchart.
Numbers in the parentheses indicate number of unique patients. ACEIs/ARBs indicates angiotensin-converting enzyme inhibitors/angiotensin II receptor blockers; aCGRP, anti-calcitonin gene-related peptide; CDW, Corporate Data Warehouse; CMS, Centers for Medicare & Medicaid Services; VA, Department of Veterans Affairs.

Figure 2. Intention-to-Treat Cumulative Incidence of Major Adverse Cardiovascular Events.
Cumulative incidence from the target trial comparing (a) lisinopril/candesartan (LSP/CAN) vs topiramate (TPM); (b) Cumulative incidence from the target trial comparing anti-calcitonin gene-related peptide (aCGRP) treatment vs TPM. The shaded ribbons indicate 95% CIs based on 500 bootstrapping samples.

Figure 3. Intention-to-Treat Cumulative Incidence of Secondary Outcomes.
Cumulative incidence of (a) myocardial infarction (MI), (b) ischemic stroke, and (c) intracerebral/subarachnoid hemorrhage (ICH/SAH) from the target trial comparing lisinopril/candesartan (LSP/CAN) vs. topiramate (TPM); and (d) MI, (e) ischemic stroke, and (f) ICH/SAH from the target trial comparing anti-calcitonin gene-related peptide (aCGRP) treatment vs. TPM. The shaded ribbons represent 95% CIs based on 500 bootstrapping samples.

Oct 2007 — Incident migraine patients in the VA — Sep 2023

Jun 2018 — Migraine preventives — Sep 2024

Oct 1999 — Characteristics from VA CDW, Community Care, and CMS

| Lisinopril Candesartan | Topiramate | aCGRP |

N = 290,629 (168,833) — Eligible — N = 229,282 (163,321)

Exclusion

- Initiated topiramate prior before migraine, n = 26,706
- Initiated lisinopril/candesartan before migraine, n = 31,877
- Treatment initiation after death, n = 5
- Co-prescribing or use of lisinopril/candesartan and topiramate in the past 3 months, n = 16,601
- Ever prescribed/co-prescribing other ACEIs/ARBs, n = 19,761
- No primary care visit in the past 12 months, n = 32,022

- Initiated topiramate prior before migraine, n = 26,706
- Treatment initiation after death, n = 2
- Co-prescribing or use of preventive aCGRP and topiramate in the past 3 months, n = 10,974
- Co-prescribing of abortive aCGRP and topiramate, n = 702
- No primary care visit in the past 12 months, n = 29,523

Analyzed

Target trial (i)
N = 163,657
(104,413)

Target trial (ii)
N = 161,375
(100,629)

a. Lisinopril/Candesartan vs. Topiramate

b. aCGRP vs. Topiramate

Num at risk

LSP/CAN 48610	44291	33748	24243	15875	9061
TPM 115047	107352	84770	63303	43274	24706

aCGRP 25635	22115	15238	9795	5601	1385
TPM 135740	125813	98836	73604	50271	28724

a. Myocardial infraction

b. Ischemic stroke

c. Intracerebral/subarachnoid hemorrhage

Num at risk						
LSP/CAN	48610	44539	34104	24608	16166	9271
TPM	115047	107521	85027	63598	43538	24899

Num at risk						
LSP/CAN	48610	44637	34257	24760	16329	9375
TPM	115047	107509	85042	63599	43550	24902

Num at risk						
LSP/CAN	48610	44860	34572	25108	16613	9571
TPM	115047	107639	85250	63844	43784	25080

d. Myocardial infraction

e. Ischemic stroke

f. Intracerebral/subarachnoid hemorrhage

Num at risk						
aCGRP	25635	22192	15329	9883	5662	1401
TPM	135740	126126	99296	74095	50711	29044

Num at risk						
aCGRP	25635	22184	15333	9879	5673	1403
TPM	135740	126121	99320	74132	50756	29066

Num at risk						
aCGRP	25635	22235	15396	9948	5718	1419
TPM	135740	126376	99705	74556	51134	29356