Genetically determined ancestry associates with morphological and molecular carotid plaque features

Authors

Nima Fahim[1*], Tim R. Sakkers[2], Joost Hoekstra[3], Dominique P.V. de Kleijn[3], Michal Mokry[1], Jose Verdezoto Mosquera[4,5], Gerard Pasterkamp[1], Hester M. den Ruijter[2], Clint L. Miller[4,5], Jessica van Setten[6], Sander W. van der Laan[1,4*].

These authors contributed equally.

Affiliations

Central Diagnostic Laboratory, Division Laboratories, Pharmacy, and Biomedical genetics, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands.

Laboratory of Experimental Cardiology, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands.

Department of Vascular Surgery, Division of Surgical Specialties, University Medical Center Utrecht, University of Utrecht, Utrecht, the Netherlands.

Department of Genome Sciences, University of Virginia, Charlottesville, VA, USA

Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA, USA

Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands.

Corresponding author(s)

dr. Sander W. van der Laan

Central Diagnostic Laboratory, Division Laboratories, Pharmacy, and Biomedical genetics, University Medical Center Utrecht, Utrecht University, Utrecht

Heidelberglaan 100, 3508 GA, Utrecht, The Netherlands

T: +31 (0) 88 75 565 73

E: s.w.vanderlaan-2@umcutrecht.nl

Manuscript contents

Manuscript of 914 words (max ±800-1000), 1 figure.

Journal subject terms

Genetics, Atherosclerosis, Plaque, Ancestry

Abstract

Atherosclerosis, the main driver of cardiovascular disease (CVD), is influenced by a plethora of risk factors, including age, sex, and diabetes, that correlate with socio-economic status and may vary across ethnicities. These factors fail to fully explain observed ethnic disparities in CVD burden. For example, coronary artery calcification increases with age regardless of ethnicity, yet CAC is more prevalent in individuals of European descent. As these findings may be confounded by self-reported ethnicity, genome-informed ancestry offers a more accurate lens through which to study these ancestral differences. Yet, the biological basis of atherosclerotic plaque development and composition across ancestries remains essential underexplored.

We hypothesized that genetically determined ancestry is associated with morphological and molecular features of atherosclerotic plaques. Leveraging the Athero-Express Biobank Study, an ongoing Dutch cohort with deep histological and transcriptomic profiling of plaques, we analyzed data from 1,944 patients after genotype quality control and ancestry inference using principal component analysis against 1000 Genomes. Two ancestry groups were identified, European (n=1,866) and non-European (n=51), reflecting Netherlands' migratory history.

Demographics were largely comparable between groups, however, ordinal logistic regression showed non-European ancestries had higher odds of increase plaque vulnerability (OR=1.67, 95% CI 1.01-2.77, p = 0.0450), a finding that remained robust after down sampling. Differential gene expression analysis highlighted *NLGN4X* and *SERPINF1*, genes linked to synaptic adhesion and vascular homeostasis, among the top differentially expressed genes. Pathway and single-cell enrichment analyses further revealed consistent enrichment of inflammation-related biological processes and diseases, including through integration with genome-wide association study data.

Our findings support that genetic ancestry correlates with morphological and molecular plaque composition, with non-European patients showing more inflammatory, higher-risk plaque features, including inflammatory signatures. Increased ancestral diversity in vascular biology research is critical for understanding atherosclerotic pathophysiology and develop equitable and personalized therapeutic strategies.

Letter

Cardiovascular disease (CVD) is the primary global cause of death, accounting for nearly 17.9 million deaths annually, with over 75% occurring in low- and middle-income countries[1] . Atherosclerosis, its main driver, is influenced by sex, aging, smoking, hypertension, family history and genetics, type 2 diabetes, lifestyle, and dyslipidemia – risk factors that correlate with socio-economic status and may vary across ethnicities[2–4]. However, they fail to fully explain observed ethnic disparities in relative rates of CVD[3,4]. For example, coronary artery calcification (CAC) – a marker of subclinical atherosclerosis – steadily increases with age, irrespective of gender or ethnicity, yet CAC is more prevalent in individuals of European descent, and more so among men compared to women[3–6]. These results may be confounded by self-reported ethnicity, however, genome-informed ancestry has been shown to correlate with subclinical atherosclerosis[5]. Thus, investigating the biological basis of atherosclerotic plaque development across ancestries remains essential.

Histological analysis is currently the gold standard method to evaluate atherosclerotic plaque composition[7,8]. High-risk lesions are characterized by variable calcific, inflammatory, lipid, and thrombotic components increasing risk of rupture and ischemic events[7,9]. While pathological studies revealed a higher extent of calcified coronary lesions in individuals of European ancestry[8], it remains unclear whether this extends to other morphological[9] or molecular plaque features. We hypothesized that genetically determined ancestry is associated with plaque composition, independent of risk factors.

We leveraged the Athero-Express (AE) biobank study[7] – a Dutch cohort with deep histological and molecular profiling[10–12] of plaques. After quality control and ancestry inference using principal component analysis (PCA, relative to the 1000G phase 3 reference), data for 1,944 patients remained, forming two groups: European (n=1,866) and non-European (n=51), reflecting Netherlands' migratory history[13–15] (**Figure 1**). Demographics were largely similar between Europeans (n=1,866) and non-Europeans (n=51), including age, sex, diabetes prevalence, hypertension, BMI, renal function, cholesterol and hsCRP levels, symptom prevalence and smoking behaviour.

To assess plaque vulnerability, we used the plaque vulnerability index (PVI, a histological score incorporating collagen, macrophages, smooth muscle cells, and lipid content[7]. Each trait was scored as stable or unstable, yielding a cumulative vulnerability score from 0 to 4. Those with non-European ancestries had higher odds of increased PVI (OR=1.67, 95% CI 1.01-2.77, p = 0.045, **Figure 1**), even after adjusting for age, gender, year of surgery, and genotyping platform in ordinal logistic regression (OR=1.69, 95% CI 1.01-2.82, p = 0.047). Sensitivity analyses using down-sampling (20-fold) of Europeans revealed a median OR=1.13, 95% CI 1.03-1.25, 18/20, $p_{binomial}$ < 0.001) indicating our results are robust with regards to the association with PVI. Further dissection of the role of individual plaque composition features revealed that non-Europeans showed lower odds of calcification (OR=0.42, 95% CI 0.21-0.80, p = 0.009) and higher odds of macrophage-rich plaques (OR=2.18, 95% CI 1.11-4.26, p = 0.023). Which was especially true for male sex: OR=0.75 95% CI 0.61-0.93 p=0.009 for calcification, and 1.4 95% CI 1.41-2.15 for macrophages, p=1.9x10^{-7}, respectively. This is consistent with earlier work that men of European descent have increased risk for CAC, underscoring that our study is sufficiently powered to detect established sex-related differences in plaque composition[6]. Our findings suggest a potential ancestry-related difference in inflammatory versus calcified plaque

phenotypes, where non-European patients exhibit higher inflammation (macrophages) and lower mineralization (calcification), consistent with the elevated overall plaque vulnerability observed.

Transcriptomic analyses of 1,028 plaques revealed 1,305 genes differentially expressed ($p_{nominal}<0.05$) out of 14,612 ($p_{binomial}$ = $7.11x10^{-87}$) between ancestry groups, including *NLGN4X* and *SERPINF1* (log_2FC=+0.61 and +0.78, respectively, FDR < 0.09) linked to synaptic adhesion[16] and vascular homeostasis[17,18]. *MMP1*, *EGR2*, *WHAMM*, and *SERPINF1* were upregulated in non-Europeans, whereas *NLGN4X*, *NRXN2*, *NR4A2*, and *ZNF432* were markedly downregulated (**Figure 1**). To further investigate the underlying mechanisms, we performed gene-set enrichment analyses using fgsea[19] and MSigDB[20] hallmark pathways revealing inflammation-related signatures including epithelial-to-mesenchymal transition, inflammatory response, apoptosis, and coagulation ($p_{adjusted}$ < 0.05). To contextualize the ancestry-associated transcriptional changes we used Enricher[21] to assess enrichment of gene-disease associations (DisGeNET[22]) showing overrepresentation of genes affecting diseases where the immune-driven diseases, such as inflammatory bowel disease, ulcerative colitis, neoplasms, arthritis, and abdominal aortic aneurysms ($p_{adjusted}$ < 0.05). Assessing the enrichment of the DEGs in plaque-derived single-cell RNAseq (n=39) data suggest a role for inflammatory cells, specifically macrophages. Enrichment against GWAS Catalog[23] further highlighted associations with circulating myeloid cell traits such as neutrophil-to-lymphocyte ratio, eosinophil percentage and mean corpuscular hemoglobin ($p_{adjusted}$ < 0.05). To assess cardiovascular relevance, we tested for enrichment of ancestry-associated genes against those linked to coronary artery disease[24], CAC[25], and carotid IMT[26] using MAGMA based gene-level GWAS summary statistics[27]. All traits showed higher-than-expected overlap via binomial testing (CAC: 106 nominally significant in CAC GWAS/1190 nominally significant in our DGEA; CAD: 149/1189; cIMT: 124/1190; all $p < 1.0 \times 10^{-16}$). These complimentary enrichment analyses underscore the robustness of our findings and suggest biological relevance of the ancestry-associated transcriptional difference to atherosclerotic disease.

Despite the strengths of this study – including histological analyses, genetic inference and transcriptomic data – several limitations should be noted. While this is the largest ancestry-focused multimodal study of carotid plaques to date, the small non-European sample (n=51) limits generalizability. The cohort reflects advanced disease, limiting the extrapolation to earlier subclinical stages of atherosclerosis. Socio-economic status was not directly assessed and may confound ancestry-related differences, although demographics did not differ between ancestry groups.

Our findings support that genetic ancestry is associated with plaque composition with non-European patients showing higher-risk plaque features. Increased representation of diverse ancestries in vascular biology research is critical for understanding atherosclerotic pathophysiology and to improve the design of equitable and personalized therapeutic strategies.

CREDiT authorship contributions

Acknowledgements

Dr. Sander W. van der Laan is funded through EU H2020 TO_AITION (grant number: 848146), EU HORIZON MIRACLE (grant number: 101115381), and Health~Holland PPP Allowance 'Getting the Perfect Image'. Dr. Clint L. Miller is funded by National Institutes of Health grants (R01HL148239, R01HL164577, and U01DK142283), Leducq Foundation 'COMET' (24CVD02) network, and AHA Transformational Project Award (24TPA1300556). Dr. Sander W. van der Laan and Dr. Clint L. Miller is funded by CZI Data Insights grant 'MetaPlaq' and EU HORIZON NextGen (grant number: 101136962).

We are thankful for the support of the Leducq Fondation 'PlaqOmics' (18CDV-02) and 'AtheroGen', and the Chan Zuckerberg Initiative 'MetaPlaq'. The research for this contribution was made possible by the AI for Health working group of the EWUU alliance (https://aiforhealth.ewuu.nl/). The collaborative project 'Getting the Perfect Image' was co-financed through use of PPP Allowance awarded by Health~Holland, Top Sector Life Sciences & Health, to stimulate public-private partnerships.

Disclosures

Dr. Sander W. van der Laan and Gerard Pasterkamp received Roche funding for unrelated work. Roche had no part in this study, neither in the conception, design, and execution of this study, nor in the preparation and contents of this manuscript. Dr. Clint L. Miller received grant support from AstraZeneca for work unrelated to the current study.
ChatGPT for macOS was used to check the text for spelling, grammar and syntax.

Figure 1: Genetically determined ancestry is associated with morphological and molecular plaque features.

References

1. Cardiovascular diseases. https://www.who.int/health-topics/cardiovascular-diseases/.
2. Lusis, A. J. Atherosclerosis. *Nature* **407**, 233–241 (2000).
3. Kronmal, R. A. *et al.* Risk factors for the progression of coronary artery calcification in asymptomatic subjects: results from the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* **115**, 2722–2730 (2007).
4. Budoff, M. J. *et al.* Ethnic differences of the presence and severity of coronary atherosclerosis. *Atherosclerosis* **187**, 343–350 (2006).
5. Gebreab, S. Y. *et al.* Genetic ancestry is associated with measures of subclinical atherosclerosis in African Americans: the Jackson Heart Study: The Jackson Heart Study. *Arterioscler. Thromb. Vasc. Biol.* **35**, 1271–1278 (2015).
6. Bild, D. E. *et al.* Ethnic differences in coronary calcification: the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* **111**, 1313–1320 (2005).
7. Verhoeven, B. A. N. *et al.* Athero-express: Differential atherosclerotic plaque expression of mRNA and protein in relation to cardiovascular events and patient characteristics. Rationale and design. *Eur. J. Epidemiol.* **19**, 1127–1133 (2004).
8. Otsuka, F., Sakakura, K., Yahagi, K., Joner, M. & Virmani, R. Has our understanding of calcification in human coronary atherosclerosis progressed? *Arterioscler. Thromb. Vasc. Biol.* **34**, 724–736 (2014).
9. van Lammeren, G. W. *et al.* Atherosclerotic Plaque Vulnerability as an Explanation for the Increased Risk of Stroke in Elderly Undergoing Carotid Artery Stenting. *Stroke* **42**, 2550–2555 (2011).
10. van der Laan, S. W. *et al.* Genetic Susceptibility Loci for Cardiovascular Disease and Their Impact on Atherosclerotic Plaques. *Circ Genom Precis Med* **11**, e002115 (2018).
11. Mokry, M. *et al.* Transcriptomic-based clustering of human atherosclerotic plaques identifies subgroups with different underlying biology and clinical presentation. *Nature Cardiovascular Research* **1**, 1140–1155 (2022).
12. Siemelink, M. A. *et al.* Smoking is Associated to DNA Methylation in Atherosclerotic Carotid Lesions. *Circ Genom Precis Med* **11**, e002030 (2018).
13. Hoeveel inwoners hebben een herkomst buiten Nederland. *Centraal Bureau voor de Statistiek* https://www.cbs.nl/nl-nl/dossier/dossier-asiel-migratie-en-integratie/hoeveel-inwoners-hebben-een-herkomst-buiten-nederland.
14. Boomsma, D. I. *et al.* The Genome of the Netherlands: design, and project goals. *Eur. J. Hum. Genet.* **22**, 221–227 (2014).
15. Deelen, P. *et al.* Improved imputation quality of low-frequency and rare variants in European samples using the "Genome of The Netherlands." *Eur. J. Hum. Genet.* **22**, 1321–1326 (2014).
16. Lehr, A. W. *et al.* Phosphorylation of NLGN4X regulates spinogenesis and synaptic function. *eNeuro* **12**, ENEURO.0278-23.2025 (2025).
17. Wang, H. *et al.* Pigment epithelial-derived factor deficiency accelerates atherosclerosis development via promoting endothelial fatty acid uptake in mice with hyperlipidemia. *J. Am. Heart Assoc.* **8**, e013028 (2019).
18. Shao, X. *et al.* Integrated single-cell RNA-seq analysis reveals the vital cell types and dynamic development signature of atherosclerosis. *Front. Physiol.* **14**, 1118239 (2023).

19. Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2016) doi:10.1101/060012.

20. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst.* **1**, 417–425 (2015).

21. Xie, Z. *et al.* Gene set knowledge discovery with Enrichr. *Curr. Protoc.* **1**, e90 (2021).

22. Piñero, J. *et al.* DisGeNET: a comprehensive platform integrating information on human disease-associated genes and variants. *Nucleic Acids Res.* **45**, D833–D839 (2017).

23. Cerezo, M. *et al.* The NHGRI-EBI GWAS Catalog: standards for reusability, sustainability and diversity. *Nucleic Acids Res.* **53**, D998–D1005 (2025).

24. Nelson, C. P. *et al.* Association analyses based on false discovery rate implicate new loci for coronary artery disease. *Nat. Genet.* **49**, 1385–1391 (2017).

25. Kavousi, M. *et al.* Multi-ancestry genome-wide study identifies effector genes and druggable pathways for coronary artery calcification. *Nat. Genet.* **55**, 1651–1664 (2023).

26. Franceschini, N. *et al.* GWAS and colocalization analyses implicate carotid intima-media thickness and carotid plaque loci in cardiovascular outcomes. *Nat. Commun.* **9**, 5141 (2018).

27. Slenders, L. *et al.* Intersecting single-cell transcriptomics and genome-wide association studies identifies crucial cell populations and candidate genes for atherosclerosis. *Eur Heart J Open* (2021) doi:10.1093/ehjopen/oeab043.

Athero-Express Biobank Study
3,633 patients with arterial stenosis

3 - RNA sequencing

Exon Intron Exon

n = 1,028 **overlapping genetic** and **histology** data

2 - Genotyping

PCA

1 - Endarterectomy

plaque processing
histopathology staining

CD34, CD66b, CD68, EvG, Fibrin, GlycC, HE, SMA, collagen

European
n = 1,866

non-European
n = 51

Differential Gene Expression Analysis

NS
Log₂ FC
p-value
p – value and log₂ FC
14,612 protein-coding genes tested

Gene-Set Enrichment analyses and **single-cell RNAseq analyses**

Top 30 DEGs: Mean per Ancestry Group

Module Score

Genetically determined **ancestry** is
associated with **morphological**
and **molecular plaque features**

non-European ancestry

Histological Analysis

higher odds for **unfavorable plaque composition**

lower odds **calcific** plaque

higher odds **inflammatory** plaque