Escalating burden and mortality of carbapenem-resistant *Klebsiella pneumoniae* species complex infections in Bangladeshi infants

Yogesh Hooda[1, #], Arif Mohammad Tanmoy[1], Naito Kanon[1], Hafizur Rahman[1], Mohammad Shahidul Islam[1], Zabed Bin Ahmed[1], Afroza Akter Tanni[1], Md Mobarok Hossain[1], Md Hasanuzzaman[1], Sharmistha Goswami[1], Tasnim Jabin[1], Rajib Chandra Das[1], Md Jamal Uddin[1], Belal Hossain[1], Shampa Saha[1], Anannya Barman Jui[1], Mohammod Shahidullah[2], Nobo Krishna Ghosh[3], AKM Shamsuzzaman[3], Nahid Sultana[4], Sanjoy Kanti Biswas[4], Feroza Aktar[5], Wazir Ahmed[5], Mohammed Monir Hossain[6], Mahbubul Hoque[6], ASM Nawshad Uddin Ahmed[1,7], Samir K Saha[1,8], Senjuti Saha[1,#]

[1] Child Health Research Foundation, Dhaka, Bangladesh

[2] Department of Neonatology, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh

[3] Department of Pediatrics, Dr M R Khan Shishu Hospital & Institute of Child Health, Dhaka, Bangladesh

[4] Department of Microbiology, Chattogram Maa O Shishu Hospital Medical College, Chattogram, Bangladesh

[5] Department of Neonatology, Chattogram Maa O Shishu Hospital Medical College, Chattogram, Bangladesh

[6] Department of Neonatal Medicine, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh

[7] Department of Pediatrics, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

[8] Department of Microbiology, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

[#]Corresponding author

Senjuti Saha

Deputy Executive Director

Child Health Research Foundation

senjutisaha@chrfbd.org

AND

Yogesh Hooda

Senior Scientist

Child Health Research Foundation

yhooda@chrfbd.org

Abstract

Klebsiella pneumoniae infections in young infants are an escalating threat in low- and middle-income countries, yet robust longitudinal data integrating hospital burden, clinical outcomes, antimicrobial, and genomics remains scarce. We performed an 18-year (2004-2021) prospective, multicenter genomic epidemiology study across four hospitals in Bangladesh. Among 122,353 enrolled children from whom blood or cerebrospinal fluid cultures were performed, 1,600 (1.3%) yielded culture-confirmed *Klebsiella pneumoniae* species complex (KpSC) isolates. Positivity increased from 16 per 1,000 cases tested in 2004 to 37 per 1,000 in 2021. Hospital case-fatality rate (CFR) rose from 21.4% to 51.4% during the study, paralleling the emergence and expansion of carbapenem resistance, first detected in 2008 and reaching 81% of isolates by 2021. Neonates accounted for 80.5% of infections and experienced a CFR of 40.8%. Whole-genome sequencing of 599 representative isolates revealed four KpSC species, 145 sequence types and 92 capsular alleles. Global high-risk clones ST11, ST16 and ST147 harbouring NDM-type carbapenemases dominated recent cases. These findings document the increasing resistance and mortality associated with KpSC infections amongst neonates in Bangladesh, underscoring the urgent need for strengthened infection prevention & control, equitable access to effective combination therapies, and vaccine-based preventative strategies.

Introduction

Gram-negative bacteria of the genus *Klebsiella* are ubiquitous in the environment and form part of the commensal flora of healthy humans[1,2]. Among them, *Klebsiella pneumoniae* and six related members of the *K. pneumoniae* species complex (KpSC) can also cause a broad spectrum of infections, including soft tissue, surgical wound, urinary tract, respiratory tract, bloodstream and central nervous system infections[3], particularly in hospital settings and immunocompromised individuals. Neonates (0-27 days old) are particularly susceptible to systemic infections by KpSC[4–8].

Reports of multidrug-resistant (MDR) *K. pneumoniae* have risen sharply worldwide over the past two decades, with the steepest increases observed in low and middle income countries (LMICs)[9–11]. The emergence of carbapenem-resistant *K. pneumoniae* is associated with high mortality rates[12], and has further constricted therapeutic options, leading to it being declared a critical priority pathogen by the WHO.

With the antibiotic pipeline faltering, alternative strategies are gaining traction. Recently, combination drugs that contain a beta-lactam antibiotic and a beta-lactamase inhibitor have shown to be efficacious for treatment of carbapenem resistant *K. pneumoniae* infections, but their activity depends on the underlying carbapenemase gene as these do not work on all family of metallo-β-lactamases.[13] Passive immunotherapies and maternal vaccines targeting capsular (K) or lipopolysaccharide (O) antigens are also under development[14–17]. However, *K. pneumoniae* exhibits extraordinary antigenic heterogeneity, with at least 13 O-types and 186 K-types documented to date[18].

Selecting vaccine valences therefore demands precise, contemporary data on

circulating serotypes, resistance genes and clonal backgrounds.

Additionally, global genomic sequencing efforts have found that other KpSC members

such as *K. quasipneumoniae* and *K. variicola* cause invasive diseases in humans, yet

routine biochemical tests cannot differentiate them from *K. pneumoniae*[2]. Failing to

recognize this hidden diversity can mask transmission chains, underestimate true

burden and misguide therapy, particularly in low-resource settings that rely on

phenotypic methods.

In this study we: (i) quantify temporal trends in the hospital burden and severity of

invasive KpSC infections in Bangladeshi children; (ii) describe the evolution of

phenotypic and genotypic AMR, including carbapenemase rise and spread; and (iii)

characterize the species, sequence types, O- and K-loci of a representative subset of

isolates using whole-genome sequencing. Overall, we generate data to inform future

infection-control measures, empirical-therapy guidelines and multivalent-vaccine design

in Bangladesh and neighboring countries.

Results

The burden and severity of KpSC infections in Bangladesh is increasing.

Between 2004 and 2021, 122,353 cases had blood and/or CSF samples cultured at the

treating physician's discretion (Sup Fig. 1); all were enrolled in this study.

Bacteriological-culture and biochemical testing confirmed that 1,600 (1.3%) infections

were caused by the KpSC. A sharp rise is observed in the number of KpSC isolates

starting in 2018 (Fig. 1a). Up to 2017, the mean positivity rate for KpSC was 16 per

1,000 cases where blood cultures were performed, which climbed to 18, 34 and 37 per

1,000 suspected sepsis cases in 2019, 2020 and 2021, respectively (Fig. 1b). Seventy-

four percent (1,182/1,600) of *KpSC*-positive cases were reported from one hospital,

BSHI (Sup Fig. 2a), which also accounted for the highest positivity rate among the four

hospitals since 2015 (Sup Fig. 2b).

The median age of KpSC-positive cases at hospital admission was 5 days (IQR: 2-18)

and at blood draw was 8 days (IQR:4-21) (Fig. 1c). Overall, 1,352/1,600 (84.5%) KpSC

infections occurred in infants <2 months old, and 1,288/1,600 (80.5%) in neonates (first

28 days old). The hospital case fatality rate (CFR) was highest during the first month of

life (41.5%) and declined for infants older than three months (Fig. 1c). Overall CFR was

38.1%. Among neonates, the median age at sample collection was 6 days (IQR: 3-21),

and the CFR was consistent throughout the first month of life (Fig. 1d). Most infections

1,076/1,600 (67.3%) were detected within 72 hours of admission (Sup Fig. 3a); the

median interval between admission to sample collection was 24 hours (IQR: 0-4).

The severity of KpSC infection is highlighted by hospital outcomes: 610/1,600 (38.1%)

cases died at the hospital, 174/1,600 (10.9%) left against medical advice, and

816/1,600 (51.0%) were discharged from the hospital (Sup Table 1). The average

hospital stay was 8 days (IQR, 4-14), but 64% babies who died, succumbed within the

first 5 days of admission (Sup Fig. 3b). CFR started increasing in 2016 (Fig. 2a), with

the odds of death in 2021 being 3.9 times higher than in 2004 (95 % CI 1.5–9.8;

Fig. 2b). The proportion of patients leaving hospital against medical advice varied over

the study period, ranging from ~4% to 20% per year, without a consistent increasing

trend over time (Sup Fig. 4a). Sensitivity analyses incorporating alternative assumptions

regarding LAMA outcomes demonstrated that the temporal increase in mortality was

preserved across all scenarios (Sup Fig. 4b).

Increasing severity of KpSC infections correlates with rise in carbapenem

resistance.

We investigated the possible drivers of escalating mortality by comparing yearly trends

in clinical features (Sup Fig. 5) and antimicrobial susceptibility profiles for 10 antibiotics

(meropenem, imipenem, ceftriaxone, ampicillin, gentamicin, netilmicin, ciprofloxacin,

amikacin, chloramphenicol, and cotrimoxazole) proposed by CLSI[19] (Sup Fig. 6). KpSC

isolates from Bangladesh exhibited high resistance against most antimicrobials used,

including amikacin, cotrimoxazole, gentamicin, netilmicin, and ciprofloxacin. The first

case of carbapenem resistance was detected in 2008, and with noticeable increase

during the study period, by 2021 >80% of KpSC isolates were resistant to carbapenem

(Fig. 2c). The rising carbapenem resistance correlated with increasing mortality, with the

odds of death being 3.2-fold higher (95% CI: 2.3-4.6) relative to carbapenem

susceptible isolates (Sup Fig. 7).

To further assess the impact of carbapenem resistance, we stratified cases into two

epochs, 2004–2015 and 2016–2021, and compared CFRs for carbapenem-resistant

versus carbapenem-susceptible KpSC infections (Fig. 3d). During 2004–2015, CFRs

were similar irrespective of resistance phenotype. From 2016 onward, carbapenem-

resistant infections exhibited a markedly higher CFR than susceptible infections,

however, mortality increased for both groups. In multivariable logistic regression

analyses, carbapenem resistance remained consistently associated with increased

odds of in-hospital CFR across all models, including those with individual adjustments

for covariates of age group, sex, hospital site, timing of culture collection, and admission

epoch (Sup Fig. 8). Upon simultaneous adjusting for neonatal status, hospital site, and

admission epoch, the estimated effect size for carbapenem resistance was attenuated.

This likely reflects shared variance between carbapenem resistance, temporal trends,

and referral patterns, as well as reduced effective sample size, rather than absence of

an association. Taken together, these analyses suggest, while carbapenem resistance

is a key driver of poor outcomes, additional factors, i.e. host, pathogen and/or

environmental, have contributed to the post-2016 surge in KpSC-associated deaths.

KpSC infections exhibit high genomic and antigenic diversity

To further characterize the KpSC isolates, we sequenced a representative subset of 527

isolates (527/1461, 36.1%) collected across the entire study period from infants younger

than two months - the age group with the highest burden (Sup Fig. 9a, Sup Table 2, Sup

Data 1)[26]. Some bio banked strains from 2004-2009 could not be revived, and hence

fewer representatives from those years were available. To partially overcome this

limitation, we included 72 sequences from three other hospitals from 2004-2009 (Sup

Table 2, Sup Data 1). Among the total 599 isolates sequenced, 503 (84.0%) genomes

were *K. pneumoniae*, 72 (12.0%) were *K. quasipneumoniae* subsp *similipneumoniae*,

20 (3.3%) were *K. quasipneumoniae* subsp *quasipneumoniae*, and 4 (0.7%) were *K.*

variicola (Fig. 3a). The annual distribution of these species remained broadly consistent

throughout the study period (Sup Fig. 9b). Infections caused by *K. pneumoniae* were

associated with a 2.18 (95% CI 1.06-4.94) fold higher case fatality rate than infections

due to other KpSC species (Sup Fig. 10).

Maximum-likelihood phylogenies of the 599 sequenced KpSC genomes (Fig. 3a) and

the 503 *K. pneumoniae* genomes (Fig. 3b) demonstrated high genetic diversity amongst

KpSC isolates identified in this study. We identified 145 distinct sequence types (STs);

however, the ten most common STs accounted for 52% of all isolates. The predominant

lineage was *K. pneumoniae* ST11, 97% of which were identified between 2019 and

2021. These ST11 genomes were sampled across three of the four surveillance

hospitals in Dhaka, Chattogram and Mirzapur, yet showed minimal within-lineage

diversity, consistent with a nationwide clonal expansion (Fig. 3c, Sup Fig 11). Metadata

reinforcing this observation include that among ST11 cases, the median interval from

admission to blood draw was 2 days (IQR 0-5); 53% (36/68) of samples were collected

within 48 hrs of hospital admission (Fig 3c). Mapping of home addresses shows infants

originating from 22 districts (Fig. 3d).

The yearly distribution of the 10 most common STs for *K. pneumoniae* are shown in Fig.

4a*.* Some of the common STs like ST11, ST147 and ST16 exhibited association with

higher CFR (Sup Fig. 10), however when adjusted for carbapenem resistance, barring

ST11, no other ST exhibited statistically significant association with higher mortality

(Sup Table 3). Compared to a global database of *K. pneumoniae,* most of the common

STs in Bangladesh have been reported in other countries, and ST11, 16, and 147

belong to global problem clones (Fig. 4b).

We next analyzed genomic carbapenem-resistance and virulence determinants in the

sequenced KpSC genomes and compared these with meropenem susceptibility results,

and mortality respectively. Genomic prediction of the meropenem phenotype achieved a

sensitivity of 96.3% and a specificity of 65.4% and matched the phenotypic carbapenem

trendline (Sup Fig. 9c). The main carbapenemase alleles detected were blaNDM-5,

blaNDM-1, along with multiple other less common variants (Table 1), indicating that

different KpSC lineages in Bangladesh have acquired carbapenem-resistance genes

independently. For virulence, we specifically looked for presence of genes encoding

siderophore-uptake systems (aerobactin and salmochelin) and hypermucoid phenotype

(RmpA/RmpA2) (Sup Fig. 10). None of the known virulence factors were associated

with increased mortality, as none of the dominant carbapenem-resistant strains (ST11,

ST16 and ST147) contained these virulence factors. In 2019, we identified a single

ST43 isolate with both bla-OXA181 and aerobactin encoding genes. While this was an

isolated case, such carbapenem-resistant hypervirulent strains should be continuously

reported.

To characterize antigenic diversity relevant to vaccine design, we analyzed the genomic

regions encoding the capsular polysaccharide (K locus) and O antigen. In total, 92

unique K loci and 11 unique O loci were identified. The ten most prevalent O loci and K

loci together accounted for 99.8% (Fig. 5a, Sup Fig. 12) and 58.3% (Fig. 5c, Sup Fig.

13) of all isolates, respectively, with year-to-year variations (Fig. 5b,5d). The combined

share of the ten dominant O loci stayed above 95% every year, whereas the share of

the ten dominant K loci was highly variable, oscillating between 0% and 86.7%.

Discussion

Geographically and temporally representative epidemiological and genomic data from

high-burden LMICs are essential to guide policy and intervention against KpSC

infections. Here we report 18 years of surveillance from four Bangladeshi pediatric

hospitals, detailing trends in burden, antimicrobial resistance and genomic diversity.

From 122,353 bacteriological culture of blood and/or CSF samples conducted between

2004 and 2021, we identified 1,600 KpSC isolates. Neonates comprised 80.5% of cases

and had a hospital case-fatality rate of 40.8%, compared with an overall CFR of 38.1%.

In a multi-country study on neonatal infections, BARNARDS, conducted in seven LMICs

including Bangladesh from Nov 2015 to Dec 2017, the CFR due to KpSC infection was

22.4% (58/258), compared to 35.7% (45/126) during the same period in this study.[6] In

the NeoObs study done across 11 countries from 2018-2019 including Bangladesh,

neonatal KpSC case fatality rate was 21.1%.[7] In NeoOBS Bangladesh site, the CFR

was 22% (2/9), compared to 39% (120/307) during that period in this study. Within

neonates, infections were most commonly detected on day 5, as seen in previous

studies [BARNARDS and NeoObs][7,8]. Collectively, these studies highlight the burden of

KpSC infections amongst neonates globally. The higher mortality we observed was

driven by BSHI, which functions as the pediatric referral center for Bangladesh,

receiving the most critically ill infants from across the country; this referral bias is likely

to inflate both the apparent burden, and the CFR compared with primary or secondary

facilities included in the other studies.

Overall, the alarming rise in neonatal deaths threatens to erode vaccine-driven gains in

child survival across LMICs. Thirteen percent of infections were detected within the first

three days of life, including 3% within 24 hours of birth, and 67% of infants were culture-

positive within 72 hours of admission. These timelines point to transmission in maternity

units, delivery clinics, or the community, prior to referral to the study hospitals.

At BSHI, the KpSC positivity rate rose from 16 per 1000 suspected sepsis cases in

2004 to 37 per 1000 in 2021, and the hospital-specific case-fatality rate climbed from a

mean of 20.7% (2004–2014) to 51.4% in 2021. Carbapenem resistance was associated

with death, with an adjusted odds ratio of 3.2 (95% CI: 2.3-4.6). A positive correlation

between carbapenem resistance and mortality has reported previously for KpSC and

other bacterial infections[34]; however, the observational design of our study precludes

inferring causality. Consistent with this association, the case-fatality rate of carbapenem-

resistant KpSC infections was significantly higher than that of carbapenem-sensitive

KpSC infections (46.3% vs 25.4%). Notably, mortality among carbapenem-susceptible

infections also increased after 2015 compared with earlier (31.1% vs 20.3%).

Adjustment for neonatal status, admission time, and hospital site reduced precision of

carbapenem resistance mortality effect estimates. These factors are likely themselves

closely linked to the emergence and clinical impact of carbapenem-resistant KpSC and

thus may lie on the causal pathway rather than acting as independent confounders.

The excess mortality in carbapenem-susceptible cases after 2015 suggests that factors

beyond drug resistance are driving severity. Several, not mutually exclusive, hypotheses

warrant exploration: (i) bacterial adaptation – dominant lineages may have acquired

additional virulence determinants[3] or stress-tolerance genes that enhance survival in

neonatal wards[35]; (ii) changing obstetric practice – the sharp rise in Caesarean

deliveries in Bangladesh increases early hospital exposure and separates infants from

protective maternal microbiota[36,37]; (iii) ward overcrowding and resource strain – more

institutional births and limited isolation facilities hamper effective infection-prevention

and control[38]; (iv) referral bias – improved road networks now funnel the sickest infants

who fail first-line therapy or need advanced support to BSHI, inflating observed

mortality[39] (v) comorbidities and other host-factors **–** rising prevalence of prematurity,

low birth-weight and congenital anomalies may increase vulnerability to sepsis[40–42]; and

(vi) climate stress – warmer, more humid conditions, extreme weather events like

flooding and heat-related infrastructure strain can favor environmental persistence and

spread of KpSC within hospitals[43,44]. These possibilities highlight the need for integrated

clinical, microbiological and health-systems investigations to disentangle drivers of the

continued rise in KpSC-associated neonatal deaths.

Several genomic studies have now established that biochemically identified *K.*

pneumoniae infections can be other members of KpSC[2,26,33]. In this study, 15% were *K.*

quasipneumoniae (*subsp. simillipneumoniae* & *subsp. quasipneumoniae*) and *K.*

variicola, rather than *K. pneumoniae*. Infections due to *K. pneumoniae* carried a higher

hospital CFR (32.9% vs 18.3%) and a greater burden of carbapenem resistance genes.

These observations suggest that pharmacological interventions directed solely at *K.*

pneumoniae, whether vaccines, monoclonal antibodies or new antimicrobials, could

create ecological space for the related species to fill. Future control strategies must

therefore target the entire KpSC repertoire to avoid pathogen replacement and should

also include broad IPC.

We also observed a high diversity of carbapenem-resistance determinants in

Bangladeshi KpSC isolates, with the *blaNDM-1* gene predominating[6,45]. Because KpSC

isolates are predicted to remain susceptible to the ceftazidime-avibactam + aztreonam

combination [13], this regimen is a promising therapeutic option. Yet neither drug is widely

available in Bangladesh due to their cost, particularly in the low-income communities

served by our surveillance hospitals, creating a critical treatment gap which needs to be

addressed through changes in drug procurement recommendations/policies. Clinical

trials are therefore urgently needed in Bangladesh to evaluate the regimen's efficacy,

safety and implementation feasibility and to guide policy aimed at reducing KpSC-

associated neonatal mortality.

Overall, the Bangladeshi KpSC population is highly heterogeneous: the 599 genomes

were distributed across 145 STs, and the dominant lineages shift and vary over time.

Notably, several dominant sequence types that emerged or expanded after 2015

accounted for a large fraction of deaths, suggesting that clonal shifts may have

contributed to the observed increase in case fatality. Notably ST11, ST147 and ST16

have been isolated in various other countries[26,33]. The clinical impact is best illustrated

by ST11, that is associated with 2.41 times (95% CI:1.39–4.24, p = 0.001) increased

odds of mortality compared to other KpSC strains, even after adjusting for carbapenem

resistance. Almost all ST11 isolates (n = 68) were recovered between 2019 and 2021,

consistent with a recent clonal expansion. None of the ST11 isolated possessed any

known virulence factors such as aerobactin, salmochelin and RmpA. The median

interval from admission to blood draw was 24 hours (IQR 0–5); 53% of cultures were

taken within 72 h of hospital admission. Cases originated from 22 districts, and most

infants were admitted directly from home or local clinics rather than transferred from

other hospitals. Minimal genetic divergence, very short admission-to-culture intervals

and wide geographic scatter together suggest transmission in maternity units, delivery

clinics or the community, settings upstream of tertiary infection-control measures. Future

studies to identify host, exposure, and care-related risk factors associated with infection

by high-risk lineages and poor outcomes are urgently needed. Such integrated

approaches will be essential to inform interventions that are tailored to the evolving

clonal structure of KpSC in hospital settings, especially neonatal wards.

Efforts to protect young infants against KpSC, particularly *K. pneumoniae*, now focus on

maternal vaccines and monoclonal antibodies that target the organism's surface

polysaccharides (capsular K antigen and lipopolysaccharide O antigen)[16]. Our data

highlight the antigenic hurdles such products must clear. Whole-genome analysis

predicted 92 distinct K (capsular) types, with the ten most common covering only 58%

of isolates. O-antigen diversity was narrower: the ten most frequent O loci

encompassed 99% of isolates, and their cumulative coverage remained >50 % each

year. Antibody access to the O-antigen may be partially constrained by the thick

polysaccharide capsule, which can shield the underlying lipopolysaccharide and limit

immune recognition in heavily encapsulated strains[46]. By contrast, K-type coverage

fluctuated markedly over time, reflecting turnover of dominant sequence types and the

strong linkage between ST and capsule genotype. These patterns indicate that a

multivalent formulation combining O- and K-antigens will be required for broad

protection in Bangladesh, yet the capsule's extensive and shifting diversity will remain a

major design challenge.

The findings of this study should be viewed within the context of certain limitations. First,

we did not collect information related to gestational age, birth weight and maternal

factors, such as maternal age, nutrition, referral status (inborn vs outborn), standardized

measures of illness severity at admission (e.g. SNAPPE-II) and mode of delivery. These

variables are recognized risk factors for KpSC infection, and their absence prevents

adjustment for host susceptibility. Second, our study lacked systematic data on

comorbidities including congenital anomalies, which may have contributed to the

observed mortality; nevertheless, previous CHAMPS investigations have implicated *K.*

pneumoniae as a leading cause of neonatal death.[8] Third, no pre-hospital records were

available, so we could not ascertain prior antibiotic exposure, treatment adequacy or the

likely place of acquisition,, illness severity at admission, hospital bed occupancy, staffing

levels, and time to antibiotic administration. These factors may plausibly contribute to

rising mortality andthe lack of such data limits our ability to disentangle pathogen-

specific effects from broader system-level drivers of mortality. Fourth, we sequenced a

representative subset of KpSC isolates; rare sequence types, capsular types or O-

antigen variants present in the isolates that were not sequenced may therefore have

been missed. Fifth, automated blood culture system (BACTEC) was introduced in BSHI,

KWMCH and MRKSH in 2016 and in CMOSH in 2017. Because parallel manual and

automated cultures were not performed during the transition, we lack internal cross-

validation data, precluding calculation of a reliable adjustment factor to account for this

methodological change. Although automation is likely to improve overall pathogen

recovery and reduce time to positivity, the rise in KpSC burden and mortality was not

temporally synchronous across sites, was driven predominantly by BSHI, continued to

increase for several years after implementation rather than showing an abrupt step

change, and was selectively enriched for carbapenem resistant lineages. Moreover,

prior studies suggest that automated systems increase overall positivity primarily

through improved detection of fastidious organisms, with similar recovery of common

pathogens such as *K. pneumoniae* across methods[47]. Taken together, BACTEC is

unlikely to explain the magnitude and selective nature of the observed increase.

Despite these constraints, the dataset provides one of the most comprehensive clinical–

genomic reports of paediatric KpSC infections available from South Asia. Rising

incidence, a four-fold increase in mortality and the dominance of blaNDM-positive global

clones expose a critical threat to neonatal survival. To mitigate this risk, policies should

include: (i) expansion of neonatal intensive-care capacity and skilled staffing at birth

facilities; (ii) stringent infection-prevention and control in labour wards and neonatal

units; (iii) continuous, denominator-based genomic surveillance to detect clonal shifts

and emerging resistance; and (iv) rapid access to effective therapeutics, including

context-specific trials of ceftazidime–avibactam plus aztreonam. Maternal vaccines or

monoclonal antibodies targeting the prevalent O-antigens could add a valuable layer of

protection, but the extensive and shifting capsule diversity documented here means

immunization must be embedded in a broader strategy that compliments vaccination

with improved clinical care, antimicrobial stewardship and robust infection control.

These recommendations are likely to be relevant beyond Bangladesh and should inform

intervention design across south Asia.

Methods

Ethics Statement:

This study analyzes *KpSC* isolates, along with associated clinical and laboratory

metadata, collected as part of the PneumoADIP study (2004-2008) and the Invasive

Bacterial Disease (IBD) study (2009-2021). Ethical approvals for data collection,

analysis and pathogen sequencing were obtained from the Ethical Review Board of the

Bangladesh Institute of Child Health (now known as Bangladesh Shishu Hospital &

Institute). All procedures were conducted in accordance with the National Ethical

Guidelines for Health Research in Bangladesh issued by the Bangladesh Medical

Research Council. We obtained written informed consent from the guardians of children

admitted to the study hospitals, from whom blood and/or cerebrospinal fluid (CSF)

samples were collected.

<u>Description of the study site:</u>

From 2004-2021, blood and/or CSF specimens were collected at four hospitals in

Bangladesh: a) Bangladesh Shishu Hospital and Institute (BSHI, formerly known as

Dhaka Shishu Hospital), a 650-bed tertiary level hospital based in Dhaka; b) Dr. M R

Khan Shishu Hospital and Institute of Child Health (MRKSH), a 250-bed primary -

secondary level hospital in Dhaka; c) Chattogram Maa-O Shishu Hospital (CMOSH), a

835-bed primary - tertiary level hospital in Chattogram, and d) Kumudini Women's

Medical College and Hospital (KWMCH), a 1050-bed based primary-tertiary level

hospital based in rural town Mirzapur. BSHI and CMOSH operate Level III neonatal

intensive care units, while MRKSH and KWMCH provide Level II neonatal care.

Children hospitalized in these hospitals who matched the WHO-defined clinical criteria

for pneumonia, severe pneumonia, severe disease, and meningitis were evaluated by

trained research physicians. Blood and/or CSF was drawn for culture at the treating

physicians' discretion. Demographic (age and sex) and clinical (symptoms, hospital

outcome, admission date, date of sample collection etc.) data were obtained for all

enrolled patients (Sup Table 1). Children were enrolled upon informed written consent if

they met the inclusion criteria.

<u>Laboratory procedures:</u>

Blood and/or CSF samples from patients were incubated in tryptic-soy broth and

incubated at 37°C for overnight then sub-cultured on blood agar, chocolate agar and

MacConkey agar media from 2004-2015. We introduced BACTEC for blood cultures in

BSHI, KWMCH and MRKSH, in 2016 and in CMOSH the following year; since then,

blood specimens were incubated in BACTEC culture systems for up to 120 hours, and

positive samples were subsequently sub-cultured on blood agar, chocolate agar and

MacConkey agar media. Upon growth, biochemical tests including, indole, urease, and

citrate utilization tests were performed to identify the genus and species. Isolates with

mucoid colony morphology, lactose fermentation on MacConkey agar, and a

biochemical profile of oxidase-negative, indole-negative, urease-positive, and citrate-

positive were confirmed as KpSC and stored at -80°C in STGG media (skim milk,

tryptone, glucose, and glycerin). These stocks served as the source for all subsequent

experiments performed in the study.

Antimicrobial susceptibility testing was performed using antibiotic discs by Kirby-Bauer

testing for: meropenem, imipenem, ceftriaxone, ampicillin, gentamicin, netilmicin,

ciprofloxacin, amikacin, chloramphenicol, and cotrimoxazole. Tests were interpreted

according the Clinical and Laboratory Standards Institute (CLSI) 2025 antibiotic

susceptibility testing clinical breakpoints[19].

From 2011 to 2019, the microbiology laboratories, as part of the WHO's Invasive

Bacterial Vaccine-Preventable Diseases (IB-VPD) surveillance, underwent external

quality assurance from UK-NEQAS.

<u>Sample selection criteria for whole genome sequencing (WGS):</u>

Approximately a third of KpSC isolates from children under the age of 2 months, were

randomly selected, keeping the distribution of isolates selected per year consistent. For

each calendar year, all eligible isolates were enumerated, and a target proportion

(~33%) was selected using computer-generated random numbers, ensuring that the

temporal distribution of sequenced isolates closely reflected the annual distribution of all

culture-confirmed cases. These selected isolates were re-cultured from biobank,

biochemically re-confirmed and sub-cultured on MacConkey agar media prior to DNA

extraction and whole genome sequencing.

To contextualize the selected isolates with other KpSC isolates from Bangladesh, in

addition to the isolates recovered in this study, we randomly selected 100 of 160 (62%)

KpSC isolates for WGS that were stored in our biobank collected between 2004-2008

from three additional hospitals that followed the same inclusion criteria: a) Dhaka

Medical College and Hospital, Dhaka, b) Chattogram Medical College and Hospital,

Chattogram, and c) Sir Salimullah Medical College and Hospital, Dhaka as part of the

pneumoADIP study.

<u>Genomic DNA extraction and sequencing:</u>

Total DNA from KpSC colonies was extracted using the QIAGEN DNA Mini extraction kit

(Cat: 56304, Qiagen, Hilden, Germany). Sequencing libraries were prepared from

extracted DNA using NEBNext Ultra II FS DNA Library kit (Cat: E7805, New England

Biolabs, MA, USA). Libraries were sequenced on an Illumina NEXTSeq2000 at the

Child Health Research Foundation, Dhaka, Bangladesh (n = 528; 150bp PE) and on the

HiSeq4000 platform at the University of Oxford, Oxford, UK (n = 71; 150bp PE).

Genome assembly

Sequence data quality was assessed using FastQC v0.11.5[20] and MultiQC v3[21]. Adapter

trimming was performed with Trimmomatic v0.39[22]. *De novo* genome assembly was

conducted using Unicycler v0.4.9[23], evaluated using Quast v.5.2.0[24], and annotated with

Prokka v.1.14.6[25]. Kleborate v2.2[26] was used to identify the sequence types (STs), AMR

genes, K and O antigen types. All generated sequence data have been submitted to

ENA (BioProject: PRJEB90555).

Phylogenetic Analysis

For reference-based genome mapping, Bowtie2 v2.3.4.1 was employed to align all

genomic data against the NCBI reference genome, *Klebsiella pneumoniae* HS11286

(Genbank accession: NC_016845.1)[27]. SAMtools v1.7.2 and BCFtools v1.7.2 were

utilized for variant calling, filtering candidate SNPs for a minimum base quality of 20, a

minimum of 5 high-quality reads, and greater than 75% read support (i.e., over 75% of

the reads supporting the SNP relative to the reference)[28]. SNPs were further filtered to

remove potential biases, including strand, mapping, and end biases (p-value < 0.001).

Additionally, SNPs located within phage regions (identified by PHASTER[29]), repeat

sequences (as noted in the reference GenBank), and recombination regions (detected

by Gubbins v2.3.1) were excluded[30]. The resulting filtered SNPs were then converted

into a SNP alignment for maximum likelihood phylogenetic tree (MLT) construction using

RAxML v8.2.11 (model: GTRGAMMA; 100 bootstraps)[31]. The alignment had 82,669

SNP positions from 472 *Klebsiella pneumoniae* isolates in our study. Tree visualization

and annotation was done using iTOL v6[32].

For global comparison of the STs of the *K. pneumoniae* isolates (n = 472) sequenced in

this study, publicly available *K. pneumoniae* genomes (n = 32,656) in PathogenWatch[33]

(retrieved on November 29, 2023) were used.

<u>Statistical analysis</u>

Statistical analyses were performed using custom scripts in R (version 4.3.1). The odds

ratio was calculated with using the *oddsratio* package, while the figures were generated

using the *ggplot2* and *Rcolorbrewer* and *ggsci* packages. Global and Bangladesh-

specific maps were made using the packages *rnaturalearth, ggspatial, maps* and

scatterpie. For ST11 geographic analysis, upazila, village or union information was used

instead of the exact address obtained during data collection. Hospital outcome was

classified as died, discharged, or left against medical advice (LAMA).

Hospital case-fatality rate (CFR) was defined as the proportion of culture-confirmed

KpSC cases that died during hospitalization, thus LAMA and discharged cases were

included in the primary CFR denominator. To evaluate the potential impact of LAMA on

mortality estimates, we performed a bounded sensitivity analysis under three

assumptions: (i) a primary analysis excluding LAMA cases (death / [death +

discharged]); (ii) a lower-bound estimate, assuming all LAMA cases survived (death /

total cases); and (iii) an upper-bound estimate, assuming all LAMA cases died ([death +

LAMA] / total cases). Year-specific mortality estimates, and 95% confidence intervals

were calculated for each scenario.

To assess the association between carbapenem resistance and in-hospital mortality, we

fitted logistic regression models with death during hospitalization as the binary outcome.

Carbapenem resistance was the primary exposure of interest. Multivariable models

were adjusted for available demographic, temporal, and hospital-level covariates,

including age group (neonate [≤28 days] vs older), sex, hospital site, timing of culture

collection relative to admission, and admission year or epoch (2004-2015 vs 2016-

2021) to account for secular trends. To evaluate robustness, we fitted a series of models

adjusting for individual covariates and all possible combinations thereof. Adjusted odds

ratios and 95% confidence intervals were reported.

Data availability

All data supporting the findings of this study are available within the paper and its

Supplementary Information. All whole genome sequencing data are available on ENA

(BioProject: PRJEB90555).

Acknowledgements

We thank all study participants and their families for their willingness to contribute to this research. We are grateful to the physicians, nurses, data collectors, and laboratory personnel at the participating hospitals for their dedicated patient care and for generating the clinical and laboratory data used in this study. We thank the funder, Gates Foundation (INV-023821 to Dr. Senjuti Saha) of supporting this study. The funder had no role in the study design, data collection, analysis, interpretation, or the decision to publish the findings.

Author Contributions

SS and SKS contributed to conceptualization, funding acquisition, study design, implementation, and monitoring. SS, YH, AMT and NK contributed to data analysis, preparation of figures, drafting the initial manuscript, and manuscript editing. MSI contributed to data analysis and coordination of the study. HR and MdH coordinated the blood culture, antimicrobial susceptibility, and other laboratory testing. AAT, SG and TJ performed DNA extraction and whole genome sequencing. YH, AMT and MdMH conducted the comparative genomic analysis. ZBA, BH, ShS, MJU and ABJ coordinated and trained doctors and nurses, and assisted in the implementation of the study. RCD performed data cleaning. MS, NKG, AKMS, NS, SKB, FA, WA, MMH, MH and ASMNU contributed to study management and patient data analysis. All authors reviewed the manuscript and agreed on the contents. The corresponding authors had full access to all the data in the study and had final responsibility for the decision to submit for

558   publication. YH, NK, and AMT directly accessed and verified the data. All authors had

559   access to de-identified data.

561   **Competing Interests**

562   The authors declare no competing financial and non-financial interests.

References

1. Wyres KL, Holt KE. Klebsiella pneumoniae as a key trafficker of drug resistance
genes from environmental to clinically important bacteria. *Current Opinion in*
Microbiology 2018; **45**: 131–9.

2 Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae.
Nature Reviews Microbiology 2020; **18**: 344–59.

3 Bengoechea JA, Sa Pessoa J. Klebsiella pneumoniae infection biology: living to
counteract host defences. *FEMS Microbiology Reviews* 2019; **43**: 123–44.

4 Zaidi AK, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. Hospital-
acquired neonatal infections in developing countries. *The Lancet* 2005; **365**: 1175–88.

5 Saha SK, Schrag SJ, Arifeen SE, *et al.* Causes and incidence of community-acquired
serious infections among young children in south Asia (ANISA): an observational
cohort study. *The Lancet* 2018; **392**: 145–59.

6 Sands K, Carvalho MJ, Portal E, *et al.* Characterization of antimicrobial-resistant
Gram-negative bacteria that cause neonatal sepsis in seven low- and middle-income
countries. *Nat Microbiol* 2021; **6**: 512–23.

7 Russell NJ, Stöhr W, Plakkal N, *et al.* Patterns of antibiotic use, pathogens, and
prediction of mortality in hospitalized neonates and young infants with sepsis: A
global neonatal sepsis observational cohort study (NeoOBS). *PLoS Med* 2023; **20**:
e1004179.

8 Verani JR, Blau DM, Gurley ES, *et al.* Child deaths caused by Klebsiella pneumoniae
in sub-Saharan Africa and south Asia: a secondary analysis of Child Health and
Mortality Prevention Surveillance (CHAMPS) data. *Lancet Microbe* 2024; : S2666-
5247(23)00290-2.

9 Munoz-Price LS, Poirel L, Bonomo RA, *et al.* Clinical epidemiology of the global
expansion of Klebsiella pneumoniae carbapenemases. *The Lancet Infectious*
Diseases 2013; **13**: 785–96.

10   Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant
Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis* 2017;
215: S28–36.

11 Stewardson AJ, Marimuthu K, Sengupta S, *et al.* Effect of carbapenem resistance on
outcomes of bloodstream infection caused by Enterobacteriaceae in low-income and
middle-income countries (PANORAMA): a multinational prospective cohort study.
Lancet Infect Dis 2019; **19**: 601–10.

12   WHO | Global priority list of antibiotic-resistant bacteria to guide research,
discovery, and development of new antibiotics. WHO.

http://www.who.int/medicines/publications/global-priority-list-antibiotic-resistant-
bacteria/en/ (accessed Sept 17, 2017).

13 Tamma PD, Heil EL, Justo JA, Mathers AJ, Satlin MJ, Bonomo RA. Infectious
Diseases Society of America 2024 Guidance on the Treatment of Antimicrobial-
Resistant Gram-Negative Infections. *Clinical Infectious Diseases* 2024; : ciae403.

14 Pennini ME, De Marco A, Pelletier M, *et al.* Immune stealth-driven O2 serotype
prevalence and potential for therapeutic antibodies against multidrug resistant
Klebsiella pneumoniae. *Nature Communications* 2017; **8**: 1991.

15 Diago-Navarro E, Motley MP, Ruiz-Peréz G, *et al.* Novel, Broadly Reactive
Anticapsular Antibodies against Carbapenem-Resistant Klebsiella pneumoniae
Protect from Infection. *mBio* 2018; **9**. DOI:10.1128/mBio.00091-18.

16 Dangor Z, Benson N, Berkley JA, *et al.* Vaccine value profile for Klebsiella
pneumoniae. *Vaccine* 2024; **42**: S125–41.

17 Opoku-Temeng C, Kobayashi SD, DeLeo FR. Klebsiella pneumoniae capsule
polysaccharide as a target for therapeutics and vaccines. *Computational and
Structural Biotechnology Journal* 2019; **17**: 1360–6.

18 Lam MMC, Wick RR, Judd LM, Holt KE, Wyres KL. Kaptive 2.0: updated capsule
and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex.
Microb Genom 2022; **8**: 000800.

19 Wayne P. Performance Standards for Antimicrobial Susceptibility Testing, 28th
edn. CLSI, 2018.

20 Andrews S. s-andrews/FastQC. 2024; published online Jan 22.
https://github.com/s-andrews/FastQC (accessed Jan 27, 2024).

21 Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* 2016; **32**: 3047–8.

22 Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 2014; **30**: 2114–20.

23 Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLOS Computational Biology*
2017; **13**: e1005595.

24 Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 2013; **29**: 1072–5.

25 Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;
30: 2068–9.

26   Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic
635   surveillance framework and genotyping tool for Klebsiella pneumoniae and its related
636   species complex. *Nat Commun* 2021; **12**: 4188.

637   27   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature*
638   *Methods* 2012; **9**: 357–9.

639   28   Danecek P, Bonfield JK, Liddle J, *et al.* Twelve years of SAMtools and BCFtools.
640   *GigaScience* 2021; **10**: giab008.

641   29   Arndt D, Grant JR, Marcu A, *et al.* PHASTER: a better, faster version of the
642   PHAST phage search tool. *Nucleic Acids Res* 2016; **44**: W16-21.

643   30   Croucher NJ, Page AJ, Connor TR, *et al.* Rapid phylogenetic analysis of large
644   samples of recombinant bacterial whole genome sequences using Gubbins. *Nucleic*
645   *Acids Research* 2015; **43**: e15–e15.

646   31   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-
647   analysis of large phylogenies. *Bioinformatics* 2014; **30**: 1312–3.

648   32   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
649   developments. *Nucleic Acids Research* 2019; **47**: W256–9.

650   33   Argimón S, David S, Underwood A, *et al.* Rapid Genomic Characterization and
651   Global Surveillance of Klebsiella Using Pathogenwatch. *Clinical Infectious Diseases*
652   2021; **73**: S325–35.

653   34   Murray CJ, Ikuta KS, Sharara F, *et al.* Global burden of bacterial antimicrobial
654   resistance in 2019: a systematic analysis. *The Lancet* 2022; **399**: 629–55.

655   35   Nieto-Rosado M, Sands K, Portal EAR, *et al.* Colonisation of hospital surfaces
656   from low- and middle-income countries by extended spectrum β-lactamase- and
657   carbapenemase-producing bacteria. *Nat Commun* 2024; **15**: 2758.

658   36   Jamil S, Obaidullah Md, Alam M. Unusual increase of cesarean section delivery
659   in Bangladesh: correspondence. *Ann Med Surg (Lond)* 2023; **85**: 2242.

660   37   Sarker AR, Zabeen I, Hossain Z, Ali N, Khan JAM. Increasing rates of cesarean
661   birth in Bangladesh: A household-level pooled analysis. *Birth* 2024; **51**: 326–42.

662   38   Kabir S, Hasan MdR, Hossain MdI, *et al.* Determinants and Trends of Health
663   Facility Delivery in Bangladesh: A Hierarchical Modeling Approach. *Biomed Res Int*
664   2022; **2022**: 1359572.

665   39   Watson SI, Arulampalam W, Petrou S, *et al.* The effects of a one-to-one nurse-to-
666   patient ratio on the mortality rate in neonatal intensive care: a retrospective,
667   longitudinal, population-based study. *Archives of Disease in Childhood - Fetal and*
668   *Neonatal Edition* 2016; **101**: F195–200.

40 Aktar S, Nu UT, Rahman M, *et al.* Trends and risk of recurrent preterm birth in
pregnancy cohorts in rural Bangladesh, 1990–2019. *BMJ Glob Health* 2023; **8**:
e012521.

41 Salam SS, Rahman AE, Mhajabin S, *et al.* Perception and experiences of
adolescent mothers and communities in caring for their preterm babies: findings from
an in-depth study in rural Bangladesh. *BMC Pregnancy Childbirth* 2024; **24**: 145.

42 Milton R, Gillespie D, Dyer C, *et al.* Neonatal sepsis and mortality in low-income
and middle-income countries from a facility-based birth cohort: an international
multisite prospective observational study. *The Lancet Global Health* 2022; **10**: e661–
72.

43 Ebi KL, Capon A, Berry P, *et al.* Hot weather and heat extremes: health risks. *The*
Lancet 2021; **398**: 698–708.

44 Dimitrova A, Dimitrova A, Mengel M, Gasparrini A, Lotze-Campen H, Gabrysch S.
Temperature-related neonatal deaths attributable to climate change in 29 low- and
middle-income countries. *Nat Commun* 2024; **15**: 5504.

45 Hu Y, Yang Y, Feng Y, *et al.* Prevalence and clonal diversity of carbapenem-
resistant Klebsiella pneumoniae causing neonatal infections: A systematic review of
128 articles across 30 countries. *PLOS Medicine* 2023; **20**: e1004233.

46 Wantuch PL, Knoot CJ, Robinson LS, *et al.* Heptavalent O-Antigen Bioconjugate
Vaccine Exhibiting Differential Functional Antibody Responses Against Diverse
Klebsiella pneumoniae Isolates. *J Infect Dis* 2024; **230**: 578–89.

47 Isaac EW, Jalo I, Difa AJ, *et al.* Bacterial Blood Isolates in Children: Conventional
vs. Bactec Automated Blood Culture System in a Tertiary Health Centre in Gombe,
North East Nigeria. *Open Journal of Medical Microbiology* 2022; **12**: 101–16.

Figure 1: Number and age distribution of culture-confirmed KpSC infections. a,

Annual number of KpSC isolates, 2004 – 2021. **b,** Annual positivity rate (per 1,000

blood cultures performed). **c,** Age distribution of all KpSC-positive cases (bars, grey)

with case fatality rate by age in months (line, yellow). **d,** Age distribution of KpSC-

positive neonates (total numbers in bars, grey; and distribution of neonates in line, blue)

with case fatality rate in days (line, yellow).

Figure 2: Rising case fatality rate of KpSC infections and correlation with rising

carbapenem resistance. a, Yearly trend in case-fatality rate (in %) amongst KpSC

cases at the study hospitals. **b,** Odds ratio (95%CI) of mortality amongst KpSC positive

cases using 2004 as the base year. **c,** The yearly trends in the percentage of KpSC

isolates that were resistant to meropenem. **d,** The comparison of case-fatality rate (in

%) of meropenem-resistant and meropenem-sensitive KpSC isolates stratified for:

2004-2015 and 2016-2021.

Figure 3: Phylogenetic structure and temporal dynamics of *Klebsiella*

***pneumoniae* species complex isolates. a,** Maximum likelihood phylogenetic tree of

599 representative KpSC genomes, colored by species. Kpn: *K. pneumoniae*, Kqpn: *K.*

quasipneumoniae subsp quasipneumoniae, Kspn: *K. quasipneumoniae subsp*

similipneumoniae, Kv: *K. variicola*. **b,** Maximum likelihood phylogenetic tree of the 503

K. pneumoniae genomes, annotated with year of isolation and sequence type (ST). **c,**

Maximum-likelihood tree showing the genomes belonging to ST-11 highlighting a large

cluster of *K. pneumoniae* ST-11 in 2019-2021. The accompanying heatmap summarizes

epidemiological variables (year of isolation, hospital site, interval between admission

and blood draw) and genomic features (ST, O locus, K type) for the ST11 genomes. **d,**

Map of Bangladesh showing the home location of patients involved in the *K.*

pneumoniae ST11 cluster who presented at the study hospitals. Points are colored by

the interval between admission and blood draw, indicating a country-wide cluster.

Figure 4: Temporal distribution of dominant *Klebsiella pneumoniae* sequence

types in Bangladesh and their global prevalence. a, Annual proportions of the ten

most common sequence types (STs) detected in Bangladesh, 2004–2021; colors

correspond to individual STs. **b,** Prevalence of the ten STs among a global collection of

32,656 *K. pneumoniae* genomes. Red outlines denote the three recognized global

problem-clone lineages (ST11, ST16, ST147).

Figure 5: Frequency and temporal coverage of dominant O- and K-loci among the

KpSC isolates. a, Absolute counts of the ten most common O-loci. Colored boxes

denote the cumulative coverage provided by the five (orange) and ten (blue) leading O-

loci. **b,** Annual percentage coverage achieved by the five (orange) and ten (blue) most

common O-loci. **c,** Absolute counts of the ten most common K-loci. Colored boxes

denote the cumulative coverage provided by the five (orange) and ten (blue) leading K-

loci. **d,** Annual percentage coverage achieved by the five (orange) and ten (blue) most

common K loci.

Table 1: List of carbapenem-resistance genes present on 599 KpSC genomes in

this study

a

Tree scale: 0.1 ⊢————┤

Species
- Kpn
- Kqpn
- Kspn
- Kv

b

Tree scale: 0.1 ⊢————┤

1
2

c

| | 1 | 2 | 3 | 4 | 5 | 6 |

Tree scale: 0.001 ⊢————┤

1. ST
- ST11
- ST147
- ST15
- ST16
- ST1998
- ST336
- ST340
- ST37
- ST48
- ST502
- others

2. Year
- 2004
- 2005
- 2007
- 2009
- 2010
- 2012
- 2014
- 2015
- 2017
- 2019
- 2021

3. K-locus
- KL15
- KL122
- Others

4. O-type
- O4
- O2afg
- O2a
- O3b

5. Site
- DSH
- KWMCH
- CMOSH
- CMCH
- DMCH

6. Gap in Collection
- <3 days
- ≥3 days

d

27°N
25°N
23°N
21°N

Latitude

KWMCH
BSHI
CMOSH

88°E 90°E 92°E

Longitude

Resistance marker	STs present	Frequency	First documented
blaNDM-5	11, 147, 48, and 7 others	126	2016
blaNDM-1	502, 16, 307 and 35 others	113	2008
blaNDM-5,blaOXA-181	16 and 147	38	2018
blaNDM-7	1998, 11, 705 and 6 others	21	2012
blaNDM-5;blaOXA-232	147, 307 and 437	14	2019
blaKPC-2;blaNDM-1	1998, 231, 340 and 70	12	2015
blaOXA-181	14, 16 and 43	4	2012
blaKPC-2	1998, 273 and 334	3	2019
blaNDM-1;blaOXA-232	14	3	2016
blaNDM-4	1584-1LV	2	2019
~blaNDM-5	152	2	2019
blaOXA-232	231	2	2018
blaNDM-24; blaOXA-181	16	1	2018
blaNDM-5; blaOXA-48	437	1	2021