Escalating carbapenem-resistant *Klebsiella* burden and mortality in Bangladeshi

infants

Yogesh Hooda[1, #], Arif Mohammad Tanmoy[1], Naito Kanon[1], Hafizur Rahman[1],

Mohammad Shahidul Islam[1], Zabed Bin Ahmed[1], Afroza Akter Tanni[1], Md Mobarok

Hossain[1], Md Hasanuzzaman[1], Sharmistha Goswami[1], Tasnim Jabin[1], Rajib C. Das[1],

Md Jamal Uddin[1], Mohammad Belal Hossain[1], Shampa Saha[1], Anannya Barman Jui[1],

Mohammod Shahidullah[2], Nobo Krishna Ghosh[3], AKM Shamsuzzaman[3], Nahid

Sultana[4], Sanjoy Kanti Biswas[4], Feroza Aktar[5], Wazir Ahmed[5], Mohammed Monir

Hossain[6], Mahbubul Hoque[6], ASM Nawshad Uddin Ahmed[1,7], Samir K Saha[1,8], Senjuti

Saha[1,#]

[1] Child Health Research Foundation, Dhaka, Bangladesh

[2] Department of Neonatology, Bangabandhu Sheikh Mujib Medical University, Dhaka,

Bangladesh

[3] Department of Pediatrics, Dr M R Khan Shishu Hospital & Institute of Child Health,

Dhaka, Bangladesh

[4] Department of Microbiology, Chattogram Maa O Shishu Hospital Medical College,

Chattogram, Bangladesh

[5] Department of Neonatology, Chattogram Maa O Shishu Hospital Medical College,

Chattogram, Bangladesh

[6] Department of Neonatal Medicine, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

[7] Department of Pediatrics, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

[8] Department of Microbiology, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

[#]Corresponding author

Senjuti Saha

Deputy Executive Director

Child Health Research Foundation

senjutisaha@chrfbd.org

AND

Yogesh Hooda

Senior Scientist

Child Health Research Foundation

yhooda@chrfbd.org

Abstract

Klebsiella infections in young infants are an escalating threat in low- and middle-income

countries, yet robust hospital burden and antimicrobial-resistance data informed by

genomics remain scarce. We performed an 18-year (2004-2021) prospective,

multicenter genomics epidemiology study across four hospitals in Bangladesh. Among

122,353 enrolled children with suspected sepsis, 1,600 (1.3%) yielded culture-confirmed

Klebsiella pneumoniae species complex (KpSC) isolates, which increased from 16 per

1,000 cases tested in 2004 to 37 in 2021. Hospital case-fatality rate (CFR) increased

from 21.4% to 51.4%, following an increase in carbapenem resistance (first detected in

2008; 81% by 2021). Neonates accounted for 80.5% of infections with a CFR of 40.8%.

Whole-genome sequencing of 599 representative isolates revealed four *Klebsiella*

species, 145 sequence types and 92 capsular loci. Global high-risk clones ST11, ST16

and ST147 harbouring NDM-type carbapenemases dominated recent cases. Stronger

infection control, access to new drugs, and multivalent vaccines are urgently needed.

Main

Gram-negative bacteria of the genus *Klebsiella* are ubiquitous in the environment and form part of the commensal flora of healthy humans[1,2]. Among them, *Klebsiella pneumoniae* and related members of the *K. pneumoniae* species complex (KpSC) can also cause a broad spectrum of infections, including soft tissue, surgical wounds, urinary tract, respiratory tract, bloodstream and the nervous system[3]. Neonates (0-27 days old) are particularly susceptible to systemic infections by KpSC[4–8].

Reports of multidrug-resistant (MDR) *K. pneumoniae* have risen sharply worldwide over the past two decades, with the steepest increases observed in low and middle income countries (LMICs)[9–11]. The emergence of carbapenem-resistant *K. pneumoniae*, now designated a critical priority pathogen by the WHO, has further constricted therapeutic options, and is associated with high mortality rates[12].

With the antibiotic pipeline faltering, alternative strategies are gaining traction. Recently, combination drugs that contain a beta-lactam antibiotic and a beta-lactamase inhibitor have shown to be efficacious for treatment of carbapenem resistant *K. pneumoniae* infections, but their efficacy hinges on the specific resistance genes present.[13] Passive immunotherapies and maternal vaccines targeting capsular (K) or lipopolysaccharide (O) antigens are also under development[14–17]. However, *Klebsiella* exhibits extraordinary antigenic heterogeneity, with at least 13 O-types and 186 K-types documented to date[18]. Selecting vaccine valences therefore demands precise, contemporary data on circulating serotypes, resistance genes and clonal backgrounds.

Additionally, global genomic sequencing efforts have found that *Klebsiella* species such

as *K. quasipneumoniae*, *K. variicola* (part of the 7 species that make up *KpSC*) cause

invasive diseases in humans, yet routine biochemical tests cannot differentiate them

from *K. pneumoniae*[2]. Failing to recognize this hidden diversity can mask transmission

chains, underestimate true burden and misguide therapy, particularly in low-resource

settings that rely on phenotypic methods.

In this study we: (i) quantify temporal trends in the hospital burden and severity of

invasive KpSC infections in Bangladeshi children; (ii) describe the evolution of

phenotypic and genotypic AMR, including carbapenemase rise and spread; and (iii)

characterize the species, sequence types, O- and K-loci of a representative subset of

isolates using whole-genome sequencing. Overall, we generate data to inform future

infection-control measures, empirical-therapy guidelines and multivalent-vaccine design

in Bangladesh and neighboring countries.

Methods

Ethics Statement:

This study analyzes *KpSC* isolates, along with associated clinical and laboratory

metadata, collected as part of the PneumoADIP study (2004-2008) and the Invasive

Bacterial Disease (IBD) study (2009-2021). The protocols for these studies were

approved by the Ethical Review Board of the Bangladesh Institute of Child Health (now

known as Bangladesh Shishu Hospital & Institute). We obtained written informed

consent from the guardians of children admitted to the study hospitals, from whom

blood and/or cerebrospinal fluid (CSF) samples were collected.

Description of the study site:

From 2004-2021, blood and/or CSF specimens were collected at four hospitals in

Bangladesh: a) Bangladesh Shishu Hospital and Institute (BSHI), a 650-bed tertiary

level hospital based in Dhaka; b) Dr. M R Khan Shishu Hospital and Institute of Child

Health (MRKSH), a 250-bed primary - secondary level hospital in Dhaka; c) Chattogram

Maa-O Shishu Hospital (CMOSH), a 835-bed primary - tertiary level hospital in

Chattogram, and d) Kumudini Women's Medical College and Hospital (KWMCH), a

1050-bed based primary-tertiary level hospital based in rural town Mirzapur. Children

hospitalized in these hospitals who matched the WHO-defined clinical criteria for

pneumonia, severe pneumonia, severe disease, and meningitis were evaluated by

trained research physicians. Blood and/or CSF was drawn for culture at the treating

physicians' discretion. Demographic (age and sex) and clinical (symptoms, hospital

outcome, admission date, date of sample collection etc.) data were obtained for all

enrolled patients (Supplementary Table 1). Children were enrolled upon informed written

consent if they met the inclusion criteria.

Laboratory procedures:

Blood and/or CSF samples from patients were directly cultured on blood agar, chocolate

agar and MacConkey agar media from 2004-2015. We introduced BACTEC for blood

cultures in BSHI, KWMCH and MRKSH, in 2016 and in CMOSH the following year;

since then, blood specimens were incubated in BACTEC culture systems for up to 120

hours, and positive samples were subsequently sub-cultured on blood agar, chocolate

agar and MacConkey agar media. Upon growth, biochemical tests including, indole,

urease, and citrate utilization tests were performed to identify the genus and species.

Isolates with mucoid colony morphology, lactose fermentation on MacConkey agar, and

a biochemical profile of oxidase-negative, indole-negative, urease-positive, and citrate-

positive were confirmed as *Klebsiella* and stored at -80°C in STGG media (skim milk,

tryptone, glucose, and glycerin). These stocks served as the source for all subsequent

experiments performed in the study.

Antimicrobial susceptibility testing was performed using antibiotic discs by Kirby-Bauer

testing for: meropenem, imipenem, ceftriaxone, ampicillin, gentamicin, netilmicin,

ciprofloxacin, amikacin, chloramphenicol, and cotrimoxazole. Tests were interpreted

according the Clinical and Laboratory Standards Institute (CLSI) 2025 antibiotic

susceptibility testing clinical breakpoints[19].

Sample selection criteria for whole genome sequencing (WGS):

Approximately a third of KpSC isolates from children under the age of 2 months, were

randomly selected, keeping the distribution of isolates selected per year consistent.

These selected isolates were re-cultured from biobank, biochemically re-confirmed and

sub-cultured on MacConkey agar media prior to DNA extraction and whole genome

sequencing.

To contextualize the selected isolates with other KpSC isolates from Bangladesh, in

addition to the isolates recovered in this study, we randomly selected 100 of 160 (62%)

KpSC isolates for WGS that were stored in our biobank collected between 2004-2008

from three additional hospitals that followed the same inclusion criteria: a) Dhaka

Medical College and Hospital, Dhaka, b) Chattogram Medical College and Hospital,

Chattogram, and c) Sir Salimullah Medical College and Hospital, Dhaka as part of the

pneumoADIP study.

Genomic DNA extraction and sequencing:

Total DNA from *Klebsiella pneumoniae* colonies was extracted using the QIAGEN DNA

Mini extraction kit (Cat: 56304, Qiagen, Hilden, Germany). Sequencing libraries were

prepared from extracted DNA using NEBNext Ultra II FS DNA Library kit (Cat: E7805,

New England Biolabs, MA, USA). Libraries were sequenced on an Illumina

NEXTSeq2000 at the Child Health Research Foundation, Dhaka, Bangladesh (n = 528;

150bp PE) and on the HiSeq4000 platform at the University of Oxford, Oxford, UK (n =

71; 150bp PE).

Genome assembly

Sequence data quality was assessed using FastQC v0.11.5[20] and MultiQC v3[21]. Adapter

trimming was performed with Trimmomatic v0.39[22]. *De novo* genome assembly was

conducted using Unicycler v0.4.9[23], evaluated using Quast v.5.2.0[24], and annotated with

Prokka v.1.14.6[25]. Kleborate v2.2[26] was used to identify the sequence types (STs), AMR

genes, K and O antigen types. All generated sequence data have been submitted to

NCBI (BioProject: PRJNA845975).

<u>Phylogenetic Analysis</u>

For reference-based genome mapping, Bowtie2 v2.3.4.1 was employed to align all

genomic data against the NCBI reference genome, *Klebsiella pneumoniae* HS11286

(Genbank accession: NC_016845.1)[27]. SAMtools v1.7.2 and BCFtools v1.7.2 were

utilized for variant calling, filtering candidate SNPs for a minimum base quality of 20, a

minimum of 5 high-quality reads, and greater than 75% read support (i.e., over 75% of

the reads supporting the SNP relative to the reference)[28]. SNPs were further filtered to

remove potential biases, including strand, mapping, and end biases (p-value < 0.001).

Additionally, SNPs located within phage regions (identified by PHASTER[29]), repeat

sequences (as noted in the reference GenBank), and recombination regions (detected

by Gubbins v2.3.1) were excluded[30]. The resulting filtered SNPs were then converted

into a SNP alignment for maximum likelihood phylogenetic tree (MLT) construction using

RAxML v8.2.11 (model: GTRGAMMA; 100 bootstraps)[31]. The alignment had 82,669

SNP positions from 472 *Klebsiella pneumoniae* isolates in our study. Tree visualization

and annotation was done using iTOL v6[32].

For global comparison of the STs of the *K. pneumoniae* isolates (n=472) sequenced in

this study, publicly available *K. pneumoniae* genomes (n=32,656) in PathogenWatch[33]

(retrieved on November 29, 2023) were used.

<u>Statistical analysis</u>

Statistical analyses were performed using custom scripts in R (version 4.3.1). The odds

ratio was calculated with using the *oddsratio* package, while the figures were generated

using the *ggplot2* and *Rcolorbrewer* and *ggsci* packages. Global and Bangladesh-

specific maps were made using the packages *rnaturalearth, ggspatial, maps* and

scatterpie. For ST11 geographic analysis, upazila, village or union information was used

instead of the exact address obtained during data collection.

Results

The burden and severity of KpSC infections in Bangladesh is increasing.

Between 2004 and 2021, 122,353 cases had blood and/or CSF samples cultured at the

treating physician's discretion (Supplementary Fig. 1); all were enrolled in this study.

Bacteriological-culture and biochemical testing confirmed that 1,600 (1.3%) infections

were caused by the KpSC. A sharp rise is observed in the number of KpSC isolates

starting in 2018 (Fig. 1a). Up to 2017, the mean positivity rate for KpSC was 16 per

1,000 cases where blood cultures were performed, which climbed to 18, 34 and 37 per

1,000 suspected sepsis cases in 2019, 2020 and 2021, respectively (Fig. 1b). Seventy-

four percent (1,182/1,600) of *KpSC*-positive cases were reported from one hospital,

BSHI (Supplementary Fig. 2a), which also accounted for the highest positivity rate

among the four hospitals since 2015 (Supplementary Fig. 2b).

Figure 1: Number and age distribution of culture-confirmed KpSC infections. a,
Annual number of KpSC isolates, 2004 – 2021. **b,** Annual positivity rate (per 1,000
blood cultures performed). **c,** Age distribution of all KpSC-positive cases (bars, grey)
with case fatality rate by age in months (line, yellow). **d,** Age distribution of KpSC-
positive neonates (total numbers in bars, grey; and distribution of neonates in line, blue)
with case fatality rate in days (line, yellow).

The median age of KpSC-positive cases at hospital admission was 5 days (IQR: 2-18)

and at blood draw was 8 days (IQR:4-21) (Fig. 1c). Overall, 1,352/1,600 (84.5%) KpSC

infections occurred in infants <2 months old, and 1,288/1,600 (80.5%) in neonates (first

28 days old). The hospital case fatality rate (CFR) was highest during the first month of

life (41.5%) and declined for infants older than three months (Fig. 1c). Overall CFR was

38.1%. Among neonates, the median age at sample collection was 6 days (IQR: 3-21),

and the CFR was consistent throughout the first month of life (Fig. 1d). Most infections

1,076/1,600 (67.3%) was detected within 72 hours of admission (Supplementary Fig.

3a); the median interval between admission to sample collection was 1 day (IQR: 0-4).

The severity of KpSC infection is highlighted by hospital outcomes: 610/1,600 (38.1%)

cases died at the hospital, 174/1,600 (10.9%) left against medical advice, and

816/1,600 (51.0%) were discharged from the hospital (Supplementary Table 1). The

average hospital stay was 8 days (IQR, 4-14), but 64% babies who died, succumbed

within the first 5 days of admission (Supplementary Fig. 3b). CFR started increasing in

2016 (Fig. 2a), with the odds of death in 2021 being 3.9 times higher than in 2004

(95 % CI 1.5–9.8; Fig. 2b).

Increasing severity of KpSC infections correlates with rise in carbapenem

resistance.

We investigated the possible drivers of escalating mortality by comparing yearly trends

in clinical features (Supplementary Fig. 4) and antimicrobial susceptibility profiles for 10

antibiotics (meropenem, imipenem, ceftriaxone, ampicillin, gentamicin, netilmicin,

ciprofloxacin, amikacin, chloramphenicol, and cotrimoxazole) proposed by CLSI[19]

(Supplementary Fig. 5). KpSC isolates from Bangladesh exhibited high resistance

against most antimicrobials used, including amikacin, cotrimoxazole, gentamicin,

netilmicin, and ciprofloxacin. The first case of carbapenem resistance was detected in

2008, and with noticeable increase during the study period, by 2021 >80% of KpSC

isolates were resistant to carbapenem (Fig. 2c). The rising carbapenem resistance

correlated with increasing mortality, with the odds of death being 3.2-fold higher (95%

CI: 2.3-4.6) relative to carbapenem sensitive isolates (Supplementary Fig. 6).

Figure 2: Rising case fatality rate of KpSC infections and correlation with rising
carbapenem resistance. a, Yearly trend in case-fatality rate (in %) amongst KpSC
cases at the study hospitals. **b,** Odds ratio (95%CI) of mortality amongst KpSC positive
cases using 2004 as the base year. **c,** The yearly trends in the percentage of KpSC
isolates that were resistant to meropenem. **d,** The comparison of case-fatality rate (in
%) of meropenem-resistant and meropenem-sensitive KpSC isolates stratified for:
2004-2015 and 2016-2021.

To further assess the impact of carbapenem resistance, we stratified cases into two

epochs, 2004–2015 and 2016–2021, and compared CFRs for carbapenem-resistant

versus carbapenem-susceptible KpSC infections (Fig. 3d). During 2004–2015, CFRs

were similar irrespective of resistance phenotype. From 2016 onward, carbapenem-

resistant infections exhibited a markedly higher CFR than susceptible infections,

however, mortality increased for both groups. These suggest, while carbapenem

resistance is a key driver of poor outcomes, additional factors, i.e. host, pathogen

and/or environmental, have contributed to the post-2016 surge in KpSC-associated

deaths.

KpSC infections exhibit high genomic and antigenic diversity

To further characterize the KpSC isolates, we sequenced a representative subset of 527

isolates (527/1461, 36.1%) collected across the entire study period from infants younger

than two months - the age group with the highest burden (Supplementary Fig. 7a,

Supplementary Table 2)[26]. Some bio banked strains from 2004-2009 could not be

revived, and hence fewer representatives from those years were available. To partially

overcome this limitation, we included 72 sequences from three other hospitals from

2004-2009. Among the total 599 isolates sequenced, 503 (84.0%) genomes were *K.*

pneumoniae, 72 (12.0%) were *K. quasipneumoniae* subsp *similipneumoniae*, 20 (3.3%)

were *K. quasipneumoniae* subsp *quasipneumoniae*, and 4 (0.7%) were *K. variicola* (Fig.

3a, Supplementary Table 2). The annual distribution of these species remained broadly

consistent throughout the study period (Supplementary Fig. 7b). Infections caused by *K.*

pneumoniae were associated with a 2.18 (95% CI 1.06-4.94) fold higher case fatality

rate than infections due to other KpSC species (Supplementary Fig. 8).

Figure 3: Phylogenetic structure and temporal dynamics of *Klebsiella pneumoniae* species complex isolates. a, Maximum likelihood phylogenetic tree of 599 representative KpSC genomes, colored by species. Kpn: *K. pneumoniae*, Kqpn: *K. quasipneumoniae subsp quasipneumoniae*, Kspn: *K. quasipneumoniae subsp similipneumoniae*, Kv: *K. variicola*. **b,** Maximum likelihood phylogenetic tree of the 503 *K. pneumoniae* genomes, annotated with year of isolation and sequence type (ST). **c,** Maximum-likelihood tree showing the genomes belonging to ST-11 highlighting a large outbreak of *K. pneumoniae* ST-11 in 2019-2021. The accompanying heatmap summarizes epidemiological variables (year of isolation, hospital site, interval between admission and blood draw) and genomic features (ST, O locus, K type) for the ST11 genomes. **d,** Map of Bangladesh showing the home location of patients involved in the *K. pneumoniae* ST11 outbreak who presented at the study hospitals. Points are colored by the interval between admission and blood draw, indicating a country-wide outbreak.

Maximum-likelihood phylogenies of the 599 sequenced KpSC genomes (Fig. 3a) and

the 503 *K. pneumoniae* genomes (Fig. 3b) demonstrated high genetic diversity amongst

KpSC isolates identified in this study. We identified 145 distinct sequence types (STs);

however, the ten most common STs accounted for 52% of all isolates. The predominant

lineage was *K. pneumoniae* ST11, 97% of which were identified between 2018 and

2020. These ST11 genomes were sampled across three of the four surveillance

hospitals in Dhaka, Chattogram and Mirzapur, yet showed minimal within-lineage

diversity, consistent with a nationwide clonal expansion (Fig. 3c). Metadata reinforcing

this observation include that among ST11 cases, the median interval from admission to

blood draw was 2 days (IQR 0-5); 53% (36/68) of samples were collected within 48 hrs

of hospital admission (Fig 3c). Mapping of home addresses shows infants originating

from 22 districts (Fig. 3d).

The yearly distribution of the 10 most common STs for *K. pneumoniae* are shown in Fig.

4a. Some of the common STs like ST11 and ST16 exhibited higher CFR

312   (Supplementary Fig. 8). When compared to a global database of *K. pneumoniae,* most

313   of the common STs in Bangladesh have been reported in other countries, and ST11, 16,

314   and 147 belong to documented global problem clones (Fig. 4b).

315
316

Figure 4: Temporal distribution of dominant *Klebsiella pneumoniae* sequence types in Bangladesh and their global prevalence. a, Annual proportions of the ten most common sequence types (STs) detected in Bangladesh, 2004–2021; colors correspond to individual STs. **b,** Prevalence of the ten STs among a global collection of 32,656 *K. pneumoniae* genomes. Red outlines denote the three recognized global problem-clone lineages (ST11, ST16, ST147).

We next analyzed genomic carbapenem-resistance determinants in the sequenced KpSC genomes (Table 1, Supplementary Fig. 7c) and compared these with meropenem susceptibility results. Genomic prediction of the meropenem phenotype achieved a sensitivity of 96.3% and a specificity of 65.4%. The main carbapenemase alleles detected were NDM-5, NDM-1, along with multiple other less common variants (Table 1), indicating that different KpSC lineages in Bangladesh have acquired carbapenem-resistance genes independently.

Table 1: List of carbapenem-resistance genes present on 599 KpSC genomes in this study

Resistance marker	STs present	Frequency	First documented
NDM-5	11, 147, 48, and 7 others	126	2016
NDM-1	502, 16, 307 and 35 others	113	2008
NDM-5,OXA-181	16 and 147	38	2018
NDM-7	1998, 11, 705 and 6 others	21	2012
NDM-5;OXA-232	147, 307 and 437	14	2019
KPC-2;NDM-1	1998, 231, 340 and 70	12	2015
OXA-181	14, 16 and 43	4	2012
KPC-2	1998, 273 and 334	3	2019
NDM-1;OXA-232	14	3	2016
NDM-4	1584-1LV	2	2019
~NDM-5	152	2	2019
OXA-232	231	2	2018
NDM-24; OXA-181	16	1	2018
NDM-5; OXA-48	437	1	2021

Figure 5: Frequency and temporal coverage of dominant O- and K-loci among the KpSC isolates. a, Absolute counts of the ten most common O-loci. Colored boxes denote the cumulative coverage provided by the five (orange) and ten (blue) leading O-loci. **b,** Annual percentage coverage achieved by the five (orange) and ten (blue) most common O-loci. **c,** Absolute counts of the ten most common K-loci. Colored boxes denote the cumulative coverage provided by the five (orange) and ten (blue) leading K-loci. **d,** Annual percentage coverage achieved by the five (orange) and ten (blue) most common K loci.

To characterize antigenic diversity relevant to vaccine design, we analyzed the genomic

regions encoding the capsular polysaccharide (K locus) and O antigen. In total, 92

unique K loci and 11 unique O loci were identified. The ten most prevalent O loci and K

loci together accounted for 99.8% (Fig. 5a, Extended Fig. 9) and 58.3% (Fig. 5c,

Extended Fig. 10) of all isolates, respectively. Their year-to-year frequencies are plotted

in Fig. 5b and Fig. 5d. The combined share of the ten dominant O loci stayed above

50% every year, whereas the share of the ten dominant K loci was highly variable,

oscillating between 0% and 86.7%.

Discussion

Geographically and temporally representative epidemiological and genomic data from

high-burden LMICs are essential to guide policy and intervention against KpSC

infections. Here we report 18 years of surveillance from four Bangladeshi pediatric

hospitals, detailing trends in burden, antimicrobial resistance and genomic diversity.

From 122,353 bacteriological culture of blood and/or CSF samples conducted between

2004 and 2021, we identified 1,600 KpSC isolates. Neonates comprised 80.5% of cases

and had a hospital case-fatality rate of 40.8%, compared with an overall CFR of 38.1%.

In a multi-country study on neonatal infections, BARNARDS, conducted in seven LMICs

including Bangladesh from Nov 2015 to Dec 2017, the CFR due to KpSC infection was

22.4% (58/258), compared to 35.7% (45/126) during the same period in this study.[6] In

the NeoObs study done across 11 countries from 2018-2019 including Bangladesh,

neonatal KpSC case fatality rate was 21.1%.[7] In NeoOBS Bangladesh site, the CFR

was 22% (2/9), compared to 39% (120/307) during that period in this study. Within

neonates, infections were most commonly detected on day 5, as seen in previous

studies [BARNARDS and NeoObs][7,8]. Collectively, these studies highlight the burden of

KpSC infections amongst neonates globally. The higher mortality we observed was

driven by BSHI, which functions as the pediatric referral center for Bangladesh,

receiving the most critically ill infants from across the country; this referral bias is likely

to inflate both the apparent burden, and the CFR compared with primary or secondary

facilities included in the other studies.

Overall, the alarming rise in neonatal deaths threatens to erode vaccine-driven gains in

child survival across LMICs. Thirteen percent of infections were detected within the first

three days of life, including 3% within 24 hours of birth, and 67% of infants were culture-

positive within 72 hours of admission. These timelines point to transmission in maternity

units, delivery clinics, or the community, well before referral to our surveillance hospitals.

At BSHI, the KpSC positivity rate rose from 16 per 1000 suspected sepsis cases in

2004 to 37 per 1000 in 2021, and the hospital-specific case-fatality rate climbed from a

mean of 20.7% (2004–2014) to 51.4% in 2021. Carbapenem resistance was associated

with death, with an adjusted odds ratio of 3.2 (95% CI: 2.3-4.6). A positive correlation

between carbapenem resistance and mortality has reported previously for KpSC and

other bacterial infections[34]; however, the observational design of our study precludes

inferring causality. In addition, while CFR of carbapenem-resistant KpSC infections was

significantly higher than carbapenem-sensitive KpSC infections (46.3% vs 25.4%), we

also observed a rise in deaths among carbapenem-susceptible infections after 2015

compared to carbapenem-susceptible infections before 2015 (31.1% vs 20.3%). The

excess mortality in carbapenem-susceptible cases after 2015 suggests that factors

beyond drug resistance are driving severity. Several, not mutually exclusive, hypotheses

warrant exploration: (i) bacterial adaptation – dominant lineages may have acquired

additional virulence determinants[3] or stress-tolerance genes that enhance survival in

neonatal wards[35]; (ii) changing obstetric practice – the sharp rise in Caesarean

deliveries in Bangladesh increases early hospital exposure and separates infants from

protective maternal microbiota[36,37]; (iii) ward overcrowding and resource strain – more

institutional births and limited isolation facilities hamper effective infection-prevention

and control[38]; (iv) referral dynamics – improved road networks now funnel the sickest

infants who fail first-line therapy or need advanced support to BSHI, inflating observed

mortality (v) co-infections and comorbidities **–** rising prevalence of prematurity, low birth-

weight and congenital anomalies may increase vulnerability to sepsis[39–41]; and (vi)

climate stress – warmer, more humid conditions, extreme weather events like flooding

and heat-related infrastructure strain can favor environmental persistence and spread of

KpSC within hospitals[42,43]. These possibilities highlight the need for integrated clinical,

microbiological and health-systems investigations to disentangle drivers of the

continued rise in KpSC-associated neonatal deaths.

Several genomic studies have now established that biochemically identified *K.*

pneumoniae infections can be other members of KpSC[2,26,33]. In this study, 15% were *K.*

quasipneumoniae (*subsp. simillipneumoniae* & *subsp. quasipneumoniae*) and *K.*

variicola, rather than *K. pneumoniae*. Infections due to *K. pneumoniae* carried a higher

hospital CFR (32.9% vs 18.3%) and a greater burden of carbapenem resistance genes.

These observations suggest that pharmacological interventions directed solely at *K.*

pneumoniae, whether vaccines, monoclonal antibodies or new antimicrobials, could

create ecological space for the related species to fill. Future control strategies must

therefore target the entire KpSC repertoire to avoid pathogen replacement and should

also include broad IPC.

We also observed a high diversity of carbapenem-resistance determinants in

Bangladeshi KpSC isolates, with the *ndm-1* gene predominating[6,44]. Because KpSC

isolates are predicted to remain susceptible to the ceftazidime-avibactam + aztreonam

combination [13], this regimen is a promising therapeutic option. Yet neither drug is widely

available in Bangladesh, particularly in the low-income communities served by our

surveillance hospitals, creating a critical treatment gap. Pragmatic clinical trials are

therefore urgently needed to evaluate the regimen's efficacy, safety and implementation

feasibility and to guide policy aimed at reducing *Klebsiella*-associated neonatal

mortality.

Overall, the Bangladeshi KpSC population is highly heterogeneous: the 599 genomes

were distributed across 145 STs**,** and the dominant lineages shift and vary over time. In

the most recent surveillance years, we detected several recognized global problem

clones, notably ST11, ST147 and ST16**,** that have been described from in various other

countries[26,33]. The clinical impact is best illustrated by ST11, the predominant ST in

recent years. Almost all ST11 isolates (n = 68) were recovered between 2018 and 2020,

consistent with a recent clonal expansion. The median interval from admission to blood

draw was 1 days (IQR 0–5); 53% of cultures were taken within 72 h of arrival. Cases

originated from 22 districts, and most infants were admitted directly from home or local

clinics rather than transferred from other hospitals. Minimal genetic divergence, very

short admission-to-culture intervals and wide geographic scatter together suggest

transmission in maternity units, delivery clinics or the community, settings upstream of

tertiary infection-control measures.

Efforts to protect young infants against KpSC, particularly *K. pneumoniae*, now focus on

maternal vaccines and monoclonal antibodies that target the organism's surface

polysaccharides (capsular K antigen and lipopolysaccharide O antigen)[16]. Our data

highlight the antigenic hurdles such products must clear. Whole-genome analysis

predicted 92 distinct K (capsular) types, with the ten most common covering only 58%

of isolates. O-antigen diversity was narrower: the ten most frequent O loci

encompassed 99% of isolates, and their cumulative coverage remained >50 % each

year. By contrast, K-type coverage fluctuated markedly over time, reflecting turnover of

dominant sequence types and the strong linkage between ST and capsule genotype.

These patterns indicate that a multivalent formulation combining O- and K-antigens will

be required for broad protection in Bangladesh, yet the capsule's extensive and shifting

diversity will remain a major design challenge.

The findings of this study should be viewed within the context of certain limitations. First,

we did not collect information related to gestational age, birth weight and maternal

factors, such as maternal age, nutrition, and mode of delivery. These variables are

recognized risk factors for KpSC infection, and their absence prevents adjustment for

host susceptibility. Second, our study lacked systematic data on comorbidities including

congenital anomalies, which may have contributed to the observed mortality;

nevertheless, previous CHAMPS investigations have implicated *K. pneumoniae* as a

leading cause of neonatal death.[8] Third, no pre-hospital records were available, so we

could not ascertain prior antibiotic exposure, treatment adequacy or the likely place of

acquisition. Fourth, we sequenced a representative subset of KpSC isolates; rare

sequence types, capsular types or O-antigen variants present in the isolates that were

not sequenced may therefore have been missed. Fifth, automated blood culture system

(BACTEC) was introduced in BSHI, KWMCH and MRKSH in 2016 and in CMOSH

in 2017. However, given the increase in KpSC was seen only in BSHI, this change is not

likely to have impacted blood-culture positivity rates.

Within these constraints, the dataset provides one of the most comprehensive clinical–

genomic reports of paediatric KpSC infections available from South Asia. Rising

incidence, a four-fold increase in mortality and the dominance of NDM-positive global

clones expose a critical threat to neonatal survival. To mitigate this risk, policies should

include: (i) expansion of neonatal intensive-care capacity and skilled staffing at birth

facilities; (ii) stringent infection-prevention and control in labour wards and neonatal

units; (iii) continuous, denominator-based genomic surveillance to detect clonal shifts

and emerging resistance; and (iv) rapid access to effective therapeutics, including

context-specific trials of ceftazidime–avibactam plus aztreonam. Maternal vaccines or

monoclonal antibodies targeting the prevalent O-antigens could add a valuable layer of

protection, but the extensive and shifting capsule diversity documented here means

immunization must be embedded in a broader strategy that compliments vaccination

with improved clinical care, antimicrobial stewardship and robust infection control.

Because pathogen-circulation patterns mirror those of other pathogenic bacteria like

Streptococcus pneumoniae and other regionally important bacteria, these

recommendations are likely to be relevant beyond Bangladesh and should inform

intervention design across south Asia.

References

1. Wyres KL, Holt KE. Klebsiella pneumoniae as a key trafficker of drug resistance genes from environmental to clinically important bacteria. *Current Opinion in Microbiology* 2018; **45**: 131–9.

Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae. *Nature Reviews Microbiology* 2020; **18**: 344–59.

Bengoechea JA, Sa Pessoa J. Klebsiella pneumoniae infection biology: living to counteract host defences. *FEMS Microbiology Reviews* 2019; **43**: 123–44.

Zaidi AK, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. Hospital-acquired neonatal infections in developing countries. *The Lancet* 2005; **365**: 1175–88.

Saha SK, Schrag SJ, Arifeen SE, *et al.* Causes and incidence of community-acquired serious infections among young children in south Asia (ANISA): an observational cohort study. *The Lancet* 2018; **392**: 145–59.

Sands K, Carvalho MJ, Portal E, *et al.* Characterization of antimicrobial-resistant Gram-negative bacteria that cause neonatal sepsis in seven low- and middle-income countries. *Nat Microbiol* 2021; **6**: 512–23.

Russell NJ, Stöhr W, Plakkal N, *et al.* Patterns of antibiotic use, pathogens, and prediction of mortality in hospitalized neonates and young infants with sepsis: A global neonatal sepsis observational cohort study (NeoOBS). *PLoS Med* 2023; **20**: e1004179.

Verani JR, Blau DM, Gurley ES, *et al.* Child deaths caused by Klebsiella pneumoniae in sub-Saharan Africa and south Asia: a secondary analysis of Child Health and Mortality Prevention Surveillance (CHAMPS) data. *Lancet Microbe* 2024; : S2666-5247(23)00290-2.

Munoz-Price LS, Poirel L, Bonomo RA, *et al.* Clinical epidemiology of the global expansion of Klebsiella pneumoniae carbapenemases. *The Lancet Infectious Diseases* 2013; **13**: 785–96.

Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant Enterobacteriaceae: The Impact and Evolution of a Global Menace. *J Infect Dis* 2017; **215**: S28–36.

Stewardson AJ, Marimuthu K, Sengupta S, *et al.* Effect of carbapenem resistance on outcomes of bloodstream infection caused by Enterobacteriaceae in low-income and middle-income countries (PANORAMA): a multinational prospective cohort study. *Lancet Infect Dis* 2019; **19**: 601–10.

WHO | Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. WHO.

http://www.who.int/medicines/publications/global-priority-list-antibiotic-resistant-
bacteria/en/ (accessed Sept 17, 2017).

13   Tamma PD, Heil EL, Justo JA, Mathers AJ, Satlin MJ, Bonomo RA. Infectious
Diseases Society of America 2024 Guidance on the Treatment of Antimicrobial-
Resistant Gram-Negative Infections. *Clinical Infectious Diseases* 2024; : ciae403.

14   Pennini ME, De Marco A, Pelletier M, *et al.* Immune stealth-driven O2 serotype
prevalence and potential for therapeutic antibodies against multidrug resistant
Klebsiella pneumoniae. *Nature Communications* 2017; **8**: 1991.

15   Diago-Navarro E, Motley MP, Ruiz-Peréz G, *et al.* Novel, Broadly Reactive
Anticapsular Antibodies against Carbapenem-Resistant Klebsiella pneumoniae
Protect from Infection. *mBio* 2018; **9**. DOI:10.1128/mBio.00091-18.

16   Dangor Z, Benson N, Berkley JA, *et al.* Vaccine value profile for Klebsiella
pneumoniae. *Vaccine* 2024; **42**: S125–41.

17   Opoku-Temeng C, Kobayashi SD, DeLeo FR. Klebsiella pneumoniae capsule
polysaccharide as a target for therapeutics and vaccines. *Computational and
Structural Biotechnology Journal* 2019; **17**: 1360–6.

18   Lam MMC, Wick RR, Judd LM, Holt KE, Wyres KL. Kaptive 2.0: updated capsule
and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex.
Microb Genom 2022; **8**: 000800.

19   Wayne P. Performance Standards for Antimicrobial Susceptibility Testing, 28th
edn. CLSI, 2018.

20   Andrews S. s-andrews/FastQC. 2024; published online Jan 22.
https://github.com/s-andrews/FastQC (accessed Jan 27, 2024).

21   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* 2016; **32**: 3047–8.

22   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 2014; **30**: 2114–20.

23   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLOS Computational Biology*
2017; **13**: e1005595.

24   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 2013; **29**: 1072–5.

25   Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;
30: 2068–9.

26   Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic
surveillance framework and genotyping tool for Klebsiella pneumoniae and its related
species complex. *Nat Commun* 2021; **12**: 4188.

27   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature
Methods* 2012; **9**: 357–9.

28   Danecek P, Bonfield JK, Liddle J, *et al.* Twelve years of SAMtools and BCFtools.
GigaScience 2021; **10**: giab008.

29   Arndt D, Grant JR, Marcu A, *et al.* PHASTER: a better, faster version of the
PHAST phage search tool. *Nucleic Acids Res* 2016; **44**: W16-21.

30   Croucher NJ, Page AJ, Connor TR, *et al.* Rapid phylogenetic analysis of large
samples of recombinant bacterial whole genome sequences using Gubbins. *Nucleic
Acids Research* 2015; **43**: e15–e15.

31   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* 2014; **30**: 1312–3.

32   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research* 2019; **47**: W256–9.

33   Argimón S, David S, Underwood A, *et al.* Rapid Genomic Characterization and
Global Surveillance of Klebsiella Using Pathogenwatch. *Clinical Infectious Diseases*
2021; **73**: S325–35.

34   Murray CJ, Ikuta KS, Sharara F, *et al.* Global burden of bacterial antimicrobial
resistance in 2019: a systematic analysis. *The Lancet* 2022; **399**: 629–55.

35   Nieto-Rosado M, Sands K, Portal EAR, *et al.* Colonisation of hospital surfaces
from low- and middle-income countries by extended spectrum β-lactamase- and
carbapenemase-producing bacteria. *Nat Commun* 2024; **15**: 2758.

36   Jamil S, Obaidullah Md, Alam M. Unusual increase of cesarean section delivery
in Bangladesh: correspondence. *Ann Med Surg (Lond)* 2023; **85**: 2242.

37   Sarker AR, Zabeen I, Hossain Z, Ali N, Khan JAM. Increasing rates of cesarean
birth in Bangladesh: A household-level pooled analysis. *Birth* 2024; **51**: 326–42.

38   Kabir S, Hasan MdR, Hossain MdI, *et al.* Determinants and Trends of Health
Facility Delivery in Bangladesh: A Hierarchical Modeling Approach. *Biomed Res Int*
2022; **2022**: 1359572.

39   Aktar S, Nu UT, Rahman M, *et al.* Trends and risk of recurrent preterm birth in
pregnancy cohorts in rural Bangladesh, 1990–2019. *BMJ Glob Health* 2023; **8**:
e012521.

40   Salam SS, Rahman AE, Mhajabin S, *et al.* Perception and experiences of
adolescent mothers and communities in caring for their preterm babies: findings from
an in-depth study in rural Bangladesh. *BMC Pregnancy Childbirth* 2024; **24**: 145.

41   Milton R, Gillespie D, Dyer C, *et al.* Neonatal sepsis and mortality in low-income
and middle-income countries from a facility-based birth cohort: an international
multisite prospective observational study. *The Lancet Global Health* 2022; **10**: e661–
72.

42   Ebi KL, Capon A, Berry P, *et al.* Hot weather and heat extremes: health risks. *The
Lancet* 2021; **398**: 698–708.

43   Dimitrova A, Dimitrova A, Mengel M, Gasparrini A, Lotze-Campen H, Gabrysch S.
Temperature-related neonatal deaths attributable to climate change in 29 low- and
middle-income countries. *Nat Commun* 2024; **15**: 5504.

44   Hu Y, Yang Y, Feng Y, *et al.* Prevalence and clonal diversity of carbapenem-
resistant Klebsiella pneumoniae causing neonatal infections: A systematic review of
128 articles across 30 countries. *PLOS Medicine* 2023; **20**: e1004233.

611