Deep brain stimulation-responsive subthalamo-cortical coupling in obsessive-compulsive disorder – a case study

Lucie Winkler MSc[1], Lucy M. Werner MSc[1], Markus Butz PhD[1], Christian J. Hartmann MD[12], Alfons Schnitzler MD[12], Jan Hirschmann PhD[1]*

[1] Institute of Clinical Neuroscience and Medical Psychology, Medical Faculty and University Hospital Düsseldorf, Heinrich Heine University, 40225, Düsseldorf, Germany

[2] Center for Movement Disorders and Neuromodulation, Department of Neurology, Medical Faculty and University Hospital Düsseldorf, Heinrich Heine University, 40225, Düsseldorf, Germany

*Jan Hirschmann
Email: Jan.Hirschmann@med.uni-duesseldorf.de

Background
Deep brain stimulation (DBS) reduces beta-band coupling between the subthalamic nucleus (STN) and cortex in Parkinson's disease (PD). How DBS affects beta-band coupling in obsessive-compulsive disorder (OCD) is unknown.

Objective
To characterize the effect of DBS on STN-cortex coupling in OCD.

Methods
Local field potential recordings from the STN via a sensing-capable DBS system were combined with magnetoencephalography. We recorded one OCD patient at rest (DBS OFF and ON), who further completed a Go/NoGo task (DBS OFF). Task-related brain activity was compared between the OCD and a PD patient.

Results
Resting-state STN power and STN-cortex coherence spectra exhibited right-lateralized beta peaks in the OCD patient, which were diminished by DBS. Prominent task-related modulations of STN activity occurred in the theta band for the OCD patient, and in the beta band for the PD patient.

Conclusions
DBS reduces subthalamic beta oscillations in patients without motor symptoms.

Introduction

Beta oscillations within the subthalamic nucleus (STN) have become a central focus in understanding the pathophysiology of Parkinson's disease (PD) and the therapeutic mechanisms of deep brain stimulation (DBS). In PD patients, beta activity is pathologically enhanced in the STN and other structures of the cortico-basal ganglia loop (1-3), and is widely recognized to contribute to PD motor symptoms such as bradykinesia and rigidity (4, 5). DBS of the STN alleviates motor symptoms, presumably by reducing excessive beta activity in the STN (6-8) and sensorimotor cortex (6, 9). These findings suggest a causal role of subthalamic beta oscillations in motor slowing. This notion, however, is mostly based on observations in PD patients.

Here, we examined a patient with obsessive-compulsive disorder (OCD), a psychiatric condition marked by persistent thoughts and urges (obsessions), and repetitive actions or mental operations (compulsions) (10). OCD is characterized by pathologically enhanced overconnectivity in a network spanning limbic cortical regions, the striatum, and the STN (11-15). Altered theta activity in the STN (16, 17) and cortex (18-20) has been identified as a potential biomarker of OCD pathology. Importantly, OCD is a condition for which, despite the absence of motor slowing, DBS of the STN is being applied as a therapeutic intervention. The therapeutic benefit possibly acts through a reduction of theta oscillations in the fronto-basal ganglia pathway (21, 22). The role of beta oscillations in OCD has not been studied extensively, but some studies suggest that beta activity is altered in cortex (23) and the STN (16) and that DBS is associated with both increases (22) and decreases (24) of beta activity in the stria terminalis/anterior limb of the internal capsule and frontal cortex. However, while it has been demonstrated that beta oscillations are present in the dorsal (17) and anteromedial STN (11) in OCD, it remains unknown whether and how DBS influences these oscillations. Therefore, we aimed to examine the effect of DBS in OCD.

Methods

The detailed methods can be found in the Supplemental Materials.

Patients

A female patient in her fifties suffering from severe OCD, marked by excessive washing of the hands, participated with written informed consent. She was implanted with DBS electrodes (3389) 12 years ago, and received a new stimulator before this study. DBS reduced her Yale-Brown Compulsive Obsessive Scale score substantially, from 39/40 pre-operatively to 7 at the time of measurement. No medications were taken at the time of measurement. For comparison, we present data from a female tremor-dominant idiopathic PD patient in her sixties in the Med ON state (DBS system implanted 3 years ago; UPDRS Part III: Med OFF/DBS OFF 51, Med OFF/DBS ON: 18). Both patients were implanted with a Medtronic Percept PC (Medtronic Inc., Minneapolis, MN, USA), capable of measuring local field potentials (LFPs) from the implanted DBS leads. The study

was approved by the Ethics Committee of the Medical Faculty of Heinrich Heine University Düsseldorf.

Recordings and stimulation

MEG was measured using a 306-channel MEG system (VectorView, MEGIN, Espoo, Finland), and we additionally monitored muscular, ocular, and cardiac activity. LFPs were recorded by the Percept system in the BrainSense streaming mode. We performed a 5 min resting-state recording in DBS OFF and subsequently applied unilateral stimulation at 130 Hz for 5 min in each hemisphere (amplitude: 1.2 mA; pulse width: 60 µs).

Paradigm

Both patients completed a cued Go/NoGo task (see Supplemental Fig. 2; OCD patient: 4 blocks; PD patient: 3 bocks; 120 trials per block). Each trial began with a fixation cross, followed by a cue indicating whether the upcoming target stimulus is likely Go or NoGo, the Go or NoGo stimulus, and feedback. Patients were instructed to press a button with their right index finger in case of a Go stimulus and to withhold any response in case of a NoGo stimulus.

Signal processing and data analysis

Following data cleaning and preprocessing (Supplemental Methods) the resting-state data was divided into non-overlapping 1 s segments. LFP power and LFP-MEG coherence were computed using Welch's method with a Hanning taper (25). For LFP power, we isolated the oscillatory components from the aperiodic background (26). Source-level coherence between the STN and cortex was computed in the beta band (13-30 Hz) by means of *Dynamic Imaging of Coherent Sources* (27).

For the Go/NoGo task data, we extracted trials centered on the Go/NoGo stimulus (-1.5 - 2 s). We source-reconstructed the activity of left and right primary motor cortices (M1, hand-knob) using Linearly Constrained Minimum Variance beamforming (28). Time-frequency spectra (2-45 Hz) were computed for STN and M1 bilaterally. As baseline, we used power averaged over all trial types (i.e. congruent/incongruent Go/NoGo trials) and all time points within those trials. Differences in power and coherence between DBS ON/OFF and Go/NoGo trials were identified through cluster-based permutation tests (29).

Results

DBS reduced STN beta power and STN-cortex beta coherence in OCD

When analyzing the resting-state data of the OCD patient, we found a pattern resembling the PD literature. Specifically, we observed a prominent peak in the beta-band for right STN power (Fig. 1A) and for coherence between right STN and right sensorimotor cortex (Fig. 1B). The beta power peak was reduced by stimulation of the left ($t_{clustersum} \leq -10.775$, $p < 0.001$) and particularly

of the right ($t_{\text{clustersum}} \le -95.168$, $p < 0.001$) STN (Fig. 1C). Similarly, the beta peak in STN-sensorimotor cortex coherence was suppressed by DBS of the right ($t_{\text{clustersum}} \le -14.258$, $p \le 0.02$) or left STN ($t_{\text{clustersum}} \le -14.704$, $p \le 0.002$; Fig. 1D). Detailed statistical results can be found in Supplemental Tables 1-2.

Figure 1. DBS reduced beta power and beta coherence in the OCD patient. Log10-transformed resting-state power spectra (aperiodic fit subtracted) and topographies of beta (13-30 Hz) STN-MEG sensor coherence for the right **(A)** and left **(B)** STN. **(C; left)** Log10-transformed resting-state power spectra (aperiodic fit subtracted) of the right STN during DBS OFF and during right and left DBS ON. Significant differences for right DBS ON *vs.* OFF: blue shade; left DBS ON *vs.* OFF: red shade; overlapping clusters: purple shade. **(C; right)** Topography of coherence between the right STN and the MEG sensors during right DBS ON *vs.* OFF and left DBS ON *vs.* OFF. Channels significantly modulated by right DBS are marked in black (left DBS: white, overlap: grey). **(D)** The source-localized contrast between DBS ON and DBS OFF is presented below (left and right DBS averaged).

Task-related modulations of subthalamic oscillations distinguish OCD and PD

Surprised by how closely the resting-state patterns of the OCD patient resembled those reported for PD, we wondered whether we would find a more distinct oscillatory signature in a task. Thus, we had the OCD patient perform a Go/NoGo task and compared the recordings to a PD patient measured with the same setup.

On the cortical level, the responses were rather similar (Fig. 2). In Go trials, we observed movement-related beta suppression after the Go stimulus, followed by a beta rebound. In NoGo trials, the suppression was interrupted by an early increase in beta power, differentiating response inhibition from execution (OCD M1: $t_{clustersum}$ = -561.202, p < 0.001; PD M1: $t_{clustersum}$ = -711.0559, p < 0.001). On the level of the STN, the beta-dominated pattern seen in cortex repeated in the PD patient ($t_{clustersum}$ = -312.723, p < 0.001), but appeared to be shifted in frequency in the OCD patient, with differences arising in the theta band in left (Fig. 2) and right STN (Supplemental Fig. 3). Specifically, NoGo trials were associated with higher theta power than Go trials ($t_{clustersum}$ = -333.027, p < 0.001; Fig. 2A). Details are reported in Supplemental Tables 3-4.

Figure 2. Modulations of left subthalamic power associated with response inhibition differed between the OCD and the PD patient. Top: time-frequency power spectra, contrast between Go and NoGo trials (pooled over cue types), for the OCD patient **(A)** and the PD patient **(B)**. The difference between baseline-corrected Go trials (% change) and baseline-corrected NoGo trials (% change) is color-coded. Significant differences are marked by hatched lines within contours. Bottom: band-average power time course, for beta (13-30 Hz), and theta (3-8 Hz) frequencies.

Discussion

DBS in OCD is rare and its effects on neural oscillations and their synchronization across basal ganglia cortex loops are underexplored. Here, we demonstrate the existence of prominent beta oscillations, synchronized across STN and motor cortex in the resting-state, in a single OCD patient. Interestingly, DBS suppressed these oscillations, as reported previously for PD patients (7, 9, 30).

Notably, we did not observe any DBS-entrained gamma activity in the OCD patient at half the stimulation frequency, in line with the idea that it is related to dopaminergic medication intake (31).

By providing a non-movement disorder control, this paper makes an important contribution to the discussion on the link between two major effects of DBS: the dampening of beta oscillations and the concurrent improvement of motor symptoms. Our results clearly indicate that these two effects can (but need not) dissociate, with beta-suppression occurring in the absence of any changes in motor performance. We thus conclude that DBS-responsive beta oscillations are not necessarily a sign of PD. The fact that they do occur in several diseases suggests that they might relate to physiological functions (32), such as somatosensory processing, the integration of sensory feedback with existing knowledge (33), and the maintenance of current motor/cognitive output (34). Alternatively, they might represent a common feature of PD and OCD, such as a high level of inhibition, arising either as a consequence of neurodegeneration (PD) or from volitional processes such as withstanding compulsions (OCD). The STN is likely involved in either process, as it integrates cognitive, limbic and motor processes (32), with beta oscillations occurring in both motor- (17) and non-motor (11, 35) regions.

In contrast to the resting-state recordings, the Go/NoGo task revealed an oscillatory pattern not familiar from the PD literature, which emphasizes the involvement of beta oscillations (36). In the OCD patient, however, we found the strongest responses in the theta band. This finding aligns with previous research linking altered theta activity to OCD symptoms (16, 17), such as the inability to inhibit compulsive behaviors or deal with conflict, an established correlate of theta activity (37). To ensure that the deviation was not due to the methodology applied here, we repeated the experiment in a single PD patient. As expected, we observed marked modulations of STN beta power, possibly reflecting the disease's characteristic overactivity of STN-cortical pathways, presumably due to insufficient shielding of the STN from motor cortical drive (3), which exhibited task-related beta modulations in both patients. Although beta and theta oscillations have been observed in the non-motor area of the STN in OCD (35) and in the motor STN in PD (38), respectively, we cannot rule out that a more anteromedial vs. dorsolateral electrode placement effected these results (Supplemental Figure 1).

In summary, this case study illustrates how a task may uncover disease-specific STN oscillations that are not apparent in resting-state. This aligns with functional MRI research, demonstrating that task-based connectivity patterns contain more behaviorally relevant information than resting-state connectivity (39, 40).

Funding Sources and Conflict of Interest

This project was funded by Brunhilde Moll Stiftung. The authors declare that there are no conflicts of interest relevant to this work.

Financial Disclosures for the Previous 12 Months

AS has been serving as a consultant for Abbott, Zambon, Medtronic Inc, and AbbVie and has received speaker's honoraria from BSH Medical Communication, Abbott, Kyowa Kirin, Novartis, AbbVie, Alexion, and GE Healthcare. He received royalties from Georg Thieme Verlag and receives funding through his institution for his research from the German Research Council (Project 4247788381-TRR 295), Abbott, and Brunhilde Moll Stiftung. JH also receives support from Brunhilde Moll Stiftung. CJH received honoraria from Abbott, AbbVie, Alexion, and Univar. LW, LMW, and MB declare that there are no additional disclosures to report.

Ethical compliance

The study was approved by the Ethics Committee of the Medical Faculty of Heinrich Heine University Düsseldorf (reference number: 2021-1587_1-andere Forschung erstvotierend). Informed consent was obtained from both participants.

Data Availability Statement

Data can be made available in anonymized form upon reasonable request.

1.   Brown P, Oliviero A, Mazzone P, Insola A, Tonali P, Di Lazzaro V. Dopamine dependency of oscillations between subthalamic nucleus and pallidum in Parkinson's disease. The Journal of Neuroscience. 2001;21(3):1033-8.

2.   Hirschmann J, Özkurt TE, Butz M, Homburger M, Elben S, Hartmann CJ, et al. Distinct oscillatory STN-cortical loops revealed by simultaneous MEG and local field potential recordings in patients with Parkinson's disease. NeuroImage. 2011;55(3):1159-68.

3.   Oswal A, Cao C, Yeh C-H, Neumann W-J, Gratwicke J, Akram H, et al. Neural signatures of hyperdirect pathway activity in Parkinson's disease. Nature Communications. 2021;12(1):5185.

4.   Kühn AA, Tsui A, Aziz T, Ray N, Brücke C, Kupsch A, et al. Pathological synchronisation in the subthalamic nucleus of patients with Parkinson's disease relates to both bradykinesia and rigidity. Experimental Neurology. 2009;215(2):380-7.

5.   Lofredi R, Tan H, Neumann W-J, Yeh C-H, Schneider G-H, Kühn AA, et al. Beta bursts during continuous movements accompany the velocity decrement in Parkinson's disease patients. Neurobiology of Disease. 2019;127:462-71.

6.   Muthuraman M, Bange M, Koirala N, Ciolac D, Pintea B, Glaser M, et al. Cross-frequency coupling between gamma oscillations and deep brain stimulation frequency in Parkinson's disease. Brain. 2020;143(11):3393-407.

7.   Mathiopoulou V, Lofredi R, Feldmann LK, Habets J, Darcy N, Neumann WJ, et al. Modulation of subthalamic beta oscillations by movement, dopamine, and deep brain stimulation in Parkinson's disease. npj Parkinson's Disease. 2024;10(1):77.

8.   Wilkins KB, Petrucci MN, Lambert EF, Melbourne JA, Gala AS, Akella P, et al. Beta burst-driven adaptive deep brain stimulation improves gait impairment and freezing of gait in Parkinson's disease. medRxiv. 2024.

9.   Abbasi O, Hirschmann J, Storzer L, Özkurt TE, Elben S, Vesper J, et al. Unilateral deep brain stimulation suppresses alpha and beta oscillations in sensorimotor cortices. NeuroImage. 2018;174:201-7.

10.   American Psychiatric Association. Diagnostic and statistical manual of mental disorders 5ed: American Psychiatric Publishing; 2013.

11.   Wojtecki L, Hirschmann J, Elben S, Boschheidgen M, Trenado C, Vesper J, et al. Oscillatory coupling of the subthalamic nucleus in obsessive compulsive disorder. Brain. 2017;140(9):e56.

12.   Qing X, Gu L, Li D. Abnormalities of localized connectivity in obsessive-compulsive disorder: A voxel-wise meta-analysis. Frontiers in Human Neuroscience. 2021;15:739175.

13.   Beucke JC, Sepulcre J, Talukdar T, Linnman C, Zschenderlein K, Endrass T, et al. Abnormally high degree connectivity of the orbitofrontal cortex in obsessive-compulsive disorder. JAMA Psychiatry. 2013;70(6):619-29.

14.	Smith EE, Schüller T, Huys D, Baldermann JC, Andrade P, Allen JJB, et al. A brief demonstration of frontostriatal connectivity in OCD patients with intracranial electrodes. NeuroImage. 2020;220:117138.

15.	Horn A, Li N, Meyer GM, Gadot R, Provenza NR, Sheth SA**.** Deep Brain Stimulation response circuits in Obsessive Compulsive Disorder. Biological Psychiatry.

16.	Bastin J, Polosan M, Piallat B, Krack P, Bougerol T, Chabardès S, et al. Changes of oscillatory activity in the subthalamic nucleus during obsessive-compulsive disorder symptoms: Two case reports. Cortex. 2014;60:145-50.

17.	Rappel P, Marmor O, Bick AS, Arkadir D, Linetsky E, Castrioto A, et al. Subthalamic theta activity: a novel human subcortical biomarker for obsessive compulsive disorder. Translational Psychiatry. 2018;8(1):118.

18.	Desarkar P, Sinha VK, Jagadheesan K, Nizamie SH**.** Subcortical functioning in obsessive-compulsive disorder: an exploratory EEG coherence study. The World Journal of Biological Psychiatry. 2007;8(3):196-200.

19.	Lee I, Kim KM, Lim MH**.** Theta and gamma activity differences in obsessive-compulsive disorder and panic disorder: Insights from resting-state eeg with eLORETA. Brain Sciences. 2023;13(10).

20.	Perera MPN, Mallawaarachchi S, Bailey NW, Murphy OW, Fitzgerald PB**.** Obsessive-compulsive disorder (OCD) is associated with increased electroencephalographic (EEG) delta and theta oscillatory power but reduced delta connectivity. Journal of Psychiatric Research. 2023;163:310-7.

21.	Figee M, Luigjes J, Smolders R, Valencia-Alfonso C-E, van Wingen G, de Kwaasteniet B, et al. Deep brain stimulation restores frontostriatal network activity in obsessive-compulsive disorder. Nature Neuroscience. 2013;16(4):386-7.

22.	Schwabe K, Alam M, Saryyeva A, Lütjens G, Heissler HE, Winter L, et al. Oscillatory activity in the BNST/ALIC and the frontal cortex in OCD: Acute effects of DBS. Journal of Neural Transmission. 2021;128(2):215-24.

23.	Koh MJ, Seol J, Kang JI, Kim BS, Namkoong K, Chang JW, et al. Altered resting-state functional connectivity in patients with obsessive–compulsive disorder: A magnetoencephalography study. International Journal of Psychophysiology. 2018;123:80-7.

24.	Xiong B, Wen R, Gao Y, Wang W**.** Longitudinal changes of local field potential oscillations in nucleus accumbens and anterior limb of the internal capsule in obsessive-compulsive disorder. Biological Psychiatry. 2023;93(11):e39-e41.

25.	Oostenveld R, Fries P, Maris E, Schoffelen JM**.** FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Computational Intelligence and Neuroscience. 2011;2011(1):156869.

26. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al. Parameterizing neural power spectra into periodic and aperiodic components. Nature Neuroscience. 2020;23(12):1655-65.

27. Gross J, Kujala J, Hämäläinen M, Timmermann L, Schnitzler A, Salmelin R. Dynamic imaging of coherent sources: Studying neural interactions in the human brain. Proceedings of the National Academy of Sciences. 2001;98(2):694-9.

28. Van Veen BD, Buckley KM. Beamforming: A versatile approach to spatial filtering. IEEE ASSP Magazine. 1988;5(2):4-24.

29. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. Journal of Neuroscience Methods. 2007;164(1):177-90.

30. Oswal A, Beudel M, Zrinzo L, Limousin P, Hariz M, Foltynie T, et al. Deep brain stimulation modulates synchrony within spatially and spectrally distinct resting state networks in Parkinson's disease. Brain. 2016;139(5):1482-96.

31. Colombo A, Bernasconi E, Alva L, Sousa M, Debove I, Nowacki A, et al. Finely tuned γ tracks medication cycles in Parkinson's disease: An ambulatory brain-sense study. Movement Disorders. 2025.

32. Accolla EA, Horn A, Herrojo-Ruiz M, Neumann W-J, Kühn AA. Reply: Oscillatory coupling of the subthalamic nucleus in obsessive compulsive disorder. Brain. 2017;140(9):e57-e.

33. Barone J, Rossiter HE. Understanding the role of sensorimotor beta oscillations. Frontiers in Systems Neuroscience. 2021;15(51).

34. Engel A, Fries P. Beta-band oscillation - signalling the status quo? Current Opinion in Neurobiology. 2010;20(2):156-65.

35. Accolla EA, Herrojo Ruiz M, Horn A, Schneider GH, Schmitz-Hübsch T, Draganski B, et al. Brain networks modulated by subthalamic nucleus deep brain stimulation. Brain. 2016;139(Pt 9):2503-15.

36. Kühn AA, Williams D, Kupsch A, Limousin P, Hariz M, Schneider GH, et al. Event-related beta desynchronization in human subthalamic nucleus correlates with motor performance. Brain. 2004;127(4):735-46.

37. Zavala B, Brittain J-S, Jenkinson N, Ashkan K, Foltynie T, Limousin P, et al. Subthalamic nucleus local field potential activity during the Eriksen Flanker Task reveals a novel role for theta phase during conflict monitoring. The Journal of Neuroscience. 2013;33(37):14758.

38. Olson JW, Nakhmani A, Irwin ZT, Edwards LJ, Gonzalez CL, Wade MH, et al. Cortical and subthalamic nucleus spectral changes during limb movements in Parkinson's disease patients with and without dystonia. Movement Disorders. 2022;37(8):1683-92.

39. Zhao W, Makowski C, Hagler DJ, Garavan HP, Thompson WK, Greene DJ, et al. Task fMRI paradigms may capture more behaviorally relevant information than resting-state functional connectivity. NeuroImage. 2023;270:119946.

40.   Elliott ML, Knodt AR, Cooke M, Kim MJ, Melzer TR, Keenan R, et al. General functional connectivity: Shared features of resting-state and task fMRI drive reliable and heritable individual differences in functional brain networks. NeuroImage. 2019;189:516-32.