Effects of Soluble Corn Fiber Consumption on Executive Functions and Gut Microbiota in Middle to Older Age Adults: A Randomized Controlled Crossover Trial

David A. Alvarado[1,4*], Tori A. Holthaus[1*], Shelby Martell[2], Nicole L. Southey[1], Marco Atallah[4], Rhea Sarma[3], David Revilla[4], Marina Brown[4], Twinkle Mehta[3], Naiman A. Khan[1,2,3,5], Hannah D. Holscher[1,4**]

[1]Division of Nutritional Sciences, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

[2]Neuroscience Program, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

[3]Health and Kinesiology, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

[4]Department of Food Science and Human Nutrition, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

[5]Beckman Institute of Advanced Science and Technology, Urbana-Champaign, Urbana, IL 61801, USA

[*]Shared first-authorship

[**]**Corresponding Author:** Hannah D. Holscher, 1201 West Gregory Drive, Urbana, IL 61801 (217) 300-2512, hholsche@illinois.edu

Abbreviations: *Parabacteroides distasonis* (*P. distasonis*), Cambridge Neuropsychological Test Automated Battery (CANTAB), carbohydrate-active enzymes (CAZyme), Event-related potential (ERP), FDR (False Discovery Rate), Maltodextrin (CON, control), Perceived Stress Score (PSS), Positive and Negative Affect Schedule (PANAS), Profile of Moods Survey (POMS), Reaction Time (RT), SE (Standard Error), SEM (Standard Error of the Mean), Short-chain fatty acids (SCFA), Branched-chain fatty acids (BCFA), Soluble corn fiber/ PROMITOR® Soluble Fibre™ (SCF), Accuracy (ACC)

Abstract

Background: Dietary fiber may support cognition through gastrointestinal-microbiota mechanisms, but clinical evidence is limited.

Objectives: We aimed to determine whether soluble corn fiber (SCF) improved cognition and altered fecal microbiota and fermentation end products in adults.

Methods: In a randomized, double-blind, crossover trial, 42 healthy adults (45-75y) consumed SCF (18g/d) or a maltodextrin placebo control (CON: 22g/d) for 4 weeks, separated by a washout. Cognitive outcomes included executive function with event-related potentials, relational memory, neuropsychological performance, and mood. Secondary outcomes included fecal microbiota, metabolomics, and gastrointestinal tolerance. Tertiary analyses related microbial and metabolite changes to cognitive improvements using correlation, mediation, and moderation models, and explored SCF fermentation pathways with 16S-predicted functional profiling, shotgun metagenomics and *in vitro* culturing.

Results: SCF improved reaction times (RT) during congruent (β = -9.8 ms, 95% CI: [-18.4, -1.2], FDR P = 0.01) and incongruent (β = -14.2 ms, 95% CI: [-22.8, -5.6], FDR P = 0.003) flanker trials and increased *Parabacteroides* (~4-fold, β = 1.44 log, 95% CI [1.01, 1.88], FDR P < 0.001). At the SCF endpoint, congruent RT tended to be inversely associated with fecal acetate (ρ = -0.33) and propionate (ρ = -0.36)*,* while *Parabacteroides* was marginally positively associated with acetate (ρ = 0.34) (all FDR P < 0.1). Moderation analyses indicated that SCF-RT relation varied by *Parabacteroides* magnitude change. At endpoint, SCF increased predicted functional potential of carbohydrate-related KOs and pathways (FDR P < 0.05). *In vitro* culturing confirmed *P. distasonis* ferments SCF.

Conclusion: SCF consumption improved attentional inhibition, altered the gut microbiota, and

selectively enriched *Parabacteroides*. Although mediation analyses did not support a direct

microbiota-to-cognition pathway, moderation analyses suggested that SCF-related cognitive

effects may depend in part on *Parabacteroides* abundance. Collectively, these findings suggest

that certain cognitive benefits of SCF consumption may be partly underpinned by the gut

microbiota.

Keywords: Reaction time; Cognition; *Parabacteroides*; Gut-Brain; Dietary Fiber

Introduction

A healthy diet is a key determinant of metabolic health, and emerging evidence extends these benefits to brain function and cognition [1–3]. Dietary fiber is associated with reduced risk of obesity, cardiovascular disease, type 2 diabetes, and colon cancer [4,5], yet nearly 95% of Americans fail to meet recommended intake levels [6]. Further, dietary fiber provides a primary substrate for microbes residing in the gastrointestinal (GI) tract, shaping ecological community diversity and metabolic capacity [7]. Preclinical studies show that dietary fibers improve cognition [8,9], but human trials remain limited [10–12], particularly those linking the microbiota, GI-derived metabolites, and cognitive outcomes [13]. Therefore, further evidence is necessary to comprehensively characterize the effect of dietary fiber on general and specific domains of cognitive function.

Soluble corn fiber (SCF) is a resistant maltodextrin with an average degree of polymerization (DP) between 10-12 units [14], that supports glycemic control, calcium absorption, and gastrointestinal health [15–17]. Its non-viscous, water-soluble structure lacks β-linkages, yet branching of the α-linkages confers resistance to complete digestion in the small intestine (<30%) [18,19]. The undigested fraction undergoes colonic fermentation, enriching saccharolytic taxa such as *Bifidobacterium*, *Lactobacillus*, and *Parabacteroides* [20,21], and increasing short-chain fatty acids (SCFA) concentrations across age groups [22], while reducing deleterious fermentation end products (phenols, indoles, ammonia) [19,23]. SCFAs are increasingly recognized as key mediators of GI-brain communication by binding free fatty acid receptors (FFAR/FFAR3) to regulate immune [24] and enteroendocrine signaling [25], modulate microglial activity and systemic inflammation [26], and stimulate vagal and serotonergic pathways [27–30].

Executive functions, including inhibitory control, are central to attention, working memory, and flexible behavior [31]. These top-down processes can be assessed through computer-

based tasks that measure a subject's ability to maintain attention while faced with distracting

stimuli, hold and manipulate information in mind, and require multi-tasking and set-shifting, while

electrophysiological markers can provide complementary insight into neural activity during such

tasks [32,33]. Given the growing evidence that microbially derived metabolites influence brain

activity [27,34–39], assessing both behavioral and neural indices provides a comprehensive overview

to evaluate dietary fiber's cognitive impact.

To address these gaps, we conducted a randomized, double-blind, placebo-controlled,

crossover trial to test the impact of daily SCF consumption in middle- to older-aged adults. The

primary aim was to determine whether four weeks of SCF consumption improved executive

function, changes in the amplitude (size) and latency (timing) of the P3 event-related potential

(ERP), as well as additional behavioral tasks assessing relational memory and neuropsychological

function. We hypothesized that SCF would enhance cognitive control and reduce the number of

neural resources required to accomplish these cognitive processes. The secondary aim examined

the effects of SCF on fecal microbiota composition and fermentation metabolites, hypothesizing

that SCF consumption would enrich health-associated taxa and effect microbial-derived

metabolites. Finally, the tertiary aim tested whether the SCF-induced changes in microbial or

metabolite profiles explained changes in cognitive outcomes, hypothesizing that these changes

would contribute to the cognitive benefits induced by SCF.

Methods

Study design and intervention

This clinical trial, registered as the SCOPE study (Soluble Corn fiber for Promoting

Executive function; NCT05066425), included two 4-week intervention periods separated by an

~4-week washout. Participants consumed soluble corn fiber (SCF; PROMITOR® Soluble Fibre™,

22g/d containing 18g fiber) or the maltodextrin placebo control (CON; 22g/d) in a counterbalanced

order. The two interventions were provided by Tate & Lyle in identical, pre-measured sachets that

were indistinguishable in taste and appearance. Sachets were labeled as 789 or 030, and

participants and research staff remained blinded to the code assignment until data analysis was

complete to ensure blinding. Participants were given 35 sachets per condition and were instructed

to consume one daily, dissolved in a fluid of their choice. Participants recorded intake of the sachet

on daily compliance logs. Unused sachets and logs were collected at the end of each period to

assess study adherence.

All study visits and testing sessions were conducted at the University of Illinois Urbana-

Champaign (Supplemental **Figure 1**). Anthropometric measures, physical activity level,

demographic information, and medical history were collected at baseline and visits 4, 5, and 6 as

potential covariates.

Participants criteria

Adults aged 45-75 years were recruited from the East Central Illinois area using email

newsletters, social media, community postings using the postal service, and flyers posted on public

transport. All participants were informed of all procedures and provided their written informed

consent before any data was collected. The SCOPE study was approved by the University of

Illinois Institutional Review Board (May 2021) and conducted following the principles of the

Declaration of Helsinki.

Exclusion criteria included physician-diagnosed gastrointestinal or metabolic disease

(inflammatory bowel disease and diabetes), neurological disorder (dementia, Parkinson's disease,

multiple sclerosis, brain tumors, autism), colorblindness or vision not corrected to 20/20, antibiotic

use within the past month, tobacco use, food allergies, pregnancy, or lactating, BMI <18.5 or >34.9

kg/m^2, Mini-Mental State Examination <24, or if habitual dietary fiber intake was >70% of the

daily recommended intake (≥18 g/d for women, ≥27 g/d for men) using the Dietary History

Questionnaire (*DHQ3*) food frequency questionnaire.

Eligible participants agreed to maintain habitual dietary patterns and physical activity, to

avoid probiotic or prebiotics supplements, and to report any changes in medications during the

SCOPE study.

Enrolled participants (**Table 1**) were randomized via coin toss to start one of the two

intervention conditions. Height and weight were measured in triplicate, without shoes and in light

clothing, to determine BMI using a *SECA* stadiometer (model 240; *SECA*, Hamburg, Germany)

and a Tanita WB-300 Plus digital scale (Tanita, Tokyo, Japan). Physical activity was assessed

using the Godin-Shephard Leisure-Time Exercise Questionnaire (*Godin*), which records the

frequency of strenuous, moderate, and light physical activities performed during a typical week,

which are totaled into a final score using a weighted formula [40].

Aim 1: Cognitive Outcomes

To minimize learning effects, participants completed practice sessions at the start of each

visit until at least 70% accuracy was met before experimental cognitive sessions were

administered. The order of cognitive task administration at each testing visit (visits 3-6) was

counterbalanced using a 2x2 Latin square design for each participant,

Executive function

Attentional inhibition was assessed using a modified Eriksen flanker task as previously

described [41]. Participants were asked to respond (*Current Designs*, Philadelphia, PA, USA) quickly

and accurately to the directionality of a central stimulus (3-cm tall white arrow) presented with

two flanking stimuli (arrows) on each side (83s stimulus presentation; 100ms response window;

jittered intertrial intervals of 1100, 1200, 1500ms). In congruent trial types, the flanking stimuli

matched the direction of the central arrow (<<<<<, >>>>>); during incongruent trials, flanking

stimuli opposed the central arrow (>><>>, <<><<), requiring greater inhibitory control. The task

began with 40 practice trials followed by two blocks of 200 randomized experimental trials.

Behavioral measures were accuracy (ACC) and reaction time (RT) for correct responses.

Interference effects (flanker effect) for ACC and RT were calculated, respectively, by subtracting

incongruent from congruent measures (ACC) and congruent from incongruent measures (RT).

These measures index the ability to maintain task performance when resolving conflicting

information.

Neuroelectric indices

EEG activity was recorded simultaneously during the flanker task via a Neuro-scan Quik-

cap with 64 passive, wet, sintered Ag/AgCl scalp electrodes arranged in the international 10–10

system, with electrodes placed near the outside of each eye and above and below the left eye [42]. A

midline sensor placed between CZ and CPZ served as a reference, and AFZ served as the ground.

Continuous EEG data were digitized using a Neuroscan SynampsRT amplifier at a sampling rate

of 500 Hz, amplified 500 times to an online low-pass 70-Hz filter with a direct current and a 60-

Hz notch filter. Electrode impedance values were maintained at $\leq$20 kohms.

Signal processing was performed in MATLAB (MathWorks, Natick, MA, version 2021b)

using EEGLab and ERPlab [43,44]. Continuous data were re-referenced to the average of two mastoid

electrodes. A trained research team member visually inspected all channels for excessive noise

during pre-processing. Channels with excessive noise were interpolated with an average of the

adjacent electrodes. Data were submitted to a 0.1 Hz high-pass Butterworth filter before eye-blink

artifacts were rejected using independent component analysis (ICA). ICA and vertical eye sensor

channel correlations greater than 0.35 were considered eye-blinks. A time window of -200 to

1200 ms around stimulus presentation was used for creating stimulus-locked epochs with a

baseline correction of -200 ms to stimulus onset. A 30-Hz low-pass filter was used. Artifacts were

detected and rejected if a moving window peak-to-peak amplitude exceeded 100 µV, using a 100-

ms window width and step. Only correct trials with >80% of artifact-free trials in all flanker task

trials were retained, and participants with excessive artifacts (>20%) were excluded ($n = 3$ CON,

$n = 4$ SCF), consistent with previous literature [45,46].

Electrodes of interest were chosen based on a collapsed localizer technique [47]. Evidence

for selection was observed from maximal voltage localization of combined baseline and post-

testing trials collapsed across condition and trial type. *Post-hoc* topographic average plots were

constructed using a stylized topographic map plugin for EEGLAB/ERPLAB [48]. The P3 component

was defined as the region of interest comprised of C1, CZ, C2, CPZ, CP1, and CP2 with a localized

peak and corresponding latency between 300 and 600ms post-stimulus onset. Amplitude and

latency were extracted to complement the analysis of the behavioral Flanker task.

Relational memory

Hippocampal-dependent relational memory was assessed using the computerized spatial

reconstruction task presentation software (*iPosition*; Neurobehavioral Systems, Berkeley, CA) [49].

Participants studied arrays of six abstract shapes on the computer screen in a randomized array for

6 seconds. The shapes were then masked by black squares for 18 seconds and could be revealed

individually by clicking on them. After a 2-second delay, the stimuli reappeared at the top of the

screen, and participants were asked to reconstruct the original array. Each participant completed

20 trials with randomized arrays. Two error metrics were used to assess performance:

misplacement error (original misplacement), calculated as the average measure of distance

between the actual and reconstructed stimuli locations, and object-location binding (accurate

single item placement), the number of times the participant correctly placed a stimulus within a

pre-defined radius.

Neuropsychological battery

Additional cognitive performance across multiple domains was assessed using the

Cambridge Neuropsychological Test Automated Battery (*CANTAB*, Cambridge Cognition Ltd)

[50,51], administered on a touchscreen tablet. The battery included the Intra-Extra Dimensional Set

Shift (IED), Stop Signal Task (SST), Rapid Visual Information Processing (RVP), and Spatial

Working Memory (SWM). The IED is a test which assesses rule acquisition and reversal,

attentional set formation, maintenance, and shifting. The RVP task assesses sustained attention,

and the SST measures response inhibition. The SWM task assesses spatial working memory.

Mood and Stress

233   Mood and stress were assessed using validated self-report instruments, including the

234 Profile of Mood States (*POMS*), the Positive and Negative Affect Schedule (*PANAS*), and the

235 Perceived Stress Scale (*PSS*). The *POMS* questionnaire includes 65-items that evaluates transient

236 mood states across six subscales (tension/anxiety, depression, anger, vigor, fatigue, and

237 confusion), with ratings from 0 ("not at all") to 5 ("extremely") [52,53]. *PANAS* includes 20 items

238 divided evenly into positive and negative affect subscales, scored from 1 ("very slightly or not at

239 all") to 5 ("extremely likely") [53,54]. *PSS-10* is a 10-item measure of stress over the past month,

240 scored from 0 ("never") to 4 ("very often") [55].

242 **Aim 2: Gut Microbiota and Metabolite Outcomes**

243 *Fecal collection and processing*

244   Participants provided two fecal samples at each of the four collection timepoints (baseline

245 and end of each of the two study conditions) for a total of 8 samples per participant. Samples were

246 collected during the five days preceding the first intervention and the final five days of each

247 treatment and washout period. Samples were delivered within 15 minutes of defecation using

248 provided kits (Commode Specimen Collection Systems, *Sage Products*) with ice packs and

249 coolers. Upon arrival at the laboratory, samples were homogenized, aliquoted into cryovials for

250 sequencing, flash-frozen in liquid nitrogen, and stored at -80°C. Additional aliquots were prepared

251 for metabolite analysis: SCFAs, BCFAs, ammonia, and dry matter basis (DMB; pre-weighed

252 aluminum tins). These aliquots were stored at -20°C until analysis. Fecal pH was measured

253 (*Denver Instrument*) after samples had been aliquoted.

255 *Fecal DNA isolation and 16S amplicon analyses*

256   Fecal microbial DNA was extracted using the Power-Lyzer PowerSoil DNA Isolation Kit

257 (*QIAGEN*). DNA was verified using 1% agarose gel electrophoresis, concentrations measured

258 using a Qbit 3.0 Fluorometer (*ThermoFisher Scientific*), quantified from two samples collected

259 per timepoint per participant, and pooled in equimolar amounts before sequencing at the University

260 of Illinois W.M Keck Center for Biotechnology. 16S Amplicons were generated with V4 primers

261 (*F*: 5'-GTGYCAGMGCCGCGGTAA, *R*: 5'-GGACTACNVGGGTWTCTAAT) [56] and

262 sequenced on an Illumina NovaSeq 6000 SP (2x250 nt), producing ~1 billion reads. Sequences

263 with quality scores >25 were processed with *DADA2* [57] integrated in *QIIME2* (v2023.7) [58].

264 Sequences were denoised and resolved into amplicon sequence variants (ASVs) using the *SILVA*

265 138-99 (515F/806R) naïve Bayes reference classifier [57,59]. Samples were normalized to 350,000

266 reads; curves plateaued at this depth, indicating adequate coverage (**Supplemental Figure 2**).

267 Rarefaction was performed with *vegan* v2.6.8 [60], filtering 1,521 ASVs (13%, primarily singletons

268 or zero-count features) and retaining 10,373 ASVs. Alpha diversity and beta diversity were

269 calculated with *phyloseq* (v1.48.0) [61], and Faith's phylogenetic diversity (Faith's PD) with *picante*

270 (v1.8.2). Taxonomic summaries were created at the genus and phylum levels for downstream

271 analyses.

273 *Targeted Metabolites*

274   Fecal dry matter was measured according to the Association of Official Analytical

275 Chemists, 1984, methods [62]. Ammonia concentrations were measured following the procedure of

276 Chaney and Marbach [63]. Aliquots for SCFA and BCFA were weighed, acidified with 2N

277 hydrochloric acid, and stored at -20°C until analysis. Briefly, SCFA (acetate, propionate, butyrate),

278 BCFA (isovalerate, valerate, and isobutyrate) concentrations were determined on the supernatant

fraction of the acidified fecal samples using gas chromatography mass spectroscopy (GC-MS).

Concentrations were calculated using standards[64]. Phenol and indole were quantified, according to

Flickinger et al.[65] Fecal metabolites are reported on the normalized DMB scale.

Digestive health

GI symptoms, bowel function, and stool characteristics were assessed using 7-day self-

reported records, questionnaires, and laboratory measures for each timepoint [66]. Participants rated

GI symptom severity (burping, cramping, bloating, gas, nausea, reflux, rumbling) on a 4-point

scale from 1 ("absent") to 4 ("severe") [67]. Bowel movements and consistency were scored using a

7-point Bristol stool scale (1- "separate hard lumps"; 2- "sausage-shaped but lumpy"; 3- "sausage-

like but with cracks on the surface"; 4- "sausage-like, smooth, and soft"; 5- "soft blobs with clear

cut edges"; 6- "fluffy pieces, mushy"; 7- "entirely liquid"). And ease of passage rated on a 5-point

scale from 1 ("very easy") to 5 ("very difficult"). In the laboratory, fresh fecal samples were

measured with a pH meter (Denver Instrument) [66], and a trained research team member recorded

stool consistency using the Bristol scale.

Aim 3: Exploratory Integrative and Functional Analyses

Mediation and moderation

Selected microbial and metabolite features for integrative analyses were prioritized from

Aim 1 and 2 outcomes that met the statistical significance threshold and were also included when

they showed consistent directional trends and biological plausibility based on prior literature [27,34–

39]. Associations between these candidates and cognitive outcomes were examined using Spearman

correlations. We then used nonparametric bootstrapped causal mediation models to estimate

indirect and direct effects within a causal inference framework [68–70]. Moderation analyses were

conducted using the Johson-Neyman framework with Aiken & West's simple-slopes procedures

[71,72]. Together, these complementary approaches will provide further clarity on whether microbes

or metabolites acted as intermediaries, transmitting the treatment (mediation), or as variables that

altered the strength or direction of the treatment's effect (moderation).

In vitro isolated culturing

Parabacteroides distasonis (*P. distasonis*), a species within the genus *Parabacteroides*

(strain 31_2, deposited as *Porphyromonas* sp. HM-169), was provided from the Biodefense and

Emerging Infections Research Resources Repository (BEI Resources, NIAID, NIH) and

maintained anaerobically in brain heart infusion (BHI) and chopped-meat broth. Growth assays

followed an adaptation of Garcia-Bayona et al. [73] using M9 supplemented with 50 mg/L L-

cysteine, 5 mg/L hemin, 2.5 µg/L vitamin K1, 2 mg/L $FeSO_4·7H2O$, and 5 µg/L vitamin B12.

Treatments included glucose (0.25% wt/vol), CON (0.25% wt/vol), or SCF (0.4% wt/vol). BHI

served as a positive control; for negative controls, M9 with no carbon (no growth control), and

blanks (no inoculum) for each treatment (media contamination control). Stock solutions were

sterile-prepared, filter-sterilized (0.22 µm), and stored at 4 °C.

Three separate *P. distasonis* overnight cultures were inoculated into M9 solutions prepared

in a 10 x 10 honeycomb plate (1:10 dilution; starting OD_{600} = 0.1). For each treatment,

measurements were performed in technical triplicate across the 3 independent biological

replicates. Growth was monitored with the Bioscreen-C system (600nm, every 30 min, for 25.5h,

37°C, shaking before/after each read) (**Supplemental Figure 3**). All samples reached the

stationary phase by 14h; therefore, analyses were restricted to the first 14h. OD values were blank-

corrected and growth rates (μ) calculated at 2 h intervals.

16S-inferred functional potential

Predicted functional potential for the whole cohort was inferred from 16S data using

PICRUSt2 (v2.5.3) [74] with default parameters (*picrust2_pipeline.py*) [74,75]. Predicted functions were

summarized as KEGG Orthologs (KOs) and pathways. *KEGGREST* (v112.1) was used to extract

carbohydrate metabolism pathways (map00010, 00030, 00500, 00620, 00650) and carbohydrate-

active KOs (KO01176, 01223, 01210, 01190, 01187, 012348). Stratified KO outputs were used to

quantify genus-level contributions to predicted functional differences across conditions.

Responder shotgun metagenomics

Participants from the whole cohort were classified based on their *Parabacteroides* response

to SCF using 16S genus-level relative abundance change (Δ = POST-PRE during the SCF period).

For an exploratory responder analysis, we selected a subset of six participants comprising the two

extremes in *Parabacteroides* responses to the SCF intervention period: three participants that

experienced the largest increases in *Parabacteroides* after the SCF intervention period

(responders, n=3) and the three participants that experienced the smallest changes in

Parabacteroides after the SCF intervention period (null or negative; non-responders, n=3).

Shotgun metagenomic sequencing of the extracted fecal DNA from the 3 responders and 3 non-

responders was conducted to assess species-level taxonomic profiles and exploratory beta-

diversity and functional patterns by responder status (**Supplemental Figure 4**).

Shotgun metagenomic sequencing was performed at the W. M. Keck Center for

Biotechnology (University of Illinois Urbana-Champaign). Libraries were pooled, quantified by

qPCR, and sequenced on half of a 10B lane for 151 cycles from both ends of the fragments using

a NovaSeq X Plus with V1.0 sequencing kits. FastQ files were demultiplexed (*bcl2fastq* v2.20),

adapters (CTGTCTCTTATACACATCT) were trimmed from the 3'-end of the reads, and base

quality scores were scored using Sanger encoding (*ASCII offset* of 33). Data pre-processing was

performed on the University cluster (Carl R. Woese Institute for Genomic Biology, University of

Illinois at Urbana-Champaign) using *KneadData* (v0.12.0) for quality control and host DNA

removal (*KneadData* 20230405 release). *Trimmomatic* (v0.39) removed low-quality bases,

Tandem Repeat Finder (TRF) removed noise from repetitive sequences, and *FastQC* (v0.11.9)

generated quality reports.

Following Huttenhower BioBakery pipelines [76–78], taxonomic profiling was performed

using *MetaPhlAn* (v4.0.6) [77], with *Bowtie2* (v2.5.0) aligning reads against the

*ChocoPhlAnSGB*_202212 reference. Functional profiles were generated using *HUMAnN* (v3.7) [76]

with the *ChocoPhlAn* nucleotide database (v201901b) and *UniRef90* protein database (Nov 2024).

Profiles were merged (*merge_metaphlan_tables.py*) to produce relative abundance and read counts

[77]. Descriptive analyses compared the log fold change between responders and non-responder's

delta values, retaining gene families with >2-fold change-targeted filtering for carbohydrate

metabolism-related terms yielded 217 candidate families. Carbohydrate-active enzymes

(CAZymes) were annotated using the CAZy database [79,80] and *run_dbcan* (v4.0.0)[81], incorporating

HMMER (*dbCAN_HMMdb), DIAMOND* (CAZy protein sequences), and *dbCAN-sub* for substrate

prediction validated against PUL (Polysaccharide Utilization Loci) datasets. Reads were pre-

processed (*Trim Galore* v0.6.0, *Kraken2* v2.1.1), assembled into contigs ≥1000 bp (*MEGAHIT*

v1.2.9), and annotated with *Prokka* (v1.4) before CAZyme assignment. Clean reads were mapped

back to coding sequences (*BWA* v0.7.17-r1188, *Samtools* v1.7, *Bedtools* v2.27.1), and abundances

were normalized to transcripts per million (TPM) using *dbcan_utils* scripts [81]. Workflow was

conducted on a Linux computer (128 GB RAM) utilizing *Anaconda* (v23.7.3) environment.

Statistical analyses

General approach. Target enrollment was determined using *a priori* power calculations

(*G-Power* v3.1.9.2) [82], estimating that 36 participants would provide 80% power ($\beta = 0.80$) to

detect a small effect size ($r = 0.20$) at $\alpha = 0.05$ for Aims 1 and 2. Due to the exploratory nature of

Aim 3, results should be interpreted as hypothesis-generating and to inform effect size estimates

for future studies. To allow for attrition, 42 participants were recruited. Analyses were conducted

in *RStudio* (v2025.05.0+496; *R* v4.4.3) [83]. The significance threshold was set at $\alpha = 0.05$ with false

discovery rate (FDR) adjustments applied where relevant.

Data preprocessing and assumptions. Data distributions were checked using Shapiro-

Wilk's test (*stats* v4.4.3; $W > 0.9$), Q-Q plots (*ggplot2* v3.5.2) [84], skewness/kurtosis metrics

(*DescTools* v0.99.6, *e1071* v2.6), and residuals inspected visually (*performance* v0.13.0). Extreme

outliers (>3 SD; *rstatix* v0.7.20) were confirmed using a Bonferroni *outlierTest* (*car* v3.1-3), then

winsorized [85,86]. To meet model assumptions, BCFAs POMS, PANAS, and Godin were square

root transformed; SCFAs were log-transformed. Data are reported as mean $\pm$ SEM unless

otherwise noted.

Modeling strategy. Outcomes were analyzed using linear (*lme4* v1.1.37) or generalized

(*glmmTMB* v1.1.11) mixed-effect models with treatment, time, and their interaction

(treatment:time) as fixed effects, and participant ID as a random intercept [*Outcome ~*

Treatment*Time + (1|ID)]. Order effects were tested and did not improve model fit, demonstrating

adequate washout time, therefore excluded. Missing data were not imputed; the mixed-effect

model provides valid estimates under the missing-at-random assumption [60]. We reported *p*-values,

estimates (β), standard errors (SE), and 95% confidence intervals (CI; *stats*) for the treatment effect

– the level of response attributable to the treatment – estimated by the treatment:time interaction

term, in the corresponding **Tables**. Fixed effects were evaluated using analysis of variance

(ANOVA; *car* v3.1-3). When the treatment:time term met the significance threshold, we

summarized simple-effects contrasts of estimated marginal means (*emmeans* v1.11.0), including

between-treatment comparisons at endpoint with Benjamini-Hochberg (FDR) corrections.

Baseline between-treatment contrasts were null across analyses. For outcomes with non-normal

residuals or otherwise not meeting model assumptions, we used paired Wilcoxon signed-rank tests.

Aim 1. All measures were analyzed with mixed-effects models (*lme4* v1.1.37). Flanker task

measures (ACC, RT, and interference scores) were tested using treatment-by-time models with

congruent and incongruent trial types evaluated separately. Two flanker ACC and two P3

amplitude outcomes were winsorized; *CANTAB* delta values were compared using paired

Wilcoxon Signed-rank tests.

Aim 2. 16S analyses utilized rarefied counts for normalization. Beta diversity was assessed

by PERMANOVA (*vegan* v2.6.10; 5,000 permutations) using participant ID as the permutation

stratum, with homogeneity of dispersion (*vegan*, 5,000 permutations) tested in parallel.

Differential abundance was assessed with negative binomial mixed-effect models (*glmmTMB*

v1.1.11) comparing a null [*Taxa ~ (1|ID)*] and full [*Taxa ~ Treatment*Time + (1|ID)*] model by a

likelihood ratio test across all included taxa [87–89]. Alpha diversity and fecal metabolites were

analyzed with mixed-effects models (*lme4* v1.1.37). Gastrointestinal symptoms were compared

with a paired Wilcoxon Signed-rank test.

Aim 3. In vitro growth rates across treatments were modeled using a linear model (*lm*) with

treatment, timepoint (every 2h), and the interaction as fixed effects, followed by pairwise contrasts

[90]. Predicted KO and pathway differences were compared with *MaAsLin2* (v1.18.0) [91] and

ggpicrust2 (v2.0.0) [92]. Correlations between cognitive and microbial outcomes were assessed

using Spearman's rank correlation test (*stats* v4.4.3). Mediation analysis used baseline-adjusted

models to (1) predict the mediator from the Treatment [$\Delta Mediator \sim Treatment + Mediator_{Baseline}$]

and (2) predict the outcome from both the Treatment and the Mediator [$\Delta RT \sim Treatment *$

$\Delta Mediator + RT_{PRE}$]. Causal inference was estimated using nonparametric bootstrapping

(*mediation* v4.5.0; 8,000 simulations) with sensitivity analyses. Reported are the average causal

mediation effect (ACME), average direct effect (ADE), total effect, and proportion mediated.

Moderation models included treatment, a centered-scaled moderator (center="TRUE",

scale="TRUE"), and its interaction [$\Delta RT \sim Treatment * \Delta Moderator_{Z\text{-}Scaled}$]. Participant ID was

included as a random intercept, which caused model singularity; therefore, it was excluded.

Comparisons were probed using simple-slopes (*interactions* v1.2.0) and provided confidence

intervals (*emmeans* v1.11.0) at low (-1 SD), mean (0 SD), and high (+1 SD) changes of the

moderator.

Results

Participant characteristics

The CONSORT diagram (**Figure 1**) summarizes recruitment, enrollment, and retention.

Forty-two participants were randomized, of whom 41 completed the trial (**Table 1**). All

participants completed at least one cognitive testing visit, including one who withdrew after the

first intervention period ($n = 1$ CON). Per-protocol analyses included compliant participants who

consumed >80% of intervention sachets during at least one intervention period (Aim 1; $n = 38$

CON, 39 SCF). Ten participants had unusable P3 amplitude data ($n=3$ CON; 1 at pre, 2 at post

testing) and seven for the SCF ($n = 2$ PRE; $n = 5$ POST). Two participants could not provide fecal

samples but completed the primary cognitive aims (Aim 2; $n = 38$ CON, 37 SCF).

Aim 1: Cognitive outcomes

Executive function

Flanker task ACC remained high across the SCOPE study ($\geq 92\%$ scores), with ANOVA

showing no treatment effect (treatment:time) during congruent ($P = 0.42$) or incongruent trial

types ($P = 0.53$) (**Supplemental Figure 5**). RT exhibited a significant treatment effect during both

trials (congruent $P = 0.01$, incongruent $P = 0.01$). At endpoint, RT was faster following SCF in

both congruent ($\beta = -9.8$ ms, SE = 4.4, 95% CI: [-18.4, -1.2], FDR $P = 0.01$) and incongruent trials

($\beta = -14.2$ ms, SE = 4.4, 95% CI: [-22.8, -5.6], FDR $P = 0.003$) (**Figure 2**). P3 distributions were

visualized (**Supplemental Figure 6**), no treatment effects were observed for P3 amplitudes

(congruent $P = 0.46$; incongruent $P = 0.33$) or peak latency (congruent $P = 0.63$; incongruent P

$= 0.14$). And no treatment effects were observed for interference scores of ACC ($P = 0.10$) or RT

($P = 0.39$).

Mood and Stress

No treatment effects were observed for mood subscales and total mood disturbance scores

(*POMS*; all $P > 0.53$), relational memory performance (*iPosition*; all $P > 0.45$), affect scores

(*PANAS*; all $P > 0.45$), and stress (*PSS*-10; $P = 0.70$) (**Supplemental Table 1**). Similarly, no

treatment effects were observed across *CANTAB* measures (all $P > 0.13$) (**Supplemental Table**

2-5).

Aim 2: Gut Microbiota and Metabolite Outcomes

16S Fecal microbiota

Alpha diversity showed a treatment effect for observed richness (FDR $P = 0.03$), a

marginal effect for Faith's PD (FDR $P = 0.09$) and none for Shannon (FDR $P = 0.26$). Observed

richness was reduced following SCF at endpoint ($\beta = -29.9$ counts, SE = 8.55, 95% CI [-47.1, -

12.6], FDR $P = 0.01$) (**Table 5**; **Supplemental Figure 7**). Beta diversity showed a small but

statistically significant treatment effect measured with Bray-Curtis (F = 0.81, $R^2 = 0.005$, FDR P

< 0.001), unweighted UniFrac (F = 0.47, $R^2 = 0.003$, FDR $P = 0.002$), and weighted UniFrac (F

= 1.07, $R^2 = 0.007$, FDR $P < 0.001$) (**Supplemental Figure 8**). Endpoint comparisons with each

metric showed differences between SCF and CON for Bray-Curtis (F = 1.39, $R^2 = 0.019$, FDR P

< 0.001), unweighted UniFrac (F = 0.64, $R^2 = 0.010$, FDR $P = 0.005$), and weighted UniFrac (F

= 1.69, $R^2 = 0.023$, FDR $P < 0.001$).

At the phylum level, none met the significance threshold (**Supplemental Table 6**). A total

of 260 genera were detected, 128 passed filtering criteria ($\geq$40% prevalence, >10 reads), and 14

met the significance threshold (all FDR $P < 0.05$); 11 were classified as low-abundance (<0.5%)

or as a group of bacteria without taxonomical assignment (incertae_sedis) (**Supplemental Table**

7). *Parabacteroides* enrichment was consistently observed across 90% of participants during the

SCF arm (**Figure 3**), demonstrating a treatment effect ($\chi^2 = 42.0$, FDR $P < 0.001$)**,** with a ~4.1-

fold increase following SCF compared to CON ($\beta = 1.44$ log, SE = 0.16, 95% CI [1.01, 1.88], FDR

P < 0.001). *Fusicatenibacter* also showed a treatment effect (χ^2 = 17.94, FDR *P* < 0.001) and

~2.7-fold increase following SCF (β = 1.10 log, SE: 0.60, 95% CI: [0.59, 1.61], FDR *P* < 0.001).

Similarly, *Dialister* had a treatment effect (χ^2 = 7.25, FDR *P* = 0.02) and increased by ~2.4-fold

following SCF (β = 1.06 log, SE: 0.39, 95% CI: [0.29, 1.83], FDR *P* = 0.03) (**Supplemental**

Figure 9).

Targeted metabolites

Fecal SCFAs (**Supplemental Figure 10**) and BCFAs did not statistically differ between

treatments (all *P* > 0.14; **Table 4**). Pooled total indoles (indole, 2-methylindole, 3-methylindole,

7-methylindole), indole, pooled total phenols (4-methylphenol, 4-ethylphenol, phenol), 4-

methylindole, and ammonia likewise did not show no treatment effect (all *P* > 0.32; **Supplemental**

Table 8).

Gastrointestinal symptoms

Symptom severity scores were generally absent to mild across conditions. No differences

were observed between SCF and CON for changes in cramping, nausea, reflux, rumbling, or

burping (all *P* > 0.17; **Table 5**). Flatulence (gas) severity scores increased following SCF (*P* =

0.01), although median severity remained <2 (mild) by endpoint (1.93 score). Bloating was

marginally increased following SCF (*P* = 0.05), though median scores remained 1.00 (absent). No

treatment effect was observed for self-reported feelings of consistency, ease of passage, bowel

frequency, or in the measured fecal pH (all *P* > 0.22; **Supplemental Table 9**).

Aim 3: Exploratory Integrative and Functional Analyses

Aim 3 analyses were restricted to primary cognitive outcomes that differed by treatment

(flanker RTs) and on the microbiota/metabolite features prioritized from Aim 2 based on either

evidence of SCF-responsiveness (*Parabacteroides*, *Fusicatenibacter*, and *Dialister*) or a

consistent directional trend with biological plausibility (acetate, propionate).

Correlation analyses

Spearman correlations at the SCF endpoint showed marginal inverse associations between

congruent RT and acetate (ρ = -0.33, FDR P = 0.09) and propionate (ρ = -0.36, FDR P = 0.09).

Parabacteroides was marginally associated with acetate (ρ = 0.34, FDR P = 0.09). No taxa were

associated with congruent RT (all FDR P > 0.14). For incongruent trials, no significant

correlations were detected with any candidate (all ρ < 0.23 and FDR P > 0.30) **(Supplementary**

Figure 11).

Mediation analyses

The proportion mediated was estimated at ≤10.0% across candidates for congruent trials

and <4.1% for incongruent trials (**Table 6**). However, across candidates, indirect effects (ACME)

were not detected for either congruency (all $p \geq 0.19$), indicating no evidence of mediation.

Moderation analyses

Moderation models provided no evidence that candidate changes modified the treatment

effect on RT (all P > 0.49), except for *Parabacteroides*, which showed a moderation effect for

congruent (P = 0.05) and incongruent RTs (P = 0.04). Adding *Parabacteroides* increased

explained variance from ΔR^2 = 0.07 (base model) to ΔR^2 = 0.12 (adjusted), whereas other

candidates contributed <1% (**Table 7**).

For congruent trials, SCF-associated RT improvement depended on *Parabacteroides*

change (z-scored; 0 SD = average change): RT was faster when *Parabacteroides* changes were

small (-1SD: β = -29.5 ms, SE = 8.76, t = -3.37, 95% CI: [-47.0, -12.03], P < 0.001), but not when

Parabacteroides change was near the mean (0 SD; β = -9.7 ms, SE = 8.27, t = -1.18, 95% CI: [-

26.2, 6.8], P = 0.24) or much larger (+1 SD: β = 10.0 ms, SE = 16.27, t = 0.62, 95% CI: [-22.5,

42.5], P = 0.54). For incongruent trials, the treatment effect on RT similarly depended on

Parabacteroides change. RT was faster when changes were small (β = -32.4 ms, SE = 10.59, t = -

3.06, 95% CI: [-53.5, -11.2], P < 0.001), but not when change was near the mean (β = -7.30 ms,

SE = 10.0, t = -0.73, 95% CI: [-27.3, 12.7], P = 0.47) or much larger (β = 17.8 ms, SE = 19.68, t

= 0.90, 95% CI: [-21.5, 57.0], P = 0.37) (**Figure 4**).

In vitro culturing of P. distasonis

P. distasonis was confirmed to grow with SCF in isolation displaying a distinct growth

kinetics with significant media effect (P < 0.001), time (P < 0.001), and interaction (P < 0.001).

Growth on glucose and CON showed a short lag phase (<1h), and a rapid log phase reaching

stationary phase within 5-6h (glucose: OD_{600} = 0.59, 5.5h; CON: OD_{600} = 0.54, 5h). Growth on

SCF exhibited a two-phase diauxic growth pattern: a slower and smaller initial phase reaching

OD_{600} = 0.31 at 6h, followed by a ~2h lag, then a second log phase that peaked at OD_{600} = 0.61 by

14h. *Post-hoc* comparisons showed that SCF had significantly lower growth rates at 2h (all FDR

P < 0.001) and 4h (vs. glucose, FDR P = 0.03; vs. CON, FDR P = 0.01) relative to controls. By

6h, no differences were detected across treatments. At 8-10h, SCF had higher growth rates

compared to controls (all FDR P < 0.001), and remained higher at 12h (vs. glucose, FDR P =

0.002). By 14h, growth stabilized with no differences detected between treatments (**Figure 5**).

16S-Predicted functional potential analyses

Predicted microbial pathway profiles from the full cohort indicated higher *N*-glycan biosynthesis and α-linolenic acid metabolism pathways following SCF at endpoint compared to control (all FDR $P < 0.01$). Predicted enzyme profiles (ECs) identified 23 functions differing by treatment at endpoint (all FDR $P < 0.05$) (**Supplemental Figure 12**). MaAsLin2 detected 2,701 carbohydrate-metabolism KOs associated with SCF at endpoint (FDR $P < 0.05$). Among the top SCF-associated KOs, 18 showed contribution by *Parabacteroides* (including glycan degradation and complex carbohydrate utilization) and two by *Fusicatenibacter* (**Supplemental Figure 13**).

Responder shotgun metagenomic analyses

Metagenomic profile comparisons between responder vs. non-responded indicated no differences in alpha diversity (all $P > 0.09$) or beta diversity. Responder status explained ~17% of variance using Bray-Curtis PERMANOVA ($R^2 = 0.167$, $P = 0.05$), with no main effect of time ($R^2 = 0.083$, $P = 0.44$) and no difference in the interaction ($R^2 = 0.079$, $P = 0.49$) (**Figure 6**). Metagenomic genus-level profiles focused on the treatment-responsive genera identified in 16S analyses and confirmed that *Parabacteroides* was enriched in responders ($\Delta = 13.6\%$) and slightly reduced in non-responders ($\Delta = -0.4\%$), consistent with 16S results. At the species level, *P. distasonis* increased in responders ($\Delta = 10.9\%$) and remained non-detectable in non-responders, while *P. merdae* also increased in responders ($\Delta = 2.7\%$) and decreased in non-responders ($\Delta = -0.4\%$). *F. saccharivorans* negligibly increased in responders ($\Delta = 0.1\%$), compared to a larger increase in non-responders ($\Delta = 2.2\%$). *D. hominis* ($\Delta = 1.4\%$) and *D. invisus* (0.3%) increased in non-responder, while only *D. invisus* was detected and decreased in responders ($\Delta = -0.3\%$).

Functional analyses with *HUMAnN* showed an increase in carbohydrate metabolism-

related KOs among responders (UDP-glucose 6-dehydrogenase, α-amylases, hexokinase), while

non-responders demonstrated a lack of presence in these KOs (**Supplemental Figure 14**). Genus-

(**Supplemental Figure 15)** and species-level time comparisons (**Figure 10**) in responders

indicated increased contributions from *P. distasonis* to multiple carbohydrate-active gene families,

including glycoside hydrolases (GHs), α-amylases, and hexokinase. CAZy annotations provided

further resolution of GH families and subfamilies between groups. Of 228 GH families detected,

those with positive log fold changes in responders were isolated (**Supplemental Figure 16**).

Several GH families were enriched (GH43_8, GH43_33, GH25, GH24, GH179, GH13_9,

GH130_2, GH12) and positively correlated with *Parabacteroides* abundances (**Supplemental**

Figure 17).

Responders demonstrated faster RT across the SCF intervention for both congruent (Δ = -

0.90ms) and incongruent (Δ = 2.7ms) trials. In contrast, non-responders exhibited slower baseline

RT but demonstrated larger SCF-induced improvements (congruent Δ = -32.1ms, incongruent Δ

= -31.8ms). Parallel changes were observed in SCFAs; responders had higher baseline SCFA

levels and was increased in both groups for acetate (responders Δ = 66.4 vs. non-responders Δ =

71.6 µmol/g DMB) and propionate (responders Δ = 20.9 vs. non-responders Δ = 28.6 µmol/g

µmol/g DMB). (**Table 8**).

Discussion

In this randomized, double-blind, placebo-controlled, crossover trial, consuming SCF daily

for 4-weeks improved attentional inhibition, reflected by faster RT without compromising

accuracy on both congruent and incongruent flanker task trial types compared to control. Effects

were domain-specific to attentional inhibition, with no changes observed in relational memory,

rule acquisition and reversal, set shifting, sustained attention, response inhibition, or spatial

working memory observed (*Aim 1*). SCF consumption also shifted fecal microbial diversity,

selectively enriched *Parabacteroides*, numerically increased fecal acetate concentrations, and

produced absent-to-mild GI symptoms (*Aim 2*). SCF consumption had no effect on P3 amplitude

or latency, mood, or stress. Exploratory analyses did not identify any mediators but moderation

analyses suggested that SCF-related RT improvements varied by the magnitude of

Parabacteroides change. Consistent with 16S results identifying *Parabacteroides* as SCF-

responsive at the genus level, shotgun metagenomics identified *P. distasonis* as the primary SCF-

responsive species, while *in vitro* culturing independently confirmed its capacity to metabolize

SCF independent of cross-feeding (*Aim 3*).

Attentional inhibition is a core domain of cognitive control that can naturally and

insidiously decline with aging [31], underscoring the need for dietary strategies to preserve healthy

cognitive aging. Few clinical trials have assessed the impact of dietary fiber consumption on

cognition while integrating microbiome outcomes. The present work demonstrated that SCF

consumption selectively supported attentional inhibition, reflected by faster RT in congruent and

incongruent flanker trials, with no changes in other cognitive outcomes. This domain-specific

pattern reinforces the concept that gut-brain communication can influence higher-order cognitive

processes. Another randomized trial among 84 adults with overweight and obesity reported

improved accuracy, but no change in RT, during a modified flanker task following 12 weeks of

daily avocado consumption (12.8-16 grams fiber) relative to a control providing 3.2 – 4 grams

fiber daily [93]. Notably, avocado provides a mixed plant cell-wall fiber profile that is predominantly

insoluble (cellulose/hemicellulose) with soluble pectin (along with other nutrients) [94], whereas

SCF is a digestion-resistant maltodextrin dextrinized to produce a mixture of α-linkages (α-1,2, -

1,3, -1,4, and -1,6) [18,19], which may yield distinct fermentation kinetics and downstream signaling.

Specific to the microbiota, the avocado intervention increased α-diversity and

enriched *Faecalibacterium, Lachnospira*, and *Alistipes* and resulted in 18% greater fecal acetate,

concentrations due to fiber-rich avocado consumption [95]. Another nutrition trial reported mood

benefits with 16 g/day inulin for 12 weeks versus 16 g maltodextrin among adults with obesity,

particularly among individuals exhibiting a specific microbial signature, however, no effects were

observed on inhibitory control, working memory, or cognitive flexibility regardless of changes in

the gut microbiota [39]. Inulin is a $\beta(2,1)$-linked fructan (fructose polymer), whereas SCF is glucose

based [96,97]. Other lifestyle interventions, such as increased physical activity, have been shown to

improve flanker RT after a single exercise session, with one randomized trial among 106 healthy-

weight adults showing improvements in both congruent (458 ms vs 475 ms) and incongruent (528

ms vs 547 ms) paradigms immediately following 30 min of high-intensity aerobic exercise

compared with seated rest [98]. Collectively, these studies suggest that cognition effects are domain-

and context-specific, shaped by factors such as fiber type, population characteristics, physical

activity status, and intervention duration, which complicates direct cross-study comparisons but

underscores nutrition as a potential modifiable factor for cognitive health. However, the fiber type

(e.g., SCF, pectin), host and microbiome characteristics, and exposure duration needed to exact

cognitive benefits is yet to be delineated, highlighting the need for additional robust clinical trials.

Our work demonstrated that *Parabacteroides* was enriched following SCF consumption,

which is consistent with previous SCF supplementation trials. Specifically, a randomized dose-

responsive trial with free-living pubertal females (0, 10, and 20g fiber/d, 4 weeks per dose), a

randomized double-blind trial with healthy adults (20 g/d; 10 days per period), and a randomized

double-blind trial with only healthy adult men (21 g/d; 21 days per period) all reported enrichment

of *Parabacteroides* [21,99,100]. In the present study, SCF consumption reduced observed richness at

endpoint, which has been previously reported with fermentable fibers when selective utilization

leads to an increase of a narrower set of taxa and a more dominated community structure, without

necessarily indicating an adverse effect [101] as shown by its consistent role as a fermentable and

generally well tolerated fiber [97].

The convergence of 16S with the full cohort and shotgun metagenomes on responder subset

supports the robustness of the *Parabacteroides* enrichment across sequencing approaches. In

responders, *P. distasonis* abundance was inversely related to *Bacteroides vulgatus*, suggesting

possible ecological competition under SCF exposure. Predicted functional profiles and shotgun

metagenomics indicated increased carbohydrate metabolism pathways and glycan degradation

capacity, with CAZy highlighting an increased in *Parabacteroides*-associated GHs (GH35, GH43,

and GH73) [102,103]. *P. distasonis*-specific GHs (GH144 and GH13) involved in the hydrolysis α-

glycosidic bonds were also detected [102,103]. CAZymes serving similar functions are typically

organized within PULs, gene clusters that coordinate sensing, import, and degradation of dietary

glycans [104]. The enrichment of these CAZymes likely reflects an activation of SCF-targeting PULs,

enabling *P. distasonis* to maintain a competitive niche in the large intestine. Our *in vitro*

experiment supported this metabolic capacity, showing that *P. distasonis*, with no prior SCF

exposure, fermented SCF in a diauxic pattern, consistent with the mixture of fiber and non-fiber

starches in the formula. Together, these findings position *Parabacteroides* as a primary degrader

of SCF, although mechanistic insights remain limited.

These findings are consistent with the ISAPP definition of a prebiotic: "a substrate that is

selectively utilized by host microorganisms, conferring a health benefit" [105]. SCF induced small

shift in the GI microbiota, selectively increasing *Parabacteroides*, and provided a cognitive benefit

in healthy older adults. *Parabacteroides* responded to SCF in 90% of the healthy cohort, consistent

with the 'Health-Associated Core Keystone' (HACK) index which identified *P. distasonis, P.*

merdae, and *F. saccharivorans* as core and stability-associated taxa [106]. Prior work links *P.*

distasonis with anti-inflammatory effects, gut barrier protection [107–109], and bioactive metabolite

production (e.g., GABA)[110], relevant to gut-brain communication and health outcomes [102,111,112].

Although *Parabacteroides* did not directly mediate cognitive benefits in the SCOPE study, its

consistent response to SCF suggests its potential role as a keystone taxon in healthy older adults.

In this study, fecal SCFA concentration differences did not reach statistical significance;

however, their directional changes align with prior dietary fiber interventions and remain plausible

candidates for gut-brain signaling, given their ability to cross and modulate the blood-brain

barrier's structural integrity [34,35,113–116]. Notably, fecal SCFA concentrations in the present study

were lower than those reported in a younger, male-only SCF trial using a higher dose (21g/d) [19],

which may reflect differences in participant characteristics such transit time, absorption, and/or

dilution with greater stool mass. Prior human studies show that colonic SCFA delivery influences

anticipatory brain reward responses and attenuates cortisol reactivity to psychosocial stress [35].

Despite biological plausibility, in the present study, mediation and moderation models did not

support fecal SCFAs as either mediators or moderators of the SCF effect.

Moderation analyses suggested that the SCF effect on RT varied with the magnitude of

change in *Parabacteroides*. RT improvements were greatest at smaller *Parabacteroides* changes

(-1SD), whereas larger *Parabacteroides* changes were not associated with faster RTs.

Furthermore, responders consistently performed quicker at baseline and endpoints, while non-

responders exhibited larger improvements from SCF for RT speeds. These exploratory findings

suggest that *Parabacteroides* may exert a context-dependent influence, but its enrichment may not

be linearly beneficial for cognitive performance.

Secondary taxa provided additional exploratory context for the SCF-RT association.

Although mediation models did not support Dialister or *Fusicatenibacter* as intermediaries of the

SCF effect on RT, their descriptive patterns were informative. Bacillota members such as

Fusicatenibacter are capable of cross-feeding on substrates released by members of the

Bacteroidota phylum [117] (e.g., *Parabacteroides*), producing beneficial metabolites including

acetate, lactate, formate, and succinate [118]. Potentially, *Fusicatenibacter* may have had the role of

the main secondary consumer working with *Parabacteroides* in response to SCF. Responders had

higher baseline levels of *F. saccharivorans* which showed negligible enrichment with SCF,

whereas non-responders exhibited a relatively larger enrichment in *F. saccharivorans*. In contrast,

D. invisus and *D. hominis* were enriched only in non-responders and remained below detectable

levels in responders. A prior study linked *Fusicatenibacter saccharivorans* with executive and

visuospatial functions with community-dwelling older adults [119], whereas *Dialister invisus* was

reported to be more abundant in older adults with mild cognitive impairment [119]. Together, these

patterns suggest that baseline community composition and differential responsiveness may

contribute to cognitive variability, though mechanistic inferences remain preliminary.

The SCOPE study has several limitations. First, the cohort was predominantly white,

female, and highly educated, which may restrict generalizability. Mediation and moderation

analysis were exploratory and did not fully account for the crossover design in its statistical

framework and utilized change and baseline adjustments. Shotgun metagenomic analyses were

limited to a small subset of responders and non-responders, reducing power to detect subtle effects.

The absence of blood samples prevented assessment of circulating SCFAs or other metabolites,

while the targeted fecal metabolomics panel may have overlooked additional relevant compounds

(e.g., tryptophan catabolites). Furthermore, fecal SCFAs can be influenced by colonic transit time

and may only reflect a fraction *in vivo* production. Systemic inflammatory markers and neuroactive

metabolites were not assessed and should be incorporated in future mechanistic trials. Although

participants agreed to maintain their habitual diets, inter-individual dietary variation may have

introduced unmeasured noise into microbiome and metabolite outcomes. Despite these limitations,

the SCOPE study provides novel evidence linking SCF to selective cognitive benefits, supported

by exploratory insights into microbial and SCFA response.

In conclusion, daily consumption of SCF for four weeks improved RT in middle aged and

older adults and induced a microbiota shift dominated by *Parabacteroides*. Exploratory

moderation suggested a threshold effect, with RT benefits most evident at smaller changes in

Parabacteroides. Notably, cognitive improvements were also observed in non-responders with

negligible *Parabacteroides*, indicating that cognitive benefits are unlikely to hinge on a single

taxon but may instead depend on the ecological context and balance between health associated

keystone microbes (e.g., *Parabacteroides* and *Fusicatenibacter*) for contributing complementary

routes to induce GI-brain interactions [120]. Host factors may also contribute to interindividual

variability in cognitive response to SCF. These findings position SCF as a dietary fiber with

potential cognitive relevance, while underscoring the need for multi-omics, untargeted

metabolomics, dosing trials, and systemic biomarker assessments to refine mechanistic models

and advance precision nutrition strategies for cognitive health.

Acknowledgments

We thank Dr. Michael J. Miller for his expertise, equipment, and facility to conduct the *in vitro*

culturing experiment. We also thank Dr. Pratik Banerjee and his graduate students (Papri Suraya

Rahman and Edwin Valenzuela De Leon) for providing the *P. distasonis* strain and assisting with

the equipment set-up. We are also grateful to the HDH graduate students: Alexis Baldeon, Marahi

Perez-Tamayo, Nadine Veasly, and Maria Sanabria-Veaz; HDH staff: Eva Cornman, Myra Esmail,

Karen Gutierrez, and Tehila Abdiel; and NAK graduate students Shelby Keyes and John Kim, as

well as the undergraduate research assistants from HDH and NAK laboratories, for their assistance

during the appointments, sample processing, and data management.

Author Contributions

HDH, NAK, DAA and TAH: designed the research; MDB and TM: recruited and enrolled

participants and coordinated study procedures; DAA: conducted the *in vitro* experiment; DAA,

TAH, SM, TM and RS: conducted the experiments and testing appointments; MA, DR and DAA:

managed biological samples collection and storages; DAA and TAH: managed study data; DAA

and NS: prepared data for bioinformatic pipelines; DAA and TAH: analyzed the data: DAA, TAH,

NAK and HDH: wrote the manuscript; NAK and HDH: had primary responsibility for final

content. All authors read and approved the final manuscript.

Author Disclosures: The authors report no conflict of interest.

Funding: This work was supported by Tate & Lyle Ingredients Americas LLC (to NAK) and the

USDA NIFA, Hatch Project (1026591, to HDH).

Data Availability: Data described in the manuscript, code book, and analytic code will be made

available upon request.

References

1. Morris MC, Wang Y, Barnes LL, Bennett DA, Dawson-Hughes B, Booth SL. Nutrients and bioactives in green leafy vegetables and cognitive decline: Prospective study. *Neurology*. 2018;90(3):E214-E222. doi:10.1212/WNL.0000000000004815
2. Perim Baldo M, Aboussaleh Y, Turrini A, et al. Adherence to a Mediterranean-Style Diet and effects on Cognition in Adults: A Qualitative evaluation and Systematic Review of Longitudinal and Prospective Trials. 2016;3:1. doi:10.3389/fnut.2016.00022
3. Khan NA, Raine LB, Donovan SM, Hillman CH. IV. The cognitive implications of obesity and nutrition in childhood. *Monogr Soc Res Child Dev*. 2014;79(4):51-71. doi:10.1111/mono.12130
4. Cho SS, Qi L, Fahey GC, Klurfeld DM. Consumption of cereal fiber, mixtures of whole grains and bran, and whole grains and risk reduction in type 2 diabetes, obesity, and cardiovascular disease. *American Journal of Clinical Nutrition*. 2013;98(2):594-619. doi:10.3945/ajcn.113.067629
5. Trock B, Lanza E, Greenwald P, Lanza E, Greenwald P. *REVIEW Dietary Fiber, Vegetables, and Colon Cancer: Critical Review and Meta-Analyses of the Epidemiologic Evidence*. https://academic.oup.com/jnci/article/82/8/650/882222
6. Gov D. *Dietary Guidelines for Americans Make Every Bite Count With the Dietary Guidelines*. https://www.
7. Marchesi JR, Ravel J. The vocabulary of microbiome research: a proposal. *Microbiome*. 2015;3(1). doi:10.1186/s40168-015-0094-5
8. Fleming SA, Monaikul S, Patsavas AJ, Waworuntu R V., Berg BM, Dilger RN. Dietary polydextrose and galactooligosaccharide increase exploratory behavior, improve recognition memory, and alter neurochemistry in the young pig. *Nutr Neurosci*. 2019;22(7):499-512. doi:10.1080/1028415X.2017.1415280
9. Gronier B, Savignac HM, Miceli M Di, et al. Increased cortical neuronal responses to NMDA and improved attentional set-shifting performance in rats following prebiotic (B-GOS s) ingestion. doi:10.1016/j.euroneuro.2017.11.001
10. Khan NA, Raine LB, Drollette ES, Scudder MR, Kramer AF, Hillman CH. Dietary Fiber Is Positively Associated with Cognitive Control among Prepubertal Children. *J Nutr*. 2015;145(1):143-149. doi:10.3945/JN.114.198457
11. Hassevoort KM, Lin AS, Khan NA, Hillman CH, Cohen NJ. Added sugar and dietary fiber consumption are associated with creativity in preadolescent children. *Nutr Neurosci*. 2020;23(10):791-802. doi:10.1080/1028415X.2018.1558003
12. Naveed S, Venäläinen T, Eloranta AM, et al. Associations of dietary carbohydrate and fatty acid intakes with cognition among children. *Public Health Nutr*. 2020;23(9):1657-1663. doi:10.1017/S1368980019003860
13. Cryan JF, O KJ, M Cowan CS, et al. The Microbiota-Gut-Brain Axis. *Physiol Rev*. 2019;99:1877-2013. doi:10.1152/physrev.00018.2018.-The
14. Stewart ML, Nikhanj SD, Timm DA, Thomas W, Slavin JL. Evaluation of the effect of four fibers on laxation, gastrointestinal tolerance and serum markers in healthy humans. *Ann Nutr Metab*. 2010;56(2):91-98. doi:10.1159/000275962
15. Shuan W, Tan K, Fen P, et al. The Role of Soluble Corn Fiber on Glycemic and Insulin Response. doi:10.3390/nu12040961

804   16.   Jakeman SA, Henry CN, Martin BR, et al. Soluble corn fiber increases bone calcium
805   retention in postmenopausal women in a dose-dependent manner: A randomized crossover
806   trial. *American Journal of Clinical Nutrition*. 2016;104(3):837-843.
807   doi:10.3945/ajcn.116.132761
808   17.   Klosterbuer AS, Hullar MAJ, Li F, et al. Gastrointestinal effects of resistant starch,
809   soluble maize fibre and pullulan in healthy adults. doi:10.1017/S0007114513000019
810   18.   LinAmy AHM, Nichols BL, Quezada-Calvillo R, et al. Unexpected high digestion rate of
811   cooked starch by the Ct-maltase-glucoamylase small intestine mucosal α-glucosidase
812   subunit. *PLoS One*. 2012;7(5). doi:10.1371/journal.pone.0035473
813   19.   Vester Boler BM, Rossoni Serao MC, Bauer LL, et al. Digestive physiological outcomes
814   related to polydextrose and soluble maize fibre consumption by healthy adult men. *British*
815   *Journal of Nutrition*. 2011;106(12):1864-1871. doi:10.1017/S0007114511002388
816   20.   Maathuis A, Venema K, Hoffman A, Evans A, Sanders L. The effect of the undigested
817   fraction of maize products on the activity and composition of the microbiota determined in
818   a dynamic in vitro model of the human proximal large intestine. *J Am Coll Nutr*.
819   2009;28(6):657-666. doi:10.1080/07315724.2009.10719798
820   21.   Whisner CM, Martin BR, Nakatsu CH, et al. Soluble corn fiber increases calcium
821   absorption associated with shifts in the gut microbiome: A randomized dose-response trial
822   in free-living pubertal females. *Journal of Nutrition*. 2016;146(7):1298-1306.
823   doi:10.3945/jn.115.227256
824   22.   Arroyo MC, Laurie I, Rotsaert C, Marzorati M, Risso D, Karnik K. Age-Dependent
825   Prebiotic Effects of Soluble Corn Fiber in M-SHIME® Gut Microbial Ecosystems. *Plant*
826   *Foods for Human Nutrition*. 2023;78(1):213-220. doi:10.1007/s11130-023-01043-z
827   23.   Holscher HD, Gregory Caporaso J, Hooda S, Brulc JM, Fahey GC, Swanson KS. Fiber
828   supplementation influences phylogenetic structure and functional capacity of the human
829   intestinal microbiome: Follow-up of a randomized controlled trial. *American Journal of*
830   *Clinical Nutrition*. 2015;101(1):55-64. doi:10.3945/ajcn.114.092064
831   24.   Brown AJ, Goldsworthy SM, Barnes AA, et al. The orphan G protein-coupled receptors
832   GPR41 and GPR43 are activated by propionate and other short chain carboxylic acids.
833   *Journal of Biological Chemistry*. 2003;278(13):11312-11319.
834   doi:10.1074/jbc.M211609200
835   25.   Cani PD, Lecourt E, Dewulf EM, et al. Gut microbiota fermentation of prebiotics
836   increases satietogenic and incretin gut peptide production with consequences for appetite
837   sensation and glucose response after a meal. *American Journal of Clinical Nutrition*.
838   2009;90(5):1236-1243. doi:10.3945/ajcn.2009.28095
839   26.   Corrêa-Oliveira R, Fachi JL, Vieira A, Sato FT, Vinolo MAR. Regulation of immune cell
840   function by short-chain fatty acids. *Clin Transl Immunology.John Wiley and Sons Inc.*
841   2016;5(4). doi:10.1038/cti.2016.17
842   27.   Dalile B, Van Oudenhove L, Vervliet B, Verbeke K. The role of short-chain fatty acids in
843   microbiota–gut–brain communication. *Nat Rev Gastroenterol Hepatol.Nature Publishing*
844   *Group*. 2019;16(8):461-478. doi:10.1038/s41575-019-0157-3
845   28.   Zhou SY, Guo ZN, Yang Y, Qu Y, Jin H. Gut-brain axis: Mechanisms and potential
846   therapeutic strategies for ischemic stroke through immune functions. *Front*
847   *Neurosci.Frontiers Media S.A.* 2023;17. doi:10.3389/fnins.2023.1081347

29.   Silva YP, Bernardi A, Frozza RL. The Role of Short-Chain Fatty Acids From Gut
Microbiota in Gut-Brain Communication. *Front Endocrinol (Lausanne).Frontiers Media*
S.A. 2020;11. doi:10.3389/fendo.2020.00025

30.   Mansuy-Aubert V, Ravussin Y. Short chain fatty acids: the messengers from down below.
Front Neurosci. 2023;17. doi:10.3389/fnins.2023.1197759

31.   Diamond A. Executive functions. *Annu Rev Psychol.Annual Reviews Inc.* 2013;64:135-
168. doi:10.1146/annurev-psych-113011-143750

32.   Duncan-Johnson CC. P300 latency: A new metric of information processing.
Psychophysiology. 1981;18(3):207-215. doi:10.1111/j.1469-8986.1981.tb03020.x

33.   Polich J. Updating P300: An integrative theory of P3a and P3b. *Clinical Neurophysiology*.
2007;118(10):2128-2148. doi:10.1016/j.clinph.2007.04.019

34.   La Torre D, Verbeke K, Dalile B. Dietary fibre and the gut–brain axis: microbiota-
dependent and independent mechanisms of action. *Gut Microbiome*. 2021;2.
doi:10.1017/gmb.2021.3

35.   Dalile B, Fuchs A, La Torre D, Vervliet B, Van Oudenhove L, Verbeke K. Colonic
butyrate administration modulates fear memory but not the acute stress response in men:
A randomized, triple-blind, placebo-controlled trial. *Prog Neuropsychopharmacol Biol*
Psychiatry. 2024;131. doi:10.1016/j.pnpbp.2024.110939

36.   Cheng J, Hu H, Ju Y, et al. Gut microbiota-derived short-chain fatty acids and depression:
deep insight into biological mechanisms and potential applications. *Gen Psychiatr.BMJ*
Publishing Group. 2024;37(1). doi:10.1136/gpsych-2023-101374

37.   Luqman A, He M, Hassan A, et al. Mood and microbes: a comprehensive review of
intestinal microbiota's impact on depression. *Front Psychiatry.Frontiers Media SA*.
2024;15. doi:10.3389/fpsyt.2024.1295766

38.   Haskell CF, Scholey AB, Jackson PA, et al. Cognitive and mood effects in healthy
children during 12 weeks' supplementation with multi-vitamin/minerals. *British Journal*
of Nutrition. 2008;100(5):1086-1096. doi:10.1017/S0007114508959213

39.   Leyrolle Q, Cserjesi R, D.G.H. Mulders M, et al. Prebiotic effect on mood in obese
patients is determined by the initial gut microbiota composition: A randomized, controlled
trial. *Brain Behav Immun*. 2021;94:289-298. doi:10.1016/j.bbi.2021.01.014

40.   Godin-Leisure-Time-Exercise-Questionnaire_070815.

41.   Edwards CG, Walk AM, Cannavale CN, et al. Dietary choline is related to neural
efficiency during a selective attention task among middle-aged adults with overweight and
obesity. *https://doi.org/101080/1028415X20191623456*. 2019;24(4):269-278.
doi:10.1080/1028415X.2019.1623456

42.   Chatrian GE, Lettich E, Nelson PL. American Journal of EEG Technology Ten Percent
Electrode System for Topographic Studies of Spontaneous and Evoked EEG Activities
Ten Percent Electrode System for Topographic Studies of Spontaneous and Evoked EEG
Activities. Published online 2015. doi:10.1080/00029238.1985.11080163

43.   Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *J Neurosci Methods*. 2004;134(1):9-
21. doi:10.1016/J.JNEUMETH.2003.10.009

44.   Lopez-Calderon J, Luck SJ. ERPLAB: An open-source toolbox for the analysis of event-
related potentials. *Front Hum Neurosci*. 2014;8(1 APR):213.
doi:10.3389/FNHUM.2014.00213/BIBTEX

45.   Kamijo K, Pontifex MB, Khan NA, et al. The association of childhood obesity to
894   neuroelectric indices of inhibition. *Psychophysiology*. 2012;49(10):1361-1371.
895   doi:10.1111/J.1469-8986.2012.01459.X

896   46.   Walk AM, Edwards CG, Baumgartner NW, et al. The role of retinal carotenoids and age
897   on neuroelectric indices of attentional control among early to middle-aged adults. *Front*
898   *Aging Neurosci*. 2017;9(JUN):183. doi:10.3389/FNAGI.2017.00183/BIBTEX

899   47.   Luck SJ, Gaspelin N. How to get statistically significant effects in any ERP experiment
900   (and why you shouldn't). *Psychophysiology*. 2017;54(1):146-157.
901   doi:10.1111/PSYP.12639

902   48.   Pontifex MB. Stylized Topographic Map Plugin for EEGLAB/ERPLAB.

903   49.   Horecka KM, Dulas MR, Schwarb H, Lucas HD, Duff M, Cohen NJ. Reconstructing
904   relational information. *Hippocampus*. 2018;28(2):164-177. doi:10.1002/hipo.22819

905   50.   Langley C, Sahakian B, Robbins T. Cambridge neuropsychological test automated battery
906   (cantab). *SAGE Publications Ltd*. 2023;0:435-468.

907   51.   Wild K V., Musser ED. The cambridge neuropsychological test automated battery in the
908   assessment of executive functioning. In: *Handbook of Executive Functioning*. Springer
909   New York; 2014:171-190. doi:10.1007/978-1-4614-8106-5_11

910   52.   Stephen J. Gibson. The Measurement of Mood States in Older Adults. *The Journals of*
911   *Gerontology: Series B*. 1997;52B(4):167-174.

912   53.   Boyle GJ, Helmes E, Matthews G, Izard CE. Measures of Affect Dimensions. In:
913   *Measures of Personality and Social Psychological Constructs*. Elsevier Inc.; 2015:190-
914   224. doi:10.1016/B978-0-12-386915-9.00008-5

915   54.   Humboldt S von, Monteiro A, Leal I. Validation of the PANAS: A Measure of Positive
916   and Negative Affect for Use with Cross-National Older Adults. *Rev Eur Stud*.
917   2017;9(2):10. doi:10.5539/res.v9n2p10

918   55.   Lee EH. Review of the psychometric evidence of the perceived stress scale. *Asian Nurs*
919   *Res (Korean Soc Nurs Sci)*. 2012;6(4):121-127. doi:10.1016/j.anr.2012.08.004

920   56.   Walters W, Hyde ER, Berg-Lyons D, et al. Improved Bacterial 16S rRNA Gene (V4 and
921   V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial
922   Community Surveys. *mSystems*. 2016;1(1). doi:10.1128/msystems.00009-15

923   57.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
924   High-resolution sample inference from Illumina amplicon data. *Nat Methods*.
925   2016;13(7):581-583. doi:10.1038/nmeth.3869

926   58.   Bolyen E, Rideout JR, Dillon MR, et al. Reproducible, interactive, scalable and extensible
927   microbiome data science using QIIME 2. *Nat Biotechnol.Nature Publishing Group*.
928   2019;37(8):852-857. doi:10.1038/s41587-019-0209-9

929   59.   Yilmaz P, Parfrey LW, Yarza P, et al. The SILVA and "all-species Living Tree Project
930   (LTP)" taxonomic frameworks. *Nucleic Acids Res*. 2014;42(D1). doi:10.1093/nar/gkt1209

931   60.   Morton JT, Marotz C, Washburne A, et al. Establishing microbial composition
932   measurement standards with reference frames. *Nat Commun*. 2019;10(1).
933   doi:10.1038/s41467-019-10656-5

934   61.   McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis
935   and Graphics of Microbiome Census Data. *PLoS One*. 2013;8(4).
936   doi:10.1371/journal.pone.0061217

937   62.   Kenneth Helrich. *Official Methods for Analysis of the Association of Official Analytical*
938   *Chemists*.; 1990.

939   63.   Chaney AL, MEP. Modified Reagents for Determination of Urea and Ammonia. *Clin*
940   *Chem*. 1962;8(2):130-132.

941   64.   Holscher HD, Doligale JL, Bauer LL, et al. Gastrointestinal tolerance and utilization of
942   agave inulin by healthy adults. *Food Funct*. 2014;5(6):1142-1149.
943   doi:10.1039/c3fo60666j

944   65.   Flickinger EA, Schreijen EMWC, Patil AR, et al. Nutrient digestibilities, microbial
945   populations, and protein catabolites as affected by fructan supplementation of dog diets
946   Get access Arrow. *J Anim Sci*. 2003;81(8):2008-2018.

947   66.   Holscher HD, Chumpitazi BP, Dahl WJ, et al. Perspective: Assessing Tolerance to
948   Nondigestible Carbohydrate Consumption. *Advances in Nutrition*. 2022;13(6):2084-2097.
949   doi:10.1093/advances/nmac091

950   67.   Maki K, Rains T, Kelley K, Cook C, Schild A, Gietl E. Fibermalt is well tolerated in
951   healthy men and women at intakes up to 60 g/d: a randomized, double-blind, crossover
952   trial. *Int J Food Sci Nutr*. 2013;63(3):274-281.

953   68.   Tingley D, Yamamoto T, Hirose K, Keele L, Imai K. *Journal of Statistical Software*
954   *Mediation: R Package for Causal Mediation Analysis*.; 2014. http://www.jstatsoft.org/

955   69.   Chi WE, Huang S, Jeon M, Park ES, Melguizo T, Kezar A. A Practical Guide to Causal
956   Mediation Analysis: Illustration With a Comprehensive College Transition Program and
957   Nonprogram Peer and Faculty Interactions. *Front Educ (Lausanne)*. 2022;7.
958   doi:10.3389/feduc.2022.886722

959   70.   Sohn MB, Li H. Compositional mediation analysis for microbiome studies. *Annals of*
960   *Applied Statistics*. 2019;13(1):661-681. doi:10.1214/18-AOAS1210

961   71.   Aiken LS, West SG. Interaction Effects. In: *Encyclopedia of Statistics in Behavioral*
962   *Science*. Wiley; 2005. doi:10.1002/0470013192.bsa306

963   72.   Preacher KJ, Curran PJ, Bauer DJ. *Computational Tools for Probing Interactions in*
964   *Multiple Linear Regression, Multilevel Modeling, and Latent Curve Analysis*. Vol 31.;
965   2006. http://www.quantpsy.org/.437

966   73.   García-Bayona L, Comstock LE. Streamlined genetic manipulation of diverse bacteroides
967   and parabacteroides isolates from the human gut microbiota. *mBio*. 2019;10(4).
968   doi:10.1128/mBio.01762-19

969   74.   Douglas GM, Maffei VJ, Zaneveld JR, et al. PICRUSt2 for prediction of metagenome
970   functions. *Nat Biotechnol.Nature Research*. 2020;38(6):685-688. doi:10.1038/s41587-
971   020-0548-6

972   75.   Langille MGI, Zaneveld J, Caporaso JG, et al. Predictive functional profiling of microbial
973   communities using 16S rRNA marker gene sequences. *Nat Biotechnol*. 2013;31(9):814-
974   821. doi:10.1038/nbt.2676

975   76.   Beghini F, McIver LJ, Blanco-Míguez A, et al. Integrating taxonomic, functional, and
976   strain-level profiling of diverse microbial communities with biobakery 3. *Elife*. 2021;10.
977   doi:10.7554/eLife.65088

978   77.   Blanco-Míguez A, Beghini F, Cumbo F, et al. Extending and improving metagenomic
979   taxonomic profiling with uncharacterized species using MetaPhlAn 4. *Nat Biotechnol*.
980   2023;41(11):1633-1644. doi:10.1038/s41587-023-01688-w

981   78.   Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation.
982   *Genome Biol*. 2011;12(6). doi:10.1186/gb-2011-12-6-r60

79. Drula E, Garron ML, Dogan S, Lombard V, Henrissat B, Terrapon N. The carbohydrate-
active enzyme database: Functions and literature. *Nucleic Acids Res*. 2022;50(D1):D571-
D577. doi:10.1093/nar/gkab1045
80. Cantarel BI, Coutinho PM, Rancurel C, Bernard T, Lombard V, Henrissat B. The
Carbohydrate-Active EnZymes database (CAZy): An expert resource for glycogenomics.
Nucleic Acids Res. 2009;37(SUPPL. 1). doi:10.1093/nar/gkn663
81. Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y. DbCAN3: Automated carbohydrate-
Active enzyme and substrate annotation. *Nucleic Acids Res*. 2023;51(W1):W115-W121.
doi:10.1093/nar/gkad328
82. Erdfelder E, FAul F, Buchner A, Lang AG. Statistical power analyses using G*Power 3.1:
Tests for correlation and regression analyses. *Behav Res Methods*. 2009;41(4):1149-1160.
doi:10.3758/BRM.41.4.1149
83. R Core Team. R: A language and environment for statistical computing. *Foundation for
Statistical Computing*. Preprint posted online 2013.
84. H. Wickham. ggplot2: Elegant Graphics for Data Analysis. *Springer-VerlagNew York*.
Published online 2016.
85. Kwak SK, Kim JH. Statistical data preparation: Management of missing values and
outliers. *Korean J Anesthesiol.Korean Society of Anesthesiologists*. 2017;70(4):407-411.
doi:10.4097/kjae.2017.70.4.407
86. Kafadar K. John Tukey and Robustness. *Statistical Science*. 2003;3(18(3)):319-331.
doi:1076102419
87. Zhang X, Mallick H, Tang Z, et al. Negative binomial mixed models for analyzing
microbiome count data. *BMC Bioinformatics*. 2017;18(1). doi:10.1186/s12859-016-1441-
7
88. Sweeny AR, Lemon H, Ibrahim A, et al. A mixed-model approach for estimating drivers
of microbiota community composition and differential taxonomic abundance. *mSystems*.
Published online July 25, 2023. doi:10.1128/msystems.00040-23
89. Jamieson PE, Smart EB, Bouranis JA, et al. Gut enterotype-dependent modulation of gut
microbiota and their metabolism in response to xanthohumol supplementation in healthy
adults. *Gut Microbes*. 2024;16(1). doi:10.1080/19490976.2024.2315633
90. Zwietering MH, Jongenburger I, Rombouts FM, Van ' K, Riet T. *Modeling of the
Bacterial Growth Curve PUm*.; 1990. https://journals.asm.org/journal/aem
91. Mallick H, Rahnavard A, McIver LJ, et al. Multivariable association discovery in
population-scale meta-omics studies. *PLoS Comput Biol*. 2021;17(11).
doi:10.1371/journal.pcbi.1009442
92. Yang C, Mai J, Cao X, Burberry A, Cominelli F, Zhang L. ggpicrust2: an R package for
PICRUSt2 pre-dicted functional profile analysis and visualiza-tion. *Bioinformatics,
YYYY*.:0-0. doi:10.1093/bioinformatics/xxxxx
93. Edwards CG, Walk AM, Thompson S V., et al. Effects of 12-week avocado consumption
on cognitive function among adults with overweight and obesity. *International Journal of
Psychophysiology*. 2020;148:13-24. doi:10.1016/j.ijpsycho.2019.12.006
94. Ford NA, Spagnuolo P, Kraft J, Bauer E. Nutritional Composition of Hass Avocado Pulp.
Foods.Multidisciplinary Digital Publishing Institute (MDPI). 2023;12(13).
doi:10.3390/foods12132516
95. Thompson S V., Bailey MA, Taylor AM, et al. Avocado Consumption Alters
Gastrointestinal Bacteria Abundance and Microbial Metabolite Concentrations among

1029   Adults with Overweight or Obesity: A Randomized Controlled Trial. *Journal of Nutrition.*
1030   2021;151(4):753-762. doi:10.1093/jn/nxaa219

1031   96.   Fda. *Review of the Scientific Evidence on the Physiological Effects of Certain Non-*
1032   *Digestible Carbohydrates.*; 2018.

1033   97.   Mysonhimer AR, Holscher HD. Gastrointestinal Effects and Tolerance of Nondigestible
1034   Carbohydrate Consumption. *Advances in Nutrition.* 2022;13(6):2237-2276.
1035   doi:10.1093/advances/nmac094

1036   98.   Mehren A, Luque CD, Brandes M, et al. Intensity-dependent effects of acute exercise on
1037   executive function. *Neural Plast.* 2019;2019. doi:10.1155/2019/8608317

1038   99.   Timm DA, Thomas W, Boileau TW, Williamson-Hughes PS, Slavin JL. Polydextrose and
1039   soluble corn fiber increase five-day fecal wet weight in healthy men and women. *Journal*
1040   *of Nutrition.* 2013;143(4):473-478. doi:10.3945/jn.112.170118

1041   100.   Hooda S, Vester Boler BM, Rossoni Serao MC, et al. 454 pyrosequencing reveals a shift
1042   in fecal microbiota of healthy adult men consuming polydextrose or soluble corn fiber.
1043   *Journal of Nutrition.* 2012;142(7):1259-1265. doi:10.3945/jn.112.158766

1044   101.   Cantu-Jungles TM, Hamaker BR. Tuning Expectations to Reality: Don't Expect Increased
1045   Gut Microbiota Diversity with Dietary Fiber. *Journal of Nutrition.* 2023;153(11):3156-
1046   3163. doi:10.1016/j.tjnut.2023.09.001

1047   102.   Shimizu H, Nakajima M, Miyanaga A, et al. Characterization and Structural Analysis of a
1048   Novel exo-Type Enzyme Acting on β-1,2-Glucooligosaccharides from Parabacteroides
1049   distasonis. *Biochemistry.* 2018;57(26):3849-3860. doi:10.1021/acs.biochem.8b00385

1050   103.   Ducarmon QR, Karcher N, Tytgat HLP, et al. Large-scale computational analyses of gut
1051   microbial CAZyme repertoires enabled by Cayman. Preprint posted online January 8,
1052   2024. doi:10.1101/2024.01.08.574624

1053   104.   Lapébie P, Lombard V, Drula E, Terrapon N, Henrissat B. Bacteroidetes use thousands of
1054   enzyme combinations to break down glycans. *Nat Commun.* 2019;10(1).
1055   doi:10.1038/s41467-019-10068-5

1056   105.   Gibson GR, Hutkins R, Sanders ME, et al. Expert consensus document: The International
1057   Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the
1058   definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol.Nature Publishing*
1059   *Group.* 2017;14(8):491-502. doi:10.1038/nrgastro.2017.75

1060   106.   Goel A, Shete O, Goswami S, et al. Toward a health-associated core keystone index for
1061   the human gut microbiome. *Cell Rep.* 2025;44(3):115378.
1062   doi:10.1016/j.celrep.2025.115378

1063   107.   Cui Y, Zhang L, Wang X, et al. Roles of intestinal Parabacteroides in human health and
1064   diseases. *FEMS Microbiol Lett.Oxford University Press.* 2022;369(1).
1065   doi:10.1093/femsle/fnac072

1066   108.   Cuffaro B, Boutillier D, Desramaut J, et al. Characterization of Two Parabacteroides
1067   distasonis Candidate Strains as New Live Biotherapeutics against Obesity. *Cells.*
1068   2023;12(9). doi:10.3390/cells12091260

1069   109.   Ezeji JC, Sarikonda DK, Hopperton A, et al. Parabacteroides distasonis: intriguing
1070   aerotolerant gut anaerobe with emerging antimicrobial resistance and pathogenic and
1071   probiotic roles in human health. *Gut Microbes.Bellwether Publishing, Ltd.* 2021;13(1).
1072   doi:10.1080/19490976.2021.1922241

110. Liang Z, Di N, Li L, Yang D. Gut microbiota alterations reveal potential gut–brain axis
changes in polycystic ovary syndrome. *J Endocrinol Invest*. 2021;44(8):1727-1737.
doi:10.1007/s40618-020-01481-5
111. Wei W, Wong CC, Jia Z, et al. Parabacteroides distasonis uses dietary inulin to suppress
NASH via its metabolite pentadecanoic acid. *Nat Microbiol*. 2023;8(8):1534-1548.
doi:10.1038/s41564-023-01418-7
112. Liu D, Zhang S, Li S, et al. Indoleacrylic acid produced by Parabacteroides distasonis
alleviates type 2 diabetes via activation of AhR to repair intestinal barrier. *BMC Biol*.
2023;21(1). doi:10.1186/s12915-023-01578-2
113. Keaney J, Campbell M. The dynamic blood-brain barrier. *FEBS Journal*.
2015;282(21):4067-4079. doi:10.1111/febs.13412
114. Van Dokkum W, Wezendonk B, Srikumar TS, Van Den Heuvel EGHM. Effect of
nondigestible oligosaccharides on large-bowel functions, blood lipid concentrations and
glucose absorption in young healthy male subjects. *Eur J Clin Nutr*. 1999;53(1):1-7.
doi:10.1038/sj.ejcn.1600668
115. Pedersen A, Sandström B, Van Amelsvoort JMM. The effect of ingestion of inulin on
blood lipids and gastrointestinal symptoms in healthy females. *British Journal of
Nutrition*. 1997;78(2):215-222. doi:10.1079/bjn19970141
116. Dalile B, La Torre D, Kalc P, et al. Extruded Wheat Bran Consumption Increases Serum
Short-Chain Fatty Acids but Does Not Modulate Psychobiological Functions in Healthy
Men: A Randomized, Placebo-Controlled Trial. *Front Nutr*. 2022;9.
doi:10.3389/fnut.2022.896154
117. Wang S, Mu L, Yu C, et al. Microbial collaborations and conflicts: unraveling interactions
in the gut ecosystem. *Gut Microbes*.*Taylor and Francis Ltd.* 2024;16(1).
doi:10.1080/19490976.2023.2296603
118. Takada T, Kurakawa T, Tsuji H, Nomoto K. Fusicatenibacter saccharivorans gen. nov.,
sp. nov., isolated from human faeces. *Int J Syst Evol Microbiol*. 2013;63(PART10):3691-
3696. doi:10.1099/ijs.0.045823-0
119. Li J, Zhu S, Wang Y, et al. Metagenomic association analysis of cognitive impairment in
community-dwelling older adults. *Neurobiol Dis*. 2023;180.
doi:10.1016/j.nbd.2023.106081
120. Loh JS, Mak WQ, Tan LKS, et al. Microbiota–gut–brain axis and its therapeutic
applications in neurodegenerative diseases. *Signal Transduct Target Ther*.*Springer
Nature*. 2024;9(1). doi:10.1038/s41392-024-01743-1

1107

Tables

Table 1. Baseline demographic and clinical characteristics of participants randomized to treatment counterbalanced.

	Overall N = 42	**CON** N = 21	**SCF** N = 21
Education			
High School Graduate	2 (4.8%)	0 (0%)	2 (9.5%)
Some college	6 (14%)	3 (14%)	3 (14%)
Bachelor's Degree	13 (31%)	8 (38%)	5 (24%)
Advanced Degree	21 (50%)	10 (48%)	11 (52%)
Ethnicity			
Asian	6 (14%)	3 (14%)	3 (14%)
Black or African American	3 (7.1%)	1 (4.8%)	2 (9.5%)
Mixed or Other	1 (2.4%)	0 (0%)	1 (4.8%)
White (Caucasian)	32 (76%)	17 (81%)	15 (71%)
Latino	1 (2.4%)	1 (4.8%)	0 (0%)
Sex			
Female	28 (67%)	15 (71%)	13 (62%)
Male	14 (33%)	6 (29%)	8 (38%)
Age (years)	56.2 ± 7.3	54.8 ± 6.2	57.5 ± 8.2
Fiber per 1000kcal	11.4 ± 3.4	12.0 ± 3.3	10.8 ± 3.5
BMI (kg/m²)	25.9 ± 4.0	25.4 ± 3.7	26.3 ± 4.2
Income			
<$31,000-40,000	3 (9.5%)	1 (4.8%)	2 (14.3%)
$41,000-50,000	3 (7.1%)	1 (4.8%)	2 (9.5%)
$51,000-60,000	3 (7.1%)	2 (9.5%)	1 (4.8%)
$61,000-70,000	4 (9.5%)	1 (4.8%)	3 (14%)
$71,000-80,000	5 (12%)	4 (19%)	1 (4.8%)
$81,000-90,000	1 (2.4%)	1 (4.8%)	0 (0%)
$91,000-100,000	7 (16.7%)	4 (19%)	3 (14%)
Greater than $100,000	15 (35.7%)	7 (33%)	8 (38%)

Baseline characteristics of participants enrolled and randomized into the clinical trial (N = 42), grouped by initial treatment assignment. Dietary fiber (g/1000 kcal) values reflect baseline intake assessed using DHQ3 prior to the first intervention period. Height and weight were measured at the baseline visit and used to calculate body mass index (BMI; kg/m^2). Continuous variables are reported as mean ± standard deviation; categorical variables are reported as number of participants and percentage within each group [n (%)].

1129 **Table 2.** Behavioral and ERP measures during the flanker task across treatment and time.
1130

| Behavioral Flanker | Maltodextrin | | Soluble Corn Fiber | | Treatment Effect (ΔSCF - ΔCON) | | | |
	PRE	POST	PRE	POST				
	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	β	SE	95% CI	p
Accuracy (%)								
Congruent	96.9 ± 0.52 (39)	95.5 ± 1.37 (39)	97.0 ± 0.48 (40)	97.6 ± 0.45 (40)	0.65	0.8	[-0.93, 2.23]	0.42
Incongruent	92.4 ± 1.00 (39)	92.0 ± 1.53 (39)	93.6 ± 0.79 (40)	94.4 ± 0.70 (40)	-0.68	1.1	[-2.82, 1.46]	0.53
Reaction Time (ms)								
Congruent	395. ± 5.79 (39)	**398. ± 6.69[a] (39)**	400.71 ± 6.22 (40)	**389. ± 5.52[b] (40)**	-18.4	4.2	[-18.4, -1.2]	**0.01***
Incongruent	431. ± 6.62 (39)	**437. ± 7.77[a] (39)**	435.43 ± 6.88 (40)	**424. ± 6.35[b] (40)**	-14.2	4.4	[-22.8, -5.6]	**0.003****
ERP Flanker								
Amplitude (μV)								
Congruent	5.96 ± 1.73 (39)	6.78 ± 0.68 (36)	6.64 ± 0.60 (39)	7.07 ± 0.77 (37)	0.88	1.18	[-2.22, 1.01]	0.46
Incongruent	6.36 ± 1.64 (39)	6.77 ± 0.69 (36)	7.14 ± 0.65 (39)	7.50 ± 0.79 (37)	1.08	1.10	[-1.06, 3.24]	0.33
Peak Latency (ms)								
Congruent	386. ± 6.71 (39)	399. ± 10.18 (36)	394. ± 7.29 (39)	398. ± 7.99 (37)	-6.16	0.32	[-31.1, 18.7]	0.67
Incongruent	424. ± 8.34 (39)	443. ± 11.70 (36)	438. ± 9.90 (39)	429. ± 8.27 (37)	-23.0	15.4	[-53.2, 6.81]	0.14

1131
Values are means ± standard error of the mean (SEM) and sample size (n) across congruencies at each timepoint. Measures include accuracy (%), reaction time
(ms), P3 amplitude (μV), and P3 peak latency (ms). Linear mixed-effect models included participant ID as a random effect, and the fixed effects of treatment, time,
and their interaction. Reported are the treatment-by-time interaction estimates (β, Standard Error (SE), 95% confidence interval (CI), p-values). Pairwise
comparisons were conducted within each congruency when a treatment-by-time interaction met the significance threshold ($P < 0.05$), with p-values adjusted using
the Benjamin-Hochberg procedure. Superscripts (a, b) denote significant *pairwise* differences within congruency trials between treatments and time. Significance
threshold p: '**' <0.01, '*' <0.05, '†' <0.1.

Table 3. 16S Alpha diversity and differential abundance of genera across intervention periods.

| 16S ASVs | Maltodextrin | | Soluble Corn Fiber | | Treatment Effect (ΔSCF - ΔCON) | | | |
| | PRE | POST | PRE | POST | | | | |
	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	β	SE	95% CI	FDR P
Genera (% of sequence)								
Bacteroides	25.2 ± 1.77 (38)	23.7 ± 1.90 (38)	27.3 ± 2.06 (37)	22.6 ± 2.07 (37)	-0.15	0.10	[-0.34, 0.05]	0.11
Faecalibacterium	10.2 ± 1.02 (38)	10.1 ± 1.11 (38)	8.66 ± 1.15 (37)	11.0 ± 1.30 (37)	0.23	0.21	[-0.18, 0.64]	0.47
Parabacteroides	2.52 ± 0.35[a] (38)	**2.06 ± 0.24[a]** (38)	2.09 ± 0.26[a] (37)	**8.37 ± 0.98[b]** (37)	1.44	0.22	[1.01, 1.88]	**<0.001***
Alistipes	3.46 ± 0.51 (38)	2.82 ± 0.35 (38)	3.88 ± 0.59 (37)	3.96 ± 0.69 (37)	0.08	0.20	[-0.31, 0.46]	0.50
Blautia	3.76 ± 0.50 (38)	4.52 ± 0.61 (38)	3.54 ± 0.42 (37)	3.24 ± 0.50 (37)	-0.27	0.13	[-0.52, -0.02]	0.10†
Agathobacter	4.16 ± 0.88 (38)	5.97 ± 1.17 (38)	3.99 ± 0.85 (37)	2.80 ± 0.58 (37)	-0.46	0.23	[-0.91, 0.01]	0.11
Subdoligranulum	2.31 ± 0.29 (38)	2.54 ± 0.38 (38)	3.24 ± 0.58 (37)	2.69 ± 0.49 (37)	-0.24	0.22	[-0.68, 0.19]	0.83
Prevotella	0.87 ± 0.41 (38)	1.64 ± 0.74 (38)	1.60 ± 0.89 (37)	2.13 ± 1.04 (37)	-0.22	0.30	[-0.81, 0.37]	0.53
Dialister	1.34 ± 0.44[a] (38)	**1.46 ± 0.51[a]** (38)	0.93 ± 0.34[a] (37)	**2.10 ± 0.63[b]** (37)	1.06	0.39	[0.29, 1.83]	**0.02***
Bifidobacterium	2.50 ± 0.53 (38)	2.69 ± 0.53 (38)	2.12 ± 0.52 (37)	2.09 ± 0.41 (37)	0.09	0.31	[-0.51, 0.70]	0.92
Fusicatenibacter	0.60 ± 0.10[a] (38)	**0.67 ± 0.09[a]** (38)	0.53 ± 0.11[a] (37)	**1.79 ± 0.28[b]** (37)	1.10	1.10	[0.59, 1.61]	**<0.001***
Alpha Diversity Metrics								
Richness	439.9 ± 19.8[a] (38)	**438.7 ± 18.2[a]** (38)	443.6 ± 17.6[a] (37)	**413.7 ± 18.6[b]** (37)	-0.76	0.29	[-1.32, -0.19]	**0.03***
Shannon	3.89 ± 0.06 (38)	3.88 ± 0.06 (38)	3.81 ± 0.06 (37)	3.71 ± 0.07 (37)	-0.08	0.07	[-0.22, 0.06]	0.26
Faith's PD	104.7 ± 1.62 (38)	103.9 ± 1.54 (38)	105.23 ± 1.54 (37)	101.9 ± 1.80 (37)	-2.43	1.28	[-4.93, 0.06]	0.09†

1139
Values are means ± standard error of the mean (SEM) and sample size (*n*) across each timepoint. For differential abundance of 16S genera, a generalized linear
mixed-effect negative binomial model included participant ID as a random effect, and the fixed effects of treatment, time, and their interaction. Reported are the
treatment-by-time interaction log-scaled estimates (β, Standard Error (SE), 95% confidence interval (CI), FDR-adjusted *p*-values). For alpha diversity metrics,
linear mixed-effect model. Reported are the treatment-by-time interaction estimates. Pairwise comparisons was conducted if an interaction *(treatment effect)* met
the significance threshold ($P < 0.05$), *p*-values adjusted using the Benjamin-Hochberg procedure. Superscripts (a, b) denote significant *pairwise* treatment effect.
Significance threshold *p*: '***' <0.001, '**' <0.01, '*' <0.05, '†' <0.1.
1146

1147 **Table 4.** Fecal volatile fatty acids across the intervention periods.

| Fermentation end-products (μmol/g DMB) | Maltodextrin | | Soluble Corn Fiber | | Treatment Effect (ΔSCF - ΔCON) | | | |
| | PRE | POST | PRE | POST | | | | |
	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	Mean ± SEM (n)	β	SE	95% CI	P
SCFAs								
Butyrate	43.3 ± 4.56 (38)	43.9 ± 5.26 (37)	38.8 ± 4.58 (37)	40.1 ± 3.18 (37)	0.10	0.13	[-0.16, 0.37]	0.44
Acetate	176. ± 13.9 (38)	182. ± 13.5 (37)	165. ± 13.2 (37)	202. ± 14.9 (37)	0.16	0.11	[-0.04, 0.36]	0.14
Propionate	48.1 ± 3.98 (38)	50.8 ± 4.40 (37)	47.5 ± 4.32 (37)	55.6 ± 4.72 (37)	0.14	0.10	[-0.09, 0.36]	0.23
Total SCFAs	267. ± 21.5 (38)	273. ± 21.97 (37)	255. ± 20.6 (37)	298. ± 22.3 (37)	0.15	0.11	[-0.05, 0.36]	0.15
BCFAs								
Valerate	6.44 ± 0.59 (38)	5.99 ± 0.47 (37)	6.00 ± 0.47 (37)	5.68 ± 0.48 (37)	-0.06	0.16	[-0.36, 0.24]	0.71
Isobutyrate	6.62 ± 0.54 (38)	5.97 ± 0.39 (36)	6.29 ± 0.52 (37)	5.22 ± 0.33 (37)	-0.08	0.11	[-0.30, 0.14]	0.48
Isovalerate	9.49 ± 0.82 (38)	8.30 ± 0.57 (36)	8.66 ± 0.69 (37)	7.35 ± 0.54 (37)	-0.04	0.14	[-0.31, 0.22]	0.75
Total BCFAs	22.6 ± 1.83 (38)	20.4 ± 1.31 (37)	21.0 ± 1.56 (37)	18.3 ± 1.14 (37)	-0.03	0.20	[-0.43, 0.36]	0.87

1148
1149 Values are descriptive means ± standard error of the mean (SEM) and sample size (n) across each timepoint. Total SCFAs and BCFAs are pooled measures, within
1150 respective category, concentrations are on a dry matter basis (DMB). Linear mixed-effect models included participant ID as a random effect, and the fixed effects
1151 of treatment, time, and their interaction (Treatment Effect). Reported are the treatment-by-time interaction estimates (β, Standard Error (SE), 95% confidence
1152 interval (CI), *p*-values).

1153 **Table 5.** Self-reported gastrointestinal symptom severity scores across the intervention.

| GI Symptoms | Maltodextrin | | Soluble Corn Fiber | | |
| | PRE | POST | PRE | POST | |
	Median [IQR] (n)	Median [IQR] (n)	Median [IQR] (n)	Median [IQR] (n)	P
Gas	1.57 [0.84] (39)	**1.43 [1.00]** (39)	1.57 [0.97] (38)	**1.93 [1.00]** (38)	**0.01***
Cramping	1.00 [0.00] (39)	1.00 [0.00] (39)	1.00 [0.00] (38)	1.00 [0.14] (38)	0.91
Bloating	1.00 [0.14] (39)	1.00 [0.14] (39)	1.00 [0.14] (38)	1.00 [0.32] (38)	0.05†
Nausea	1.00 [0.00] (39)	1.00 [0.00] (39)	1.00 [0.00] (38)	1.00 [0.00] (38)	0.17
Reflux	1.00 [0.00] (39)	1.00 [0.00] (39)	1.00 [0.03] (38)	1.00 [0.00] (38)	0.84
Rumbling	1.00 [0.16] (39)	1.00 [0.16] (39)	1.00 [0.17] (38)	1.00 [0.41] (38)	0.68
Burping	1.00 [0.16] (39)	1.00 [0.00] (39)	1.00 [0.28] (38)	1.00 [0.17] (38)	0.70

1154
1155 Values are median [interquartile range (IQR)] and sample size (n) at each timepoint. Symptom were rated on a 4-point
1156 scale (1 = Absent, 2 = Mild, 3 = Moderate, 4 = Severe). Reported values include *p*-values from the paired Wilcoxon
1157 Signed-Rank testing the treatment effect on change over time using delta values. Significance threshold *p*: '*' <0.05,
1158 '†' <0.1.
1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

Table 6. Mediation of SCF-related changes in Flanker reaction times through fecal SCFAs and
16S genera.

		Nonparametric Bootstrap Confidence Intervals with Percentile			
Mediator	**Flanker Task**	**Effect**	**β**	**[95% CI]**	**P**
Acetate	Congruent RT	ACME	-0.57	[-2.51, 0.87]	0.48
		ADE	-13.42	[-23.7, -3.23]	0.01*
		Total effect	-13.99	[-24.4, -14.0]	0.009**
		Proportion Mediated	4.09%	[-0.08, 0.21]	0.48
	Incongruent RT	ACME	-0.44	[-2.87, 1.61]	0.72
		ADE	-17.03	[-29.4, -4.38]	0.008**
		Total effect	-17.47	[-30.1, -4.63]	0.007**
		Proportion Mediated	2.51%	[-0.11, 0.19]	0.72
Propionate	Congruent RT	ACME	-0.64	[-2.39, 0.81]	0.43
		ADE	-13.39	[-23.9, -3.30]	0.01*
		Total effect	-14.03	[-24.4, -3.78]	0.008**
		Proportion Mediated	4.57%	[-0.28, 0.08]	0.43
	Incongruent RT	ACME	-0.38	[-4.32, 0.78]	0.22
		ADE	-17.11	[-23.0, -2.36]	0.03*
		Total effect	-17.49	[-24.3, -3.77]	0.01*
		Proportion Mediated	2.22%	[-0.11, 0.54]	0.23
Parabacteroides	Congruent RT	ACME	-0.15	[-9.31, 4.27]	0.93
		ADE	-13.85	[-25.9, 0.39]	0.06
		Total effect	-14.00	[-24.3, -3.83]	0.008**
		Proportion Mediated	1.06%	[-0.36, 1.02]	0.93
	Incongruent RT	ACME	-0.48	[-12.64, 4.53]	0.86
		ADE	-17.00	[-30.5, 0.20]	0.03*
		Total effect	-17.48	[-30.0, -4.80]	0.01*
		Proportion Mediated	2.75%	[-0.34, 0.96]	0.86
Fusicatenibacter	Congruent RT	ACME	-1.40	[-4.98, 0.65]	0.19
		ADE	-12.63	[-23.0, -2.28]	0.02*
		Total effect	-14.03	[-24.3, -3.84]	0.007**
		Proportion Mediated	10.0%	[-0.06, 0.50]	0.19
	Incongruent RT	ACME	-0.72	[-4.98, 2.16]	0.59
		ADE	-16.75	[-29.1, -3.85]	0.01*
		Total effect	-17.47	[-30.0, -4.79]	0.007**
		Proportion Mediated	4.1%	[-0.18, 0.35]	0.59
Dialister	Congruent RT	ACME	0.44	[-0.89, 3.47]	0.52
		ADE	-14.44	[-25.4, -4.22]	0.006**
		Total effect	-14.00	[-24.2, -3.75]	0.007**
		Proportion Mediated	-3.15%	[-0.33, 0.08]	0.53
	Incongruent RT	ACME	0.67	[-1.26, 4.95]	0.52
		ADE	-18.15	[-31.8, -5.12]	0.007**
		Total effect	-17.49	[-30.0, -4.70]	0.007**
		Proportion Mediated	-3.8%	[-0.35, 0.11]	0.53

Mediation analysis (*n* = 37) was conducted using nonparametric bootstrapped linear regression models (R; *mediation*)
to estimate the average causal mediation effect (ACME), average direct effect (ADE), and total effects of treatment
on changes in reaction times (RT) Reported are the model estimates (β), Standard Error (SE), 95% confidence interval
(CI). Proportion mediated indicates the percentage of the total effect explained by the ACME pathway. Significance
threshold *p*: '***' <0.001, '**' <0.01, '*' <0.05, '†' <0.1.

1176 **Table 7.** Moderation of the SCF effect on change in flanker reaction time by change in 16S
1177 genera and fecal metabolites

Moderator	Flanker Task	Treatment x Moderator (Model Summary)						
		ΔR^2	Adj. ΔR^2	β	SE	t	[95% CI]	*P*
Acetate	Congruent RT	0.11	0.07	3.13	5.62	0.56	[-8.08, 14.3]	0.58
	Incongruent RT	0.10	0.06	2.21	6.80	0.33	[-11.4, 15.8]	0.75
Propionate	Congruent RT	0.11	0.07	0.43	5.60	0.08	[-10.76, 11.6]	0.94
	Incongruent RT	0.10	0.06	0.40	6.80	0.06	[-13.17, 14.0]	0.95
Parabacteroides	Congruent RT	**0.16**	0.12	**-19.8**	10.12	-1.95	[-40.0, 0.43]	**0.05***
	Incongruent RT	**0.15**	0.12	**-25.1**	12.24	-2.05	[-49.5, 0.63]	**0.04***
Fusicatenibacter	Congruent RT	0.10	0.06	-1.31	13.48	-0.10	[-28.2, 25.6]	0.92
	Incongruent RT	0.11	0.07	11.49	16.20	0.71	[-20.9, 43.8]	0.49
Dialister	Congruent RT	0.11	0.07	-4.49	7.39	-0.61	[-19.2, 10.3]	0.55
	Incongruent RT	0.10	0.06	0.23	8.98	0.03	[-17.7, 18.2]	0.98

1178
1179 Moderation analysis (*n* = 37) was tested using linear regression models testing if changes in moderators explain
1180 variability in flanker reaction times (RT). Each model includes treatment, Δ moderator, and its interaction. Reported
1181 are the treatment-by-moderator estimates (β, Standard Error (SE), 95% confidence interval (CI), *p*-values). The ΔR^2
1182 and adjusted ΔR^2 columns show the variance explained by the full model. Statistical threshold for *p*: '***' <0.001,
1183 '**' <0.01, '*' <0.05, '†' <0.1.
1184

1185

1186

1187

1188

1189

Table 8. Responder vs. non-responder descriptive summary during the SCF intervention period

Pilot Analyses	Non-Responder		Responder	
	PRE	POST	PRE	POST
	Mean ± SD *(n)*	Mean ± SD *(n)*	Mean ± SD *(n)*	Mean ± SD *(n)*
Genera (% of Sequences)				
Parabacteroides	0.43 ± 0.64 *(3)*	*ND (3)*	1.04 ± 0.33 *(3)*	14.61 ± 2.82 *(3)*
Dialister	1.50 ± 0.79 *(3)*	3.18 ± 3.25 *(3)*	0.25 ± 0.43 *(3)*	*ND (3)*
Fusicatenibacter	0.48 ± 0.45 *(3)*	2.69 ± 2.77 *(3)*	1.24 ± 1.65 *(3)*	1.35 ± 1.34 *(3)*
Species (% of Sequences)				
P. distasonis	0.01 ± 0.01 *(3)*	*ND (3)*	0.49 ± 0.17 *(3)*	11.38 ± 1.47 *(3)*
P. merdae	0.41 ± 0.63 *(3)*	*ND (3)*	0.55 ± 0.50 *(3)*	3.23 ± 3.67 *(3)*
P. faecis	0.01 ± 0.01 *(3)*	*ND (3)*	*ND (3)*	*ND (3)*
D. invisus	0.73 ± 0.73 *(3)*	1.00 ± 1.74 *(3)*	0.25 ± 0.43 *(3)*	*ND (3)*
D. hominis	0.77 ± 1.34 *(3)*	2.17 ± 3.75 *(3)*	*ND (3)*	*ND (3)*
F. saccharivorans	0.48 ± 0.45 *(3)*	2.69 ± 2.77 *(3)*	1.24 ± 1.65 *(3)*	1.35 ± 1.34 *(3)*
Flanker Reaction Time (ms)				
Congruent	428.9 ± 24.4 *(3)*	396.7 ± 11.6 *(3)*	363.0 ± 50.2 *(3)*	362.1 ± 42.7 *(3)*
Incongruent	456.6 ± 3.9 *(3)*	424.9 ± 14.4 *(3)*	403.9 ± 60.2 *(3)*	401.1 ± 50.2 *(3)*
SCFAs (μmol/g DMB)				
Acetate	153.8 ± 60.9 *(3)*	225.4 ± 88.9 *(3)*	180.2 ± 95.1 *(3)*	246.6 ± 59.1 *(3)*
Propionate	28.47 ± 12.2 *(3)*	57.02 ± 23.7 *(3)*	51.89 ± 23.6 *(3)*	72.81 ± 27.2 *(3)*
Demographics				
Age *(year)*	53.7 ± 7.2 *(3)*		54.3 ± 2.5 *(3)*	

Responders (n=3) and non-responders (n=3) following the SCF intervention. Responders were the three participants with the largest increase in *Parabacteroides* (16S) following SCF; non-responders were the three participants with the smallest change following SCF (total N = 6 participants). Data are presented as mean ± standard deviation (SD) and sample size (*n*). Genus and species relative abundances (%) were derived from shotgun metagenomic profiling using MetaPhlAn4. Species initials correspond as genus followed by species name: *P* = *Parabacteroides*, *D* = *Dialister*, and *F* = *Fusicatenibacter*. ND = not detected (no reads assigned at the reported taxonomic level.)

 Figure Legend

1202 **Figure 1. CONSORT flow diagram of recruitment, randomization, and retention.**
1203 Participants were randomized (n=42) to start with either the maltodextrin control (CON) or soluble
1204 corn fiber (SCF) in Period 1, then crossed over to the alternate condition in Period 2 following
1205 washout. Boxes show the "*n*" indicating the number of participants at each stage and reasons for
1206 exclusion or missing data of the SCOPE study. (*Figure created using BioRender)*

1207

Figure 2. Soluble corn fiber (SCF) consumption improved flanker reaction times. Observed
reaction times (RT, millisecond) are shown at both time points for the maltodextrin control (CON)
and SCF intervention periods for (**A**) congruent trials and (**B**) incongruent trials. Right panels
display model-estimated marginal means (coefficients and the 95% confidence intervals for both
congruent and incongruent trials (FDR corrected pairwise contrasts; symbols shown in right
panels). Statistical significance derived from pairwise comparisons. FDR-adjusted *p*-values
annotated as follows: ** < 0.01; * < 0.05.

1215

Figure 3. 16S microbiota composition across condition and participants. (n=38 CON, n=37
SCF). (**A**) Mean relative abundance (% of sequence) of the top 5 for phyla (*left*) and top 10 for
genera (*right*) at PRE and POST for each intervention. Genus colors correspond to their parent
phylum. (**B**) Differential abundance of genera from a generalized linear mixed model showing the
treatment effect estimates with effect size increasing from the center dashed vertical line. The y-
axis corresponds to the significance of the genera, higher meaning more significant, with a
horizontal dashed line indicating the FDR P = 0.05 threshold. Individual participant-sample
relative abundances are shown as a (**C**) waterfall of the distribution of SCF-responsive features.
Incertae sedis is group of bacteria without taxonomical assignment.

1225

Figure 4. Interaction between 16S-derived Δ*Parabacteroides* and the SCF effect on flanker
RT. Panels show moderation results for (**A**) for congruent and (**B**) incongruent trials. The outcome
is ΔRT (POST-PRE, ms), where negative values indicate faster RT. The moderator is
Δ*Parabacteroides* (POST-PRE), mean-centered and z-scaled (negative values reflect smaller-
than-average changes). Left panels show observed data and model-predicted simple slopes for
CON (blue) and SCF (orange); vertical reference lines indicate -1 SD (standard deviation), average
(0 SD), and +1 SD. Right panels show estimated SCF-CON emmeans contrasts in ΔRT at -1,0,
and +1SD of Δ*Parabacteroides*; the vertical line at 0 indicates no difference. Statistical
significance threshold *p* * <0.05.

1235

Figure 5. Growth rates of *Parabacteroides distasonis* cultured in M9 supplement with SCF
compared to controls. The relative log growth curve of *P. distasonis* (**A**) in M9 minimal medium
supplemented with different carbohydrate sources. OD_0 represents the initial optical density
(OD_{600}) at time zero. Each curve represents a mean biological replicate (N = 3), and error bars
indicate propagated standard deviations. The mean growth rate (**B**), calculated as the slope of
ln(OD) over sequential 2-hour intervals, cut off at 10-hrs. Bars represent average growth rate
across replicates. Significance threshold *p*: '***' < 0.001, '**' < 0.01, '*' < 0.05, '†' < 0.1, 'NS'
> 0.1.

1244

Figure 6. Exploratory responder analysis during the SCF period with MetaPhlAn4. N = 6.
Responders (n=3) were the participants with the largest *Parabacteroides* increase after SCF
consumption, and non-responders (n=3) were the three with the smallest change (null/negative).
The cognitive outcomes from the Flanker task demonstrate the responders (**A**) congruent and (**B**)
incongruent reaction times (RT). Shotgun sequencing and MetaPhlan4 yielded (**C**) species
resolution of microbial hits from 16S genus analysis. The bottom row demonstrates the diversity
metrics for Alpha (**D**) richness and (**E**) Shannon evenness, and the principal coordinate analysis of
(**F**) Beta diversity is shown using Bray-Curtis with adonis2 PERMANOVA values.