Original Article

Initial proteinuria reduction and adverse kidney outcomes in IgA nephropathy: An

analysis from the Japan IgA nephropathy cohort study (J-IGACS)

Takaya Sasaki[1], Nobuo Tsuboi[1]*, Kentaro Koike[1], Hiroyuki Ueda[1], Masahiro Okabe[1], Shinya

Yokote[1], Akihiro Shimizu[1], Keita Hirano[1], Tetsuya Kawamura[1], Takashi Yokoo[1], Yusuke

Suzuki[2] and the J-IGACS working group.

1 Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei

University School of Medicine, Tokyo, Japan

2 Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan.

Running title: Early Proteinuria Reduction and Kidney Outcomes in IgAN

Word counts: abstract 250/250 words; manuscript body 2903/4000 words including

References

*Correspondence to: Nobuo Tsuboi, M.D., Ph.D.

Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei

University School of Medicine, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo, Japan

Tel: 81-3-3433-1111, Fax: 81-3-3433-4297, E-mail: tsuboi-n@jikei.ac.jp

Keywords: IgA nephropathy, proteinuria, eGFR slope, kidney outcome

Abstract

Background: Proteinuria reduction is considered a potential surrogate endpoint predictive of

reflecting long-term kidney prognosis in IgA nephropathy (IgAN). However, its association

with adverse kidney outcomes and IgAN-related decline in estimated glomerular filtration

rate (eGFR) remains uncertain.

Methods: Patients with biopsy-proven IgAN from the Japan IgA Nephropathy Cohort Study

(J-IGACS) were analyzed. Participants were categorized into tertiles based on their 12-month

proteinuria-to-baseline proteinuria ratios. The primary outcome was a composite of ≥40%

eGFR decline or initiation of kidney-replacement therapy. Associations between proteinuria

ratio and outcomes were assessed using Cox proportional hazards models and restricted cubic

splines. Multivariable analyses adjusted for age, sex, baseline eGFR, log-transformed

proteinuria, Oxford classification scores, and use of corticosteroids and renin-angiotensin-

aldosterone system inhibitors within 12 months.

Results: Among 793 patients, those in the lowest tertile (greatest proteinuria reduction) had

significantly lower risk of the primary endpoint (P for trend <0.001) and a more favorable

eGFR slope. Spline analysis showed a continuous, dose-response association between

proteinuria ratio and improved outcomes. These findings remained robust in sensitivity

analyses restricted to patients likely qualifying for clinical trials. The results showed that

patients with lower proteinuria ratios tended to have slower rates of eGFR decline (P for

trend <0.001), even after multivariable adjustment.

Conclusion: Proteinuria reduction within the first-year post-diagnosis is independently

associated with lower risk of adverse kidney outcomes and a slower decline in kidney

function in patients with IgAN. These results support the use of proteinuria reduction as a

surrogate endpoint in both clinical trials and disease management.

1 Introduction

Immunoglobulin A nephropathy (IgAN) is the most prevalent primary glomerulonephritis

globally and a major contributor to progressive chronic kidney disease (CKD) and kidney

failure [1-4]. Despite advances in understanding its pathophysiology and the development of

histopathologic classification systems such as the Oxford Classification [5, 6], treatment

strategies remain largely supportive, with limited disease-specific therapies approved to date.

Proteinuria is a well-established risk factor in IgAN and is frequently utilized in

clinical practice as a target for therapeutic intervention [7, 8]. Several studies have proposed

proteinuria reduction as a potential surrogate endpoint for kidney outcomes, particularly in

the context of clinical trials [9, 10]. However, while short-term reduction in proteinuria is

generally considered favorable, the extent to which this improvement translates into long-

term preservation of kidney function, particularly as reflected by the slope of the estimated

glomerular filtration rate (eGFR), remains incompletely elucidated, especially in Asian

countries, including Japan, where high-quality evidence is still limited.

To address this knowledge gap, we analyzed data from a well-characterized, biopsy-

confirmed cohort of IgAN patients enrolled in the Japan IgA Nephropathy Cohort Study (J-

IGACS) [11, 12]. We aimed to investigate the association between the magnitude of

proteinuria reduction at 12 months post-diagnosis and both a composite kidney outcome and

the longitudinal trajectory of eGFR. We further explored whether this association persists

across clinically relevant subgroups, including those with characteristics commonly

considered in the context of interventional clinical trial eligibility.

Methods

Study population

This study represents a *post-hoc* analysis of the Japan IgA Nephropathy Cohort Study (J-IGACS), a prospective, multicenter registry enrolling patients with primary IgAN across Japan between April 1, 2005, and August 31, 2015 [11, 12]. Eligibility required a biopsy-confirmed diagnosis of IgAN and a kidney biopsy specimen containing at least 10 glomeruli for primary study. The study population consisted of individuals from the primary cohort after excluding those with missing values in any covariates included in the multivariable adjustment. Ethical approval was granted by The Jikei University School of Medicine (approval no. 37-069 [12706]). This study was conducted in accordance with the Declaration of Helsinki and reported in compliance with STROBE guidelines.

Data collection and definitions

Clinical data were collected at the time of biopsy and subsequently at 6-month intervals, including age, sex, blood pressure, serum creatinine, estimated glomerular filtration rate (eGFR), serum uric acid, 24-hour urinary protein excretion (UPE) (the urinary protein-to-creatinine ratio [UPCR] was permitted in place of 24-hour urinary protein excretion at all time points except baseline), and urinary red blood cell count. eGFR was calculated using Uemura's equation [13] for individuals under 20 years of age and Matsuo's formula [14] thereafter. Treatments initiated within 1 year after biopsy were considered initial interventions and included corticosteroid therapy, tonsillectomy, and renin-angiotensin-aldosterone system (RAAS) inhibitor use, including any prescriptions ongoing at the time of biopsy. Kidney biopsy histopathological findings were scored using the Oxford MEST-C classification, based on standardized definitions for mesangial (M) and endocapillary (E)

hypercellularity, segmental sclerosis (S), tubular atrophy/interstitial fibrosis (T), and

crescents (C), in accordance with prior literature [5, 6].

Exposure and outcome measures

Proteinuria reduction was assessed using the proteinuria ratio, defined as the ratio of

proteinuria at 12 months to baseline proteinuria. It was categorized into tertiles or log-

transformed for analyses as needed. The primary endpoint was defined as either a ≥40%

decrease in eGFR or the initiation of kidney replacement therapy.

Statistical analysis

Baseline patient characteristics were presented as mean (standard deviation) or median

[interquartile range] for continuous variables, and as frequencies and proportions for

categorical variables. The Jonckheere–Terpstra test and Cochran–Armitage test were used to

evaluate trends in baseline characteristics across tertiles of the proteinuria ratio.

For kidney outcomes, the cumulative incidence of the composite endpoint, defined

as a 40% decline in eGFR or initiation of kidney replacement therapy, was estimated using

Kaplan–Meier survival curves with the log-rank test for trend across tertiles of proteinuria

ratio. Cox proportional hazards models were used to estimate hazard ratios (HRs) and 95%

confidence intervals (CIs) for the composite endpoint. These models were adjusted for

established risk factors, namely, age, sex, systolic blood pressure, baseline eGFR, baseline

log-transformed proteinuria, Oxford MEST-C scores, tonsillectomy, and initiation of RAAS

inhibitors and corticosteroids within one year after diagnosis. In addition, to explore the

association of continuous values of proteinuria ratio to HR of the composite endpoint, spline

regression models were also performed with the median value of the lowest tertile as the

reference point and knots at the 5th, 35th, 65th, and 95th percentiles of the proteinuria ratio

distribution. Additional analyses were conducted by restricting the study population to a

subgroup with baseline eGFR $\geq$30 and <120 mL/min/1.73 m^2 and baseline proteinuria $\geq$0.5

and <3.5 g/day, a range often considered clinically relevant in therapeutic research, to

evaluate the robustness of the main findings.

To assess longitudinal changes in kidney function, a linear mixed-effects model for

repeated measures was employed to estimate the annual eGFR slope based on the relationship

between follow-up time and eGFR levels, where a variance components matrix for the

random intercepts to account for individual variability and a first-order autoregressive

covariance structure for modeling within-subject correlation across repeated measures were

employed. The linearity between follow-up time and eGFR was visually validated using a

generalized additive model with smoothing functions.

A sensitivity analysis was performed using the proteinuria ratio at 6 months, a

shorter time period compared to the main analysis timeframe, as the explanatory variable.

A two-sided P-value <0.05 was considered statistically significant. All statistical

analyses, except for generalized additive model fitting, were performed using SAS version

9.4 (SAS Institute Inc., Cary, NC, USA). Generalized additive model analyses were

conducted using R version 4.5.0 (R Foundation for Statistical Computing, Vienna, Austria).

Results

Baseline characteristics and follow-up

After excluding individuals with missing values in any of the covariates from the original

cohort of 991 patients, a total of 793 patients with complete baseline variables were included

in the analysis. The proteinuria ratio, defined as the ratio of proteinuria at 12 months to

baseline proteinuria, was categorized into tertiles (Tertile 1, <0.194; Tertile 2, 0.194–0.625;

and Tertile 3, ≥0.625). Baseline characteristics showed that the proteinuria ratio was

significantly associated with several clinical parameters, including urinary protein excretion,

microscopic hematuria, the presence of E, S, and C lesions according to the Oxford

classification, initiation of steroid therapy within 1 year, and tonsillectomy within 1 year.

Association between initial proteinuria reduction and primary outcome

Kaplan–Meier analysis was used to analyze the associations between the kidney composite

endpoint, which was defined as a 40% decline in eGFR or initiation of kidney replacement

therapy, and the proteinuria ratio. The analysis demonstrated that higher proteinuria ratios

were significantly associated with an increased cumulative incidence of the composite

endpoint, as indicated by a significant trend in the log-rank test (P for trend <0.001; **Figure**

1). Moreover, in multivariable Cox proportional hazards models adjusted for age, sex,

systolic blood pressure, Oxford MEST-C scores, baseline eGFR, baseline log-transformed

proteinuria, tonsillectomy, initiation of RAAS inhibitor therapy, and initiation of steroid

therapy, higher proteinuria ratios remained independently associated with an elevated risk of

the incident composite endpoint (**Table 2**). Additional analyses using spline models, with the

median value of Tertile 1 used as the reference point, revealed that the risk of the incident

composite endpoint increased continuously as the proteinuria ratio rose above the median

value (**Figure 2**).

Restricted cohort analysis

To further assess the robustness of the findings, we performed an analysis limited to patients with baseline eGFR ≥30 and <120 mL/min/1.73 m^2 and baseline proteinuria ≥0.5 and <3.5 g/day, a range often considered relevant in clinical and research contexts. These analyses demonstrated results consistent with the main findings, and the association between a higher proteinuria ratio and adverse kidney outcomes appeared even stronger within this restricted cohort (**Figure 3**).

Association of initial proteinuria reduction and eGFR slope

A multivariable-adjusted mixed-effects model for repeated measures demonstrated that higher proteinuria ratios were significantly associated with a steeper decline in eGFR slope (P for trend <0.001; **Figure 4**).

Sensitivity analysis

As a sensitivity analysis, we conducted an additional analysis using the proteinuria ratio at 6 months, a shorter time period compared to the main analysis timeframe, as an explanatory variable. Both multivariable Cox proportional hazards models and mixed-effects models for repeated measures yielded significant associations consistent with the primary analyses, although the associations were attenuated (**Supplementary Table S1, Supplementary Figure S1**).

Discussion

In this post hoc analysis of a large-scale, nationwide, prospective cohort of biopsy-confirmed IgAN patients, we found that early reduction in proteinuria, quantified at 12 months after histopathological diagnosis using the proteinuria ratio, was independently and consistently associated with the long-term clinical endpoints, defined as a 40% or greater decline in eGFR or initiation of kidney replacement therapy. Moreover, early reduction in proteinuria was found to be consistently associated with the intermediate-term trajectories of kidney function decline assessed by the eGFR slope. The latter association remained significant even after multivariable adjustment for demographic, clinical, and histopathological covariates, and across sensitivity analyses. These results reinforce the association between early proteinuria response and long-term kidney outcomes, highlighting the potential of early proteinuria response as a clinically actionable, pathophysiologically relevant, and dynamically measurable surrogate endpoint. Incorporating proteinuria reduction into risk stratification algorithms and treatment response evaluation may enhance individualized clinical decision-making in the management of IgAN.

From a mechanistic perspective, proteinuria serves as a surrogate indicator of structural injury to the glomerular filtration barrier and reflects ongoing immunologically mediated inflammation [15, 16]. A reduction in proteinuria, therefore, is likely indicative of favorable modulation of underlying pathogenic pathways. Importantly, proteinuria possesses several practical advantages over other biomarkers: it is routinely quantified in clinical practice, exhibits prompt responsiveness to therapeutic intervention, and is readily interpretable at the point of care. These attributes collectively render it highly amenable to incorporation into dynamic, real-time clinical decision-making frameworks. Our data provide compelling evidence that, even within the constraints of a real-world, non-randomized setting, early amelioration of proteinuria is independently associated with long-term kidney

function preservation [7]. In addition to its established association with definitive clinical

endpoints, our analysis revealed that a lower proteinuria ratio was significantly correlated

with a more gradual subsequent decline in eGFR, consistent with previous meta-regression

analyses that evaluated the relationship between treatment-induced reductions in proteinuria

and improvements in eGFR decline [17, 18]. This observation supports the hypothesis that

proteinuria may function as an upstream surrogate marker, reflecting both functional

impairment and structural progression of kidney disease [19, 20]. These findings lend further

biological and clinical plausibility to the use of proteinuria as a comprehensive indicator of

disease activity and therapeutic response in IgAN.

From a clinical perspective, these findings offer practical and actionable insights for

the early stratification of patient risk in IgAN [7, 15, 21, 22]. Individuals who achieve a

marked reduction in proteinuria within the first year following diagnosis may be identified as

having a more favorable prognosis. This subgroup may be appropriate for de-escalation of

immunosuppressive regimens, reduction in monitoring frequency, and provision of

reassurance based on observed associations with favorable outcomes. In contrast, patients

demonstrating insufficient proteinuria reduction may warrant intensified therapeutic

strategies, closer clinical surveillance, or consideration for inclusion in interventional clinical

trials. Owing to its therapeutic modifiability, ease of measurement, and interpretability in

routine practice, proteinuria represents a particularly valuable parameter for dynamic risk

assessment and response-adaptive treatment strategies, principles that lie at the core of

precision nephrology.

To further validate the clinical applicability of our findings, we conducted an

additional analysis restricted to patients with baseline proteinuria between 0.5 and 3.5 g/day,

a range often considered clinically relevant in therapeutic contexts [23]. This subgroup

reflects a population at moderate to high risk, in whom treatment decisions are frequently

guided by proteinuria levels [24]. Additionally, we performed a subgroup analysis limited to patients with baseline eGFR between 30 and 120 mL/min/1.73 m^2, representing a population for whom kidney function is preserved enough to permit intervention. The consistent and even strengthened associations observed in both subgroups support the robustness of proteinuria reduction as a meaningful marker and highlight its significance for clinical trial design and real-world decision-making in IgAN.

Several limitations of this study should be acknowledged. First, although the cohort was prospectively assembled, this was a *post hoc* analysis and is therefore subject to inherent limitations of observational research, including potential residual confounding. Second, proteinuria measurements were obtained according to institutional protocols rather than centralized standardization, potentially introducing inter-site variability. Third, the 12-month window used for assessing proteinuria reduction, while clinically pragmatic, may not fully capture heterogeneity in early treatment responses [25]. However, our sensitivity analysis using the 6-month proteinuria ratio yielded consistent results, suggesting that evaluation at 6 to 12 months may be acceptable in clinical and research settings. Lastly, our cohort consisted exclusively of Japanese patients. While this homogeneity offers unique insights into a high-prevalence population [26], it may limit the generalizability of our findings to other ethnic and geographic groups with differing genetic predispositions, environmental exposures, and healthcare systems.

In conclusion, early reduction in proteinuria serves as a reliable, pathophysiologically meaningful, and clinically applicable marker associated with both long-term and intermediate kidney outcomes in patients with IgAN. Its availability, responsiveness to therapy, and consistent association with clinical outcomes highlight its value as a surrogate endpoint in both clinical practice and clinical research. Future studies should aim to refine

1   optimal thresholds, examine disease- and treatment-specific responsiveness, and validate

2   these findings across more diverse populations and settings.

References

1. Wyatt, R.J. and B.A. Julian, *IgA nephropathy.* N Engl J Med, 2013. **368**(25): p. 2402-14.
2. Moriyama, T., et al., *Prognosis in IgA nephropathy: 30-year analysis of 1,012 patients at a single center in Japan.* PLoS One, 2014. **9**(3): p. e91756.
3. Koyama, A., M. Igarashi, and M. Kobayashi, *Natural history and risk factors for immunoglobulin A nephropathy in Japan. Research Group on Progressive Renal Diseases.* Am J Kidney Dis, 1997. **29**(4): p. 526-32.
4. D'Amico, G., *Natural history of idiopathic IgA nephropathy and factors predictive of disease outcome.* Semin Nephrol, 2004. **24**(3): p. 179-96.
5. Haas, M., et al., *A Multicenter Study of the Predictive Value of Crescents in IgA Nephropathy.* J Am Soc Nephrol, 2017. **28**(2): p. 691-701.
6. Trimarchi, H., et al., *Oxford Classification of IgA nephropathy 2016: an update from the IgA Nephropathy Classification Working Group.* Kidney Int, 2017. **91**(5): p. 1014-1021.
7. Reich, H.N., et al., *Remission of proteinuria improves prognosis in IgA nephropathy.* J Am Soc Nephrol, 2007. **18**(12): p. 3177-83.
8. Philibert, D. and D. Cattran, *Remission of proteinuria in primary glomerulonephritis: we know the goal but do we know the price?* Nat Clin Pract Nephrol, 2008. **4**(10): p. 550-9.
9. Mathur, M., et al., *A Phase 2 Trial of Sibeprenlimab in Patients with IgA Nephropathy.* N Engl J Med, 2024. **390**(1): p. 20-31.
10. Perkovic, V., et al., *Alternative Complement Pathway Inhibition with Iptacopan in IgA Nephropathy.* N Engl J Med, 2025. **392**(6): p. 531-543.
11. Kawamura, T., et al., *Associations of corticosteroid therapy and tonsillectomy with kidney survival in a multicenter prospective study for IgA nephropathy.* Sci Rep, 2023. **13**(1): p. 18455.
12. Koike, K., et al., *Clinicopathological prognostic stratification for proteinuria and kidney survival in IgA nephropathy: a Japanese prospective cohort study.* Clin Kidney J, 2024. **17**(1): p. sfad294.
13. Uemura, O., et al., *Creatinine-based equation to estimate the glomerular filtration rate in Japanese children and adolescents with chronic kidney disease.* Clin Exp Nephrol, 2014. **18**(4): p. 626-33.
14. Matsuo, S., et al., *Revised equations for estimated GFR from serum creatinine in Japan.* Am J Kidney Dis, 2009. **53**(6): p. 982-92.
15. El Karoui, K., F.C. Fervenza, and A.S. De Vriese, *Treatment of IgA Nephropathy: A Rapidly Evolving Field.* J Am Soc Nephrol, 2024. **35**(1): p. 103-116.
16. Daehn, I.S. and J.S. Duffield, *The glomerular filtration barrier: a structural target for novel kidney therapies.* Nat Rev Drug Discov, 2021. **20**(10): p. 770-788.
17. Inker, L.A., et al., *Early Change in Urine Protein as a Surrogate End Point in Studies of IgA Nephropathy: An Individual-Patient Meta-analysis.* Am J Kidney Dis, 2016. **68**(3): p. 392-401.
18. Inker, L.A., et al., *Association of Treatment Effects on Early Change in Urine Protein and Treatment Effects on GFR Slope in IgA Nephropathy: An Individual Participant Meta-analysis.* Am J Kidney Dis, 2021. **78**(3): p. 340-349 e1.
19. Tryggvason, K. and E. Pettersson, *Causes and consequences of proteinuria: the kidney filtration barrier and progressive renal failure.* J Intern Med, 2003. **254**(3): p. 216-24.

20. Remuzzi, G., A. Benigni, and A. Remuzzi, *Mechanisms of progression and regression of renal lesions of chronic nephropathies and diabetes.* J Clin Invest, 2006. **116**(2): p. 288-96.

21. Bartosik, L.P., et al., *Predicting progression in IgA nephropathy.* Am J Kidney Dis, 2001. **38**(4): p. 728-35.

22. Barbour, S.J., et al., *Evaluating a New International Risk-Prediction Tool in IgA Nephropathy.* JAMA Intern Med, 2019. **179**(7): p. 942-952.

23. Shen, X., et al., *Long-term outcomes of IgA nephropathy in China.* Nephrol Dial Transplant, 2024.

24. Okonogi, H., et al., *A grading system that predicts the risk of dialysis induction in IgA nephropathy patients based on the combination of the clinical and histological severity.* Clin Exp Nephrol, 2019. **23**(1): p. 16-25.

25. Canney, M., et al., *Quantifying Duration of Proteinuria Remission and Association with Clinical Outcome in IgA Nephropathy.* J Am Soc Nephrol, 2021. **32**(2): p. 436-447.

26. Ghaddar, M., M. Canney, and S.J. Barbour, *IgA Nephropathy: Epidemiology and Disease Risk Across the World.* Semin Nephrol, 2024. **44**(5): p. 151564.

Authors' Contributions

Conceptualization: Takaya Sasaki, Nobuo Tsuboi, Takashi Yokoo, Yusuke Suzuki.

Data curation: Takaya Sasaki.

Formal analysis: Takaya Sasaki.

Funding acquisition: Yusuke Suzuki.

Investigation: Takaya Sasaki, Nobuo Tsuboi.

Methodology: Takaya Sasaki.

Project administration: Nobuo Tsuboi, Takashi Yokoo, Yusuke Suzuki.

Supervision: Takashi Yokoo, Yusuke Suzuki.

Visualization: Takaya Sasaki.

Writing – original draft: Takaya Sasaki, Nobuo Tsuboi.

Writing – review & editing: Kentaro Koike, Hiroyuki Ueda, Masahiro Okabe, Shinya Yokote, Akihiro Shimizu, Keita Hirano, Tetsuya Kawamura, Takashi Yokoo, Yusuke Suzuki.

Funding

This study was partly supported by a Grant-in-Aid for Progressive Renal Diseases Research, Research on Rare and Intractable Disease, from the Ministry of Health, Labour and Welfare of Japan. This research was supported by the Japan Agency for Medical Research and Development under grant JP19ek0109261.

Financial Disclosure

All the authors have no competing interest in regard to this study.

Acknowledgments

1 This study was approved by the ethics review board of the Jikei University School of

2 Medicine (37-069[12706]).

4 **Data availability statement**

5 All data produced in the present study are available upon reasonable request to the authors.

Figure legends

Figure 1. Kaplan–Meier curves for the kidney composite endpoint according to

proteinuria ratio tertiles.

Kaplan–Meier survival curves show the cumulative incidence of the kidney composite

endpoint, defined as a 40% decline in estimated glomerular filtration rate or initiation of

kidney replacement therapy, stratified by tertiles of the proteinuria ratio at 12 months (Tertile

1,<0.194; Tertile 2, 0.194–0.625; Tertile 3, ≥0.625).

Figure 2. Adjusted hazard ratios for the kidney composite endpoint by proteinuria ratio

tertiles.

Forest plot displaying adjusted hazard ratios for the kidney composite endpoint according to

proteinuria ratio tertiles, using a multivariable Cox proportional hazards model. Adjustment

variables included age, sex, systolic blood pressure, initiation of renin-angiotensin-

aldosterone inhibitor therapy, initiation of steroid therapy, tonsillectomy, Oxford MEST-C*

scores, baseline estimated glomerular filtration rate, and log-transformed proteinuria. The risk

of adverse outcomes increased with higher proteinuria ratios.

*Oxford MEST-C indicates mesangial hypercellularity, endocapillary hypercellularity,

segmental sclerosis, tubular atrophy/interstitial fibrosis, and cellular/fibrocellular crescents.

Figure 3. Association between continuous proteinuria ratio and hazard ratio (HR) of the

kidney composite endpoint using spline models.

Spline curves depict the continuous relationship between the proteinuria ratio at 12 months

and the multivariable-adjusted HR for the kidney composite endpoint, where the median

proteinuria ratio of the 1st tertile was treated as the reference point and the 5th, 35th, 65th,

and 95th points were used as knots. The model was adjusted for age, sex, systolic blood

pressure, and initiation of renin-angiotensin-aldosterone system inhibitors or corticosteroids

within one year after diagnosis, tonsillectomy, baseline estimated glomerular filtration rate,

baseline log-transformed proteinuria, and Oxford MEST-C scores*.

*Oxford MEST-C indicates mesangial hypercellularity, endocapillary hypercellularity,

segmental sclerosis, tubular atrophy/interstitial fibrosis, and cellular/fibrocellular crescents.

Figure 4. Annual slope of the estimated glomerular filtration rate (eGFR) according to

proteinuria ratio tertiles.

The association between eGFR levels and follow-up time was analyzed using a mixed-effects

model for repeated measures across proteinuria ratio tertiles (Tertile 1, <0.194; Tertile 2,

0.194–0.625; Tertile 3, ≥0.625). The model was adjusted for age, sex, systolic blood pressure,

and initiation of renin-angiotensin-aldosterone system inhibitors or corticosteroids within one

year after diagnosis, tonsillectomy, baseline estimated glomerular filtration rate, baseline log-

transformed proteinuria, and Oxford MEST-C scores*.

* Oxford MEST-C indicates mesangial hypercellularity, endocapillary hypercellularity,

segmental sclerosis, tubular atrophy/interstitial fibrosis, and cellular/fibrocellular crescents.

Table 1. Baseline characteristics and initial treatment according to proteinuria ratio tertiles.

	Proteinuria ratio			
	Tertile 1 (<0.194) n = 264	Tertile 2 (0.194–0.625) n = 265	Tertile 3 (≥0.625) n = 264	P for trend
Clinical characteristics				
Age, year	37.9 (16.6)	38.9 (16.0)	40.2 (17.0)	0.13
Male sex, n (%)	140 (53.0)	138 (52.1)	129 (48.5)	0.30
Height, cm	161.2 (12.5)	161.9 (11.0)	162.4 (10.1)	0.54
Weight, kg	58.5 (14.5)	59.1 (13.9)	59.1 (12.6)	0.70
Systolic blood pressure, mmHg	122.0 (18.0)	121.4 (17.4)	123.2 (17.7)	0.37
eGFR, mL/min/1.73 m^2	75.5 (29.2)	75.4 (26.9)	76.3 (29.4)	0.46
UPE, g/day	1.1 [0.5–2.1]	0.6 [0.4–1.1]	0.4 [0.2–0.8]	<0.001
Microhematuria*, n %	239 (90.5)	226 (85.3)	208 (78.8)	<0.001
Pathological characteristics†				
M1, n (%)	93 (35.2)	70 (26.4)	68 (25.8)	0.017
E1, n (%)	125 (47.4)	94 (35.5)	69 (26.1)	<0.001
S1, n (%)	216 (81.8)	203 (76.6)	177 (67.1)	<0.001
T1 or T2, n (%)	62 (23.5)	50 (19.3)	65 (24.2)	0.83
C1 or C2, n (%)	150 (56.8)	94 (35.5)	67 (25.4)	<0.001
Initial treatment				
RAAS inhibitor, n (%)	155 (58.7)	154 (58.1)	165 (62.5)	0.37
Corticosteroid, n (%)	222 (84.1)	190 (71.7)	114 (42.8)	<0.001
Tonsillectomy, n (%)	138 (39.4)	121 (45.7)	91 (34.5)	<0.001

Abbreviations: eGFR, estimated glomerular filtration rate; RAAS, renin-angiotensin-aldosterone system; UPE, urinary protein excretion.

*Defined as 5 or more red blood cells / high-power field

†Indicates Oxford MEST-C scores: M, mesangial hypercellularity; E, endocapillary hypercellularity; S, segmental sclerosis; T, tubular atrophy/interstitial fibrosis; and C, cellular/fibrocellular crescents.

Table 2. Hazard ratios (HRs) for kidney composite endpoint according to proteinuria ratio tertiles.

Tertile	No. of outcomes	No. of participants	Age- and sex-adjusted		Multivariable adjusted*	
			HR (95% CI)	P for trend	HR (95% CI)	P for trend
Tertile 1 (<0.194)	48	264	1 (ref)		1 (ref)	
Tertile 2 (0.194–0.625)	43	265	0.93 (0.61–1.40)	<0.001	1.00 (0.65–1.54)	0.021
Tertile 3 (≥0.625)	85	264	1.65 (1.30–2.63)		1.65 (1.09–2.52)	

Abbreviation: CI, confidence interval.

*Adjusted for age, sex, systolic blood pressure, baseline estimated glomerular filtration rate, baseline log-transformed proteinuria, and Oxford MEST-C scores†, initiation of renin-angiotensin-aldosterone system inhibitors and corticosteroids within one year after diagnosis, tonsillectomy.

†Oxford MEST-C indicates mesangial hypercellularity, endocapillary hypercellularity, segmental sclerosis, tubular atrophy/interstitial fibrosis, cellular/fibrocellular crescents.

Figure 1

Figure 2

Figure 3

Figure 4