Practice effects persist over two decades of cognitive testing: Implications for longitudinal research

Jeremy A. Elman[a,b], Erik Buchholz[a,b], Rouhui Chen[c], Mark Sanderson-Cimino[d], Tyler R. Bell[a,b], Nathan Whitsel[a,b], Katherine J. Bangen[a,e], Alice Cronin-Golomb[f], Anders M. Dale[g,h], Lisa T. Eyler[a], Nathan A. Gillespie[i], Eric L. Granholm[a], Daniel E. Gustavson[j], Donald J. Hagler Jr.[g], Richard L. Hauger[a,b,k], Diane M. Jacobs[h], Amy J. Jak[a], Mark W. Logue[l,m,n,o], Ruth E. McKenzie[p], Michael C. Neale[i], Robert A. Rissman[q], Rosemary Toomey[f], Arthur Wingfield[r], Hong Xian[s,t], Christine Fennema-Notestine[a,b], Carol E. Franz[a,b], Michael J. Lyons[f], Chandra A. Reynolds[j], Xin M. Tu[u], William S. Kremen[a,b,*], and Matthew S. Panizzon[a,b,*]

* Joint senior authors
[a] Department of Psychiatry, University of California San Diego, La Jolla, CA, USA
[b] Center for Behavior Genetics of Aging, University of California San Diego, La Jolla, CA, USA
[c] Department of Preventive Medicine, Feinberg School of Medicine, Northwestern University, Evanston, IL, USA
[d] Memory and Aging Center, Weill Institute for Neurosciences, San Francisco, CA, USA,
[e] VA San Diego Healthcare System, San Diego, CA, USA
[f] Department of Psychological and Brain Sciences, Boston University, Boston, MA, USA
[g] J. Craig Venter Institute, La Jolla, CA, USA
[h] Department of Neurosciences, University of California San Diego, La Jolla, CA, USA
[i] Virginia Institute for Psychiatric and Behavior Genetics, Richmond, Virginia, USA.
[j] Institute for Behavioral Genetics and Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder, CO, USA,
[k] Center of Excellence for Stress and Mental Health (CESAMH), VA San Diego Healthcare System, San Diego, CA, USA
[l] Department of Psychiatry, Boston University School of Medicine, Boston, MA, USA.
[m] Biomedical Genetics, Boston University School of Medicine, Boston, MA, USA.
[n] Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA.
[o] Department of Psychological and Brain Sciences, Boston University, Boston, MA, USA.
[p] School of Education and Social Policy, Applied Human Development and Community Studies, Merrimack College, North Andover, MA, USA
[q] Department of Physiology and Neuroscience, Alzheimer's Therapeutic Research Institute of the Keck School of Medicine of the University of Southern California, San Diego, CA, USA
[r] Department of Psychology and Volen National Center for Complex Systems, Brandeis University, Waltham, MA, USA
[s] Department of Epidemiology and Biostatistics, Saint. Louis University, St. Louis, Missouri, USA
[t] Research Service, VA St. Louis Healthcare System, St. Louis, Missouri, USA
[u] Herbert Wertheim School of Public Health & Human Longevity Science, University of California San Diego, La Jolla, CA, USA

Correspondence:
Jeremy A. Elman
Department of Psychiatry
University of California, San Diego
9500 Gilman Dr. (MC0738)
La Jolla, CA 92093
Phone: (858) 534-6842
E-mail: jaelman@health.ucsd.edu

ABSTRACT

INTRODUCTION: Repeated cognitive testing can boost scores due to practice effects (PEs). It remains unclear whether PEs persist across multiple follow-ups and long durations. We examined PEs across multiple assessments from midlife to old age in a nonclinical sample.

METHOD: Men (N=1,608) in the Vietnam Era Twin Study of Aging (VETSA) underwent neuropsychological assessment across 4 waves from mean age 56 to 74. We leveraged age-matched attrition-replacement (AR) participants to estimate PEs at each wave. We compared cognitive trajectories and prevalence of mild cognitive impairment (MCI) using unadjusted versus PE-adjusted scores.

RESULTS: Across follow-ups, a range of 7-12 out of 30 measures demonstrated significant PEs, especially in episodic memory and visuospatial domains. Adjusting for PEs resulted in steeper cognitive decline with up to 29% higher MCI prevalence.

DISCUSSION: PEs persist across multiple assessments and decades. The AR-participant method provides accurate sample-specific PE estimates that enable significantly earlier detection of MCI.

1. INTRODUCTION

Longitudinal designs are critical for understanding cognitive development and decline [1,2]. In the context of studies on Alzheimer's disease and Alzheimer's disease-related dementias (AD/ADRD), the decades-long pathological process and impact of risk factors across the lifespan further underscores the need for long-term longitudinal studies to identify modifiable risk factors, understand variation in disease progression, and evaluate efficacy of treatments [3,4]. However, it has long been acknowledged that performance at follow-up may be artifactually inflated due to practice effects (PEs) [5-7]. Nevertheless, accounting for PEs is not standard practice, which can obscure the true nature of cognitive trajectories and has implications for aging research and AD/ADRD clinical trials [8-10].

PEs are often defined as improved performance at follow-up compared to baseline [11,12]. However, even with observed decline, PEs may still be present when age- or disease-related declines are greater than the magnitude of the PEs [13]. One solution implemented by Rönnlund et al. [14] is to use attrition replacement (AR) participants, new participants recruited from the same population during follow-up study waves who are age-matched to the ongoing longitudinal cohort. This approach is conceptually similar to a randomized controlled trial. By comparing well-matched groups drawn from the same population who differ only on whether they have previously been tested, we can estimate the expected improvement (at the group level) due to practice. Critically, because both groups should experience similar levels of age-related normative decline, we can estimate PEs even when there is an observed *decrease* in scores at follow-up for returnees. This method allows us to create an adjusted follow-up score that can be more appropriately compared to norms or thresholds for impairment, which assume that test scores are unaffected by practice. We have previously shown that using PE-adjusted scores results in earlier detection of mild cognitive impairment (MCI) [13,15,16].

Studies of PEs have typically examined test-retest intervals of less than 5 years, with most being in the range of 6 months to 2 years [17,18]. There has been little investigation of PEs in

cohorts that have followed individuals for extended periods of time (i.e., over 10 years, but see [19] and [20] for notable exceptions). Extended longitudinal follow-ups that account for PEs are necessary to clarify cognitive trajectories across critical transition periods such as midlife to old age, especially with increased recognition that disease processes can begin decades prior to clinical onset. The Vietnam Era Twin Study of Aging (VETSA) presents a rare opportunity to examine PEs on cognitive performance in ~1,600 individuals assessed over 4 study waves. Here, we extend the method [14] that we previously applied to two waves of VETSA data [15] using a generalized approach that more flexibly handles complex testing schedules and missingness patterns. We examined how PEs evolve over two decades and compared cognitive trajectories and MCI prevalence using unadjusted versus PE-adjusted scores.

2. METHODS

2.1. Participants

Participants were 1,608 individuals tested at one or more of the 4 completed VETSA study waves (**Table 1**). VETSA is an on-going longitudinal study of cognitive and brain aging beginning in middle age [21-23]. Participants were members of the Vietnam Era Twin Registry, a national, community-dwelling sample of male-male twins who served in the U.S. military during the Vietnam era (1965-1975) [24]. All Registry members were invited to participate in the Harvard Drug Study [25], for which ascertainment was not based on any diagnostic or substance use criteria. VETSA participants were then randomly recruited from the Harvard Drug Study sample. At baseline, VETSA participants were similar to American men in their age cohort with respect to health, education, and lifestyle characteristics based on Center for Disease Control and Prevention data [26], and nearly 80% reported no combat exposure during their military service [27].

Of the 1,608 individuals in the current study, 1,291 were enrolled and tested during the wave 1 baseline assessment. At waves 2 and 3, attrition replacement participants age-matched to the ongoing sample were recruited from the VET Registry and tested for the first time (wave 2

n=193; wave 3 n=124). These participants were then invited for follow-up at all subsequent

waves (see **Supplemental Figure S1A** for all patterns of assessments). On average,

participants were 56 years of age (range 51-61) at wave 1, 62 years (range 56-67) at wave 2,

68 years (range 61-73) at wave 3, and 74 (range 67-79) at wave 4 (see **Supplemental Figure**

S1B for age distributions by wave). The average time between wave 1 and 2 was 5.7 years,

with 5.9 years between waves 2 and 3, and 5.6 years between waves 3 and 4.

The study was performed in accordance with the ethical standards per the 1964 Declaration

of Helsinki and later amendments. Informed consent was obtained from all participants and

institutional review boards at both sites approved all study procedures.

2.2. Cognitive tests and measures

Cognitive measures included in the current analysis were a set of 30 component scores

from neuropsychological tests covering multiple cognitive domains (see **Table 2** and

Supplemental Material for full list of tests and measures). For consistency, scores from tests

where lower values indicate better performance (e.g., reaction time) were reverse-coded so that

higher scores uniformly indicate better performance. The same tests were assessed at all

waves with 2 exceptions: spatial span (waves 1, 2, and 3 only) and Boston Naming Test (waves

3 and 4 only). Therefore, all 30 measures have 1 follow-up, with 29 and 28 measures having

two and three follow-ups, respectively. Additionally, young adult general cognitive ability (GCA)

was assessed with the validated Armed Forces Qualification Test (AFQT) [28], which is highly

correlated with standard IQ scales (r=.84) [29]. AFQT was administered during military induction

at average age 20 (hereafter referred to as "age 20 GCA").

2.3. Estimation of practice effects

PEs were estimated for each of the 30 measures using generalized estimating equations

(GEE). We opted for this class of semiparametric regression models because it requires no

assumptions about data distributions, such as normality, to provide valid inferences for virtually

all data distributions arising in practice [30].

Let Y_{iaw} denote the score on a cognitive measure at the a^{th} assessment of subject i occurring at wave w:

$$\begin{aligned}
Y_{\text{iaw}} = \beta_0 &+ \beta_{age}(age) + \beta_{afqt}(age\ 20\ AFQT) + \beta_{s1}I(s = 1) + \beta_{s2}I(s = 2) + \beta_1I(w = 2) \\
&+ \beta_2I(w = 3) + \beta_3I(w = 4) + \beta_4I(w = 2\ \&\ a = 2) + \beta_5I(w = 3\ \&\ a = 2) \\
&+ \beta_6I(w = 3\ \&\ a = 3) + \beta_7I(w = 4\ \&\ a = 3) + \beta_8I(w = 4\ \&\ a = 4)
\end{aligned}$$

Here, $I(\cdot)$ denotes an indicator function that takes the value 1 if the condition inside is true, and 0 otherwise. Wave w and Assessment a can take values from 1 to 4. Importantly, the inclusion of AR participants means there are groups of individuals at each follow-up wave that differ in number of prior assessments. Therefore, additional indicator variables capture the interaction between Wave and Assessment (with the necessary condition that $a \leq w$). Participants may miss a study wave but then return for follow-up at subsequent waves. To account for the potential impact of longer intervals between assessments on PEs, we include the Skip variable s that can take the values 1 or 2 to indicate whether they missed 1 or 2 waves immediately prior to the current assessment (only missed assessments after enrollment are counted).

We additionally adjusted for age and age 20 GCA. Adjusting for age allows us to estimate PEs independent of normative age-related decline. Previously, we found that the attrition replacement group at Wave 2 had significantly lower young adult (age 20) GCA scores than the group enrolled at Wave 1, which could result in artificially inflated PEs. Given that participants were randomly recruited from the same population (i.e., the VET Registry), this is likely due to random sampling variation. However, this adjustment helps account for potential long-standing differences in performance. Details on interpretation of each model coefficient are provided in **Supplemental Methods** and **Supplemental Table S1.**

2.4. Adjusting test scores for practice effects

For a given individual, we can calculate the expected PE for a measure by taking the coefficients corresponding to the relevant wave and assessment number, and whether they have skipped any previous assessments. The resulting "adjustment value" is then subtracted

from their observed score. This results in a score that we would expect if they had been taking the test for the first time. As an example, consider two individuals at Wave 4. The first has completed all four waves of testing, so we can adjust their score on a given measure at Wave 4 by subtracting the value of β_8 (i.e., the PE for someone completing a 4^{th} assessment at Wave 4) from their observed score. The other individual missed Wave 3, so it would be their third assessment. For this person, we would instead use β_7 and add to that the estimated effect of skipping one prior assessment (i.e., β_{s1}). See **Supplemental Methods** for further discussion of selecting coefficients for practice effect adjustment.

2.5. Cognitive factor score composites

We have previously shown that composites such as cognitive factor scores can improve reliability and prediction of cognitive decline [31]. Therefore, we were interested in understanding how adjusting individual scores could impact cognitive composites. We calculated composites for 5 cognitive abilities from both adjusted and unadjusted scores: episodic memory, executive function, verbal fluency, processing speed, and visuospatial ability. Details of the factor models from which these composites were derived are described in previous publications [31-34] and in **Supplemental Methods**. Higher values reflect better performance in each domain. We used paired t-tests to compare cognitive composites calculated from adjusted and unadjusted scores at each wave.

2.6. Classification of mild cognitive impairment

Classification of MCI was compared both before and after adjusting for PEs. We defined MCI according to the Jak/Bondi approach as described previously [35,36] and in **Supplemental Methods**. To ensure that MCI classification captured cognitive decline rather than lifelong low ability, neuropsychological scores were adjusted using early adult GCA (age 20 AFQT) as a covariate. Impairment was defined as having 2+ measures within a domain >1.5 SD below age-based normative means. Individuals with an impaired memory domain were classified as amnestic MCI (aMCI), and those with impairments in domains other than memory were

classified as non-amnestic MCI (naMCI). Differences in MCI classification based on adjusted or unadjusted scores were assessed at each wave with McNemar's χ^2 test with 1 degree of freedom.

3. RESULTS

3.1. Practice effects on cognitive tests across 4 study waves.

Models included all data and patterns of assessments, but here we focus on reporting estimates for individuals who attended all 4 waves. This was the most common pattern of participation and allowed us to examine the evolution of PEs over the longest follow-up period and greatest number of assessments. **Figure 1** presents PE estimates and confidence intervals for each of the 3 follow-up assessments for those that completed all waves. The PE estimates can be interpreted as the expected boost in performance an individual receives from having taken the test a given number of times previously compared to someone of a similar age and young adult GCA taking the test for the first time. At the first follow-up, 12 of 30 measures demonstrated significant PEs ranging in magnitude from 0.14 SD units to 0.29 SD units. At the second follow-up, 7 of 29 measures demonstrated significant PEs ranging from 0.16 SD units to 0.34 SD units. At the third follow-up, 8 of the 28 measures demonstrated significant PEs ranging from 0.23 SD units to 0.35 SD units. PEs. Although the number of significant PEs was smaller at later follow-ups, this is likely due to smaller sample size at these visits. Visuospatial and episodic memory measures consistently showed the strongest effects, and measures that showed significant PEs at later follow-ups typically, but not always, showed significant PEs at earlier follow-ups.

3.2. Impact on cognitive factor scores.

PE-adjustment led to significantly lower composite scores across all follow-up waves for all domains (all ps < 0.05), indicating that unadjusted scores can overestimate cognitive performance in these domains. Consistent with results from individual test measures,

differences in performance were most notable for composites of episodic memory (ranging between -0.21 to -0.25 SD units) and visuospatial ability (ranging between -0.19 to -0.22 SD units), and were weakest in the fluency domain (ranging between -0.04 to -0.07 SD units). See **Figure 2** and **Supplemental Table S3** for full results of comparisons across waves.

3.3. Impact on classification of MCI

We next examined the impact of adjusting for PEs on the rate of MCI at follow-up assessments (**Figure 3**). Adjusting for PEs significantly increased the prevalence of MCI at all follow-up waves (wave 2: 12.3% vs. 15.6%, χ^2=32.60, p<0.001; wave 3: 15.1% vs. 18.1%, χ^2=23.67, p<0.001; wave 4: 16.3% vs. 21.0%, χ^2=33.23, p<0.001). These increases suggest that failure to adjust for PEs masks clinically meaningful cognitive decline, particularly in long-term follow-up. Consistent with findings that measures in the memory domains exhibited the strongest PEs, the increased rates of MCI were largest for amnestic MCI. After PE adjustment, the rate of amnestic MCI was significantly higher at all follow-up waves (wave 2: 8.5% vs 11.4%, χ^2=26.036, p<0.001; wave 3: 9.6% vs 13.2%, χ^2=25.289, p<0.001; wave 4: 12.6% vs 16.6%, χ^2=22.042, p<0.001). Rates of non-amnestic MCI were significantly higher at waves 2 and 4, but not wave 3 (wave 2: 4.6% vs 5.2%, χ^2=5.786, p=0.016; wave 3: 6.6% vs 6.5%, χ^2=0.125, p=0.724; wave 4: 4.9% vs 6.3%, χ^2=9.600, p=0.002).

4. DISCUSSION

These results demonstrate the persistence of PEs on multiple cognitive measures over extended periods of time and multiple assessments. Our findings are broadly consistent with other studies in that PEs were most significant at first follow-up with fewer significant PEs at later follow-ups [12,19,37-40]. However, this was not due to a consistent decrease in effects sizes and may be explained by smaller sample sizes at these waves, thus PEs at later follow-ups should not be discounted. Importantly, even when PEs were modest and non-significant, we found that they could have meaningful impacts on downstream analyses. Adjusting for PEs

revealed significantly greater decline in cognitive trajectories as well as increased rates—and hence, earlier detection—of MCI at all waves. Here, we address 5 issues that are relevant to the present findings and are frequently raised in regard to PEs.

First, which domains are most susceptible to PEs and how does the magnitude of PEs compare with other studies? Consistent with prior studies, we found PEs were most apparent and most persistent in the memory domain, whereas fluency, executive function, and speed showed the smaller effects [11,12,17,37,41]. We additionally found strong PEs in the visuospatial domain, which is consistent with some studies [12] but not others [17,37]. However, direct comparisons across studies may not be entirely appropriate. PEs vary depending on the specific tests used, length of retest interval, number of follow-ups, and participant characteristics such as general cognitive ability, age, sex, race/ethnicity, diagnostic status, and presence of pathology [11,37,42,43]. Underscoring this point, we found that the magnitude of PEs differed across individual measures within the same domain. Because estimates from one study may not apply to another, it is ideal to estimate study-specific PEs. Estimating PEs using an AR-participant approach builds in a mechanism for deriving valid estimates that are specific to the sample characteristics (e.g., age and other demographics) and study design (e.g., test battery, retest interval).

Second, how long do practice effects last? Our results provide evidence of PEs across testing intervals of at least 5-6 years. Although this testing interval is comparatively long, other studies have found evidence of PEs after 7 years [14,17,19,44]. Regarding how PEs persist across number of follow-ups, our study provides evidence that PEs continued to influence performance through Wave 4. One of the few studies examining PEs for multiple assessments over a comparable period of time also found PEs on an intelligence test up to the fourth wave [19,40]. It should be noted that most prior studies finding a lack of PEs at follow-up did not use AR-participant approaches, which may underestimate or fail to detect PEs (i.e., they may be obscured by greater decline at later timepoints). Moreover, performance on certain tests may

have plateaued at later follow-ups as participant performance reached ceiling on the administered tests, preventing any further practice-related boosts [12,37-39].

Third, what are the mechanisms underlying PEs? There are likely to be multiple mechanisms, with varying contributions depending on the test. Participants may explicitly remember certain content from prior assessments (e.g., aspects of the story given during Logical Memory), which aids their performance. Yet on some tests such as Digit Span, it is highly unlikely participants are remembering specific sequences of digits. In these scenarios, PEs may be driven by greater familiarity with the *context.* This could include reduced test anxiety due to familiarity with the testing environment, procedural memory for tasks with a motor component, or identifying effective test-taking strategies. These context-related factors may explain why even individuals with severe episodic memory impairment can sometimes still benefit from practice [45]. It is worth noting, however, that the AR-based approach used here to estimate and adjust for PEs here does not hinge on knowing their underlying causes, but quantifies their aggregate effect.

Fourth, how do PEs affect detection of MCI? We found that adjusting for PEs resulted in significant increases in the rate of MCI at all waves, driven primarily by increases in amnestic MCI. This supports our prior findings that not accounting for PEs underestimates or delays diagnosis of MCI [13,15,16,46]. As shown elsewhere, accounting for PEs to identify MCI earlier can reduce the required sample size of clinical trials that use progression to impairment as an endpoint, resulting in multi-million dollar cost savings and allowing for more accurate estimate of treatment effects [8,9,16]. Of course, adjusting scores downward necessarily means that more people will be below the impairment threshold, but in an independent sample we showed that PE-adjusted diagnoses resulted in lower rates of reversion and increased concordance with biomarker-positivity [16,46], providing evidence that the increased prevalence of MCI was not driven by false positive diagnoses. Finally, earlier detection is critical to enabling early intervention, which is likely to improve treatment effectiveness.

Fifth, how does the AR-participant approach compare with other PE methods? We utilized an AR-based approach to estimate group-level PEs representing the average increase in score expected from having taken a given test once or several previous times. Our study extends prior AR-based methods [14-16,46,47] to better handle larger numbers of follow-up assessments and patterns of missingness. Some approaches employ multiple tests within short time frames (e.g., 1 week test-retest interval) [18]. Although bearing a similar name, these short-term PE studies have a different purpose than our approach, namely, for prognosis or predicting future decline. Other methods, such as the Reliable Change Index (RCI) [48] or Standardized Regression Based (SRB) change indices [49], focus on analysis of *change scores*. In contrast, our approach focuses on obtaining *adjusted follow-up scores* that, when compared to norm-based thresholds, enable earlier identification of impairment. Despite the shared use of the term "practice effects," these approaches are therefore not comparable nor are they in competition as they have different goals. Importantly, the AR-based approach can disentangle PEs from normative decline, which avoids underestimation of PEs. Moreover, the estimates are study- and sample-specific, which enable earlier and more accurate identification of MCI.

We note some limitations of the current study. The VETSA is an all-male sample and primarily non-Hispanic White, limiting its generalizability to other samples. Similar results have been found over 1- to 5-year intervals in mixed-sex samples [14,16,46], but as noted, the AR approach is meant to be applied within individual studies to obtain study-specific PEs. The AR-approach does add costs and time to complete a study, yet a relatively small number of replacement participants (~10% of the sample) was sufficient to estimate PEs specific to our study. Importantly, we have previously described an approach to obtain "pseudo-replacements" [16] that can also be used in studies when dedicated AR participants were not part of an initial study design. The fact that the method determines PEs at the group, rather than the individual level may also be a limitation. However, as described above, approaches that estimate individual-level PEs typically have a different goal (i.e., prognosis or prediction of decline). It is

not clear that there is an approach to obtain individual-level PEs for the purposes of adjusting

follow-up scores that can properly disentangle differential practice from differential decline.

In summary, with the AR-participant approach we were able to demonstrate that PEs

arising from repeated cognitive testing may persist across multiple assessments and over

periods as long as 20 years, with substantial impacts on cognitive trajectories and detection of

MCI. This approach uniquely allows for detection of PEs even when observed scores decline

and extends our prior work to handle multiple follow-ups and patterns of missingness. Adjusting

follow-up scores enables earlier detection of progression to MCI, which has the potential for

substantial cost savings in clinical trials [10,16] and allows for earlier and perhaps more effective

intervention. These results underscore the importance of accounting for PEs in longitudinal

studies of aging and clinical trials aimed at slowing cognitive decline.

5. REFERENCES

[1] Schaie KW. The course of adult intellectual development. Am Psychol. 1994;49:304–13.

[2] Nesselroade JR, Baltes PB. Longitudinal research in the study of behavior and development: Academic Press New York; 1979.

[3] Miller JB, Cummings J, Nance C, Ritter A. Neuroscience learning from longitudinal cohort studies of Alzheimer's disease: Lessons for disease-modifying drug programs and an introduction to the Center for Neurodegeneration and Translational Neuroscience. Alzheimers Dement (N Y). 2018;4:350–6.

[4] Galasko D, Corey-Bloom J, Thal LJ. Monitoring progression in Alzheimer's disease. J Am Geriatr Soc. 1991;39:932–41.

[5] McCaffrey RJ, Ortega A, Haase RF. Effects of repeated neuropsychological assessments. Archives of clinical neuropsychology : the official journal of the National Academy of Neuropsychologists. 1993;8:519–24.

[6] McCaffrey RJ, Ortega A, Orsillo SM, Nelles WB, Haase RF. Practice effects in repeated neuropsychological assessments. Clinical Neuropsychologist. 1992;6:32–42.

[7] Thorndike EL. Practice Effects in Intelligence Tests. J Exp Psychol. 1922;5:101–7.

[8] Jacobs DM, Ard MC, Salmon DP, Galasko DR, Bondi MW, Edland SD. Potential implications of practice effects in Alzheimer's disease prevention trials. Alzheimers Dement (N Y). 2017;3:531–5.

[9] Goldberg TE, Harvey PD, Wesnes KA, Snyder PJ, Schneider LS. Practice effects due to serial cognitive assessment: Implications for preclinical Alzheimer's disease randomized controlled trials. Alzheimers Dement (Amst). 2015;1:103–11.

[10] Duehring JA, Jacobs DM, Thomas ML, Dodge HH, Feldman HH, Edland SD. Implications of practice effects for the design of Alzheimer clinical trials. Alzheimer's & Dementia: Translational Research & Clinical Interventions. 2025;11:e70154.

[11] Mitrushina M, Satz P. Effect of repeated administration of a neuropsychological battery in the elderly. J Clin Psychol. 1991;47:790–801.

[12] Machulda MM, Pankratz VS, Christianson TJ, Ivnik RJ, Mielke MM, Roberts RO, et al. Practice effects and longitudinal cognitive change in normal aging vs. incident mild cognitive impairment and dementia in the Mayo Clinic Study of Aging. Clin Neuropsychol. 2013;27:1247–64.

[13] Sanderson-Cimino M, Chen R, Tu XM, Elman JA, Jak AJ, Kremen WS. Misinterpreting cognitive change over multiple timepoints: When practice effects meet age-related decline. Neuropsychology. 2023;37:568–81.

[14] Ronnlund M, Nyberg L, Backman L, Nilsson LG. Stability, growth, and decline in adult life span development of declarative memory: cross-sectional and longitudinal data from a population-based study. Psychol Aging. 2005;20:3–18.

[15] Elman JA, Jak AJ, Panizzon MS, Tu XM, Chen T, Reynolds CA, et al. Underdiagnosis of mild cognitive impairment: A consequence of ignoring practice effects. Alzheimers Dement (Amst). 2018;10:372–81.

[16] Sanderson-Cimino M, Elman JA, Tu XM, Gross AL, Panizzon MS, Gustavson DE, et al. Cognitive practice effects delay diagnosis of MCI: Implications for clinical trials. Alzheimers Dement (N Y). 2022;8:e12228.

[17] Calamia M, Markon K, Tranel D. Scoring higher the second time around: meta-analyses of practice effects in neuropsychological assessment. Clin Neuropsychol. 2012;26:543–70.

[18] Jutten RJ, Grandoit E, Foldi NS, Sikkes SAM, Jones RN, Choi SE, et al. Lower practice effects as a marker of cognitive performance and dementia risk: A literature review. Alzheimers Dement (Amst). 2020;12:e12055.

[19] Rabbitt P, Diggle P, Holland F, McInnes L. Practice and drop-out effects during a 17-year longitudinal study of cognitive aging. J Gerontol B Psychol Sci Soc Sci. 2004;59:P84–97.

[20] Schaie KW. Developmental influences on adult intelligence: The Seattle longitudinal study: Oxford University Press; 2005.

[21] Kremen WS, Franz CE, Lyons MJ. VETSA: the Vietnam Era Twin Study of Aging. Twin Res Hum Genet. 2013;16:399–402.

[22] Kremen WS, Franz CE, Lyons MJ. Current Status of the Vietnam Era Twin Study of Aging (VETSA). Twin Res Hum Genet. 2019;22:783–7.

[23] Kremen WS, Thompson-Brenner H, Leung YM, Grant MD, Franz CE, Eisen SA, et al. Genes, environment, and time: the Vietnam Era Twin Study of Aging (VETSA). Twin Res Hum Genet. 2006;9:1009–22.

[24] Henderson WG, Eisen S, Goldberg J, True WR, Barnes JE, Vitek ME. The Vietnam Era Twin Registry: a resource for medical research. Public Health Rep. 1990;105:368–73.

[25] Tsuang MT, Bar JL, Harley RM, Lyons MJ. The Harvard Twin Study of Substance Abuse: what we have learned. Harv Rev Psychiatry. 2001;9:267–79.

[26] Schoenborn CA, Heyman KM. Health characteristics of adults aged 55 years and over: United States, 2004-2007. Natl Health Stat Report. 2009;16:1–31.

[27] Eisen S, True W, Goldberg J, Henderson W, Robinette CD. The Vietnam Era Twin (VET) Registry: method of construction. Acta Genet Med Gemellol (Roma). 1987;36:61–6.

[28] Bayroff A, Anderson AA. Development of the Armed Forces Qualification Tests 7 and 8 (Technical Research Report 1122). In: Institute. UAR, editor. Alexandria, VA1963.

[29] Lyons MJ, York TP, Franz CE, Grant MD, Eaves LJ, Jacobson KC, et al. Genes determine stability and the environment determines change in cognitive ability during 35 years of adulthood. Psychol Sci. 2009;20:1146–52.

[30] Tang W, He H, Tu XM. Applied categorical and count data analysis: Chapman and Hall/CRC; 2023.

[31] Gustavson DE, Elman JA, Sanderson-Cimino M, Franz CE, Panizzon MS, Jak AJ, et al. Extensive memory testing improves prediction of progression to MCI in late middle age. Alzheimers Dement (Amst). 2020;12:e12004.

[32] Gustavson DE, Panizzon MS, Elman JA, Franz CE, Beck A, Reynolds CA, et al. Genetic and Environmental Influences on Verbal Fluency in Middle Age: A Longitudinal Twin Study. Behav Genet. 2018;48:361–73.

[33] Gustavson DE, Panizzon MS, Franz CE, Friedman NP, Reynolds CA, Jacobson KC, et al. Genetic and environmental architecture of executive functions in midlife. Neuropsychology. 2018;32:18–30.

[34] Sanderson-Cimino M, Panizzon MS, Elman JA, Gustavson DE, Franz CE, Reynolds CA, et al. Genetic and environmental architecture of processing speed across midlife. Neuropsychology. 2019;33:862–71.

[35] Bondi MW, Edmonds EC, Jak AJ, Clark LR, Delano-Wood L, McDonald CR, et al. Neuropsychological criteria for mild cognitive impairment improves diagnostic precision, biomarker associations, and progression rates. Journal of Alzheimer's disease : JAD. 2014;42:275–89.

[36] Jak AJ, Bondi MW, Delano-Wood L, Wierenga C, Corey-Bloom J, Salmon DP, et al. Quantification of five neuropsychological approaches to defining mild cognitive impairment. Am J Geriatr Psychiatry. 2009;17:368–75.

[37] Bartels C, Wegrzyn M, Wiedl A, Ackermann V, Ehrenreich H. Practice effects in healthy adults: a longitudinal study on frequent repetitive cognitive testing. BMC Neurosci. 2010;11:118.

[38] Machulda MM, Hagen CE, Wiste HJ, Mielke MM, Knopman DS, Roberts RO, et al. Practice effects and longitudinal cognitive change in clinically normal older adults differ by Alzheimer imaging biomarker status. Clin Neuropsychol. 2017;31:99–117.

[39] Collie A, Maruff P, Darby DG, McStephen M. The effects of practice on the cognitive test

performance of neurologically normal individuals assessed at brief test-retest intervals. Journal of the International Neuropsychological Society : JINS. 2003;9:419–28.

[40] Rabbitt P, Lunn M, Wong D, Cobain M. Age and ability affect practice gains in longitudinal studies of cognitive change. J Gerontol B Psychol Sci Soc Sci. 2008;63:P235–P40.

[41] Ferrer E, Salthouse TA, Stewart WF, Schwartz BS. Modeling age and retest processes in longitudinal studies of cognitive abilities. Psychol Aging. 2004;19:243–59.

[42] Reeve CL, Lam H. The Relation Between Practice Effects, Test-Taker Characteristics and Degree ofg-Saturation. International Journal of Testing. 2007;7:225–42.

[43] Salthouse TA. Influence of age on practice effects in longitudinal neurocognitive change. Neuropsychology. 2010;24:563–72.

[44] Salthouse TA, Schroeder DH, Ferrer E. Estimating retest effects in longitudinal assessments of cognitive functioning in adults between 18 and 60 years of age. Dev Psychol. 2004;40:813–22.

[45] Gross AL, Chu N, Anderson L, Glymour MM, Jones RN, Coalition Against Major D. Do people with Alzheimer's disease improve with repeated testing? Unpacking the role of content and context in retest effects. Age Ageing. 2018;47:866–71.

[46] Sanderson-Cimino M, Elman JA, Tu XM, Gross AL, Panizzon MS, Gustavson DE, et al. Practice Effects in Mild Cognitive Impairment Increase Reversion Rates and Delay Detection of New Impairments. Front Aging Neurosci. 2022;14:847315.

[47] Sanderson-Cimino M, Elman JA, Tu XM, Gross AL, Panizzon MS, Gustavson DE, et al. Cognitive Practice Effects Delay Diagnosis; Implications for Clinical Trials. medRxiv. 2020.

[48] Heaton RK, Temkin N, Dikmen S, Avitable N, Taylor MJ, Marcotte TD, et al. Detecting change: A comparison of three neuropsychological methods, using normal and clinical samples. Archives of Clinical Neuropsychology. 2001;16:75–91.

[49] McSweeny AJ, Naugle RI, Chelune GJ, Lüders H. "TScores for Change": An illustration of a regression approach to depicting change in clinical neuropsychology. Clinical Neuropsychologist. 1993;7:300–12.

ACKNOWLEDGMENTS

This work was supported by the National Institute on Aging at the National Institutes of Health (grant numbers R01s AG076838, AG022381, AG050595, AG064955; and K01 AG063805). The content of this manuscript is the responsibility of the authors and does not represent official views of NIA/NIH, or the Veterans' Administration. Numerous organizations provided invaluable assistance in the conduct of the VET Registry, including: U.S. Department of Veterans Affairs, Department of Defense; National Personnel Records Center, National Archives and Records Administration; Internal Revenue Service; National Opinion Research Center; National Research Council, National Academy of Sciences; the Institute for Survey Research, Temple University. The authors gratefully acknowledge the continued cooperation of the twins and the efforts of many staff members.

CONFLICTS

AMD is a founder and holds equity in CorTechs Laboratories, Inc., and serves on its Scientific Advisory Board. He is a member of the Scientific Advisory Board of Human Longevity, Inc., and receives funding through research agreements with GE HealthCare and Medtronic. The terms of this arrangement have been reviewed and approved by the University of California San Diego in accordance with its conflict of interest policies. All other authors report no financial interests or potential conflicts of interest.

FUNDING SOURCES

This work was supported by the National Institute on Aging at the National Institutes of Health (grant numbers R01s AG076838, AG022381, AG050595, AG064955; and K01 AG063805). The funding sources had no role in the study design; in the collection, analysis and interpretation of data; or in the writing of the report.

CONSENT STATEMENT

Informed consent was obtained from all participants and institutional review boards at the University of California San Diego and Boston University approved all study procedures. data; or in the writing of the report.

KEYWORDS

practice effects; repeat testing; serial testing; longitudinal testing; mild cognitive impairment; cognitive change

DATA AVAILABILITY

Instructions for data access requests are available on the VETSA website (https://psychiatry.ucsd.edu/research/programs-centers/vetsa/researchers.html). Access to data from military induction can be requested from the Vietnam Era Twin Registry (https://www.seattle.eric.research.va.gov/VETR/Investigator_Access.asp).

LEGENDS

Figure 1. Practice effect estimates across follow-up assessments. The forest plot presents practice effect estimates and 95% confidence intervals for practice effect estimates at each follow-up. The first, second and third follow-ups occurred at waves 2, 3, and 4, respectively, for all tests except the Boston Naming Test. The Boston Naming Test was introduced at wave 3, so the first follow-up occurred at wave 4. Models included all participants and assessment patterns but only estimates corresponding to individuals that participated in all four study waves are presented in this figure (i.e., column 1 = β_4, column 2 = β_6, column 3 = β_8). Coefficients can be interpreted as the expected boost in performance in standard deviation units expected for returnees at a given wave and given number of prior assessments compared to a test-naïve participant of a similar age and young adult general cognitive ability level. All items were coded such that higher values reflect better performance. See Table 2 for full names of measures in each domain. AFQT=Armed Forces Qualifying Test.

* Boston Naming Test was administered at waves 3 and 4 only.

† Spatial Span was administered at waves 1, 2, and 3 only.

Figure 2. Plots of cognitive factor score trajectories. Means and within-subject standard errors for cognitive factor score composites calculated from unadjusted (triangles and dashed lines) versus practice effect-adjusted (dots and solid lines) scores. Scores were standardized using the sample means and standard deviations at wave 1.

Figure 3. Rates of mild cognitive impairment. Bar plots present prevalence of any mild cognitive impairment (left panel), amnestic MCI (middle panel) and non-amnestic MCI (right panel) at each wave based on unadjusted (orange) and practice effect-adjusted (blue) test scores. Asterisks indicate significant (p<0.05) differences in prevalence between unadjusted and adjusted rates within a wave.

Table 1. Demographic characteristics of the sample.

N Total	1608
Age (years); mean (SD)	
Wave 1	55.94 (2.43)
Wave 2	61.72 (2.45)
Wave 3	67.56 (2.53)
Wave 4	73.09 (2.09)
Education (years); mean (SD)	13.85 (09)
Age 20 AFQT (percentile); mean (SD)	59.95 (23.11)
Race; n (%)	
American Indian or Alaskan Native	5 (0.3%)
Native Hawaiian or Pacific Islander	2 (0.1%)
Black or African-American	88 (5.5%)
White	1492 (92.8%)
More than one race	21 (1.3%)
Decline to answer	2 (0.1%)
Ethnicity; n (%)	
Hispanic	44 (2.7%)
Non-Hispanic	1564 (97.3%)
Testing interval (years); mean (SD)	
Wave 1 to 2	5.69 (0.69)
Wave 2 to 3	5.91 (0.76)
Wave 3 to 4	5.63 (0.60)

Note: AFQT = Armed Forces Qualifying Test

Table 2. Neuropsychological test scores included in practice effects estimation. Individual scores are grouped based on the cognitive composite that they contribute to. Measures listed in the "Other" category were not used to calculate cognitive factor scores but were used to classify mild cognitive impairment.

Cognitive Domain	Measure
Episodic Memory	CVLT-II Total of Trials 1-5
	CVLT-II Short Delay Free Recall
	CVLT-II Long Delay Free Recall
	WMS-III Logical Memory - Immediate Recall Story Units Total Score
	WMS-III Logical Memory - Delayed Recall Story Units Total Score
	WMS-III Visual Reproduction: Immediate Recall Total Score
	WMS-III Visual Reproduction: Delayed Recall Total Score
Executive function	WMS-III Digit Span: Forward + Backward Score
	WMS-III Letter-Number Sequencing: Total Score for Trials Passed
	D-KEFS Trails 4 time
	D-KEFS Category switching accuracy (# correct switches)
	Stroop: Interference Score
	Reading Span: Total Score Ascending
Verbal Fluency	D-KEFS Letter Fluency FAS total correct
	D-KEFS Category Fluency Animal/Boys total correct
	D-KEFS Category Switching total correct
Processing Speed	D-KEFS Trails 2 time
	D-KEFS Trails 3 time
	Stroop: Raw Word Score
	Stroop: Raw Color Score
	Simple Reaction Time
	Choice Reaction Time
Visuospatial	Mental Rotation: Total Correct Part 1
	Gottschaldt Hidden Figures Total Correct All Parts
	AFQT – Box Folding Subtest
Other	D-KEFS Trails 4 time
	WMS-III Spatial Span: Total Trials Passed Forward + Backward*
	WASI Matrix Reasoning Raw Score
	WMS-III Visual Reproduction: Copy Total Score
	Boston Naming Test[†]

Note: AFQT = Armed Forces Qualifying Test; CVLT = California Verbal Learning Test-II; WMS-III = Wechsler Memory Scale-III; D-KEFS = Delis-Kaplan Executive Function System; WASI = Wechsler Abbreviated Scale of Intelligence
*Spatial span was administered at waves 1, 2, and 3.
†Boston Naming Test was administered at waves 3 and 4.