Time-lagged Cognitive Impairment from Long-term Occupational Exhaust Exposure: A NHANES Analysis Using Distributed Lag Nonlinear Models

Lili Liang[a], Perry E Sheffield[b], Rose Saint Fleur-Calixte[c], Tianxu Xia[d], Spencer Xinyi Zhang[d], Jenny J Lin[e]

[a] Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA.
[b] Departments of Pediatrics and Environmental Medicine and Climate Science, Icahn School of Medicine at Mount Sinai, New York, New York, USA.
[c] Department of Epidemiology and Biostatistics, SUNY Downstate Medical Center, New York, New York, USA.
[d] Departments of Neurology and Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York, USA.
[e] Division of General Internal Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

Abstract

Background: Occupational exposure to harmful substances such as exhaust fumes is a known risk factor for respiratory and cardiovascular diseases. However, its impact on cognitive function remains underexplored in older adults. As the population in the United States ages, understanding the influence of occupational exposures on cognitive health is critical for identifying modifiable risk factors for cognitive decline.

Methods: This study investigates the relationship between ~~past~~ occupational exposure to exhaust fumes and cognitive performance among older adults aged 60 years and above, utilizing data from the 2011–2012 National Health and Nutrition Examination Survey (NHANES) cohort. Cognitive performance was assessed using the Consortium to Establish a Registry for Alzheimer's Disease Word List Learning (CERAD-WL) recall test (immediate and delayed), the Animal Fluency Test (AFT), and the Digit Symbol Substitution Test (DSST). Cognitive impairment was defined as an adjusted Z-score $\leq$ -1.5 standard deviations (SD) (based on age- and education-corrected norms). Multivariable logistic regression analysis was conducted to evaluate the associations between occupational exposure and cognitive impairment, adjusting for demographic and health-related covariates. A Distributed Lag Nonlinear Model (DLNM) was applied to investigate the complex relationships between exposure levels and cognitive impairment.

Results: Among the 1,110 participants, 266 (24.0%) reported occupational exposure to exhaust fumes. Comparing to the unexposed group, exposed individuals were more likely to be male (81.6% vs. 41.8%), have lower education levels (31.2% vs. 24.4% with less than 12[th] grade level), and exhibit higher rates of smoking (65.0% vs. 48.6%) and excessive alcohol use (15.7% vs. 7.0%). Cognitive performance was significantly poorer in the exposed group, after controlling for age, sex and education. Occupational exhaust fumes exposure was associated with substantial impairments in delayed memory (CERAD-WL Delayed recall: OR = 2.55, 95% CI: 1.61–4.05), verbal fluency (AFT: OR = 2.41, 95% CI: 1.48–3.94), and processing speed (DSST: OR = 1.95, 95% CI: 1.19–3.18). Duration of occupational exposure exhibited biphasic, time-lagged effects on cognitive function, with low levels of exposure showing minimal positive impact, and higher levels leading to significant declines in immediate and delayed recall, semantic memory, and processing speed, particularly over prolonged exposure periods (20-30 years).

Conclusion: Occupational exposure to exhaust fumes is significantly associated with poorer cognitive performance in older adults, particularly in domains of memory, semantic fluency, and processing speed. The DLNM analysis further indicates a significant time-dependent decline in cognitive scores at higher exposure levels. While further research is needed to explore the mechanisms linking occupational exposure to cognitive decline and to explore long-term brain health outcomes, these findings highlight the need for targeted interventions to reduce occupational exposures and mitigate their cognitive impacts. The current trend to use of electric vehicles may aid in reducing exposures, and populations with higher exposure risk should be monitored closely for cognitive changes.

Keywords

Occupational Exposure; Exhaust Fumes; Cognitive Function; National Health and Nutrition Examination Survey (NHANES); Distributed lag nonlinear models (DLNM)

1. Introduction

Occupational exposure to exhaust fumes, a key contributor to traffic-related air pollution (TRAP),[1] is an important source of environmental exposure with significant implications for human health. Exhaust fumes consist of fine particulate matter (PM2.5), polycyclic aromatic hydrocarbons (PAHs), metals, and reactive compounds capable of penetrating deep into the respiratory system and crossing the blood-brain barrier.[1] While TRAP has long been associated with cardiovascular and respiratory diseases,[2-5] increasing evidence highlights its potential neurotoxic effects and contributions to neurodegenerative diseases.[6-11]

A growing body of epidemiological evidence has linked chronic exposure to TRAP with cognitive decline, dementia, and Alzheimer's disease.[12-32] Observational studies suggest that long-term exposure to PM2.5 and other traffic-related pollutants accelerates cognitive dysfunction in older adults.[16,33] Proposed mechanisms include direct pathways, where inhaled ultrafine particles (<0.1 μm) enter the brain through the olfactory bulb, and indirect pathways, where systemic inflammation and oxidative stress affect brain function.[34,35] Animal studies have demonstrated acute and subchronic neuroinflammation, oxidative damage, and elevated markers of neurodegeneration following exhaust fumes exposure.[8,9,10,11] Behavioral experiments have further shown impairments in learning, memory, and spatial cognition in mice exposed to exhaust fumes.[36-40] Experimental studies investigating cognitive effects of sub-chronic or acute exhaust fumes exposures in humans provide further insight. For instance, retrofitting school buses to reduce exhaust fumes resulted in improved cognitive performance among students, including increased English and math test scores.[41,42] Further studies have shown decrements in cognitive function following short-term TRAP exposure, with diminished attention, processing speed, and executive function observed in commuters and office workers exposed to PM2.5.[43,44] These findings suggest that even short-term exposure to exhaust fumes can affect cognitive domains critical for decision-making and occupational safety.

Despite substantial evidence from environmental exposure studies, the impacts of occupational exposure to exhaust fumes —characterized by higher intensity and prolonged duration—remain underexplored. Workers in industries such as transportation, construction, and heavy machinery are frequently exposed to high concentrations of exhaust fumes, potentially compounding risks of systemic inflammation, hypertension, and metabolic syndrome—conditions associated with cognitive decline.[45,46] Occupational settings also often involve co-exposures to noise and stress, which may exacerbate cognitive impairments.[47,48] However, limited research has examined the impact of occupational fumes exposure on cognitive function.

This study aims to address the gap in understanding the effects of occupational exhaust fumes exposure on cognitive function and how this relationship may differ by race and ethnicity. This study uses data from the 2011–2012 National Health and Nutrition Examination Survey (NHANES). Key exposure metrics, including occupational exhaust exposure history and exposure duration, were analyzed to explore dose-response relationships and assess specific cognitive outcomes, such as delayed memory and processing speed.

2. Materials and methods

2.1 Study population

We used NHANES data collected in the 2011–2012 cycle.[49] This analysis focused on older adults aged 60 years and above to explore the association between occupational exposure to exhaust fumes and cognitive function. Participants were eligible for inclusion if they had completed cognitive function tests, such as the Consortium to Establish a Registry for Alzheimer's Disease Word List Learning Test (CERAD-WL) (immediate and delayed), the Animal Fluency Test (AFT), and the Digit Symbol Substitution Test (DSST) and complete data on occupational exposure to exhaust fumes, demographic characteristics, and relevant covariates (e.g., educational attainment, household income, race/ethnicity, smoking and alcohol use). Following the inclusion criteria outlined in Figure 1, a total of 1110 adults were included in the study.

2.2 Measurement of occupational exposure history

The Occupation Questionnaire section of the NHANES 2011-2012 study includes interview data regarding employment and work environment-related variables. A subset of individuals aged 16-79 were asked about their lifetime history of exposure to mineral and organic dusts, gases, fumes, and vapors in the workplace, specifically, participants were first asked whether they had ever been exposed to exhaust fumes at work (OCQ550), defined as breathing in fumes or having a lingering smell on their clothes, skin, or hair. These exhaust fumes could originate from sources such as trucks, buses, heavy machinery, and diesel engines in any job. Additionally, participants who answered 'yes' were then asked to report the total number of years they had been exposed to exhaust fumes at work from all jobs (OCQ560). OCQ550 (binary, exposed or unexposed) was used to categorize participants. OCQ560 (years of exposure) was treated as a continuous variable.

2.3 Cognitive function test

The CERAD-WL recall test measures both immediate and delayed recall abilities for new verbal information.[50,51] This assessment includes three sequential learning trials followed by a delayed recall test. In each learning trial, participants are asked to read aloud a list of 10 unrelated words. Immediately after hearing the list, participants are instructed to recall and recite as many words as they can remember. The delayed recall test takes place roughly 10 minutes after the start of the initial word learning trials. Each trial allows for a maximum score of 10, with a total possible score of 40, which includes the combined scores from all three trials plus the delayed recall.

The AFT evaluates verbal category fluency, which is an aspect of executive function, alongside other cognitive functions like semantic memory and processing speed.[52] Participants are required to name as many different animals as they can within a one-minute period, earning one point for each correctly named animal.

The DSST, which serves as an overall indicator of cognitive health, assesses skills such as processing speed, visual scanning ability, sustained attention, and short-term memory.[53] This test involves a paper form where participants are given a key displaying nine numbers, each paired with a unique symbol. Over the course of two minutes, participants must accurately copy the corresponding symbols into 133 boxes aligned with the numbers.

According to the guidelines of the Uniform Data Set (UDS),[54, 55] cognitive impairment is defined as an adjusted Z-score ≤ -1.5 standard deviations (SD) (based on age- and education-corrected norms), with cognitive performance dichotomized as "impaired" or "unimpaired" to identify mild cognitive impairment (MCI).

2.4 Covariates

Demographic variables included age (either 60–69 or 70–79 years), gender (male, female), race or ethnicity (Hispanic, non-Hispanic White, non-Hispanic Black, non-Hispanic Asian and other race), and educational attainment (< HSG: less than 12th grade, including 12th grade with no diploma; HSG: high school graduate or GED; >HS: some college or associate degree, and college graduate or higher). Health-related variables included body mass index (BMI), smoking history, alcohol use, high blood pressure, abnormal cholesterol, heart disease, depression, sleep problems, and hyperglycemia. BMI (kg/m²) was calculated using height and weight measured by trained staff. Obesity was defined as a BMI ≥30 kg/m². Smoking history was divided into two groups: those who smoked at least 100 cigarettes in their life and those who did not.[56] Excessive alcohol use was defined as more than 350 g/week for females or more than 420 g/week for males.[57] High blood pressure was defined by a measured blood pressure of 140/90 mmHg or higher or if the person reported taking prescription for hypertension. Dyslipidemia was defined by triglyceride levels of 1.70 mmol/L (150 mg/dL) or higher, low-density lipoprotein cholesterol of 3.4 mmol/L (130 mg/dL) or higher, or high-density lipoprotein cholesterol less than 1.0 mmol/L (40 mg/dL) in men and less than 1.3 mmol/L (50 mg/dL) in women. History of cardiovascular diseases (CVD) was reported by participants and included history of heart failure, coronary heart disease, angina, heart attack or stroke. Depression was measured using the PHQ-9 questionnaire, with scores of 10 or higher indicating depression.[58] Sleep problems were identified if participants said they slept 1–4 hours per day or said "yes" to the questions, "Did a doctor ever say you had trouble sleeping?" or "Did a doctor ever say you had a sleep disorder?"[59] Hyperglycemia was defined by fasting glucose of 5.6 mmol/L (100 mg/dL) or higher, hemoglobin A1c of 5.7% (39 mmol/mol) or higher, or if the person said they had type 2 diabetes or were being treated for it.

2.5 Statistical analysis

This study utilized data from NHANES, which employs a complex sampling design. Descriptive statistics were calculated for demographic and health-related variables. Chi-square tests were used to compare categorical variables (e.g., age group, gender, race/ethnicity, education levels, smoking history, and excessive alcohol consumption, obesity, hypertension, dyslipidemia, hyperglycemia, history of CVD, depression and sleep disorders) between exposed and unexposed groups.

Weighted survey logistic regression models were used to evaluate the association between occupational exposure to exhaust fumes and cognitive impairment. Multivariable logistic regression models were adjusted for potential confounding variables, including age group, gender, race/ethnicity, educational attainment, smoking history, and excessive alcohol consumption, obesity, hypertension, dyslipidemia, hyperglycemia, history of CVD, depression and sleep disorders. Covariates were selected based on their association with cognitive

impairment. Adjusted analyses were conducted to examine associations between cognitive impairment and gender, age, and education. NHANES-recommended sampling weights, strata, and primary sampling units were incorporated to account for the complex survey design and ensure representative results. All analyses accounted for the NHANES complex sampling design using survey procedures in SAS software version 9.4 (SAS Institute Inc., Cary, NC). The significance level for all statistical tests was set at $p < 0.05$.

We employed a Distributed Lag Nonlinear Model (DLNM) to investigate the complex relationships between years of exposure (OCQ560, continuous variable) and cognitive function scores (continuous variables). Cross-basis function was constructed to quantify the exposure-response relationship and lag effects, allowing non-linear characteristics in both the time and dose dimensions. Both exposure and lag dimensions were modeled using natural splines (df = 3), with a maximum lag duration of 10 years. The model adjusted for potential confounders such as age, gender, race, education, depression, hyperglycemia and history of CVD. with the optimal parameter configuration selected using the Akaike Information Criterion (AIC). All analyses were conducted in R (v4.4.1) using the *dlnm* and *splines* packages. Effect estimates were visualized via the *crosspred* function, with effect sizes represented by β coefficients and 95% confidence intervals, and statistical significance set at a two-tailed α = 0.05.

3. Results

3.1 Characteristics of participants by exposure history

Based on Table 1, a total of 1,110 participants aged 60 and older were analyzed, comprising 266 individuals (24.0%) in the exposed group and 844 (76.0%) in the unexposed group. Significant differences were observed between the two groups. The exposed group had a higher proportion of males (81.6% vs. 41.8%, $P < 0.0001$) and individuals with less than high school education (31.2% vs. 24.4%, $P = 0.017$). Smoking history differed notably, with 65.0% of the exposed group reporting smoking >100 cigarettes compared to 48.6% in the unexposed group ($P < 0.0001$). Excessive alcohol use was more common in the exposed group (15.7% vs. 7.0%, $P = 0.001$).

3.2 Cognitive Function Test Scores and Occupational Exposure

Participants with occupational exposure to exhaust fumes demonstrated poorer cognitive performance across all tests (CERAD-WL recall test (immediate and delayed); AFT; DSST). For the CERAD-WL delayed recall test, the association between occupational exposure and cognitive function was statistically significant (OR = 2.55, 95% CI: 1.61–4.05), indicating a substantial negative impact on delayed memory. Similar trends were observed for the AFT and DSST. Occupational exposure was significantly associated with poorer verbal fluency in the AFT (OR = 2.41, 95% CI: 1.48–3.94) and lower processing speed in the DSST (OR = 1.95, 95% CI: 1.19–3.18). However, this association weakened and became non-significant for the CERAD-WL immediate recall test (OR = 1.45, 95% CI: 0.91–2.30), suggesting that immediate memory may be less affected by occupational exposure. These results indicate that occupational exposure is associated with cognitive decline, particularly in domains related to delayed memory, verbal fluency, and processing speed (Figure 2).

3.3 Exposure-Response Curves and Time-lagged Effect Analysis

The exposure-response curves and biphasic time-lagged effects of occupational exposure (OCQ560) on cognitive function scores are shown in Figure 3.

For CERAD-WL immediate recall, the exposure-response curve shows an initial effect size of 0.5 at OCQ560 = 10. This decreases to 0.0 at OCQ560 = 30. This means that low exposure levels may have a small positive effect, but as exposure levels increase, the effect quickly drops to zero. The 3D lag effect analysis revealed a significant time-dependent decline in cognitive function scores with increasing exposure levels (OCQ560). At lower exposure levels (OCQ560 < 20), the effect size remained close to zero ($\beta \approx 0$). However, at higher exposure levels (OCQ560 > 30) and longer lag times (20-30 lags), the effect size became significantly negative ($\beta = -0.6$), suggesting a pronounced decline in cognitive performance over time.

For CERAD-WL delayed recall, the exposure-response curve shows a threshold effect. There is no significant association at OCQ560 < 20 ($\beta \approx 0$). But above this threshold, the effect decreases linearly ($\beta = -0.02$ per unit increase). This means that a significant negative impact on delayed recall only happens above a certain exposure level. The 3D lag effect plot shows significant negative effects between lags 15 and 25 ($\beta = -0.4$). These effects get worse at higher exposure levels (OCQ560 > 30), ranging from $\beta = -0.4$ to -0.6. This means that prolonged high exposure levels have a time-dependent harmful effect on delayed recall.

The effect on AFT is highest at lags 0-10 and keeps decreasing at high exposure levels (OCQ560 > 40). The exposure-response curve shows a negative linear relationship. The effect size increases by $\beta = -0.5$ for every 10-unit rise in OCQ560 ($p < 0.01$). This shows a clear dose-response relationship between exposure levels and semantic memory impairment. The 3D lag effect plot demonstrated the same clear negative trend. At higher exposure levels (OCQ560 > 30) and longer lag times (20-30 lags), the effect size dropped significantly ($\beta = -1.0$), indicating a substantial decline in semantic fluency. This suggests that prolonged exposure at higher levels has a time-dependent negative impact on semantic memory.

The exposure-response analysis for the DSST showed a dose-dependent relationship between exposure levels (OCQ560) and cognitive performance. At low exposure levels (OCQ560 < 10), the effect size was positive ($\beta \approx 2$), indicating no decline in performance. However, as exposure levels increased (OCQ560 > 30), the effect size decreased linearly ($\beta = -0.2$ per unit increase), reaching a significant negative value ($\beta = -0.6$) at the highest exposure levels (OCQ560 = 50). The 3D lag effect plot shows changes in the impact on DSST across different exposure levels and lag times. At lower exposure levels (OCQ560 < 10), there is a short-term positive effect within 0-5 lags. But at higher exposure levels (OCQ560 > 30), this effect changes significantly, with a β value of -0.2. This demonstrates that higher exposure levels are associated with a progressive decline in processing speed and working memory.

3.4 Adjusted Analysis of Gender, Age and Education Levels

As shown in the Supplementary Table 1, differences in cognitive performance were observed across gender, age, and education levels. Women generally scored higher than men on all

cognitive tests, with the most pronounced differences observed in the DSST and AFT. Declines in cognitive function were associated with advancing age and lower educational attainment. These associations were more pronounced among men, whose cognitive scores showed greater variability in response to age and education.

3.5 Racial Disparities

Significant racial disparities were observed in cognitive performance across all tests, with non-Hispanic Black participants consistently scoring lower than other racial groups (Supplementary Figure 1). These differences persisted even after adjusting for age, gender, and education levels. In the DSST, non-Hispanic Black participants had notably higher odds of poorer performance compared to non-Hispanic White participants (OR=7.13, 95% CI: 4.31–11.79). Similar trends were observed in CERAD-WL recall test (immediate and delayed). Participants of Mexican American or Other Hispanic backgrounds also demonstrated worse cognitive performance compared to non-Hispanic White participants, though the disparities were less pronounced (DSST: OR=5.39, 95% CI: 2.65–10.93).

4. Discussion

This study evaluated the relationship between occupational exposure to exhaust fumes and cognitive function among older adults using data from the 2011-2012 NHANES. The findings revealed significant associations between occupational exposure to exhaust fumes and declines in cognitive performance, particularly in delayed memory, verbal fluency, and processing speed. A dose-response relationship with exposure duration highlighted the cumulative impact of prolonged occupational exposure. These results contribute to the growing body of evidence linking air pollution and occupational hazards to neurodegeneration and cognitive decline. By focusing on occupational exposures, this research investigates how prolonged, high-intensity exposure to exhaust fumes impacts cognitive domains, contributing to growing evidence of environmental and occupational risk factors for neurodegeneration.

The findings are consistent with previous research on air pollution and cognitive outcomes. Carey et al reported a 16% increased risk of dementia with NO☐ and a 7% increase with PM2.5 among older adults in London.[60] Similarly, the Adult Changes in Thought (ACT) study observed a 16% higher hazard of all-cause dementia with a 10-year average PM2.5 exposure.[17] Gao et al identified a 10.4% increased risk of cognitive impairment per 10 μg/m³ increase in PM2.5 in a Chinese cohort.[15] While these studies focused on environmental exposures, our findings extend this research to occupational settings, where exposure intensity is often higher. The results also parallel Colicino et al results that link a traffic-related air pollution marker to declines in verbal memory, recognition, and executive function.[61,62] However, the lack of association with immediate memory in this study suggests domain-specific vulnerabilities that warrant further exploration.

Our DLNM analysis revealed novel insights into the neurotoxic effects of occupational exhaust fume exposure through two key mechanistic findings. First, we observed a biphasic dose-response relationship where low-level exposure (<20 years) transiently improved processing speed (DSST β≈2), while higher exposures (>30 years) caused significant cognitive impairment.

This pattern, potentially mediated by Nrf2 pathway activation,[63] contrasts with the monotonic cognitive decline typically reported for environmental pollutants like PM2.5 and occupational solvents.[64,65] While similar low-dose neuroadaptive responses were documented in cadmium-exposed animals,[66] this phenomenon remains rarely reported in human occupational studies. We acknowledge potential confounding from healthier work environments in low-exposure groups, a limitation noted in air pollution research. Second, our DLNM analysis identified critical time windows (15-25 lag years) for cognitive decline, suggesting cumulative damage thresholds analogous to β-amyloid dynamics in Alzheimer's disease. [67] This finding advance beyond traditional occupational studies that used static exposure metrics or single lag periods. Furthermore, our domain-specific approach addresses important gaps in occupational neurotoxicology, [68] revealing distinct vulnerability patterns: DSST's sensitivity to low exposure (likely reflecting prefrontal cortex adaptability) versus AFT's susceptibility to chronic exposure (possibly indicating temporal lobe vulnerability), partially aligning with previous research.[69] These results underscore the need for dynamic exposure assessment and cognitive domain stratification in future occupational health studies.

Gender differences in cognitive outcomes were significant, with females consistently outperforming males. This aligns with evidence that estrogen confers neuroprotective effects through antioxidative and anti-inflammatory mechanisms.[70] However, this advantage diminishes with age due to declining estrogen levels during menopause. Mitochondrial health further explains this disparity. Females exhibit greater mitochondrial repair capacity and antioxidant defenses, offering resilience against oxidative damage. In contrast, males showed pronounced deficits in processing speed and executive function, likely due to greater susceptibility to oxidative stress and neurodegeneration.[71]

This study also uncovered significant socioeconomic and racial disparities. Non-Hispanic Black and Hispanic participants scored consistently lower on cognitive tests compared to non-Hispanic White participants, even after adjusting for confounders such as education and age. These disparities reflect systemic inequities in education, occupational opportunities, and healthcare access. Lower educational attainment, overrepresented in high-risk occupational groups, exacerbates cognitive decline risks by reducing cognitive reserve, a factor that enhances resilience to neurodegeneration.[12] Environmental exposures further amplify these disparities. Minority populations often face disproportionate exposure to harmful pollutants like diesel exhaust, coupled with limited access to healthcare for managing associated conditions such as cardiovascular disease.[73] These findings emphasize the need for targeted interventions in high-risk communities, including educational programs and stricter regulatory policies to minimize workplace hazards.

Age was another critical factor, with participants aged 70–79 years experiencing greater cognitive declines than those aged 60–69 years. This finding supports evidence that cumulative exposure and age-related systemic inflammation amplify susceptibility to neurodegeneration.[15] The overrepresentation of lower-educated individuals in high-risk occupations further underscores the intersection of occupational hazards and socioeconomic inequities.

Several important limitations of this study should be acknowledged. First, the cross-sectional design prevents us from establishing causal relationships between exposure and outcomes, as we

cannot rule out the possibility of reverse causality. For instance, individuals with pre-existing cognitive impairments may have been more likely to select certain occupations, potentially biasing our results. Second, our reliance on self-reported exposure metrics introduces the possibility of recall bias, which could be mitigated in future studies through objective measures such as workplace air quality monitoring or industrial hygiene records. Third, while we adjusted for major confounders, we were unable to account for other occupational co-exposures that may influence cognitive outcomes, including noise pollution and psychosocial stressors, which are known to interact with chemical exposures. Fourth, the lack of detailed temporal data on participants' job histories and residential locations limited our ability to reconstruct complete exposure trajectories over time. Finally, the absence of biological markers of neurodegeneration or neuroinflammation represents a significant constraint, as such biomarkers could provide direct evidence of the pathological mechanisms linking exhaust fume exposure to cognitive decline.

Despite these limitations, our findings have important implications for both public health practice and future research. The identification of occupational exhaust exposure as a modifiable risk factor for cognitive decline suggests several actionable strategies. Public health initiatives should focus on implementing technological interventions in high-risk industries, including the adoption of cleaner engine technologies and enhanced workplace air filtration systems. Occupational health programs should consider incorporating regular cognitive screening for workers with significant exposure histories, as this could facilitate early detection of impairment and timely intervention. Furthermore, our results underscore the need for policies that address systemic socioeconomic inequities, as these factors appear to compound both exposure risks and cognitive vulnerabilities.

For future direction, longitudinal study designs would provide stronger evidence for causal relationships by tracking both exposure and cognitive changes over time. Future investigations should incorporate biomarkers of neuroinflammation and oxidative stress, along with neuroimaging modalities, to better elucidate the biological pathways underlying these associations. There is also a need to examine how occupational exposures interact with lifestyle factors such as diet and physical activity, as well as genetic predispositions including APOEε4 status, to better understand the multifactorial nature of cognitive decline. Additionally, our observed exposure thresholds should be validated across diverse occupational cohorts to inform more precise regulatory standards for workplace air quality.

5. Conclusion

By focusing on occupational exposure, this research investigates how prolonged, high-intensity exposure to exhaust fumes impacts cognitive domains in American adults, contributing to growing evidence of environmental and occupational risk factors for neurodegeneration. Vulnerable populations, particularly those in high-risk industries, are disproportionately affected. The dose-response relationship underscores the cumulative impact of exposure, with early declines suggesting the need for timely interventions. Addressing systemic disparities and implementing targeted workplace safety measures are critical steps toward mitigating these risks. By bridging gaps in knowledge and policy, future research and public health initiatives can advance efforts to protect cognitive health in at-risk populations.

Data availability statement
Publicly available datasets were analyzed in this study. This data can be found here: the datasets for this study can be found in the NHANES repository. Please see the https://wwwn.cdc.gov/nchs/nhanes/Default.aspx for more details.

Ethics statement
The studies involving human participants were reviewed and approved by the Research Ethics Review Board (ERB) of the US National Center for Healthcare Statistics (NCHS) authorized the 2011–2012 NHANES (Protocol Number: protocol#2011-12 and continuation of protocol #2011-12). The patients/participants provided their written informed consent to participate in this study. Written informed consent was obtained from the individual(s) for the publication of any potentially identifiable images or data included in this article.

CRediT authorship contribution statement
Lili Liang: conceptualization, formal analysis, and writing original draft.
Perry E Sheffield: conceptualization, methodology, editing and review.
Rose Saint Fleur-Calixte: software, and formal analysis.
Tianxu Xia: methodology and review.
Spencer Xinyi Zhang: validation and writing original draft.
Jenny J Lin: methodology, review and editing.

Funding
This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest
The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments
We express our special appreciation to Dr. John Doucette for his invaluable support on statistical consultation. We are grateful for the NHANES team and all participants.

Appendix A. Supplementary data
Supplementary data to this article can be found online.

References

1. Ghio, A.J., Smith, C.B., Madden, M.C., 2012. Diesel exhaust particles and airway inflammation. Curr. Opin. Pulm. Med. 18, 144–150. https://doi.org/10.1097/MCP.0b013e32834f0e2a

2. Brook, R.D., Rajagopalan, S., Pope, C.A. 3rd, Brook, J.R., Bhatnagar, A., Diez-Roux, A.V., Holguin, F., Hong, Y., Luepker, R.V., Mittleman, M.A., Peters, A., Siscovick, D., Smith, S.C. Jr., Whitsel, L., Kaufman, J.D.; American Heart Association Council on Epidemiology and Prevention, Council on the Kidney in Cardiovascular Disease, and Council on Nutrition, Physical Activity and Metabolism, 2010. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. Circulation 121, 2331–2378. https://doi.org/10.1161/CIR.0b013e3181dbece1

3. Pope, C.A. 3rd, Ezzati, M., Dockery, D.W., 2009. Fine-particulate air pollution and life expectancy in the United States. N. Engl. J. Med. 360, 376–386. https://doi.org/10.1056/NEJMsa0805646

4. Zhang, K., Batterman, S., 2013. Air pollution and health risks due to vehicle traffic. Sci. Total Environ. 450–451, 307–316. https://doi.org/10.1016/j.scitotenv.2013.01.074

5. Schraufnagel, D.E., Balmes, J.R., De Matteis, S., Hoffman, B., Kim, W.J., Perez-Padilla, R., Rice, M., Sood, A., Vanker, A., Wuebbles, D.J., 2019. Health Benefits of Air Pollution Reduction. Ann. Am. Thorac. Soc. 16, 1478–1487. https://doi.org/10.1513/AnnalsATS.201907-538CME

6. de Crom, T.O.E., Ginos, B.N.R., Oudin, A., Ikram, M.K., Voortman, T., Ikram, M.A., 2023. Air Pollution and the Risk of Dementia: The Rotterdam Study. J. Alzheimers Dis. 91, 603–613. https://doi.org/10.3233/JAD-220804

7. Costa, L.G., Cole, T.B., Coburn, J., Chang, Y.C., Dao, K., Roqué, P.J., 2017. Neurotoxicity of traffic-related air pollution. Neurotoxicology 59, 133–139. https://doi.org/10.1016/j.neuro.2015.11.008

8. Levesque, S., Surace, M.J., McDonald, J., Block, M.L., 2011. Air pollution & the brain: Subchronic diesel exhaust exposure causes neuroinflammation and elevates early markers of neurodegenerative disease. J. Neuroinflammation 8, 105. https://doi.org/10.1186/1742-2094-8-105

9. Coburn, J.L., Cole, T.B., Dao, K.T., Costa, L.G., 2018. Acute exposure to diesel exhaust impairs adult neurogenesis in mice: prominence in males and protective effect of pioglitazone. Arch. Toxicol. 92, 1815–1829. https://doi.org/10.1007/s00204-018-2180-5

10. Ehsanifar, M., Tameh, A.A., Farzadkia, M., Kalantari, R.R., Zavareh, M.S., Nikzaad, H., Jafari, A.J., 2019. Exposure to nanoscale diesel exhaust particles: Oxidative stress, neuroinflammation, anxiety and depression on adult male mice. Ecotoxicol. Environ. Saf. 168, 338–347. https://doi.org/10.1016/j.ecoenv.2018.10.090

11. Milani, C., Farina, F., Botto, L., Massimino, L., Lonati, E., Donzelli, E., Ballarini, E., Crippa, L., Marmiroli, P., Bulbarelli, A., Palestini, P., 2020. Systemic Exposure to Air Pollution Induces Oxidative Stress and Inflammation in Mouse Brain, Contributing to Neurodegeneration Onset. Int. J. Mol. Sci. 21, 3699. https://doi.org/10.3390/ijms21103699

12. Weuve, J., Bennett, E.E., Ranker, L., Gianattasio, K.Z., Pedde, M., Adar, S.D., Yanosky, J.D., Power, M.C., 2021. Exposure to Air Pollution in Relation to Risk of Dementia and Related Outcomes: An Updated Systematic Review of the Epidemiological Literature. Environ. Health Perspect. 129, 96001. https://doi.org/10.1289/EHP8716

13. Chen, M.C., Wang, C.F., Lai, B.C., Hsieh, S.W., Chen, S.C., Hung, C.H., Kuo, C.H., 2021. Air Pollution Is Associated with Poor Cognitive Function in Taiwanese Adults. Int. J. Environ. Res. Public Health 18, 316. https://doi.org/10.3390/ijerph18010316

14. Duchesne, J., Gutierrez, L.A., Carrière, I., Mura, T., Chen, J., Vienneau, D., de Hoogh, K., Helmer, C., Jacquemin, B., Berr, C., Mortamais, M., 2022. Exposure to ambient air pollution and cognitive decline: Results of the prospective Three-City cohort study. Environ. Int. 161, 107118. https://doi.org/10.1016/j.envint.2022.107118

15. Gao, Q., Zang, E., Bi, J., Dubrow, R., Lowe, S.R., Chen, H., Zeng, Y., Shi, L., Chen, K., 2022. Long-term ozone exposure and cognitive impairment among Chinese older adults: A cohort study. Environ. Int. 160, 107072. https://doi.org/10.1016/j.envint.2021.107072

16. He, F., Tang, J., Zhang, T., Lin, J., Li, F., Gu, X., Chen, A., Nevill, A., Chen, R., 2022. Impact of air pollution exposure on the risk of Alzheimer's disease in China: A community-based cohort study. Environ. Res. 205, 112318. https://doi.org/10.1016/j.envres.2021.112318

17. Shaffer, R.M., Blanco, M.N., Li, G., Adar, S.D., Carone, M., Szpiro, A.A., Kaufman, J.D., Larson, T.V., Larson, E.B., Crane, P.K., Sheppard, L., 2021. Fine Particulate Matter and Dementia Incidence in the Adult Changes in Thought Study. Environ. Health Perspect. 129, 87001. https://doi.org/10.1289/EHP9018

18. Aretz, B., Janssen, F., Vonk, J.M., Heneka, M.T., Boezen, H.M., Doblhammer, G., 2021. Long-term exposure to fine particulate matter, lung function and cognitive performance: A prospective Dutch cohort study on the underlying routes. Environ. Res. 201, 111533. https://doi.org/10.1016/j.envres.2021.111533

19. Åström, D.O., Adolfsson, R., Segersson, D., Forsberg, B., Oudin, A., 2021. Local Contrasts in Concentration of Ambient Particulate Air Pollution (PM2.5) and Incidence of Alzheimer's Disease and Dementia: Results from the Betula Cohort in Northern Sweden. J. Alzheimers Dis. 81, 83–85. https://doi.org/10.3233/JAD-201538

20. Ilango, S.D., Gonzalez, K., Gallo, L., Allison, M.A., Cai, J., Isasi, C.R., Hosgood, D.H., Vasquez, P.M., Zeng, D., Mortamais, M., Gonzalez, H., Benmarhnia, T., 2021. Long-Term Exposure to Ambient Air Pollution and Cognitive Function Among Hispanic/Latino Adults in San Diego, California. J. Alzheimers Dis. 79, 1489–1496. https://doi.org/10.3233/JAD-200766

21. Mortamais, M., Gutierrez, L.A., de Hoogh, K., Chen, J., Vienneau, D., Carrière, I., Letellier, N., Helmer, C., Gabelle, A., Mura, T., Sunyer, J., Benmarhnia, T., Jacquemin, B., Berr, C., 2021. Long-term exposure to ambient air pollution and risk of dementia: Results of the prospective Three-City Study. Environ. Int. 148, 106376. https://doi.org/10.1016/j.envint.2020.106376

22. Park, S.Y., Han, J., Kim, S.H., Suk, H.W., Park, J.E., Lee, D.Y., 2022. Impact of Long-Term Exposure to Air Pollution on Cognitive Decline in Older Adults Without Dementia. J. Alzheimers Dis. 86, 553–563. https://doi.org/10.3233/JAD-215120

23. Parra, K.L., Alexander, G.E., Raichlen, D.A., Klimentidis, Y.C., Furlong, M.A., 2022. Exposure to air pollution and risk of incident dementia in the UK Biobank. Environ. Res. 209, 112895. https://doi.org/10.1016/j.envres.2022.112895

24. Ran, J., Zhang, Y., Han, L., Sun, S., Zhao, S., Shen, C., Zhang, X., Chan, K.P., Lee, R.S., Qiu, Y., Tian, L., 2021. The joint association of physical activity and fine particulate matter exposure with incident dementia in elderly Hong Kong residents. Environ. Int. 156, 106645. https://doi.org/10.1016/j.envint.2021.106645

25. Ji, J.S., Liu, L., Zeng, Y., Yan, L.L., 2022. Effect of FOXO3 and Air Pollution on Cognitive Function: A Longitudinal Cohort Study of Older Adults in China From 2000 to 2014. J. Gerontol. A Biol. Sci. Med. Sci. 77, 1534–1541. https://doi.org/10.1093/gerona/glac016

26. Li, J., Wang, Y., Steenland, K., Liu, P., van Donkelaar, A., Martin, R.V., Chang, H.H., Caudle, W.M., Schwartz, J., Koutrakis, P., Shi, L., 2022. Long-term effects of PM2.5 components on incident dementia in the northeastern United States. Innovation (Camb) 3, 100208. https://doi.org/10.1016/j.xinn.2022.100208

27. Semmens, E.O., Leary, C.S., Fitzpatrick, A.L., Ilango, S.D., Park, C., Adam, C.E., DeKosky, S.T., Lopez, O., Hajat, A., Kaufman, J.D., 2023. Air pollution and dementia in older adults in the Ginkgo Evaluation of Memory Study. Alzheimers Dement. 19, 549–559. https://doi.org/10.1002/alz.12654

28. Sullivan, K.J., Ran, X., Wu, F., Chang, C.H., Sharma, R., Jacobsen, E., Berman, S., Snitz, B.E., Sekikawa, A., Talbott, E.O., Ganguli, M., 2021. Ambient fine particulate matter exposure and incident mild cognitive impairment and dementia. J. Am. Geriatr. Soc. 69, 2185–2194. https://doi.org/10.1111/jgs.17188

29. ßTan, J., Li, N., Wang, X., Chen, G., Yan, L., Wang, L., Zhao, Y., Li, S., Guo, Y., 2021. Associations of particulate matter with dementia and mild cognitive impairment in China: A multicenter cross-sectional study. Innovation (Camb) 2, 100147. https://doi.org/10.1016/j.xinn.2021.100147

30. Power, M.C., Weisskopf, M.G., Alexeeff, S.E., Coull, B.A., Spiro, A. 3rd, Schwartz, J., 2011. Traffic-related air pollution and cognitive function in a cohort of older men. Environ. Health Perspect. 119, 682–687. https://doi.org/10.1289/ehp.1002767

31. Block, M.L., Calderón-Garcidueñas, L., 2009. Air pollution: mechanisms of neuroinflammation and CNS disease. Trends Neurosci. 32, 506–516. https://doi.org/10.1016/j.tins.2009.05.009

32. Lin, M.T., Beal, M.F., 2006. Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. Nature 443, 787–795. https://doi.org/10.1038/nature05292

33. Zare Sakhvidi, M.J., Yang, J., Lequy, E., Chen, J., de Hoogh, K., Letellier, N., Mortamais, M., Ozguler, A., Vienneau, D., Zins, M., Goldberg, M., Berr, C., Jacquemin, B., 2022. Outdoor air pollution exposure and cognitive performance: findings from the enrolment phase of the CONSTANCES cohort. Lancet Planet. Health 6, e219–e229. https://doi.org/10.1016/S2542-5196(22)00001-8

34. Shi, L., Steenland, K., Li, H., Liu, P., Zhang, Y., Lyles, R.H., Requia, W.J., Ilango, S.D., Chang, H.H., Wingo, T., Weber, R.J., Schwartz, J., 2021. A national cohort study (2000-2018) of long-term air pollution exposure and incident dementia in older adults in the United States. Nat. Commun. 12, 6754. https://doi.org/10.1038/s41467-021-27049-2

35. Yao, Y., Wang, K., Xiang, H., 2022. Association between cognitive function and ambient particulate matters in middle-aged and elderly Chinese adults: Evidence from the China Health and Retirement Longitudinal Study (CHARLS). Sci. Total Environ. 828, 154297. https://doi.org/10.1016/j.scitotenv.2022.154297

36. Yokota, S., Sato, A., Umezawa, M., Oshio, S., Takeda, K., 2015. In utero exposure of mice to diesel exhaust particles affects spatial learning and memory with reduced N-methyl-D-aspartate receptor expression in the hippocampus of male offspring. Neurotoxicology 50, 108–115. https://doi.org/10.1016/j.neuro.2015.08.009

37. Hullmann, M., Albrecht, C., van Berlo, D., Gerlofs-Nijland, M.E., Wahle, T., Boots, A.W., Krutmann, J., Cassee, F.R., Bayer, T.A., Schins, R.P.F., 2017. Diesel engine exhaust accelerates plaque formation in a mouse model of Alzheimer's disease. Part. Fibre Toxicol. 14, 35. https://doi.org/10.1186/s12989-017-0213-5

38. Garrick, J.M., Cole, T.B., Dao, K., Phillips, A., Costa, L.G., 2023. Perinatal diesel exhaust exposure causes persistent changes in the brains of aged mice: An assessment of behavioral and biochemical endpoints related to neurodegenerative disease. Environ. Toxicol. 38, 899–913. https://doi.org/10.1002/tox.23733

39. Li, X., Zhang, Y., Li, B., Yang, H., Cui, J., Li, X., Zhang, X., Sun, H., Meng, Q., Wu, S., Li, S., Wang, J., Aschner, M., Chen, R., 2020. Activation of NLRP3 in microglia exacerbates diesel exhaust particles-induced impairment in learning and memory in mice. Environ. Int. 136, 105487. https://doi.org/10.1016/j.envint.2020.105487

40. Win-Shwe, T.T., Yamamoto, S., Fujitani, Y., Hirano, S., Fujimaki, H., 2008. Spatial learning and memory function-related gene expression in the hippocampus of mouse exposed to nanoparticle-rich diesel exhaust. Neurotoxicology 29, 940–947. https://doi.org/10.1016/j.neuro.2008.09.007

41. Lutz, K.L., Rakowska, S.B., Adams, M.D., 2024. Examining the impacts of school bus travel on students' academic performance in two major cities. Can. Geogr. / Géogr. Can. 68, 603–614. https://doi.org/10.1111/cag.12957

42. Pedde, M., Szpiro, A., Hirth, R.A., Adar, S.D., 2024. School Bus Rebate Program and Student Educational Performance Test Scores. JAMA Netw. Open 7, e243121. https://doi.org/10.1001/jamanetworkopen.2024.3121

43. Mallach, G., Shutt, R., Thomson, E.M., Valcin, F., Kulka, R., Weichenthal, S., 2023. Randomized Cross-Over Study of In-Vehicle Cabin Air Filtration, Air Pollution Exposure, and Acute Changes to Heart Rate Variability, Saliva Cortisol, and Cognitive Function. Environ. Sci. Technol. 57, 3238–3247. https://doi.org/10.1021/acs.est.2c06556

44. Yang, R., He, J., Zhong, K.K., Fan, M.X., Bao, T., Liu, X.C., Zhong, Q., 2024. Relationship of solid fuels use with cognitive function and efficacy of switching to cleaner fuels or using ventilation: A systematic review and meta-analysis. Environ. Res. 249, 118314. https://doi.org/10.1016/j.envres.2024.118314

45. Yaffe, K., Kanaya, A., Lindquist, K., Simonsick, E.M., Harris, T., Shorr, R.I., Tylavsky, F.A., Newman, A.B., 2004. The metabolic syndrome, inflammation, and risk of cognitive decline. JAMA 292, 2237–2242. https://doi.org/10.1001/jama.292.18.2237

46. Costache, A.D., Ignat, B.E., Grosu, C., Mastaleru, A., Abdulan, I., Oancea, A., Roca, M., Leon, M.M., Badescu, M.C., Luca, S., Jigoranu, A.R., Chetran, A., Mitu, O., Costache, I.I., Mitu, F., 2023. Inflammatory Pathways in Overweight and Obese Persons as a Potential Mechanism for Cognitive Impairment and Earlier Onset Alzeihmer's Dementia in the General Population: A Narrative Review. Biomedicines 11, 3233. https://doi.org/10.3390/biomedicines11123233

47. Yu, Y., Su, J., Jerrett, M., Paul, K.C., Lee, E., Shih, I.F., Haan, M., Ritz, B., 2023. Air pollution and traffic noise interact to affect cognitive health in older Mexican Americans. Environ. Int. 173, 107810. https://doi.org/10.1016/j.envint.2023.107810

48. Wright, B., Peters, E., Ettinger, U., Kuipers, E., Kumari, V., 2014. Understanding noise stress-induced cognitive impairment in healthy adults and its implications for schizophrenia. Noise Health 16, 166–176. https://doi.org/10.4103/1463-1741.134917

49. NHANES III https://www.cdc.gov/nchs/nhanes/

50. Fillenbaum, G.G., van Belle, G., Morris, J.C., Mohs, R.C., Mirra, S.S., Davis, P.C., Tariot, P.N., Silverman, J.M., Clark, C.M., Welsh-Bohmer, K.A., Heyman, A., 2008. Consortium to Establish a Registry for Alzheimer's Disease (CERAD): the first twenty years. Alzheimers Dement. 4, 96–109. https://doi.org/10.1016/j.jalz.2007.08.005

51. Morris, J.C., Heyman, A., Mohs, R.C., Hughes, J.P., van Belle, G., Fillenbaum, G., Mellits, E.D., Clark, C., 1989. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer's disease. Neurology 39, 1159–1165. https://doi.org/10.1212/wnl.39.9.1159

52. Clark, L.J., Gatz, M., Zheng, L., Chen, Y.L., McCleary, C., Mack, W.J., 2009. Longitudinal verbal fluency in normal aging, preclinical, and prevalent Alzheimer's disease. Am. J. Alzheimers Dis. Other Demen. 24, 461–468. https://doi.org/10.1177/1533317509345154

53. Wechsler D. WAIS Manual - Third Edition. New York: Psychological Corporation. 1997.

54. Dodge, H.H., Goldstein, F.C., Wakim, N.I., et al., 2020. Differentiating among stages of cognitive impairment in aging: Version 3 of the Uniform Data Set (UDS) neuropsychological test battery and MoCA index scores. Alzheimers Dement. (NY) 6, e12103. https://doi.org/10.1002/trc2.12103

55. Albert, M.S., DeKosky, S.T., Dickson, D., et al., 2011. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimers Dement. 7, 270–279. https://doi.org/10.1016/j.jalz.2011.03.008

56. Zhu, S., Ji, L., He, Z., Zhang, W., Tong, Y., Luo, J., Hong, Z., Zhang, Y., Yu, D., Zhang, Q., Bi, Q., 2023. Association of smoking and osteoarthritis in US (NHANES 1999-2018). Sci. Rep. 13, 3911. https://doi.org/10.1038/s41598-023-30644-6

57. Kalligeros, M., Vassilopoulos, A., Vassilopoulos, S., Victor, D.W., Mylonakis, E., Noureddin, M., 2024. Prevalence of Steatotic Liver Disease (MASLD, MetALD, and ALD) in the United States: NHANES 2017-2020. Clin. Gastroenterol. Hepatol. 22, 1330–1332.e4. https://doi.org/10.1016/j.cgh.2023.11.003

58. Kroenke, K., Spitzer, R.L., Williams, J.B., 2001. The PHQ-9: validity of a brief depression severity measure. J. Gen. Intern. Med. 16, 606–613. https://doi.org/10.1046/j.1525-1497.2001.016009606.x

59. Lee, C.L., Tzeng, H.E., Liu, W.J., Tsai, C.H., 2021. A cross-sectional analysis of the association between sleep duration and osteoporosis risk in adults using 2005-2010 NHANES. Sci. Rep. 11, 9090. https://doi.org/10.1038/s41598-021-88739-x

60. Carey, I.M., Anderson, H.R., Atkinson, R.W., Beevers, S.D., Cook, D.G., Strachan, D.P., Dajnak, D., Gulliver, J., Kelly, F.J., 2018. Are noise and air pollution related to the incidence of dementia? A cohort study in London, England. BMJ Open 8, e022404. https://doi.org/10.1136/bmjopen-2018-022404

61. Colicino, E., Power, M.C., Cox, D.G., Weisskopf, M.G., Hou, L., Alexeeff, S.E., Sanchez-Guerra, M., Vokonas, P., Spiro, A. 3rd, Schwartz, J., Baccarelli, A.A., 2014. Mitochondrial haplogroups modify the effect of black carbon on age-related cognitive impairment. Environ. Health 13, 42. https://doi.org/10.1186/1476-069X-13-42

62. Colicino, E., Wilson, A., Frisardi, M.C., Prada, D., Power, M.C., Hoxha, M., Dioni, L., Spiro, A., Vokonas, P.S., Weisskopf, M.G., Schwartz, J.D., Baccarelli, A.A., 2017. Telomere Length, Long-Term Black Carbon Exposure, and Cognitive Function in a Cohort of Older Men: The VA Normative Aging Study. Environ. Health Perspect. 125, 76–81. https://doi.org/10.1289/EHP241Benjamin, D.I., Brett, J.O., Both, P., Benjamin, J.S., Ishak, H.L., Kang, J., Kim, S., Chung, M., Arjona, M., Nutter, C.W., Tan, J.H., Krishnan, A.K., Dulay, H., Louie, S.M., de Morree, A., Nomura, D.K., Rando, T.A., 2023. Multiomics reveals glutathione metabolism as a driver of bimodality during stem cell aging. Cell Metab. 35, 472–486.e6. https://doi.org/10.1016/j.cmet.2023.02.001

63. Chen, G., Qian, Z.M., Zhang, J., Wang, X., Zhang, Z., Cai, M., Arnold, L.D., Abresch, C., Wang, C., Liu, Y., Fan, Q., Lin, H., 2024. Associations between Changes in Exposure to Air Pollutants due to Relocation and the Incidence of 14 Major Disease Categories and All-Cause Mortality: A Natural Experiment Study. Environ. Health Perspect. 132, 97012. https://doi.org/10.1289/EHP14367

64. Lim, C., Oh, H., 2023. Organic solvent exposure for the chronic kidney disease: updated systematic review with meta-analysis. Ann. Occup. Environ. Med. 35, e11. https://doi.org/10.35371/aoem.2023.35.e11

65. Lalouette, A., Chaumot, A., Lepeule, L., Gaget, K., Delorme, N., Garnero, L., Calevro, F., Esposti, D.D., 2025. Cadmium tolerance is associated with tissue-specific plasticity of metallothionein gene expression in Gammarus fossarum field populations. Sci. Rep. 15, 13913. https://doi.org/10.1038/s41598-025-98937-6

66. Fowler, S.L., Behr, T.S., Turkes, E., O'Brien, D.P., Cauhy, P.M., Rawlinson, I., Edmonds, M., Foiani, M.S., Schaler, A., Crowley, G., Bez, S., Ficulle, E., Tsefou, E., Fischer, R., Geary, B., Gaur, P., Miller, C., D'Acunzo, P., Levy, E., Duff, K.E., Ryskeldi-Falcon, B., 2025. Tau filaments are tethered within brain extracellular vesicles in Alzheimer's disease. Nat. Neurosci. 28, 40–48. https://doi.org/10.1038/s41593-024-01801-5

67. Cromerg Cromberg, J.L., Schettgen, T., Willmes, K., Lang, J., Kraus, T., Fimm, B., 2024. Occupational exposure to polychlorinated biphenyls: Development of neuropsychological functions over time. Neurotoxicology 101, 6–15. https://doi.org/10.1016/j.neuro.2024.01.001

68. López-Vicente, M., Kusters, M., Binter, A.C., Petricola, S., Tiemeier, H., Muetzel, R., Guxens, M., 2025. Long-term exposure to traffic-related air pollution and noise and dynamic brain connectivity across adolescence. Environ. Health Perspect. https://doi.org/10.1289/EHP14525

69. Bustamante-Barrientos, F.A., Méndez-Ruette, M., Ortloff, A., Luz-Crawford, P., Rivera, F.J., Figueroa, C.D., Molina, L., Bátiz, L.F., 2021. The Impact of Estrogen and Estrogen-Like Molecules in Neurogenesis and Neurodegeneration: Beneficial or Harmful? Front. Cell. Neurosci. 15, 636176. https://doi.org/10.3389/fncel.2021.636176

70. Demarest, T.G., McCarthy, M.M., 2015. Sex differences in mitochondrial (dys)function: Implications for neuroprotection. J. Bioenerg. Biomembr. 47, 173–188. https://doi.org/10.1007/s10863-014-9583-7

71. Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne, C., Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S.G., Dias, A., Fox, N., Gitlin, L.N., Howard, R., Kales, H.C., Kivimäki, M., Larson, E.B., Ogunniyi, A., Orgeta, V., Ritchie, K., Rockwood, K., Sampson, E.L., Samus, Q., Schneider, L.S., Selbæk, G., Teri, L., Mukadam, N., 2020. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. Lancet 396, 413–446. https://doi.org/10.1016/S0140-6736(20)30367-6

Figure 1. Flowchart for selecting eligible population in the study.

Table 1. Baseline characteristics of participants by exposure status to exhaust fumes at work (n=1110).

	No. (%)	Exposed group No. (%)	Never Exposed group No. (%)	*P* Value
Total	1110 (100.0)	266 (24.0)	844 (76.0)	
Age (years)				0.576
60-69	727 (65.5)	178 (16.1)	549 (49.4)	
70-79	383 (34.5)	88 (7.9)	295 (26.6)	
Gender				**0.0001**
Male	570 (51.4)	217 (19.6)	353 (31.8)	
Female	540 (48.6)	49 (4.4)	491 (44.2)	
Race/Ethnicity				**0.019**
Mexican American and Other Hispanic	214 (19.3)	51 (4.6)	163 (14.7)	
Non-Hispanic White	435 (39.2)	113 (10.2)	322 (29.0)	
Non-Hispanic Black	347 (31.2)	88 (7.9)	259 (23.3)	
Non-Hispanic Asian and Other Race	114 (10.3)	14 (1.3)	100 (9.0)	
Education Level				**0.017**
< HS	290 (26.1)	84 (7.5)	206 (18.6)	
HSG	245 (22.1)	65 (5.9)	180 (16.2)	
> HS	575 (51.8)	118 (10.6)	457 (41.2)	

	No. (%)	Exposed group No. (%)	Never Exposed group No. (%)	P Value
Smoking history (Ever smoke)				**0.0001**
>100 cigarettes	583 (52.5)	173 (15.6)	410 (36.9)	
< 100 cigarettes	527 (47.5)	93 (8.4)	434 (39.1)	
Excessive Alcohol Use[†]				**0.001**
Yes	59 (9.3)	26 (4.1)	33 (5.2)	
No	576 (90.7)	140 (22.0)	436 (68.7)	
Obesity				0.107
Yes	466 (42.0)	123 (11.1)	343 (30.9)	
No	644 (58.0)	143 (12.9)	501 (45.1)	
Hypertension				0.555
Yes	655 (59.0)	161 (14.5)	493 (44.5)	
No	455 (41.0)	105 (9.5)	350 (31.5)	
Dyslipidemia[*]				0.290
Yes	645 (62.6)	150 (14.5)	495 (48.0)	
No	386 (37.4)	101 (9.8)	285 (27.7)	
Hyperglycemia[§]				0.640
Yes	759 (71.3)	187 (17.6)	572 (53.7)	
No	305 (28.7)	71 (6.7)	234 (22.0)	

	No. (%)	Exposed group No. (%)	Never Exposed group No. (%)	*P* Value
History of CVD				0.098
Yes	204 (18.4)	58 (5.2)	146 (13.2)	
No	906 (81.6)	208 (18.7)	698 (62.9)	
Depression (PHQ9 score)				0.215
Yes ($\geq$ 10)	105 (9.5)	20 (1.8)	85 (7.7)	
No (< 10)	1005 (90.5)	246 (22.1)	759 (68.4)	
Have Sleep Problems				0.860
Yes	379 (34.3)	92 (8.3)	287 (26.0)	
No	727 (65.7)	173 (15.6)	554 (50.1)	

[*][†][§]Missing data: Dyslipidemia (79), excessive alcohol use (475), hyperglycemia (46).

Figure 2. Forest plot of weighted multivariable survey logistic regression analyses on the association between occupational exhaust fumes exposure and cognitive decline (n = 1,110). Adjusted for: Model 1, no adjustment; Model 2, age, gender, and educational level; Model 3, age, gender, educational level, race/ethnicity, History of CVD, Depression and Hyperglycemia. Note: CVD, cardiovascular diseases. CERAD-WL, the Consortium to Establish a Registry for Alzheimer's Disease Word List Learning Recall Test. AFT, the Animal Fluency Test. DSST, the Digit Symbol Substitution Test. CI, confidence interval.

Figure 3. Associations between occupational exhaust fumes exposure and cognitive decline in NHANES participants. The exposure-response curves show the dose-dependent effects of exposure on cognitive outcomes, with the effect sizes quantified as changes in cognitive scores per unit (10 years) increase in exposure. The three-dimensional contour plots show the joint effects of exposure intensity and duration (lag time) on cognitive function scores, showing biphasic pattens characterized by initial adaptive responses followed by potential neurotoxic effects at higher cumulative exposures. Logistic regression models with DLNMs were adjusted for age, gender, educational level, race/ethnicity, History of CVD, depression and hyperglycemia. The solid lines represent the estimated effect sizes from the fitted DLNM, and the shaded areas represent the 95% confidence intervals around the estimate for each exposure level. Note: NHANES, National Health and Nutrition Examination Survey. CVD, cardiovascular diseases. DLNM, distributed lag nonlinear model.