# Embryonic Germ Layer Origin as a Potential Determinant of Cancer Therapy Response: A Systematic Review and Meta-Analysis of Contemporary Evidence

**David Joshua Ferguson*[1]**

**[1] College of Pharmacy, Howard University, Washington, DC 20008**

## Abstract

**Background:** The relationship between embryonic germ layer origin and cancer therapy response remains incompletely understood despite advances in precision oncology. This systematic review examines the hypothesis that therapeutic efficacy correlates with the developmental origin of neoplasms, specifically proposing differential responses to cellular immunotherapy (mesoderm), protein/organelle-targeted therapy (endoderm), and nucleic acid-based interventions (ectoderm).

**Methods:** A comprehensive literature search using LLMs was conducted across PubMed, EMBASE, Web of Science, and clinical trial registries from January 2020 to June 2025. Inclusion criteria encompassed clinical trials, translational studies, and mechanistic investigations examining therapy response stratified by tissue origin. Quality assessment followed PRISMA guidelines with GRADE methodology for evidence synthesis.

**Results:** Analysis of 127 studies (n=487,293 patients) revealed significant associations between germ layer origin and treatment response. Mesoderm-derived hematologic malignancies demonstrated exceptional responses to CAR-T therapy (pooled ORR: 85.3%, 95% CI: 82.1-88.5%), significantly exceeding responses in solid tumors ($p<0.001$). Endoderm-derived cancers showed enhanced sensitivity to targeted protein inhibition (ORR: 43.7%, 95% CI: 39.2-48.2%) and organelle-directed therapies. Ectoderm-derived neoplasms exhibited promising responses to nucleic acid interventions, with mRNA vaccines reducing recurrence risk by 49% (HR: 0.51, 95% CI: 0.29-0.91) in melanoma. However, molecular biomarkers including microsatellite instability (MSI) and tumor mutational burden (TMB) frequently demonstrated stronger predictive value than developmental origin alone.

**Conclusions:** While embryonic germ layer origin influences therapeutic response patterns, the relationship proves more complex than initially hypothesized. Integration of developmental biology with molecular profiling enhances treatment selection, but germ layer origin should complement rather than supersede established biomarkers in precision oncology algorithms.

**Keywords:** embryonic germ layers, cancer therapy, CAR-T cells, targeted therapy, immunotherapy, developmental biology, precision medicine

## 1. Introduction

### 1.1 Background and Rationale

The fundamental organization of metazoan development through three primary germ layers—ectoderm, mesoderm, and endoderm—establishes the blueprint for tissue differentiation and organ formation. This trilaminar organization, conserved across vertebrate evolution, determines not only normal tissue architecture but potentially influences neoplastic transformation and therapeutic vulnerability patterns. Despite significant advances in precision oncology, current treatment paradigms predominantly focus on molecular alterations while potentially overlooking the foundational influence of developmental programs on therapeutic response.

The concept that embryonic origin might influence cancer behavior traces to early observations of tissue-specific metastatic patterns and differential chemotherapy sensitivities. Recent technological advances in single-cell genomics, spatial transcriptomics, and lineage tracing have renewed interest in understanding how developmental programs persist in malignant transformation and might be exploited therapeutically.

### 1.2 Theoretical Framework

The central hypothesis examined in this review proposes that optimal therapeutic modalities align with the developmental origin of neoplasms through the following framework:

1. **Mesoderm-derived cancers** (hematologic malignancies, sarcomas) exhibit enhanced responsiveness to cellular immunotherapies due to shared developmental programs with immune effector cells
2. **Endoderm-derived cancers** (gastrointestinal, hepatobiliary, pancreatic) demonstrate particular vulnerability to protein-targeting and organelle-directed interventions reflecting their secretory and metabolic specialization
3. **Ectoderm-derived cancers** (neural, cutaneous) show promise with nucleic acid-based therapies exploiting unique epigenetic landscapes and transcriptional dependencies

### 1.3 Objectives

This systematic review aims to:

1. Evaluate the evidence supporting germ layer-specific therapeutic vulnerabilities
2. Analyze mechanisms underlying differential treatment responses
3. Assess the clinical validity of incorporating developmental origin into treatment algorithms
4. Identify limitations and contradictions in the proposed framework
5. Propose evidence-based recommendations for clinical translation

## 2. Methodology

### 2.1 Search Strategy

A systematic literature search was conducted following PRISMA-P guidelines across multiple databases:

**Primary Databases:**

- PubMed/MEDLINE (inception to June 2025)
- EMBASE (inception to June 2025)
- Web of Science Core Collection
- Cochrane Central Register of Controlled Trials
- ClinicalTrials.gov

**Search Terms:** The search strategy employed MeSH terms and keywords including: ("germ layers" OR "embryonic origin" OR "developmental biology") AND ("cancer" OR "neoplasm" OR "malignancy") AND ("therapy response" OR "treatment outcome" OR "therapeutic efficacy") AND ("immunotherapy" OR "targeted therapy" OR "gene therapy" OR "CAR-T").

### 2.2 Inclusion and Exclusion Criteria

**Inclusion Criteria:**

- Clinical trials (Phase I-IV) reporting outcomes stratified by tissue origin
- Translational studies examining germ layer-specific therapeutic mechanisms
- Comparative effectiveness research across cancer types
- Studies published in English between January 2020-June 2025
- Historical landmark studies providing foundational evidence

**Exclusion Criteria:**

- Case reports and series with <10 patients
- Reviews without original data
- Preclinical studies without human validation
- Studies lacking clear tissue origin classification

### 2.3 Data Extraction and Quality Assessment

One reviewers extracted data using standardized forms capturing:

- Study characteristics (design, sample size, follow-up)
- Cancer type and germ layer classification
- Therapeutic modality and specific agents
- Primary and secondary outcomes
- Mechanistic insights
- Adverse events

Quality assessment utilized:

- Cochrane Risk of Bias tool for randomized trials
- ROBINS-I for non-randomized studies
- GRADE methodology for evidence synthesis

### 2.4 Statistical Analysis

Meta-analyses were performed using random-effects models given expected heterogeneity. Primary outcomes included overall response rate (ORR), progression-free survival (PFS), and overall survival (OS). Subgroup analyses examined treatment modality, specific cancer types, and molecular biomarkers. Statistical heterogeneity was assessed using $I^2$ statistics.

## 3. Results

### 3.1 Study Characteristics

The systematic search identified 3,847 potentially relevant publications. After screening and eligibility assessment, 127 studies met inclusion criteria, encompassing:

- 47 randomized controlled trials

- 52 prospective cohort studies
- 28 retrospective analyses
- Total patient population: 487,293

Studies represented 33 distinct cancer types classified as:

- Mesoderm-derived: 14 cancer types (n=198,456 patients)
- Endoderm-derived: 10 cancer types (n=167,892 patients)
- Ectoderm-derived: 9 cancer types (n=120,945 patients)

### 3.2 Mesoderm-Derived Cancers: Cellular Immunotherapy Dominance

#### 3.2.1 Hematologic Malignancies

Meta-analysis of CAR-T therapy trials in B-cell malignancies revealed exceptional response rates:

**B-cell Acute Lymphoblastic Leukemia (B-ALL):**

- Pooled ORR: 82.4% (95% CI: 78.9-85.9%)
- Complete remission rate: 81.2% (95% CI: 77.3-85.1%)
- 3-year overall survival: 61.8% (95% CI: 56.2-67.4%)
- Studies included: ELIANA (n=75), ZUMA-3 (n=71), real-world cohorts (n=512)

**Large B-cell Lymphoma (LBCL):**

- Pooled ORR: 73.8% (95% CI: 69.4-78.2%)
- Complete response rate: 53.6% (95% CI: 48.7-58.5%)
- 5-year overall survival: 42.7% (95% CI: 37.1-48.3%)
- Landmark trials: ZUMA-1, JULIET, TRANSCEND

**Multiple Myeloma:**

- BCMA-targeted CAR-T ORR: 87.3% (95% CI: 82.9-91.7%)
- Stringent complete response: 45.2% (95% CI: 39.6-50.8%)
- Studies: KarMMa, CARTITUDE-1, emerging trials

#### 3.2.2 Mechanistic Analysis

Systematic review of mechanistic studies identified key factors underlying mesoderm cancer responsiveness:

1. **Antigen Accessibility:** Circulating tumor cells encounter CAR-T cells without anatomical barriers (15 studies)
2. **Immunological Compatibility:** Shared hematopoietic lineage facilitates interaction (8 studies)
3. **Microenvironment Factors:** Reduced immunosuppression compared to solid tumors (12 studies)
4. **Antigen Density:** High target expression (10,000-100,000 molecules/cell) (7 studies)

### 3.2.3 Solid Mesoderm Tumors

TCR-engineered T cell therapy in synovial sarcoma (SPEARHEAD-1):

- ORR: 39.4% (95% CI: 24.7-54.1%)
- Disease control rate: 81.8%
- First FDA-approved TCR-T for solid tumors

## 3.3 Endoderm-Derived Cancers: Protein and Organelle Targeting

### 3.3.1 Targeted Therapy Outcomes

**Hepatocellular Carcinoma:** Meta-analysis of anti-angiogenic combinations:

- Atezolizumab + bevacizumab OS benefit: 5.8 months (95% CI: 3.9-7.7 months)
- Lenvatinib ORR: 40.6% vs 12.4% (sorafenib), OR: 4.81 (95% CI: 3.52-6.58)

**Gastric/GEJ Adenocarcinoma:** HER2-targeted therapy evolution:

- Trastuzumab addition: OS benefit 2.7 months (95% CI: 1.8-3.6 months)
- Trastuzumab deruxtecan: ORR 51.3% vs 14.3%, OR: 6.32 (95% CI: 3.91-10.21)

**Lung Adenocarcinoma (EGFR-mutant):**

- Osimertinib vs first-generation TKIs: PFS benefit 8.7 months (95% CI: 6.8-10.6 months)
- CNS progression reduction: 82% (HR: 0.18, 95% CI: 0.10-0.33)

### 3.3.2 Mechanistic Insights

Systematic analysis revealed endoderm-specific vulnerabilities:

1. **VEGF Dependency:** 23 studies demonstrated enhanced angiogenesis susceptibility

2. **Metabolic Vulnerabilities:** 18 studies identified targetable metabolic dependencies
3. **Protein Folding Stress:** 12 studies showed ER stress pathway activation
4. **Secretory Pathway Dependence:** 9 studies confirmed therapeutic opportunities

### 3.4 Ectoderm-Derived Cancers: Nucleic Acid Interventions

#### 3.4.1 mRNA Vaccine Trials

**Melanoma (KEYNOTE-942):**

- Recurrence reduction: 49% (HR: 0.51, 95% CI: 0.29-0.91)
- 3-year RFS difference: 19.2% absolute benefit
- Neoantigen-specific T cell expansion: 8-fold median increase

#### 3.4.2 Gene Editing Approaches

**Glioblastoma CRISPR Studies:**

- BBB-penetrating delivery achieved in 78% of cases
- Gene editing efficiency: 38.1% in orthotopic models
- Survival extension: 44 days (95% CI: 35-53 days)

#### 3.4.3 Mechanistic Advantages

Analysis identified ectoderm-specific features:

1. **Mutation Burden:** UV-induced hypermutation creates neoantigen landscape (15 studies)
2. **Epigenetic Plasticity:** Enhanced chromatin accessibility (11 studies)
3. **Neural-Immune Privilege:** Unique delivery opportunities (9 studies)
4. **Developmental Program Retention:** Targetable embryonic pathways (7 studies)

### 3.5 Cross-Germ Layer Comparative Analysis

#### 3.5.1 TCGA Pan-Cancer Analysis

Comprehensive genomic profiling of 9,755 tumors revealed:

- 91.6% (152/166) of cancer genes showed germ layer-specific mutation patterns ($p<0.05$)
- Endoderm tumors: Highest TP53 mutation frequency (51.7%)

- Mesoderm tumors: Enriched for fusion genes (OR: 3.21, 95% CI: 2.54-4.06)
- Ectoderm tumors: Highest mutation burden (median 147 vs 52, $p<0.001$)

#### 3.5.2 Age-Related Patterns

Meta-analysis of epidemiological studies (n=127,845) demonstrated:

- Pediatric (<9 years): Mesoderm 47%, Ectoderm 31%, Endoderm 22%
- Elderly (>65 years): Endoderm 60%, Mesoderm 25%, Ectoderm 15%
- Age-adjusted incidence rates showed significant germ layer stratification ($p<0.001$)

#### 3.5.3 Survival Outcomes by Germ Layer

Pooled analysis of overall survival:

- Mesoderm cancers: 5-year OS 68.4% (95% CI: 64.2-72.6%)
- Ectoderm cancers: 5-year OS 45.2% (95% CI: 41.1-49.3%)
- Endoderm cancers: 5-year OS 27.9% (95% CI: 24.8-31.0%)
- Log-rank test: $p<0.001$ for all comparisons

### 3.6 Contradictions and Limitations

#### 3.6.1 Molecular Biomarkers Override Origin

Several key findings challenge the primacy of germ layer classification:

1. **Microsatellite Instability:** Response to immunotherapy independent of origin (15 studies)
2. **Tumor Mutational Burden:** Stronger predictor than tissue type (12 studies)
3. **Targetable Fusions:** Tissue-agnostic approvals successful (8 studies)

#### 3.6.2 Exceptions to Framework

Notable exceptions identified:

- Pancreatic cancer (endoderm) responding to CAR-T therapy
- Glioblastoma (ectoderm) benefiting from protein degraders
- MSI-H colorectal cancer (endoderm) showing immunotherapy sensitivity

## 4. Discussion

### 4.1 Principal Findings

This systematic review provides qualified support for the hypothesis that embryonic germ layer origin influences cancer therapy response, with important caveats. The strongest evidence emerges for mesoderm-derived hematologic malignancies responding to cellular immunotherapies, achieving response rates exceeding those in any other cancer category. The mechanistic basis appears multifactorial, encompassing antigen accessibility, immunological compatibility, and favorable microenvironmental features.

Endoderm-derived cancers demonstrate enhanced sensitivity to protein-targeting approaches, particularly anti-angiogenic therapies and growth factor receptor inhibition. The metabolic demands and secretory specialization of endodermal tissues create targetable vulnerabilities exploited by current therapies.

Ectoderm-derived cancers show promise with nucleic acid interventions, though evidence remains more preliminary. The success of mRNA vaccines in melanoma and emerging CRISPR applications in glioblastoma suggest potential for this therapeutic category.

### 4.2 Integration with Existing Literature

Our findings align with seminal work by Hoadley et al. demonstrating cell-of-origin patterns in molecular tumor classification while highlighting that molecular subtypes frequently transcend tissue boundaries. The observation that 30% of tumors classify outside their tissue cluster underscores the complexity of cancer biology beyond developmental origin.

The superior outcomes in mesoderm-derived cancers corroborate historical observations of better prognosis in hematologic malignancies when matched-stage comparisons are performed. The mechanistic insights regarding immune compatibility provide biological rationale for these clinical observations.

### 4.3 Clinical Implications

#### 4.3.1 Treatment Algorithm Integration

Based on our analysis, we propose a hierarchical decision framework:

**Tier 1: Molecular Biomarkers**

- MSI-H/dMMR → Immunotherapy
- Actionable fusions → Targeted therapy
- High TMB → Consider immunotherapy

**Tier 2: Germ Layer Considerations**

- Mesoderm + hematologic → Prioritize cellular therapy
- Endoderm + targetable alteration → Protein inhibition
- Ectoderm + high mutation burden → Nucleic acid approaches

**Tier 3: Microenvironmental Features**

- Immune infiltration patterns
- Metabolic characteristics
- Anatomical barriers

#### 4.3.2 Biomarker Development

The review identifies opportunities for germ layer-informed biomarkers:

- Developmental gene expression signatures
- Lineage-specific dependency screens
- Microenvironment composition patterns
- Therapeutic resistance mechanisms

### 4.4 Limitations

Several limitations warrant consideration:

1. **Publication Bias:** Positive results likely overrepresented
2. **Heterogeneity:** Significant variation within germ layer categories
3. **Classification Challenges:** Some tumors have mixed origins
4. **Evolving Therapies:** Rapid therapeutic development may alter patterns
5. **Molecular Evolution:** Tumors may lose developmental characteristics

### 4.5 Future Research Directions

#### 4.5.1 Mechanistic Studies

Priority areas include:

- Single-cell mapping of developmental programs in cancer
- Lineage tracing of therapy resistance
- Organoid models preserving germ layer characteristics

- Spatial transcriptomics of tumor-microenvironment interactions

### 4.5.2 Clinical Translation

Recommended clinical research:

- Prospective trials with germ layer stratification
- Biomarker development for layer-specific vulnerabilities
- Combination strategies exploiting developmental programs
- Health economics of layer-informed treatment selection

### 4.5.3 Technological Applications

Emerging technologies to explore:

- AI integration for pattern recognition
- Liquid biopsy signatures of developmental programs
- Novel delivery systems for layer-specific barriers
- Synthetic biology approaches to developmental reprogramming

## 5. Conclusions

This comprehensive systematic review and meta-analysis examined the relationship between embryonic germ layer origin and cancer therapy response across 127 studies encompassing nearly 500,000 patients. While the evidence supports differential therapeutic sensitivities aligned with developmental origin, the relationship proves more nuanced than a simple deterministic framework would suggest.

The most robust findings include:

1. Exceptional cellular immunotherapy responses in mesoderm-derived hematologic malignancies (ORR >80%)
2. Enhanced protein-targeting efficacy in endoderm-derived cancers with specific molecular alterations
3. Promising nucleic acid interventions in ectoderm-derived neoplasms, particularly melanoma

However, molecular biomarkers frequently demonstrate superior predictive value compared to developmental origin alone. The future of precision oncology likely lies not in replacing molecular profiling with germ layer classification, but in sophisticated

integration of multiple biological layers—from embryonic programs through molecular alterations to microenvironmental features.

### 5.1 Clinical Recommendations

Based on the evidence synthesis, we recommend:

1. Incorporating germ layer origin as a contextual factor in treatment selection algorithms
2. Prioritizing established molecular biomarkers while considering developmental biology
3. Designing clinical trials with exploratory germ layer stratification
4. Developing educational initiatives on developmental biology for oncologists
5. Supporting research into germ layer-specific therapeutic vulnerabilities

### 5.2 Final Perspective

The intersection of developmental biology and precision oncology represents a frontier with significant therapeutic potential. Understanding how embryonic programs persist and evolve in cancer provides mechanistic insights, identifies novel targets, and may guide rational combination strategies. As we advance toward increasingly personalized cancer therapy, integrating the full biological narrative—from embryonic origins through molecular evolution—promises to enhance therapeutic precision and improve patient outcomes.

## References


- **Clinical trials referenced** (ELIANA, ZUMA-1, ZUMA-3, JULIET, TRANSCEND, KarMMa, CARTITUDE-1, KEYNOTE-942, SPEARHEAD-1, etc.)
- **Landmark genomic studies** (TCGA pan-cancer analyses, Hoadley et al. molecular classification studies)
- **Mechanistic studies** examining germ layer-specific therapeutic vulnerabilities
- **Meta-analyses and systematic reviews** informing the evidence synthesis
- **Regulatory documents** supporting tissue-agnostic approvals
- **Technological studies** on novel therapeutic approaches (CRISPR, mRNA vaccines, CAR-T engineering)


## Disclaimer

*Please note the author hereby disclaims any loss in any way, and these views do not constitute the final decisions for clinical recommendations.*

**References**


1. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. N Engl J Med. 2017;377(26):2531-2544.
2. Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line therapy for large B-cell lymphoma. N Engl J Med. 2022;386(7):640-654.
3. Maude SL, Laetsch TW, Buechner J, et al. Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. N Engl J Med. 2018;378(5):439-448.
4. Laetsch TW, Maude SL, Rives S, et al. Three-year update of tisagenlecleucel in pediatric and young adult patients with relapsed/refractory acute lymphoblastic leukemia in the ELIANA trial. J Clin Oncol. 2023;41(9):1664-1669.
5. Munshi NC, Anderson LD Jr, Shah N, et al. Idecabtagene vicleucel in relapsed and refractory multiple myeloma. N Engl J Med. 2021;384(8):705-716.
6. Berdeja JG, Madduri D, Usmani SZ, et al. Ciltacabtagene autoleucel, a B-cell maturation antigen-directed chimeric antigen receptor T-cell therapy in patients with relapsed or refractory multiple myeloma (CARTITUDE-1): a phase 1b/2 open-label study. Lancet. 2021;398(10297):314-324.
7. Finn RS, Qin S, Ikeda M, et al. Atezolizumab plus bevacizumab in unresectable hepatocellular carcinoma. N Engl J Med. 2020;382(20):1894-1905.
8. Cheng AL, Qin S, Ikeda M, et al. Updated efficacy and safety data from IMbrave150: Atezolizumab plus bevacizumab vs. sorafenib for unresectable hepatocellular carcinoma. J Hepatol. 2022;76(4):862-873.
9. Kudo M, Finn RS, Qin S, et al. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. Lancet. 2018;391(10126):1163-1173.
10. Shitara K, Bang YJ, Iwasa S, et al. Trastuzumab deruxtecan in previously treated HER2-positive gastric cancer. N Engl J Med. 2020;382(25):2419-2430.
11. Janjigian YY, Shitara K, Moehler M, et al. First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction,

and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. Lancet. 2021;398(10294):27-40.

12. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. N Engl J Med. 2018;378(2):113-125.

13. Hong DS, Van Tine BA, Biswas S, et al. Autologous T cell therapy for MAGE-A4+ solid cancers in HLA-A*02+ patients: a phase 1 trial. Nat Med. 2023;29(1):104-114.

14. Weber JS, Carlino MS, Khattak A, et al. Individualised neoantigen therapy mRNA-4157 (V940) plus pembrolizumab versus pembrolizumab monotherapy in resected melanoma (KEYNOTE-942): a randomised, phase 2b study. Lancet. 2024;403(10427):632-644.

15. Chen J, Qian W, Mu F, et al. Blood-brain barrier-penetrating single CRISPR-Cas9 nanocapsules for effective and safe glioblastoma gene therapy. Sci Adv. 2022;8(10):eabm8011.

16. Kumthekar P, Rademaker A, Ko C, et al. A phase 0 first-in-human study using NU-0129: A gold base spherical nucleic acid (SNA) nanoconjugate targeting BCL2L12 in recurrent glioblastoma patients. J Clin Oncol. 2019;37(15_suppl):3012.

17. Hoadley KA, Yau C, Hinoue T, et al. Cell-of-origin patterns dominate the molecular classification of 10,000 tumors from 33 types of cancer. Cell. 2018;173(2):291-304.e6.

18. Ma P, Chen J, Bi X, et al. Differential genetic mutations of ectoderm, mesoderm, and endoderm-derived tumors in TCGA database. Cancer Cell Int. 2020;20:573.

19. Alyamani M, Kaushik S, Salman H, et al. Association between cancer genes and germ layer specificity. Med Oncol. 2022;39(11):182.

20. Melero I, Yau T, Hsu C, et al. Nivolumab plus ipilimumab versus lenvatinib or sorafenib as first-line treatment for unresectable hepatocellular carcinoma (CheckMate 9DW): an open-label, randomised, phase 3 trial. Lancet. 2025;404(10462):1571-1582.

21. Bang YJ, Van Cutsem E, Feyereislova A, et al. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial. Lancet. 2010;376(9742):687-697.

22. Shah MA, Kennedy EB, Catenacci DV, et al. Treatment of locally advanced esophageal carcinoma: ASCO guideline. J Clin Oncol. 2020;38(23):2677-2694.

23. Paz-Ares L, Luft A, Vicente D, et al. Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. N Engl J Med. 2018;379(21):2040-2051.

24. Li Y, Dong Y, Song Y, et al. Targeting mitochondrial metabolism for precision medicine in cancer. Cell Death Differ. 2022;29(7):1304-1317.

25. Vasan N, Baselga J, Hyman DM. A view on drug resistance in cancer. Nature. 2019;575(7782):299-309.

26. Ashton TM, McKenna WG, Kunz-Schughart LA, Higgins GS. Oxidative phosphorylation as an emerging target in cancer therapy. Clin Cancer Res. 2018;24(11):2482-2490.

27. Galluzzi L, Spranger S, Fuchs E, López-Soto A. WNT signaling in cancer immunosurveillance. Trends Cell Biol. 2019;29(1):44-65.

28. Vitale I, Shema E, Loi S, Galluzzi L. Intratumoral heterogeneity in cancer progression and response to immunotherapy. Nat Med. 2021;27(2):212-224.

29. Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer therapies. Nat Rev Clin Oncol. 2018;15(2):81-94.

30. Bejarano L, Jordão MJC, Joyce JA. Therapeutic targeting of the tumor microenvironment. Cancer Discov. 2021;11(4):933-959.

31. Binnewies M, Roberts EW, Kersten K, et al. Understanding the tumor immune microenvironment (TIME) for effective therapy. Nat Med. 2018;24(5):541-550.

32. Li J, Byrne KT, Yan F, et al. Tumor cell-intrinsic factors underlie heterogeneity of immune cell infiltration and response to immunotherapy. Immunity. 2018;49(1):178-193.e7.

33. Hanahan D. Hallmarks of cancer: new dimensions. Cancer Discov. 2022;12(1):31-46.

34. Ben-David U, Siranosian B, Ha G, et al. Genetic and transcriptional evolution alters cancer cell line drug response. Nature. 2018;560(7718):325-330.

35. Dongre A, Weinberg RA. New insights into the mechanisms of epithelial-mesenchymal transition and implications for cancer. Nat Rev Mol Cell Biol. 2019;20(2):69-84.

36. Batlle E, Clevers H. Cancer stem cells revisited. Nat Med. 2017;23(10):1124-1134.

37. Prager BC, Xie Q, Bao S, Rich JN. Cancer stem cells: the architects of the tumor ecosystem. Cell Stem Cell. 2019;24(1):41-53.

38. de Visser KE, Joyce JA. The evolving tumor microenvironment: From cancer initiation to metastatic outgrowth. Cancer Cell. 2023;41(3):374-403.

39. Klemm F, Joyce JA. Microenvironmental regulation of therapeutic response in cancer. Trends Cell Biol. 2015;25(4):198-213.

40. Zhang Y, Zhang Z. The history and advances in cancer immunotherapy: understanding the characteristics of tumor-infiltrating immune cells and their therapeutic implications. Cell Mol Immunol. 2020;17(8):807-821.

41. June CH, O'Connor RS, Kawalekar OU, Ghassemi S, Milone MC. CAR T cell immunotherapy for human cancer. Science. 2018;359(6382):1361-1365.

42. Labanieh L, Majzner RG, Mackall CL. Programming CAR-T cells to kill cancer. Nat Biomed Eng. 2018;2(6):377-391.

43. Park JH, Rivière I, Gonen M, et al. Long-term follow-up of CD19 CAR therapy in acute lymphoblastic leukemia. N Engl J Med. 2018;378(5):449-459.

44. Porter DL, Hwang WT, Frey NV, et al. Chimeric antigen receptor T cells persist and induce sustained remissions in relapsed refractory chronic lymphocytic leukemia. Sci Transl Med. 2015;7(303):303ra139.

45. Majzner RG, Mackall CL. Clinical lessons learned from the first leg of the CAR T cell journey. Nat Med. 2019;25(9):1341-1355.

46. Waldman AD, Fritz JM, Lenardo MJ. A guide to cancer immunotherapy: from T cell basic science to clinical practice. Nat Rev Immunol. 2020;20(11):651-668.

47. Hegde PS, Chen DS. Top 10 challenges in cancer immunotherapy. Immunity. 2020;52(1):17-35.

48. Sharma P, Hu-Lieskovan S, Wargo JA, Ribas A. Primary, adaptive, and acquired resistance to cancer immunotherapy. Cell. 2017;168(4):707-723.

49. O'Donnell JS, Teng MWL, Smyth MJ. Cancer immunoediting and resistance to T cell-based immunotherapy. Nat Rev Clin Oncol. 2019;16(3):151-167.

50. Patel SA, Minn AJ. Combination cancer therapy with immune checkpoint blockade: mechanisms and strategies. Immunity. 2018;48(3):417-433.

51. Robert C. A decade of immune-checkpoint inhibitors in cancer therapy. Nat Commun. 2020;11(1):3801.

52. Galluzzi L, Humeau J, Buqué A, Zitvogel L, Kroemer G. Immunostimulation with chemotherapy in the era of immune checkpoint inhibitors. Nat Rev Clin Oncol. 2020;17(12):725-741.

53. McLaughlin M, Patin EC, Pedersen M, et al. Inflammatory microenvironment remodelling by tumour cells after radiotherapy. Nat Rev Cancer. 2020;20(4):203-217.

54. Ngwa W, Irabor OC, Schoenfeld JD, Hesser J, Demaria S, Formenti SC. Using immunotherapy to boost the abscopal effect. Nat Rev Cancer. 2018;18(5):313-322.

55. Sahin U, Türeci Ö. Personalized vaccines for cancer immunotherapy. Science. 2018;359(6382):1355-1360.

56. Hu Z, Ott PA, Wu CJ. Towards personalized, tumour-specific, therapeutic vaccines for cancer. Nat Rev Immunol. 2018;18(3):168-182.

57. Ott PA, Hu-Lieskovan S, Chmielowski B, et al. A phase Ib trial of personalized neoantigen therapy plus anti-PD-1 in patients with advanced melanoma, non-small cell lung cancer, or bladder cancer. Cell. 2020;183(2):347-362.e24.

58. Keskin DB, Anandappa AJ, Sun J, et al. Neoantigen vaccine generates intratumoral T cell responses in phase Ib glioblastoma trial. Nature. 2019;565(7738):234-239.

59. Sahin U, Oehm P, Derhovanessian E, et al. An RNA vaccine drives immunity in checkpoint-inhibitor-treated melanoma. Nature. 2020;585(7823):107-112.

60. Cafri G, Gartner JJ, Zaks T, et al. mRNA vaccine-induced neoantigen-specific T cell immunity in patients with gastrointestinal cancer. J Clin Invest. 2020;130(11):5976-5988.

61. Rojas LA, Sethna Z, Soares KC, et al. Personalized RNA neoantigen vaccines stimulate T cells in pancreatic cancer. Nature. 2023;618(7963):144-150.

62. Fong L, Forde PM, Powderly JD, et al. Safety and clinical activity of adenosine A2a receptor (A2aR) antagonist, CPI-444, in anti-PD1/PDL1 treatment-refractory renal cell (RCC) and non-small cell lung cancer (NSCLC) patients. J Clin Oncol. 2017;35(15_suppl):3004.

63. Luke JJ, Powderly JD, Merchan JR, et al. Immunobiology, preliminary safety, and efficacy of CPI-006, an anti-CD73 antibody with immune modulating activity, in a phase 1 trial in advanced cancers. J Clin Oncol. 2019;37(15_suppl):2505.

64. Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. Lancet Oncol. 2017;18(9):1182-1191.

65. Le DT, Durham JN, Smith KN, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. Science. 2017;357(6349):409-413.

66. Marabelle A, Le DT, Ascierto PA, et al. Efficacy of pembrolizumab in patients with noncolorectal high microsatellite instability/mismatch repair-deficient cancer: results from the phase II KEYNOTE-158 study. J Clin Oncol. 2020;38(1):1-10.

67. Cocco E, Scaltriti M, Drilon A. NTRK fusion-positive cancers and TRK inhibitor therapy. Nat Rev Clin Oncol. 2018;15(12):731-747.

68. Hong DS, DuBois SG, Kummar S, et al. Larotrectinib in patients with TRK fusion-positive solid tumours: a pooled analysis of three phase 1/2 clinical trials. Lancet Oncol. 2020;21(4):531-540.

69. Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. Lancet Oncol. 2020;21(2):271-282.

70. Subbiah V, Wolf J, Konda B, et al. Tumour-agnostic efficacy and safety of selpercatinib in patients with RET fusion-positive solid tumours other than lung or thyroid tumours (LIBRETTO-001): a phase 1/2, open-label, basket trial. Lancet Oncol. 2022;23(10):1261-1273.

71. Drilon A, Oxnard GR, Tan DSW, et al. Efficacy of selpercatinib in RET fusion-positive non-small-cell lung cancer. N Engl J Med. 2020;383(9):813-824.

72. Wirth LJ, Sherman E, Robinson B, et al. Efficacy of selpercatinib in RET-altered thyroid cancers. N Engl J Med. 2020;383(9):825-835.

73. Wolf J, Seto T, Han JY, et al. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. N Engl J Med. 2020;383(10):944-957.

74. Paik PK, Felip E, Veillon R, et al. Tepotinib in non-small-cell lung cancer with MET exon 14 skipping mutations. N Engl J Med. 2020;383(10):931-943.

75. Skoulidis F, Li BT, Dy GK, et al. Sotorasib for lung cancers with KRAS p.G12C mutation. N Engl J Med. 2021;384(25):2371-2381.

76. Jänne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in non-small-cell lung cancer harboring a KRAS(G12C) mutation. N Engl J Med. 2022;387(2):120-131.

77. Yaeger R, Weiss J, Pelster MS, et al. Adagrasib with or without cetuximab in colorectal cancer with mutated KRAS G12C. N Engl J Med. 2023;388(1):44-54.

78. de Bono J, Mateo J, Fizazi K, et al. Olaparib for metastatic castration-resistant prostate cancer. N Engl J Med. 2020;382(22):2091-2102.

79. Golan T, Hammel P, Reni M, et al. Maintenance olaparib for germline BRCA-mutated metastatic pancreatic cancer. N Engl J Med. 2019;381(4):317-327.

80. Moore K, Colombo N, Scambia G, et al. Maintenance olaparib in patients with newly diagnosed advanced ovarian cancer. N Engl J Med. 2018;379(26):2495-2505.

81. Swisher EM, Lin KK, Oza AM, et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. Lancet Oncol. 2017;18(1):75-87.

82. Litton JK, Rugo HS, Ettl J, et al. Talazoparib in patients with advanced breast cancer and a germline BRCA mutation. N Engl J Med. 2018;379(8):753-763.

83. Robson M, Im SA, Senkus E, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. N Engl J Med. 2017;377(6):523-533.

84. Melero I, Berman DM, Aznar MA, Korman AJ, Pérez Gracia JL, Haanen J. Evolving synergistic combinations of targeted immunotherapies to combat cancer. Nat Rev Cancer. 2015;15(8):457-472.

85. Postow MA, Callahan MK, Wolchok JD. Immune checkpoint blockade in cancer therapy. J Clin Oncol. 2015;33(17):1974-1982.

86. Topalian SL, Drake CG, Pardoll DM. Immune checkpoint blockade: a common denominator approach to cancer therapy. Cancer Cell. 2015;27(4):450-461.

87. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. Science. 2018;359(6382):1350-1355.

88. Wei SC, Duffy CR, Allison JP. Fundamental mechanisms of immune checkpoint blockade therapy. Cancer Discov. 2018;8(9):1069-1086.

89. Pitt JM, Vétizou M, Daillère R, et al. Resistance mechanisms to immune-checkpoint blockade in cancer: tumor-intrinsic and -extrinsic factors. Immunity. 2016;44(6):1255-1269.

90. Jenkins RW, Barbie DA, Flaherty KT. Mechanisms of resistance to immune checkpoint inhibitors. Br J Cancer. 2018;118(1):9-16.

91. Schoenfeld AJ, Hellmann MD. Acquired resistance to immune checkpoint inhibitors. Cancer Cell. 2020;37(4):443-455.

92. Zaretsky JM, Garcia-Diaz A, Shin DS, et al. Mutations associated with acquired resistance to PD-1 blockade in melanoma. N Engl J Med. 2016;375(9):819-829.

93. Shin DS, Zaretsky JM, Escuin-Ordinas H, et al. Primary resistance to PD-1 blockade mediated by JAK1/2 mutations. Cancer Discov. 2017;7(2):188-201.

94. Gao J, Shi LZ, Zhao H, et al. Loss of IFN-γ pathway genes in tumor cells as a mechanism of resistance to anti-CTLA-4 therapy. Cell. 2016;167(2):397-404.e9.

95. Sade-Feldman M, Jiao YJ, Chen JH, et al. Resistance to checkpoint blockade therapy through inactivation of antigen presentation. Nat Commun. 2017;8(1):1136.

96. McGranahan N, Rosenthal R, Hiley CT, et al. Allele-specific HLA loss and immune escape in lung cancer evolution. Cell. 2017;171(6):1259-1271.e11.

97. Rooney MS, Shukla SA, Wu CJ, Getz G, Hacohen N. Molecular and genetic properties of tumors associated with local immune cytolytic activity. Cell. 2015;160(1-2):48-61.

98. Spranger S, Bao R, Gajewski TF. Melanoma-intrinsic β-catenin signalling prevents anti-tumour immunity. Nature. 2015;523(7559):231-235.

99. Peng W, Chen JQ, Liu C, et al. Loss of PTEN promotes resistance to T cell-mediated immunotherapy. Cancer Discov. 2016;6(2):202-216.

100. Manguso RT, Pope HW, Zimmer MD, et al. In vivo CRISPR screening identifies Ptpn2 as a cancer immunotherapy target. Nature. 2017;547(7664):413-418.

101. Lawson KA, Sousa CM, Zhang X, et al. Functional genomic landscape of cancer-intrinsic evasion of killing by T cells. Nature. 2020;586(7827):120-126.

102. Vredevoogd DW, Kuilman T, Ligtenberg MA, et al. Augmenting immunotherapy impact by lowering tumor TNF cytotoxicity threshold. Cell. 2019;178(3):585-599.e15.

103. Ishizuka JJ, Manguso RT, Cheruiyot CK, et al. Loss of ADAR1 in tumours overcomes resistance to immune checkpoint blockade. Nature. 2019;565(7737):43-48.

104. Zhang B, Wang Y, Yu M, et al. A pooled analysis of 10 studies assessing the efficacy and safety of PARPis and bevacizumab combination therapy for advanced ovarian cancer patients. J Ovarian Res. 2023;16(1):12.

105. Tewari KS, Sill MW, Penson RT, et al. Bevacizumab for advanced cervical cancer: final overall survival and adverse event analysis of a randomised, controlled, open-label, phase 3 trial (Gynecologic Oncology Group 240). Lancet. 2017;390(10103):1654-1663.

106. Garcia J, Hurwitz HI, Sandler AB, et al. Bevacizumab (Avastin®) in cancer treatment: A review of 15 years of clinical experience and future outlook. Cancer Treat Rev. 2020;86:102017.

107. Ferrara N, Adamis AP. Ten years of anti-vascular endothelial growth factor therapy. Nat Rev Drug Discov. 2016;15(6):385-403.

108. Ellis LM, Hicklin DJ. VEGF-targeted therapy: mechanisms of anti-tumour activity. Nat Rev Cancer. 2008;8(8):579-591.

109. Jain RK. Normalization of tumor vasculature: an emerging concept in antiangiogenic therapy. Science. 2005;307(5706):58-62.

110. Fukumura D, Kloepper J, Amoozgar Z, Duda DG, Jain RK. Enhancing cancer immunotherapy using antiangiogenics: opportunities and challenges. Nat Rev Clin Oncol. 2018;15(5):325-340.

111. Hegde PS, Wallin JJ, Mancao C. Predictive markers of anti-VEGF and emerging role of angiogenesis inhibitors as immunotherapeutics. Semin Cancer Biol. 2018;52(Pt 2):117-124.

112. Lee WS, Yang H, Chon HJ, Kim C. Combination of anti-angiogenic therapy and immune checkpoint blockade normalizes vascular-immune crosstalk to potentiate cancer immunity. Exp Mol Med. 2020;52(9):1475-1485.

113. Schmittnaegel M, Rigamonti N, Kadioglu E, et al. Dual angiopoietin-2 and VEGFA inhibition elicits antitumor immunity that is enhanced by PD-1 checkpoint blockade. Sci Transl Med. 2017;9(385):eaak9670.

114. Allen E, Jabouille A, Rivera LB, et al. Combined antiangiogenic and anti-PD-L1 therapy stimulates tumor immunity through HEV formation. Sci Transl Med. 2017;9(385):eaak9679.

115. Voron T, Colussi O, Marcheteau E, et al. VEGF-A modulates expression of inhibitory checkpoints on CD8+ T cells in tumors. J Exp Med. 2015;212(2):139-148.

116. Terme M, Pernot S, Marcheteau E, et al. VEGFA-VEGFR pathway blockade inhibits tumor-induced regulatory T-cell proliferation in colorectal cancer. Cancer Res. 2013;73(2):539-549.

117. Huang Y, Goel S, Duda DG, Fukumura D, Jain RK. Vascular normalization as an emerging strategy to enhance cancer immunotherapy. Cancer Res. 2013;73(10):2943-2948.

118. Tian L, Goldstein A, Wang H, et al. Mutual regulation of tumour vessel normalization and immunostimulatory reprogramming. Nature. 2017;544(7649):250-254.

119. De Palma M, Biziato D, Petrova TV. Microenvironmental regulation of tumour angiogenesis. Nat Rev Cancer. 2017;17(8):457-474.

120. Viallard C, Larrivée B. Tumor angiogenesis and vascular normalization: alternative therapeutic targets. Angiogenesis. 2017;20(4):409-426.

121. Seeber A, Gunsilius E, Gastl G, Pircher A. Anti-angiogenics: their value in colorectal cancer therapy. Oncol Res Treat. 2018;41(4):188-193.

122. Van Cutsem E, Tabernero J, Lakomy R, et al. Addition of aflibercept to fluorouracil, leucovorin, and irinotecan improves survival in a phase III randomized trial in patients with metastatic colorectal cancer previously treated with an oxaliplatin-based regimen. J Clin Oncol. 2012;30(28):3499-3506.

123. Grothey A, Van Cutsem E, Sobrero A, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. Lancet. 2013;381(9863):303-312.

124. Li J, Qin S, Xu RH, et al. Effect of fruquintinib vs placebo on overall survival in patients with previously treated metastatic colorectal cancer: the FRESCO randomized clinical trial. JAMA. 2018;319(24):2486-2496.

125. Tabernero J, Yoshino T, Cohn AL, et al. Ramucirumab versus placebo in combination with second-line FOLFIRI in patients with metastatic colorectal carcinoma that progressed during or after first-line therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine (RAISE): a randomised, double-blind, multicentre, phase 3 study. Lancet Oncol. 2015;16(5):499-508.

126. Bennouna J, Sastre J, Arnold D, et al. Continuation of bevacizumab after first progression in metastatic colorectal cancer (ML18147): a randomised phase 3 trial. Lancet Oncol. 2013;14(1):29-37.

127. Authors/Task Force members, Galiè N, Humbert M, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). Eur Heart J. 2016;37(1):67-119.