The Effects of Yoga Nidra Practice on EEG: A Systematic Review

Vicky Kachera[a*], Dimple Patel[a], Saurabh Mishra[b], Alka Mishra[c], James Gomes[a,d], Rahul Garg[a,e,f]

[a]*National Resource Centre for Value Education in Engineering, Indian Institute of Technology, Delhi, Hauz Khas, New Delhi, India*
[b]*Department of Career Support, Guidance and Progression, Dev Sanskriti Vishwavidyalaya, Gayatrikunj-Shantikunj, Haridwar, Uttarakhand, India*
[c]*Department of Ayurveda and Holistic Health, Dev Sanskriti Vishwavidyalaya, Gayatrikunj-Shantikunj, Haridwar, Uttarakhand, India*
[c]*Department of Complementary and Alternative Medicine, Dev Sanskriti Vishwavidyalaya, Gayatrikunj-Shantikunj, Haridwar, Uttarakhand, India*
[d]*Kusuma School of Biological Sciences, Indian Institute of Technology Delhi, Hauz Khas, New Delhi, India*
[e]*Department of Computer Science and Engineering, Indian Institute of Technology, Delhi, Hauz Khas, New Delhi, India*
[f]*Amar Nath and Shashi Khosla School of Information Technology, Indian Institute of Technology, New Delhi, India*

*Corresponding author
E-mail address: vickyoctober1998@gmail.com

Abstract

Background: Yoga Nidra practice (YNP) is a systematic method of inducing physical, mental and emotional relaxation. YNP has been claimed to improve sleep quality and reduce anxiety and stress. Research has begun exploring its neural mechanisms using EEG to determine how it induces a sleep-like state or is different from sleep, but the findings remain fragmented. This review investigates the impact of YNP on EEG activity, based on existing quantitative studies.

Methods: Studies examining the effects of YNP on EEG oscillations, published in the English language, were searched in PubMed, Web of Science, and Scopus from inception to May 6, 2025. Study quality was assessed using the Mixed Methods Appraisal Tool (MMAT). Since a meta-analysis could not be done owing to heterogeneity in study designs, Power Spectral Density (PSD) and EEG-based sleep-related findings were synthesised narratively.

Results: Twelve studies with 326 participants met the inclusion criteria, employing varied designs including RCTs, as well as pre-post, within-subject and crossover studies. Only two studies met 100% of the MMAT quality criteria, and four included an active control group. The most consistent finding was increased theta power among experienced practitioners, reported in four studies. Changes in alpha, delta, beta, and gamma bands were inconsistent. A common observation was that participants with prior YNP experience remained awake during practice. Additionally, YNP was associated with improvements in sleep quality.

Conclusion: Yoga Nidra Practice shows promise in improving sleep-related outcomes, promoting deep relaxation. More well-designed studies with larger sample sizes and diverse populations are essential to better understand its impact on EEG and its underlying neural mechanisms.

PROSPERO registration number: CRD42024511747.

Keywords: Yoga nidra, meditation, NSDR, EEG, brain waves, mental health

Highlights:

- Yoga Nidra practice (YNP) has significant effects on EEG activity.
- By facilitating deep relaxation, YNP improves sleep-related outcomes.
- No consistent patterns were observed in the PSD of alpha, beta, delta and gamma bands.
- The findings elicit the need for further studies with larger sample sizes, diverse populations and longitudinal designs.

1. Introduction

The state of Yoga Nidra (YN) is an altered state of consciousness, not yet fully understood by modern science (1). According to Parker et al. (2013), the YN state resembles deep non-REM sleep, while awareness remains intact (1). Although no published scientific studies have verified this phenomenon, pilot reports documented it in accomplished yogis such as Swami Rama and his disciple Swami Veda Bharati, who have demonstrated the ability to consciously enter the YN state (2,3). The Yoga Nidra practice (YNP) is an ancient technique rooted in the Tantras and defined as a systematic method for achieving complete physical, mental, and emotional relaxation, and was revived in modern times by yogis such as Swami Satyananda Saraswati and Swami Rama, who emphasised that the Yoga Nidra state can be attained through sustained practice (3,4).

Variants of YNP include the '61-Points Relaxation Exercise' developed by Swami Rama of the Himalayan yoga tradition (5); Marmanasthanam Kriya, from the Rishiculture Ashtanga Yoga (Gitananda Tradition) (6); and Integrative Restoration (iRest), a modern adaptation developed by Richard Miller based on the ancient YNP (7). Dr. Andrew Huberman considers Yoga Nidra a form of Non-Sleep Deep Rest (NSDR), a term he coined to describe mindfulness practices that induce deep relaxation without leading to sleep (8,9).

YNP has been reported to enhance mental health, particularly by alleviating symptoms of stress, anxiety, and depression (10). Several clinical trials on YNP have shown improvements in sleep quality and reductions in symptoms of insomnia (11−14). Additionally, YNP has been shown to lower blood pressure in hypertensive patients (15) and improve the psychological well-being of women experiencing menstrual disorders (16).

Electroencephalography (EEG) is a non-invasive modality that measures brain function and can be used to characterise the electrical activity patterns associated with both YNP and the YN state. According to Parker et al. (2013), YNP can be described electrophysiologically across several levels (1). In the initial stages of practice, individuals typically experience deep relaxation marked by increased alpha wave activity, which may gradually shift toward theta waves with deeper engagement (1). Swami Satyananda Saraswati also emphasised the predominance of alpha waves during YNP, associating it with profound mental, emotional, and muscular relaxation (4).

Furthermore, both Parker et al. (2013) and Swami Satyananda propose that in the deeper YN state, practitioners may consciously enter a state characterised by delta wave activity, typically associated with deep, non-REM sleep, while remaining fully aware of the environment (1,4). To the best of our knowledge, no published studies have demonstrated the existence of the YN state under controlled conditions. Building on these theoretical perspectives and anecdotal reports, recent studies have employed EEG to objectively examine the brain wave patterns during YNP. However, drawing definitive conclusions about specific brainwave patterns remains challenging owing to methodological heterogeneity resulting from variability in YN protocols, practitioner experience, and study designs. Additionally, some studies have recorded EEG activity during the practice, while others have assessed post-practice effects.

Therefore, the objective of this systematic review is to identify and characterise studies examining brain wave activity associated with YNP.

2. Methods

This systematic review adhered to the updated 2020 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (17), and the protocol was registered in PROSPERO (CRD42024511747).

2.1 Information Sources and Search Strategy

A PROSPERO search conducted on January 7, 2024, revealed no registered systematic reviews focusing specifically on YN and its effects on brain waves using EEG. Three databases namely PubMed, Web of Science, and Scopus were searched for articles published from the inception of each database until May 6, 2025, using broad search terms such as 'Yoga Nidra,' 'Yoganidra,' 'Yogic Sleep,' 'Psychic Sleep,' 'Hypnagogic

Sleep,' 'NSDR,' 'iRest,' 'Electroencephalogram,' 'EEG,' and 'Brain Waves.' These terms were intentionally chosen to capture diverse terminology associated with YN and related neurophysiological states. Terms such as 'Yoganidra,' 'Yogic Sleep,' 'Psychic Sleep,' and 'Hypnagogic Sleep' addressed conceptual variations, while 'EEG,' 'Electroencephalogram,' and 'Brain Waves' ensured inclusion of studies involving neuroelectrical measurements. The final searches for PubMed, Web of Science and Scopus were conducted on May 6, 2025. No filters or restrictions were applied to any database. The detailed search query for all databases is provided in Supplementary Table A.

2.2 Study Selection Process:

The detailed eligibility criteria are summarised in Table 1. Findings from each database were imported into RAYYAN for deduplication and screening (18). Two reviewers (VK and DP) independently screened the articles by title and abstract according to the predetermined inclusion and exclusion criteria. Only accessible full-text articles were downloaded and thoroughly reviewed to confirm their eligibility. Additionally, the reference lists of the selected articles were examined to identify any other relevant studies. Before the final inclusion of articles, consensus was reached, and any disagreements were resolved through discussion. Study authors were contacted by email to clarify doubts, confirm details, or request additional information as needed. Referencing was managed using Zotero 6.0.36, an open-access reference management tool (19).

Table 1: Inclusion and Exclusion Criteria

Key Details	Inclusion	Exclusion
Relevant Publications	Studies assessing the effect of YN on brain waves as measured by EEG	Studies assessing the effect of YN on variables other than EEG
Publication Date	From the inception of the databases until May 2025	After May 2025
Literature	Published literature	Opinion pieces, discussion papers, reviews, unpublished literature, textbook chapters, grey literature
Study Framework	All types of quantitative studies	Qualitative studies, reviews, case series/reports
Document Availability	Full text	Abstracts only

2.3 Data Collection Process and Data Items:

The data collection process involved creating a table, with the reviewer VK extracting the data including the following aspects: study information (title, journal, author(s), year, country, study design, blinding, randomisation details); sample characteristics (inclusion-exclusion criteria, sample size (n), gender, age range); control and intervention details (control/comparator group details, experimental intervention details, duration and frequency); EEG data and methodology (EEG channels, locations analysed, impedance, sampling rate, feature extraction, EEG methodology); and key findings (EEG). This data was verified by another reviewer (DP), with any disagreements resolved by a third reviewer (RG). Data extraction was performed manually, without the use of automation tools.

2.4 Quality Assessment:

The MMAT tool can appraise heterogeneous empirical studies across five categories: Qualitative, Quantitative Randomised Controlled Trials (RCTs), Quantitative Non-Randomised, Quantitative Descriptive, and Mixed Methods studies (20). It has been implemented in various mixed-method systematic reviews (21–23). Two reviewers (VK and DP) independently evaluated the quality of all included studies by using the 2018 version of MMAT (20,24). Each question in the selected category was rated as 'Yes,' 'No,' or 'Can't tell' (when sufficient information was not provided to answer 'Yes' or 'No') (20). Any discrepancies in ratings were resolved through discussion, and a third independent reviewer (RG) was involved whenever needed. For 'Can't tell' ratings, the corresponding authors of the studies were contacted by email for more information or clarification. The results are presented as follows: a score of 100% if all five quality criteria are met, 80% if four, 60% if three, 40% if two, and 20% if one (25). No automation or AI-based tools were used to assess study quality.

2.5 Statistical Synthesis:

The heterogeneity in the YNP type, duration, results, population, and study design did not permit a meta-analysis. Instead, the results were reported in tabular form and summarised narratively.

3. Results

3.1 Search Results:

A total of 36 records were identified through our search strategy, including 10 from PubMed, 7 from Web of Science, 16 from Scopus, and 3 from reference searches. 15 duplicates were removed using RAYYAN, leaving 21 articles. After screening titles and abstracts in RAYYAN, 14 articles remained. One article's (26) full text was unavailable, and another was published in a journal not indexed in global databases, leaving 12 articles for quality appraisal. See Figure 1 for the PRISMA 2020 flow diagram.

3.2 Study Characteristics:

Tables 2a and 2b summarise the characteristics of the included studies. The review included 12 papers that met the eligibility criteria: Three were pre-post interventions (14,27,28), Three were RCTs (29–31), One was a within-subject randomised crossover study (32), Three were within-subject studies (33–35), and two were pre-post with a control group (36,37). Most studies (n = 8) were published within the last 10 years.

3.3 Participant Characteristics:

Tables 2a and 2b summarise participant characteristics across studies. A total of 326 participants were included in the 12 studies. Three studies had 50 to 60 participants, four had 20 to 49, and five had fewer than 20. Three studies involved male participants only, one involved female participants only, while three included both male and female participants. Two studies had an equal number of male and female participants, another two did not report gender information, while one mentioned both genders were involved but did not provide specific numbers. The age of participants ranged from 18 to 90 years. Two studies included participants with yoga experience ranging from 5 to 26 years (32,33), while one focused on long-term YN practitioners (35). Two studies involved healthy novices (14,27), one included migraine patients (31), and another involved novice insomnia patients (30). The remaining five studies did not report participants' yoga experience.

Figure 1: PRISMA flow diagram (17)

Table 2a: Characteristics of Studies Employing Power Spectral Density (PSD) Analysis of EEG During Yoga Nidra Practice

S.No.	Study (Year)	Country	Study Design	Population Characteristics and Sample Size (N)	Sample Characteristics [Gender (M/F), Age Range/Mean (Years)]
1.	Datta et al., 2022 (14)	India	Pre-post intervention	Healthy novices N = 30	M/F (15/15); 35.53 ± 8.69 years
2.	Kjaer et al., 2002 (32)	UK	Within-subject, randomised crossover	Meditation teachers with 7 to 26 years of experience. N = 8	Males; 31–50 years
3.	Lou et al., 1999 (33)	Not reported	Within-subjects	Yoga practitioners with over 5 years of experience in Kriya Yoga, YNP, and other yoga techniques. N = 9	M/F (6/3); 23–41 years
4.	Sharpe et al., 2023 (30)	USA	RCT	Novices with insomnia. N = 18 (YNP = 9, CG = 9)	M/F, YNP = 3/6 CG = 1/8; YN = 33 ± 7 years CG = 30 ± 6 years
5.	Zaccaro et al., 2021 (35)	France	Within-subjects	Healthy practitioners with 10 to 12 years of formal YNP. N = 6	M/F (2/4); 31–74 years

RCT: Randomised controlled trial, CG: Control group

Table 2b: EEG Outcomes: Following YNP or Interventions Combined with YNP

S.No.	Study (Year)	Country	Study Design	Population Characteristics and Sample Size (N)	Sample Characteristics [Gender (M/F), Age Range/Mean (Years)]
1.	Datta, Yadav, et al., 2022 (29)	India	RCT	Athletes N = 58 (YNP = 28, CG = 29)	Males; 22.45 ± 4.53 years, age not reported separately for YNP/CG
2.	Shashikiran et al., 2022 (31)	India	RCT	Migraine patients. N = 60 (YNP = 30, CG = 30)	M/F, 18–25 years; gender and age figures not reported separately for YNP/CG
3.	Datta et al., 2023 (27)	India	Pre-post interventional cohort	Healthy novice N= 41	Males; 25.146 ± 2.275 years
4.	Rodin et al., 2017 (34)	USA	Within-subjects	Interested individuals in mindfulness practices. N = 10	M/F (5/5); 26-90 years/41.2 years
Studies that included Yoga Nidra in combination with other interventions as the independent variable					
1.	Bharadwaj et al., 2013 (36)	India	Pre-Post with control group	Working women. N = 50 (YNP = 25, CG = 25)	Females; 25-39 years, age not reported separately for YNP/CG
2.	Anuradha R et al., 2022 (37)	India	Pre-Post with control group	Physically fit students. N = 40 (YNP = 20, CG = 20)	Gender and age figures not reported
3.	Kumar & Joshi, 2009 (28)	India	Pre-Post	College students. N = 40	Gender information not reported; 18-25 years

3.4 Results of Quality Assessment:

Table 3 summarises the quality assessment results. Of the 12 studies, two met 100% of the criteria (14,30), six met 80% (27,29,31,33,34,36), one met 60% (35), two met 40% (32,37), and one met 20% (28). None of the studies were excluded due to low methodological quality, as this is generally discouraged (20).

3.5 Intervention Characteristics:

A detailed overview of EEG acquisition parameters, including electrode locations, sampling rates, and feature extraction methods, is presented in Table 4, while the characteristics of the interventions are provided in Tables 5 and 6. Two studies used Swami Satyananda Saraswati's YNP developed for insomnia (14,30), while the other 10 did not specify the tradition. The session durations ranged from 20 minutes (27) to 2 hours (35), with the most common duration being 27-30 minutes (14,28–30,36). Two studies did not report the duration of YN sessions (31,37). The duration of intervention was largely variable among the studies. Four studies evaluated the effect of a single session of YNP (32–34,30); while others ranged from 12 sessions over 6 days (35) to daily sessions for 45 days (36). One study did not specify the exact intervention period, only stating it was between 3 and 6 months (37). All studies focused on YN as an independent variable except for three (28,36,37), which combined YNP with other yogic practices such as Nadi Shodhan Pranayama, Pranakarshan Pranayama, Nine-Centre Meditation, and Pragya Yoga.

Table 3: Assessment of Study Quality Following the Mixed Methods Appraisal Tool (MMAT)

S.No.	Study (Year)	1. Is randomisation appropriately performed?	2. Are the groups comparable at baseline?	3. Are there complete outcome data?	4. Are outcome assessors blinded to the intervention provided?	5. Did the participants adhere to the assigned intervention?	Quality (%)
1. RANDOMISED CONTROLLED TRIALS							
1.	Datta, Yadav, et al., 2022	Yes	Yes	No	Yes	Yes	80
2.	Kjaer et al., 2002	Can't tell	Yes	No	Can't tell	Yes	40
3.	Sharpe et al., 2023	Yes	Yes	Yes	Yes	Yes	100
4.	Shashikiran et al., 2022	Yes	Yes	Yes	Can't tell	Yes	80

S.No.	Study (Year)	1. Are the participants representative of the target population?	2. Are measurements appropriate regarding both the outcome and intervention (or exposure)?	3. Are there complete outcome data?	4. Are the confounders accounted for in the design and analysis?	5. During the study period, is the intervention administered (or exposure occurred) as intended?	Quality (%)
2. NON-RANDOMISED STUDIES							
1.	Anuradha R et al., 2022	Yes	Yes	Can't tell	Can't tell	Can't tell	40
2.	Bharadwaj et al., 2013	Yes	Yes	Yes	Can't tell	Yes	80
3.	Datta et al., 2022	Yes	Yes	Yes	Yes	Yes	100
4.	Datta et al., 2023	Yes	Yes	Yes	No	Yes	80
5.	Kumar & Joshi, 2009	Yes	Can't tell	Can't tell	Can't tell	Can't tell	20
6.	Lou et al., 1999	Yes	Yes	Yes	Can't tell	Yes	80
7.	Rodin et al., 2017	Yes	Yes	Yes	No	Yes	80
8.	Zaccaro et al., 2021	Yes	Yes	No	No	Yes	60

Table 4: Technical Specifications of EEG Recording and Analysis in Yoga Nidra Practice Studies

S.No.	Study (Year)	EEG Channels	Locations Analysed	Impedance	Sampling Rate	Feature Extraction
1.	Datta et al., 2022 (14)	19	C3, C4, Cz, F3, F4, F7, F8, Fz, Fp1, Fp2, P3, P4, Pz, T3, T4, T5, T6, and O1, O2.	≤ 5 kΩ	500 Hz	PSD
2 A.	Datta et al., 2023 (27)	3	F3, O1, and C3.	NR	256 Hz	PSD
2 B.	Datta et al., 2023 (27)	NR	Electrodes were positioned at O1, O2, F3, C3, C4, and F4, along with EOG and EMG sites.	< 5 kΩ	NR	PSG analysis
3.	Datta, Yadav, et al., 2022 (29)	NR	O1-M2/O2-M1, C3-M2/C4-M1, F3-M2/F4-M1, and two EOG and EMG.	NR	NR	NR
4.	Kjaer et al., 2002 (32)	16	NR	NR	NR	PSD
5.	Lou et al., 1999 (33)	16	NR	NR	NR	PSD
6.	Rodin et al., 2017 (34)	128	As per standard 10-20 system with EOG	≤ 50 kΩ	600 Hz	FFT and ICA
7.	Sharpe et al., 2023 (30)	NR	F3, O1, C3, EOG and chin locations.	NR	NR	PSD, PSG sleep scoring
8.	Shashikiran et al., 2022 (31)	NR	NR	NR	NR	NR
9.	Zaccaro et al., 2021 (35)	128	NR	< 50 kΩ	500 Hz	PSD
10.	Bharadwaj et al., 2013 (36)	NR	NR	NR	NR	NR
11.	Anuradha R et al., 2022 (37)	8	O1, O2, F3, F4, P3, P4, and C3, C4.	NR	256 Hz	DWT
12.	Kumar & Joshi, 2009 (28)	NR	NR	NR	NR	NR

O: Occipital, F: Frontal, T: Temporal, C: Central, DWT: Discrete wavelet transform, FFT: Fast fourier transformation, ICA: Independent component analysis, PSG: Polysomnography, NR: Not reported

3.6 EEG Power Analysis During Yoga Nidra Practice

Table 5 provides a detailed overview of power spectral density (PSD) changes observed during YNP across studies. All studies reported PSD analysis during YNP except the study by Rodin et al. (2017), which reported "near-DC activity that extended beyond one hour, as well as rhythmic wave durations ranging from about 10 to 35 min" during a single 65-minute YNP session" primarily in the frontal midline and inferior hemispheric regions, suggesting a deeply altered neural state (34). The following subsections provide a breakdown of results by frequency band.

3.6.1 Delta Band

Delta activity showed varied changes across studies. Datta et al. (2022) reported an increase in delta power in the central region and a decrease in the prefrontal region during YNP compared to the pre-YNP baseline (14). In another study, delta power decreased by 32% in the control group and by 17% in the YNP group; however, these changes were not statistically significant (30). Zaccaro et al. (2020) found no significant change in delta power during the YNP session (35).

3.6.2 Theta Band

Theta power generally increased during YNP. Kjaer et al. (2002) observed a 9% increase in theta power during YNP compared to the control condition (32), while Lou et al. (1999) reported an 11% increase during YNP compared to a pre-YNP resting state (33). Zaccaro et al. (2021) found that theta power increased during the mental visualisation stage of YNP but gradually declined thereafter (35). Sharpe et al. (2023) noted that theta power decreased less in the YNP group than in the control group during a post-YNP nap session; however, this difference was not statistically significant (30).

3.6.3 Alpha Band

Changes in alpha power during YNP varied across studies, with differing patterns observed depending on participant characteristics and session conditions. Kjaer et al. (2002) reported an 8% decrease in alpha power compared to the control condition (32), and Datta et al. (2022) similarly found reduced alpha power in novices, though PSG analysis confirmed participants remained awake (14). In contrast, Lou et al. (1999) reported only a 2% non-significant decrease from resting baseline, suggesting wakefulness during YNP (33). Zaccaro et al. (2021) observed an initial increase in alpha power that gradually declined over time (35). Sharpe et al. (2023) found a slight increase at O1 in the YNP group and a decrease in the control group; however, PSG analysis showed 89% of participants fell asleep during the session (30). Additionally, post-YNP nap data showed a smaller alpha power decrease in the YNP group than in controls, though the difference was not statistically significant (30).

3.6.4 Beta and Gamma Bands

Changes in beta and gamma power were reported only by Zaccaro et al. (2021) (35). Beta power initially increased during YNP and gradually declined as the session progressed. Gamma power showed an increase during the final stages of YNP (35).

3.7 EEG-Based Sleep Outcomes

As summarised in Table 6, two studies assessed the impact of YNP on objective sleep parameters using overnight EEG recordings. Datta et al. (2023) reported significant improvements in wake-after-sleep onset, sleep efficiency, and increased delta percentage during slow-wave sleep after four weeks of daily YN practice (27). Similarly, Datta, Yadav et al. (2022) found significant improvements in sleep onset latency and time in bed following YNP, with comparable improvements also observed in the progressive muscle relaxation control group (29). These findings suggest that YNP may enhance sleep continuity and quality, as evidenced by changes in electrophysiological markers of sleep.

3.8 EEG Power Analysis Following Yoga Nidra Practice or Combined Interventions

As detailed in Table 6, several studies examined power spectral density (PSD) changes following YNP, either as a standalone intervention or in combination with other yogic techniques. Datta et al. (2023) reported a significant increase in delta and theta power at the occipital site (O1) during a cognitive task after two weeks of YNP (27). Shashikiran et al. (2022) observed a significant increase in the alpha power spectrum in migraine patients following six weeks of YNP, compared to a control group receiving supine rest, with no significant changes in other frequency power bands (31). Other studies reported changes in the power of frequency bands following YNP combined with additional yogic practices. Bharadwaj et al. (2013) found a significant increase in "alpha activity" after a 45-day intervention incorporating Pragya Yoga, Nadi Shodhan Pranayama, and YNP among working women (36). Kumar and Joshi (2009) also reported increased "alpha activity" after 40 days of combined YNP and Pranakarshan Pranayama practice in college students (28). In contrast, Anuradha et al. (2022) observed reductions in both "alpha and beta frequencies" following six months of YNP combined with Nine-Centre Meditation and Nadi Shodhan Pranayama among students (37). These findings suggest that YNP, especially when practised alongside other techniques, may influence alpha and beta band powers, although the direction and extent of change appear to depend on the specific intervention protocol and participant characteristics. Overall, due to poor reporting, low methodological quality, and inconsistencies in the results related to alpha power, no definitive conclusions can be drawn.

Table 5: Power Spectral Density (PSD) Changes in EEG During Yoga Nidra Practice

S.No.	Study (Year)	Experimental Protocol	Control Group	Baseline Condition Description	Comparison Condition	Findings Across Frequency Bands					Remarks
						Δ Delta	Δ Theta	Δ Alpha	Δ Beta	Δ Gamma	
1.	Datta et al., 2022 (14)	SSS's YNP for insomnia was practised for 2 weeks with ~27-min audio; EEG was recorded during YNP after 2 weeks.	NA	Four cycles of 1-min eyes closed and 1 min eyes open (8 min total).	Baseline	Central ↑ Prefrontal ↓	No change in the central region.	Occipital and parietal alpha ↓	NR	NR	PSG data confirmed the awake state of participants.
2.	Kjaer et al., 2002 (32)	Single 72-minute YNP session. EEG was recorded during YNP for both groups.	Yes	NA	Listened to speech (same voice as YNP), eyes closed, 72 min.	NR	↑ 9% (NS)	↓ 8% (NS)	NR	NR	-
3.	Lou et al., 1999 (33)	Two-hour Tantric Kriya Yoga (preparation), followed by a single 45-minute YNP session. EEG was recorded during the entire YNP session.	NA	Supine position, no sensory input, eyes covered, pre-YNP.	Baseline	NR	Increased	↓ 2% (NS)	NR	NR	Alpha was similar to the resting state, indicating wakefulness.

No.	Study	Intervention	Blinding	Setup	Comparison						Other findings
4.	Sharpe et al., 2023 (30)	SSS's YNP for insomnia was used. Single 30-minute YNP session followed by 60-minute quiet rest (nap) at V2. EEG was recorded throughout.	Yes	Supine position, no sensory input, 90-min quiet rest at visit 1(baseline). EEG was recorded during visit 1.	Baseline and 90-minute quiet rest at V2 (CG).	O1: ↓ 32% in the CG and ↓ 17% in the YNP group between V1 and V2 (NS).	During the nap, ↓ less in the YNP group compared to the CG (NS).	Slight ↑ at O1 in the YNP group and slight ↓ in the CG from V1 to V2 (NS within & between). During the 60-minute nap after YNP, ↓ less in the YNP group than in CG (NS).	NR	NR	89% of the intervention group slept, compared to 78% of controls.
5.	Zaccaro et al., 2021 (35)	Twelve group YNP sessions over 6 days; each 2 hours. EEG was recorded during each YNP session.	NA	Supine position, spontaneous breathing, 10 min, pre-YNP.	Baseline	No change	Widespread ↑ during mental visualisation, ↓ later.	Initial widespread ↑, then ↓.	Initial widespread ↑, then ↓.	Widespread gamma ↑ in late stages;	No K-complexes or sleep spindles observed.
6.	Rodin et al., 2017[1] (34)	Single 65-min YNP session performed via audio seated with eyes closed; EEG recorded throughout.	NA	NR	NR	Rhythmic waves below the infraslow range (about 8-30 min) and a near-DC component, primarily in the frontal midline and inferior hemispheric regions.					

NA: Not available, NR: Not reported, SSS: Swami Satyananda Saraswati, ↑: Increase, ↓: Decrease, CG: Control group, V1: visit 1, V2: visit 2

---

[1] **Rodin et al., 2017 recorded EEG during YNP, but did not report PSD analysis.**

Table 6: EEG Outcomes: Following YNP or Interventions Combined with YNP

S.No.	Study (Year)	Experimental Protocol	Control Group	Comparison Condition	Key Findings (EEG)
1.	Datta et al., 2023 (27)	YNP (20 min/day) for 4 weeks; overnight PSG recorded at baseline and after 4 weeks. EEG was recorded during daytime cognitive tasks.	NA	Baseline	Significant improvements in wake-after-sleep onset, sleep efficiency, and delta (%) in slow-wave sleep.
				Control with Eyes Open (CEO) and Control with Eyes Closed (CEC)	O1: ↑ theta during VOLT, ↑ delta/beta during BART (after 2 weeks of YN) C3: ↑ theta during VOLT and MPT (after 1 week of YN) F3: ↑ delta/alpha, ↑ delta/theta during ERT (after 1 week of YN)
2.	Datta, Yadav, et al., 2022 (29)	YNP (~27.2 min/day) for 4 weeks, with one pre-sleep hygiene session; overnight EEG recorded at baseline and after 2 weeks.	PMR practised before sleep (10–15 min) for 4 weeks, preceded by a sleep hygiene session.	Baseline	Significant improvements in sleep onset latency and time in bed in both groups.
3.	Shashikiran et al., 2022 (31)	YNP for 6 weeks. EEG was recorded at baseline and after 6 weeks.	Received supine rest for 6 weeks.	Control group	A significant increase in "Alpha frequency" in the YNP group; no significant changes in the other EEG waveforms.
4.	Bharadwaj et al., 2013 (36)	Pragya Yoga, NSP, and YNP (30 min) practised together (60 min/day) for 45 days; Alpha EEG biofeedback was recorded at baseline and after 45 days.	NR	Control group	A significant increase in "alpha EEG" was observed in the intervention group.

5.	Anuradha R et al., 2022 (37)	YNP, NSP, and NCM were first practised as preparation, then individually (the duration was not explicitly reported). EEG was recorded for 40 min at baseline and after 6 months.	NR	Control group	"Alpha and beta frequencies" decreased in the YNP group.
6.	Kumar & Joshi, 2009 (28)	YNP (30 min) and PP (15 min) practised daily for 40 days; Alpha EEG biofeedback was recorded at baseline and after 40 days.	NA	Baseline	A significant increase in "alpha EEG".

O1: Occipital region, C3: Central region VOLT: Visual Object Learning Task, MPT: Motor Praxis Task, BART: Balloon Analog Risk Task, ERT: Emotion Recognition Task, ↑: Increase, ↓: Decrease, PMR: Progressive Muscle Relaxation, NSP: Nadi Shodhan Pranayama, PP: Pranakarshan Pranayama, NCM: Nine-Centre Meditation

4. Discussion

4.1 Interpretation of Results

This systematic review summarises findings from 12 studies involving 326 participants pertaining to the effects of Yoga Nidra practice (YNP) on EEG activity. The included studies span various populations, including long-term YN practitioners, healthy novices, migraine patients, and insomnia patients.

A consistent finding across studies is increased theta power during YNP, particularly among experienced practitioners (30,32,33,35). According to Swami Satyananda, increased theta activity has been linked to the release of suppressed emotions, fears and desires (4). Prior research suggests a positive correlation between theta power and meditation experience (38). Increased theta band power is a key characteristic of meditation practices and is commonly associated with enhanced cognitive functions, including improved episodic memory retrieval, learning, and spatial navigation (38,39). One study further associated increased theta activity during YNP with endogenous dopamine release (32). Dopamine plays a key role in regulating mood and the sleep–wake cycle and is also released in response to natural rewards such as food, drink, and sex (40,41).

Findings on alpha power during YNP were inconsistent: a significant decrease was reported by Datta et al. (2022) (14); non-significant decreases were shown by Kjaer et al. (2002) and Lou et al. (1999) (32,33); Sharpe et al. (2023) found a non-significant increase (30); and Zaccaro et al. (2021) observed an initial increase followed by a decrease (35). These variations may reflect differences in participant experience, session duration, or EEG protocols. Nevertheless, a recurring pattern was that experienced practitioners remained awake during YNP, as confirmed by PSG analysis (33,35). One study also found that inexperienced participants who received two weeks of YNP training remained awake, possibly as a result of the prior training (14), while another reported sleep onset in novice insomnia patients with no prior exposure (30). Overall, the non-significant changes in alpha power (increases or decreases) suggest that participants remained awake, as alpha power typically diminishes by more than 50% during stage 1 sleep (42).

Beyond theta and alpha power, findings on other frequency bands during YNP are limited and inconsistent. One study reported dynamic beta fluctuations (initial increase followed by decrease) and late-stage gamma elevations, while delta activity varied across regions, potentially reflecting heterogeneous protocols and participant characteristics. Some studies suggest that several weeks of YNP, alone or with other yogic practices, may influence alpha and beta activity post-intervention. However, inconsistent findings and poor methodological quality limit definitive conclusions.

In addition to brainwave activity during the practice itself, a few studies have explored sleep-related EEG outcomes following YNP. These include improvements such as increased sleep efficiency, reduced wake-after-sleep onset, higher delta power during slow-wave sleep, improved sleep onset latency, and increased total time in bed (27,29). Although limited, these findings suggest potential benefits of YNP for sleep, warranting further research with larger samples and standardised methods.

4.2 Evidence Limitations and Future Recommendations

The evidence included in this review has several limitations to be taken into consideration. Many of the studies had small sample sizes, with five studies involving fewer than 20 participants, which limits the generalisability of their findings. Only two studies met 100% MMAT quality assessment criteria. Participant characteristics varied widely in age, health status, and yoga experience, potentially introducing confounding factors. Methodological inconsistencies, such as differences in study design, YNP protocols, and EEG parameters, make it challenging to compare results directly or perform a meta-analysis. Incomplete reporting of key EEG technical details and poor documentation of randomisation, blinding, and control conditions also limit the reliability and reproducibility of results.

Future research should prioritise well-powered, long-term RCTs to differentiate state and trait effects of YN on brain activity and elucidate its underlying mechanisms. Deep-design studies with repeated EEG recordings in the same individuals can help capture dynamic, state-related neural changes during YNP. Standardised YN protocols, use of active control conditions, and diverse participant samples will enhance comparability and generalisability. Clear reporting of EEG specifications (e.g., sampling rates, feature extraction), pre-registration, and open data sharing are essential for methodological transparency. Additionally, explicitly stating the YNP tradition, its lineage, and the specific version used would help contextualise results, reduce variability across studies, and facilitate reproducibility.

5. Conclusion

This systematic review highlights the potential of YN to influence brain activity, provide insights into neural mechanisms, and improve sleep-related outcomes. The observed increase in theta during YN, along with improved sleep parameters including enhanced sleep efficiency, suggests an overarching trend towards deep relaxation and overall well-being. However, further studies are needed to understand these effects better.

Credit Authorship Contribution Statement:

Rahul Garg: Conceptualisation, Supervision, Writing – review and editing. Vicky Kachera: Data curation, Formal analysis, Writing – original draft, Writing – review and editing. Dimple Patel: Data curation, Formal analysis, Writing – review and editing. Alka Mishra: Writing – Final review and editing. Saurabh Mishra: Writing – Final review and editing. James Gomes: Writing – Final review and editing.

Declaration of Interests:

The authors declare that there are no competing financial or personal relationships that could inappropriately influence the work reported in this paper.

Funding Sources:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References:

1. Parker S, Bharati SV, Fernandez M. Defining Yoga-Nidra: Traditional Accounts, Physiological Research, and Future Directions. International Journal of Yoga Therapy. 2013 Jan 1;23(1):11–6.
2. Parker S. Chapter 11 - Training attention for conscious non-REM sleep: The yogic practice of yoga-nidrā and its implications for neuroscience research. In: Srinivasan N, editor. Progress in Brain Research [Internet]. Elsevier; 2019 [cited 2025 Mar 24]. p. 255–72. (Meditation; vol. 244). Available from: https://www.sciencedirect.com/science/article/pii/S0079612318301559
3. Parker S. Yoga Nidrā: An Opportunity for Collaboration to Extend the Science of Sleep States. Sleep Vigilance. 2017 Dec;1(2):57–63.
4. Saraswati S. Yoga Nidra. 6th ed. Munger, Bihar, India: Yoga Publications Trust, Ganga Darshan; 2009.
5. Dvivedi J, Dvivedi S, Mahajan KK, Mittal S, Singhal A. Effect of '61-points relaxation technique' on stress parameters in premenstrual syndrome. Indian J Physiol Pharmacol. 2008;52(1):69–76.
6. Thiruvalluvan A, Sekizhar V, Ramanathan M, Bhavanani AB, Chakravathy D, Reddy JRC. Effect of Pranayama Techniques with Marmanasthanam Kriya as Yogic Relaxation on Biopsychosocial Parameters Prior to Endodontic Therapy: A Cross Sectional Study Design. Int J Yoga. 2021;14(2):146–51.
7. iRest Institute [Internet]. About iRest. Available from: https://shop.irest.org/pages/about-irest
8. Non-Sleep Deep Rest (NSDR) [Internet]. [cited 2025 Jun 9]. Available from: https://www.hubermanlab.com/nsdr
9. Boukhris O, Suppiah H, Halson S, Russell S, Clarke A, Geneau MC, et al. The acute effects of nonsleep deep rest on perceptual responses, physical, and cognitive performance in physically active participants. Applied Psych Health & Well. 2024 Nov;16(4):1967–87.
10. Malviya S, Saoji AA, Pravalika B. Yoga nidra for mental health: A systematic review of current evidence. Journal of Spirituality in Mental Health. 2023 Dec 11;1–27.
11. Datta K, Tripathi M, Verma M, Masiwal D, Mallick HN. *Yoga nidra* practice shows improvement in sleep in patients with chronic insomnia: A randomized controlled trial. NMJI. 2021 Oct 22;34:143–50.
12. Gulia KK, Sreedharan SE. Yogic Sleep and Walking Protocol Induced Improvement in Sleep and Wellbeing in Post-menopausal Subject: A Longitudinal Case Study During COVID Lockdown. Sleep Vigilance. 2022 Jun;6(1):229–33.
13. Sharpe E, Tibbitts D, Wolfe B, Senders A, Bradley R. Qualitative Impressions of a Yoga Nidra Practice for Insomnia: An Exploratory Mixed-Methods Design. The Journal of Alternative and Complementary Medicine. 2021 Oct 1;27(10):884–92.
14. Datta K, Mallick HN, Tripathi M, Ahuja N, Deepak KK. Electrophysiological Evidence of Local Sleep During Yoga Nidra Practice. Front Neurol [Internet]. 2022 Jul 12 [cited 2024 Aug 20];13. Available from: https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2022.910794/full
15. Ahuja N, Bhardwaj P, Pathania M, Sethi D, Kumar A, Parchani A, et al. Yoga Nidra for hypertension: A systematic review and meta-analysis. Journal of Ayurveda and Integrative Medicine. 2024 Mar;15(2):100882.
16. Kim SD. Psychological effects of yoga nidra in women with menstrual disorders: A systematic review of randomized controlled trials. Complementary Therapies in Clinical Practice. 2017 Aug 1;28:4–8.
17. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021 Mar 29;372:n71.
18. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. Syst Rev. 2016 Dec 5;5:210.
19. Ahmed KKM, Al Dhubaib BE. Zotero: A bibliographic assistant to researcher. J Pharmacol Pharmacother. 2011;2(4):303–5.

20. Hong QN, Pluye P, Fàbregues S, Bartlett G, Boardman F, Cargo M, et al. Mixed Methods Appraisal Tool (MMAT), version 2018. Canadian Intellectual Property Office, Industry Canada.;

21. Peek STM, Wouters EJM, van Hoof J, Luijkx KG, Boeije HR, Vrijhoef HJM. Factors influencing acceptance of technology for aging in place: a systematic review. Int J Med Inform. 2014 Apr;83(4):235–48.

22. Puts MTE, Tu HA, Tourangeau A, Howell D, Fitch M, Springall E, et al. Factors influencing adherence to cancer treatment in older adults with cancer: a systematic review. Ann Oncol. 2014 Mar;25(3):564–77.

23. Humphries S, Stafinski T, Mumtaz Z, Menon D. Barriers and facilitators to evidence-use in program management: a systematic review of the literature. BMC Health Serv Res. 2014 Apr 14;14:171.

24. Hong QN, Fàbregues S, Bartlett G, Boardman F, Cargo M, Dagenais P, et al. The Mixed Methods Appraisal Tool (MMAT) version 2018 for information professionals and researchers. Education for Information. 2018 Jan 1;34(4):285–91.

25. Hong QN. mixedmethodsappraisaltoolpublic [licensed for non-commercial use only] / FAQ [Internet]. [cited 2024 Sep 13]. Available from: http://mixedmethodsappraisaltoolpublic.pbworks.com/w/page/71030694/FAQ

26. Ray GC. DFT ANALYSIS OF EEG PICKED UP DURING SHABASANA AND YOGANIDRA. Journal of the Institution of Engineers India Interdisciplinary and General Engineering. 1985 Feb;65:56–9.

27. Datta K, Bhutambare A, V L M, Narawa Y, Srinath R, Kanitkar M. Improved sleep, cognitive processing and enhanced learning and memory task accuracy with Yoga nidra practice in novices. PLoS One. 2023;18(12):e0294678.

28. Kumar K, Joshi B. Study on the effect of Pranakarshan pranayama and Yoga nidra on alpha EEG & GSR. INDIAN JOURNAL OF TRADITIONAL KNOWLEDGE. 2009 Jul;8(3):453–4.

29. Datta K, Yadav N, Narawade Y, Mahajan U, Joseph T, Kumar A. Sleep Strategies for Sportsmen; What can they be? A Randomised Controlled Trial. Sleep Vigilance. 2022 Jun 1;6(1):165–71.

30. Sharpe E, Butler MP, Clark-Stone J, Soltanzadeh R, Jindal R, Hanes D, et al. A Closer Look at Yoga Nidra- Early Randomized Sleep Lab Investigations. J Psychosom Res. 2023 Mar;166:111169.

31. Shashikiran HC, Shetty P, Akshay R, Venugopal A, Shetty S. Effect of Yoga Nidra on the brain activity in individuals with migraine. Yoga Mimamsa. 2022 Jan 1;54(1):18–23.

32. Kjaer TW, Bertelsen C, Piccini P, Brooks D, Alving J, Lou HC. Increased dopamine tone during meditation-induced change of consciousness. Brain Res Cogn Brain Res. 2002 Apr;13(2):255–9.

33. Lou HC, Kjaer TW, Friberg L, Wildschiodtz G, Holm S, Nowak M. A 15O-H2O PET study of meditation and the resting state of normal consciousness. Hum Brain Mapp. 1999;7(2):98–105.

34. Rodin E, Bornfleth H, Johnson M. DC-EEG recordings of mindfulness. Clin Neurophysiol. 2017 Apr;128(4):512–9.

35. Zaccaro A, Riehl A, Piarulli A, Alfì G, Neri B, Menicucci D, et al. The Consciousness State of Traditional Nidrâ Yoga/Modern Yoga Nidra: Phenomenological Characterization and Preliminary Insights from an EEG Study. Int J Yoga Therap. 2021 Jan 1;31(1):Article_14.

36. Bharadwaj I, Kulshrestha A, Anuja. Effect of Yogic Intervention on Blood pressure and Alpha- EEG level of working women. Indian Journal of Traditional Knowledge. 2013 Jul;12(3):542–6.

37. Anuradha R R, Rathi G, Krishnappa MRm, Suresh Kumar MS, Kalpana M. Detecting Stress Level of Students Using Brain Waves Reducing It Using Yoga Therapy. In: 2022 IEEE World Conference on Applied Intelligence and Computing (AIC) [Internet]. 2022 [cited 2024 Aug 20]. p. 764–8. Available from: https://ieeexplore.ieee.org/document/9848919

38. Lee DJ, Kulubya E, Goldin P, Goodarzi A, Girgis F. Review of the Neural Oscillations Underlying Meditation. Front Neurosci [Internet]. 2018 Mar 26 [cited 2025 May 30];12. Available from: https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2018.00178/full

39. Lomas T, Ivtzan I, Fu CHY. A systematic review of the neurophysiology of mindfulness on EEG oscillations. Neuroscience & Biobehavioral Reviews. 2015 Oct 1;57:401–10.

40. Bressan RA, Crippa JA. The role of dopamine in reward and pleasure behaviour – review of data from preclinical research. Acta Psychiatr Scand. 2005 Jun;111(s427):14–21.

41. Radwan B, Liu H, Chaudhury D. The role of dopamine in mood disorders and the associated changes in circadian rhythms and sleep-wake cycle. Brain Research. 2019 Jun 15;1713:42–51.

42. Berry RB, Wagner MH. Introduction. In: Sleep Medicine Pearls [Internet]. Elsevier; 2015 [cited 2025 Jun 12]. p. 17–25. Available from: https://linkinghub.elsevier.com/retrieve/pii/B9781455770519000036