Next-Generation Sequencing Methods for Sensitive Hepatitis B Viral Genome

Analysis: A European Study

Detection of HBV DNA at Low Viral Loads using NGS

Michael X. Fu[1]*, Maria F. Perdomo[2], Sheila F. Lumley[1,3], Johan Ringlander[4,5], Kai Kean[1],

Kaitlin Reid[1], Richard Mayne[1], Oscar Enrique Torres Montaguth[6], Leysa Forrest[7], Sarah

Buddle[6], Johannes C. Botha[6,8], Joakim B. Stenbäck[4,5], Zachery Dickson[2], Chris Kent[9],

Haiting Chai[1], Matthew Byott[6,8], Leo Hannolainen[2], Shannah Secret[1], George Airey[1], Klaus

Hedman[2], Monique I. Andersson[3,10], M. Azim Ansari[1], Eleni Nastouli[6,8], Judith Breuer[6],

Philippa C. Matthews[11,12], Tanya Golubchik[1,13], William L. Irving[14], Peter Simmonds[1], Heli

Harvala[10,15,16]*

1. Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

2. Department of Virology, University of Helsinki, Helsinki, Finland

3. Department of Infectious Diseases and Microbiology, Oxford University Hospitals

NHS Foundation Trust, Oxford, United Kingdom

4. Department of Clinical Microbiology, Sahlgrenska University Hospital, Gothenburg,

Sweden

5. Institute of Biomedicine, Department of Infectious Diseases, University of

Gothenburg, Gothenburg, Sweden

6. Infection, Immunity and Inflammation Department, Great Ormond Street Institute of

Child Health, University College London, London, United Kingdom

7. Genetics and Genomic Medicine Department, Great Ormond Street Institute of Child

Health, University College London, London, United Kingdom

8. Advanced Pathogen Diagnostics Unit, University College London Hospitals NHS Trust, London, United Kingdom

9. Institute of Microbiology and Infection, University of Birmingham, Birmingham, United Kingdom

10. Radcliffe Department of Medicine, University of Oxford, Oxford, United Kingdom

11. Division of Infection and Immunity, University College London, London, United Kingdom

12. The Francis Crick Institute, London, United Kingdom

13. Sydney Infectious Diseases Institute, School of Medical Sciences, Faculty of Medicine and Health, University of Sydney, Sydney, Australia

14. NIHR Nottingham Biomedical Research Centre, Nottingham University Hospitals NHS Trust and the University of Nottingham, Nottingham, United Kingdom

15. Institute of Biomedicine, University of Turku and University Hospital of Turku, Turku, Finland

16. Microbiology Services, NHS Blood and Transplant, Colindale, United Kingdom

*Corresponding authors: Michael X Fu, michael.fu@ndm.ox.ac.uk, Heli Harvala, heli.harvala@utu.fi

Abstract

Objectives: This multicentre study investigated the utility of next-generation sequencing

(NGS) to detect and generate hepatitis B virus (HBV) genomes in samples of low viral load

(from 0.2 to 6207 IU/ml).

Methods: 23 HBV DNA positive plasma samples of genotypes A-E and one HBV-negative

control sample were assayed blindly via 9 established NGS methods from 6 European

laboratories. Methods included untargeted metagenomics, pre-enrichment by probe-capture

followed by Illumina sequencing, and HBV-specific PCR pre-amplification followed by

sequencing with Nanopore or Illumina.

Results: Full HBV genomes were obtained only from samples with viral loads >1000 IU/ml

using probe-capture methods, >200 IU/ml using PCR-Illumina methods, >10 IU/ml using

PCR-Nanopore methods, and in no samples using metagenomic methods. Contamination

was observed in the negative control and samples with very low viral loads in all PCR-

based methods. Probe-capture and metagenomic methods detected additional viruses not

routinely screened in blood donations, including polyomaviruses and herpesviruses;

positive results were confirmed by PCR.

Conclusions: NGS may delineate whole-genome sequences at low viral loads if supported

by a PCR pre-amplification step. Probe-capture methods also reliably detect HBV without

pre-amplification but achieve limited genome characterisation at low viral loads; they may

additionally detect a wide range of blood-borne viruses.

Key words: high-throughput sequencing; sensitivity; virus detection; viral metagenomics;

diagnostics; clinical microbiology; Nanopore; Illumina.

Introduction

The advent of high-throughput sequencing platforms in the mid-2000s, coined 'next-

generation sequencing' (NGS), led to dramatic advances in the characterisation of host and

pathogen genomes in health and disease[1]. Metagenomics, a commonly used NGS method

to sequence all genomic material in a sample, has been instrumental in detecting broad-

spectrum microbes, surveillance of emerging infections, and identifying novel pathogens[2,3].

Compared to Sanger sequencing where sub-genomic amplicons spanning regions of

interest are sequenced, the ability to sequence complete microbial genomes using NGS in

an unbiased manner has proven to be a powerful diagnostic tool[2,3]. However, the wider

adoption of NGS in clinical microbiology and public health has been hindered by complex

methodological workflows, complications in the detection and reporting of contaminating

sequences, and a reduced sensitivity compared to real-time and nested PCR assays[2,4]. In

particular, the inadequate sensitivity of untargeted metagenomics has limited the detection

of hepatitis B virus (HBV), where many HBV-infected clinical samples contain low copy

numbers of the viral genome[5,6].

HBV remains a global public health issue despite the availability of a prophylactic vaccine

and antiviral treatments; the virus chronically infects around 246 million individuals

worldwide, accounting for 1.1 million HBV-related deaths in 2022[7]. In the blood, HBV exists

in relaxed-circular DNA (rcDNA) form: a partially double-stranded genome of approximately

3.2 kb, consisting of overlapping reading frames encoding four genes: polymerase, surface,

precore/core, and X. HBV can be classified into 10 genotypes (A to J) based upon

sequence divergence ≥8%, with inter-genotypic differences in geographic distribution,

transmission mode, and clinical outcomes[8]. Sequencing full-length HBV genomes identifies

(sub)genotypes and recombinants and provides a better understanding of mechanisms of

transmission and persistence, identifies treatment-resistant and immune-escape mutations,

and thus can aid in the development of new therapeutics[9]. Furthermore, whole-genome

sequencing (WGS) of low viral load (VL) samples enables the characterisation of occult

HBV infection and the assessment of viral clearance, persistence and reactivation.

Methodologies used in clinical diagnostics for sequencing viral genomes need to be time-

and cost-efficient, specific, and sensitive to support their integration into clinical, public

health, and research laboratories[10]. Previous evaluations of viral sequencing methods have

been limited to individual workflows and laboratories, with few coordinated comparisons,

typically using synthetic or low-diversity materials due to limited clinical sample availability[11].

Method comparisons using clinical samples of varying loads are limited, but include a

previous investigation of the effectiveness of NGS methods for the characterisation of

hepatitis C virus detection frequencies and genome coverage[12]. The value of many studies

is limited by the inclusion of samples with relatively high VLs without determining the

detection limits of different NGS methods, for example[11].

Amid calls for the HBV field to work progressively toward developing NGS methods[8], this

study brought together six European laboratories with nine established virus sequencing

protocols, encompassing four commonly used NGS methods. First, the untargeted

metagenomics method sequenced all genomic material in the samples. The second method

involved the addition of biotinylated nucleic acid probes to enrich for large numbers of

known pathogens and selectively capture target sequences before Illumina sequencing.

The final methods involved PCR amplification of HBV fragments, followed by sequencing

using the short-read Illumina or the long-read Oxford Nanopore Sequencing (ONT)

platforms. All methods except untargeted metagenomics have previously demonstrated the

ability to read and assemble whole HBV genomes from clinical samples[3,6,10,13].

The four NGS methods were evaluated using a panel of 23 HBV DNA-positive plasma

samples with low VLs. Through this comparison, we assessed the ability of NGS to detect

and generate HBV sequences, as well as to co-detect other viruses potentially present in

the samples. Overall, this study aims to provide an overview of the HBV-specific

sensitivities of currently applied NGS methodologies and to evaluate their potential

suitability for sequencing low VL samples in clinical practice.

Materials and Methods

Samples. Twenty-three large-volume samples, containing up to around 200 mL of plasma,

from individuals infected with HBV genotypes A, B, C, D, and E were obtained from blood

donors from NHS Blood and Transplant (NHSBT). To reflect the heterogeneity in the

extraction methods used by NGS protocols and account for the non-negligible stochastic

effect of measuring low VLs, geometric mean VLs were calculated across eight separate

real-time PCR measurements from eight extractions (five of which were obtained from a

previous investigation[14] and three further extractions from 5 ml of plasma). The samples

had geometric mean VLs ranging from 0.2 to 6207 IU/ml (approximately 1 to 31,000

copies/ml). Furthermore, a control plasma sample (around 200 mL) that screened negative

for HBV serological markers, HBV DNA, and other blood-borne infections was obtained

from NHSBT, where HBV DNA negativity was independently confirmed by real-time PCR

measurements. All donors provided signed consent to use their samples for research

purposes. This study was approved by the Blood Supply Clinical Audit, Risk and

Effectiveness Committee of NHSBT (numbers 22031 and 23002). All 24 samples, including

the negative control, were sent to each laboratory on dry ice, with anonymised sample

identifiers and no other sample information.

Sequencing methods. The panel of 24 blinded samples was used to evaluate the

performance of nine established NGS protocols at six expert European laboratories,

performed in the years 2023-2024. Methods included two untargeted metagenomic

protocols using Illumina (MTG-A and MTG-B), three pan-viral probe-based enrichment

protocols using Illumina (TAC-A, TAC-B, and TAC-C), two HBV-specific PCR amplification

protocols using Illumina (PCR-A and PCR-B [PCR-Illumina]) and two HBV-specific PCR

amplification protocols using ONT (PCR-1 and PCR-2 [PCR-ONT]). The protocols are

specific to each laboratory, and the impact of individual methodological variables could not

be evaluated independently within this study. Further details of protocols are found in the

supplementary material. Of note, the input plasma volumes for extractions ranged from 0.2

to 20 ml. Extracted nucleic acids underwent library preparation according to local protocols,

where input volumes ranged from 5 to 50 µl. Effective test volumes ranged from 4 to 800 µl.

Average bases sequenced per sample differed from 2.2 Mb to 5.03 Gb per sample. The

pre-capture libraries for TAC-B were sequenced for MTG-B; thus, the samples were

prepared similarly for both protocols.

Factors relating to the technical practicality and costs associated with the use of NGS

assays greatly influence their adoption for clinical diagnostics. Laboratories were asked to

provide the total time and cost of each protocol step from extraction to sequencing on a

standardised template document, considering the typical number of samples that would

normally be included per run for each protocol. Times for each stage were separated into

per-person labour time and waiting times (which included machine running times), and

costs were standardised using a 1:0.90:0.75 USD:EUR:GBP exchange rate, as of May 18,

2025.

Bioinformatic processing. Post-read filtering and genome assembly, either mapping to

the closest available reference or *de novo*, were performed using established protocol-

specific pipelines at each centre (see supplementary material). After deduplication of reads

or removal of PCR duplicates and processing of sequences to begin at the EcoR1

restriction site (the conventional origin of the genome[15]), CSV files for each assembled

genome/sample were sent from each centre, containing the number of base calls at each

nucleotide site. These files were received at the coordinating laboratory, where further

bioinformatics analyses were performed using a common set of tools on Microsoft Excel

(version 16.103) and SSE[16,17] (version 1.4) as previously described[12]. A 95% majority base

consensus sequence was calculated at each genome position possessing one or more

base reads, incorporating the calling of ambiguous bases when necessary. Constructed

assembled sequences for each sample were examined and realigned to ensure consistent

numbering, compare nucleotide similarities, and assess genome coverage and accuracy.

Within-sample consensus sequences were assembled for each sample from the different

NGS protocols by combining more than one similar assembled sequence with genome

coverage of around 100% using the program Sequence Join in the SSE package[17],

followed by visual inspection of sites with ambiguous bases. Specific sequences from each

method were compared with the within-sample consensus, and the numbers of nucleotide

differences were recorded using the program Sequence Dist[17]. Missing bases were not

included in comparisons of sequence identity. Thresholds for base-calls were set to

practically represent the realities of sequencing: a threshold of one base read per

nucleotide site was utilised for MTG and TAC methods, whilst a more stringent minimum

threshold of 10 base reads at each nucleotide site was applied for PCR-based methods.

Within-sample consensus NGS sequences were (sub)genotyped as previously described[14].

When whole-genome assemblies could not be constructed (i.e. no two complete NGS

sequences were similar), genotypes were defined from either the most complete NGS

sequence or Sanger sequence, the latter if there were no similar NGS sequences.

Individual NGS sequences from each protocol were genotyped using the Genome

Detective system[18].

Nested PCR and Sanger sequencing. Sequences of a portion of the surface gene

obtained previously from extensive nested PCR and Sanger sequencing were included for

comparison to the NGS sequences[19] (see Figure S1 for phylogenetic tree of all HBV-

positive samples obtained via Sanger sequencing of nested PCR amplicons). Using DNA

extracted from 5 ml of plasma[14], nested PCR and sequencing of two further regions were

performed: one spanning the surface and polymerase genes, and another spanning the X

and precore/core genes (see Table S2 for primer details and nucleotide positions).

Definitions. The following terms used in this manuscript have been defined to aid

readability.

Protocol: unique individual NGS workflows performed by individual laboratories.

Method: a collection of similar protocols based on sequencing technology and platform.

Methods included untargeted metagenomics followed by Illumina sequencing (MTG),

targeted metagenomics using probe capture followed by Illumina sequencing (TAC), PCR

amplification of HBV before Illumina sequencing (PCR-Illumina), and PCR amplification of

HBV before ONT sequencing (PCR-ONT).

Genotype: distinct genetic variations of HBV based on nucleotide differences of $\geq 8\%$[8].

Subgenotype: subgroups within specific HBV genotypes differing by 4-8% in viral genome

sequence[8] and distinct from the broader HBV genotypes.

Strain: a sequence that differed by >5% from other sequences from the same sample,

indicating a different (sub)genotype. A 5% difference was chosen as a practical, rounded

cutoff to reliably distinguish distinct subgenotypes.

Reference sequence: a known digital HBV DNA sequence used as a standard for genome

assembly.

Assembled sequence: individual sequences constructed from individual protocols, which

underwent read filtering and assembly via protocol-specific bioinformatic pipelines, followed

by calculation of majority base consensus sequence at the coordinating laboratory and

realignment of assembled sequences where necessary.

Within-sample consensus sequence: a more robust complete genome sequence for each

sample, derived from the most commonly observed nucleotide at each genome position

across available assembled sequences from all protocols for the same sample.

Contamination/contaminating sequence: a genetically different sequence and strain from

the within-sample consensus sequence or other assembled sequences, where applicable

(>5% nucleotide differences).

True-positive: similar to other sequences from the same sample (<5% nucleotide

differences).

False-positive sequence: detection of an HBV DNA sequence in the negative control.

HBV-specific bases read/base counts: the total number of nucleotide bases aligned to the

HBV genome, following deduplication of sequencing reads and removal of PCR duplicates

where applicable.

Genome coverage: the proportion of the viral genome with consensus bases called.

Shannon entropy: a measure of the diversity of base counts in a sequence, which may

result from technical sequencing errors or natural variability within infecting viral populations

and could also be influenced by VL. Shannon entropy value ranges from 0 (no variability) to

1 (equal frequencies of all bases).

Detection of other viruses by PCR. Since HBV was not the only possible virus sequenced

from the five TAC and MTG protocols from three laboratories, the deduplicated read counts

and genome coverage percentages were obtained for any additional viruses detected from

the sample panel, following the full bioinformatics pipelines for each centre (see

supplementary material for details of full pipelines). When a certain virus was detected by

more than one NGS method in the same sample, confirmatory PCR was performed in all

samples (see supplementary material for details of the PCR protocols).

Statistical analyses. Spearman's correlation was used to assess associations between VL

and sequencing metrics across nine protocols. *P*-values were adjusted for multiple testing

within each set of comparisons using the Benjamini-Hochberg method. α was 0.05. Data

analyses and visualisation were performed with GraphPad Prism (v10.4.2, LLC), except *p*-

value adjustment performed with Microsoft Excel (version 16.103).

Results

HBV detection and accuracy of assembled sequences. HBV amplicons were obtained

for all 23 samples by Sanger sequencing, with good concordance between sequences

generated by Sanger and NGS sequencing. Within-sample consensus NGS sequences

were constructed for 14/15 samples with VL above 50 IU/ml (Table 1). Sequences

assembled using different NGS protocols were generally identical or nearly identical to the

within-sample consensus. A few individual sequences were >5% divergent from the within-

sample consensus (shown in pink), indicating a different contaminating HBV strain.

Assembled sequences were compared for eight samples with lower VLs, for which within-

sample consensus sequences were unavailable. Different HBV genotypes were detected

by at least one protocol for all eight samples (Table S3). Three samples, VLs 1.1, 1.8, and

8.5 IU/ml, yielded discrepant sequences in all NGS protocols.

All four PCR-based protocols detected false-positive HBV sequences in the negative

control (genome coverage, 48%-100%), which were identified as different genotypes (Table

S3). Three of the four protocols also detected contaminating sequences in other samples,

including those with low VL (<50 IU/ml). These four protocols were performed in three

different laboratories, except for extraction, which the coordinating centre conducted for

three of the four protocols. Some contaminants and false positives matched sequences of

other *ex vivo* viruses (Table 1), but most were non-identical. For PCR-based methods other

than PCR-1, the proportion of contaminants or the lack of detection at the lowest VLs

deterred the ability to ascertain true-positive detection confidently. TAC methods had very

few contaminating sequences, and no contaminating sequences were assembled from

MTG methods.

Overall, true-positive HBV sequences of >5% genome coverage were detected in 5-6/23

samples using MTG, 10-17/23 samples using TAC, 9-13/23 samples using PCR-Illumina,

and 16-19/23 samples using PCR-ONT protocols. MTG-B detected 7/13 samples at lower

coverage and read numbers, which were detected by TAC-B, including the six samples with

the highest coverage detected by TAC-B; samples for both protocols were prepared in the

same run before the capture step.

HBV base counts and genome coverage. The highest number of HBV-specific bases was

read in PCR methods (up to 10^9), followed by TAC (up to 10^7) and the lowest in MTG (up to

10^4). A strong correlation was observed between base counts and VL using the PCR-ONT

and TAC methods, whereas this correlation was moderate when using the PCR-Illumina

and MTG methods (Figure 1). Base counts were similar between samples for the PCR-

Illumina methods. For PCR-ONT, bases obtained for true-positive samples with VL <50

IU/ml ranged from 1×10^3 to 5×10^7. For TAC and MTG methods, base counts were much

lower for samples with lower VL. The approximate threshold to reliably detect the sample

panel with true-positive sequences was >50 IU/ml or >10^6 bases for PCR-ONT methods

(except no threshold for PCR-1), >50 IU/ml or >10^4 bases for TAC methods, >100 IU/ml

or >10^7 bases for PCR-Illumina methods, and >3000 IU/ml or >10^3 bases for MTG methods.

Following multiple alignment of read assemblies, the completeness of assembled

sequences was analysed. Complete or near-complete genome sequences were assembled

for all samples with VL >10 IU/ml for PCR-1 protocol, >200 IU/ml for PCR-B, >1000 IU/ml

for TAC-C, and in none of the samples for MTG methods (Figure 2). Coverage thresholds of

80% and 40% could be applied to exclude contamination in the PCR-Illumina and PCR-

ONT methods, respectively, if the 100% coverage in the negative control from PCR-1 was

excluded.

Inspection of the mean genome coverage revealed that the methods yielded uneven

coverage across the genome (Figure 3). PCR-ONT methods produced the most uniform

genome coverage, with the exception of PCR-2, which showed reduced amplification

spanning the region from the start of the polymerase to the surface genes. PCR-Illumina

methods had more regions of reduced coverage, including genome positions 1100 to 2100

for PCR-A and the start of the surface gene for PCR-B. Probe capture efficiency was highly

variable across protocols, each showing areas of markedly low capture efficiency. Coverage

of the HBV genome was even sparser for MTG methods. When analysing individual

samples, there was considerable heterogeneity in the distribution of HBV genome coverage

between protocols, without clear patterns (Figure S2).

Heterogeneity. The diversity of base counts at each site was quantified through

calculations of Shannon entropy. Variability in Shannon values was observed between the

NGS methods for the same samples, which may reflect the variations in the quantities of

base counts by each method. MTG methods showed the least intra-sample HBV sequence heterogeneity, with a maximum mean Shannon entropy of 0.01, followed by the TAC methods (up to 0.04) and the PCR-Illumina methods (up to 0.09). Heterogeneity was extremely high (up to 0.32) for PCR-ONT methods (Figure 4). However, the presence of ambiguous bases in sequences from PCR-ONT methods did not affect the accuracy of consensus sequences generated from the reads. Diversity was higher when VL was greater for PCR-Illumina methods, but was lower when VL was higher for PCR-ONT methods.

Co-detection. Protocol-specific bioinformatic pipelines were employed to determine the presence, read counts, and genome coverage of other viruses for the TAC and MTG methods, which simultaneously screened for multiple viruses from generated sequences (Table 2). Altogether, anellovirus sequences were detected in 22/24 samples by the five NGS protocols. Anellovirus was detected by at least two protocols in 19/24 samples, and 2/24 samples by all five protocols. >50% genome coverage for anellovirus was observed in two samples although consensus sequence generation was hampered by the likely presence of multiple strains and types of anelloviruses within samples. Confirmatory PCR for alpha, beta, and gammatorqueviruses revealed the presence of DNA sequences of all three anellovirus genera in all 24 samples (20/24 αTTV-positive αTTV; 22/24 βTTV-positive; 17/24 γTTV-positive), all with relatively high VLs (Ct values ranged from 4.7 to 26.1). Sanger sequencing of amplicons generated by nested PCR from randomly selected anellovirus-positive samples confirmed the presence of different strains from each genus.

Various human herpesviruses were detected in 17/24 samples across the five MTG and TAC protocols at <50% genome coverage, and were detected by more than one protocol for three samples (EBV in 4/5 protocols in two laboratories for sample HBV8; HHV6 in 2/5

protocols in two laboratories for HBV9; KSHV in 3/5 protocols in three laboratories for

HBV12). Real-time PCR confirmed the presence of these herpesviruses in these three

samples, but no other sample was HHV PCR-positive, including the 13 samples where

TAC-C reported HHV-7 DNA sequences. Polyomavirus DNA sequences were detected in

11/24 samples using NGS protocols; however, they were only detected by more than one

protocol in one sample (HPyV6, detected by two protocols in two laboratories in HBV6).

HyPV6 was, however, not detected by real-time PCR. Instead, MCPyV was detected by

real-time PCR in low copy numbers in HBV6 (detected by one NGS protocol) and HBV21

(not detected by any NGS protocol). Overall, TAC methods detected a greater number of

viruses at higher read counts and coverage than MTG methods.

Time and costs. The per-person labour and waiting times varied between protocols (Figure

4a). The turnaround time was shortest for MTG and most PCR-based protocols. TAC

methods took longer, mostly due to overnight hybridisation steps. Costs were lowest for

PCR-ONT methods, higher for PCR-Illumina methods, followed by MTG methods, whilst

TAC methods had the greatest cost (Figure 4b).

Discussion

Findings from our multicentre study provide an overview of currently available NGS

methodologies for sequencing low VL HBV clinical samples. They highlight the HBV-specific

sensitivities and potential applicability in clinical practice.

While all evaluated methods demonstrated varying capacities to generate HBV sequences

and detect co-infecting viruses, protocols incorporating pre-amplification steps showed

greater sensitivity at low VLs. Reliable detection of HBV at VL ~10 IU/ml (for PCR-1 and

TAC-C) or ~100 IU/ml (for other pre-amplification protocols) would be sufficient for most

HBV infections encountered in clinical settings. For detecting and quantifying samples with

very low VL, such as those associated with occult HBV[14], conventional real-time or nested

PCR remained essential. Previous investigations of PCR-1 and PCR-2 also showed

successful amplification of full HBV genomes at VL <30 IU/ml[6,13], and alternative probe-

capture methods observed a similar 50 copies/ml limit of detection for HBV[20] and 90-100%

coverage for a range of DNA and RNA viruses at ≥1000 copies/ml[21,22]. The substantially

reduced depth of coverage for MTG methods we observed has been well-described for a

range of viruses[10–12]. However, the method retains its value in other applications due to its

ability to detect unanticipated or potentially entirely novel pathogens without a requirement

to pre-select targets in capture and PCR-assisted NGS methods[2].

Although detection sensitivity was similar, genomes generated from PCR-based methods

were near-complete compared to lower genome coverage from TAC methods for our low

VL samples. As an inexpensive method for sequencing whole HBV genomes, PCR-ONT

NGS may be a beneficial screening tool for the HBV and microbiology fields, especially in

low- and middle-income countries[23] when ONT allows preliminary analysis before a run is

completed: highly advantageous in terms of turnaround time for high VL samples. Our

findings contrast with a similar study on hepatitis C virus, which concluded that PCR-based

NGS was relatively laborious[12]; improvements in technologies and workflows could explain

the differences found. However, PCR-based NGS would be more challenging for more

divergent and/or larger viruses than HBV, as primer design for whole-genome amplification

can be difficult, and amplicon drop-out necessitating primer re-design is not uncommon.

Variables that may have affected the methods' detection capabilities in this study include

extraction from larger volumes of plasma (as previously found with PCR-1[13] and TAC-A/B

protocols[19]), having greater sample representation and fewer samples for library

preparation, and sequencing more gigabases per sample. For PCR-based methods,

sensitivity is affected less by the sequencing platform and more by the effectiveness of the

PCR for genome pre-amplification, particularly the conservation of primer-binding sites. A

nested PCR strategy with shorter amplicons would improve sensitivity if WGS is not

required. For TAC methods, sensitivity may have been affected by factors such as probe

design, hybridisation, and wash temperatures. As with the detection of viruses around the

limit of detection of conventional PCR assays, the stochastic effect around the limits of

detection of NGS protocols affects the detection of samples with low VL, albeit at a higher

detection limit than conventional PCR.

Although the sensitivity of PCR-based NGS methods is sufficient for most clinical cases of

HBV, the finding of low-level HBV DNA contamination compromised their diagnostic

reliability at the lowest VLs. Without applying a minimum threshold of 10 for consensus

base calling to account for potential background contamination or sequencing errors, HBV

sequences would have been detected in all samples (24/24) using PCR-based methods;

however, contaminant HBV sequences would also have appeared more frequently (Table

S4). The detection of contaminating sequences has been reported in other studies

comparing NGS protocols[11,21], highlighting contamination as a persistent obstacle to the

reliable characterisation of low VL genomes. Notably, one planned protocol could not be

included in this study because the background level of contamination exceeded the VLs of

the sample panel.

Investigating the exact sources and mechanisms of contamination was beyond the scope of

this study. The negative control was confirmed to be HBV-negative prior to processing,

suggesting that contamination likely occurred during library preparation, amplification, or

sample handling in each of the participating laboratories. This interpretation is supported by

the detection of identical sequences across multiple samples processed within the same

protocol or laboratory (Table 1). The variation observed in both the number and

characteristics of contaminant strains among protocols indicates that contamination events

in low VL samples are relatively frequent but highly context-dependent. In the presence of

background contamination, stricter analytical thresholds can reduce sensitivity but improve

the reliability of positive calls. Such measures may include increasing the minimum read

depth required for consensus base calls, enforcing minimum genome coverage or VL

thresholds, and setting defined read count criteria relative to negative controls (Table S5).

Importantly, minimising contamination in complex NGS workflows requires extensive

experience and rigorous laboratory practice[24], such as using separate PCR workspaces

and tubes, exercising meticulous pipetting, and including appropriate negative plasma

controls.

The substantial variation in base counts and error rates observed among ONT base-calling

algorithms used between the years 2023 and 2024 made it difficult to accurately identify

minority variants. Although these errors are unlikely to impact interpretation for most clinical

and public health purposes, we advise against performing variant calling directly from

mapped ONT reads without considerable bioinformatics expertise to exclude false

substitutions and improve variant calling[25]. Subsequent improvements in R10 flow cells and

base-calling algorithms, after our experiments were performed, are likely to enhance

sequencing accuracy and reduce base count variability[26,27].

NGS methods involving PCR amplification are inherently limited to the characterisation of one known target pathogen. Their use for whole genome characterisation is compromised for viruses with long genomes such as herpesviruses, where multiple, overlapping PCRs may be required, or for targets which are highly genetically heterogeneous and which may lack conserved sites for primer binding across the genome, such as anelloviruses. Metagenomic methods possess clear advantages in their breadth of detection, with frequent co-detections of multiple anelloviruses, human herpesviruses, and polyomaviruses in the samples. The 100% (24/24) prevalence of anellovirus DNA detection by PCR was consistent with previous studies, which have shown that most immunocompetent individuals carry the virus in their blood and at high VLs in patients with HBV infection[28]. The low detection rate by all TAC protocols may have originated from the extreme genetic heterogeneity of the anellovirus species and genera that were not adequately represented among the probes used for target capture[29]. However, this does not explain the similarly low rates of detection by MTG methods, which were not a sensitivity problem as the anelloviruses had high VLs by PCR; it is possible that their circular sequences were too divergent to map to what might be a very limited number of reference sequences used by the bioinformatic software to assign reads. Our detection rates of polyomaviruses and HHVs were comparable to those reported in population studies[30,31]. The high detection rate of HHVs and polyomaviruses with TAC-C compared to fragmented PCR could potentially be explained by the presence of incomplete or highly fragmented genome sequences derived from degraded viral DNA from disintegrating persistently infected cells[32].

Our study focused on HBV samples with low VLs; including samples with higher VLs would likely have increased the proportion achieving high base counts and complete genome coverage, while reducing the relative impact of contamination on detection reliability.

Although we centralised some of the bioinformatics to facilitate comparison, the different

pipeline-specific steps performed by each laboratory could have affected sensitivity, such as

the choice of taxonomic classifier[10]. In a follow-up study, we plan to evaluate the

performance of different bioinformatic pipelines on the raw data generated in the current

study to ascertain optimal tools for HBV characterisation.

In conclusion, this study provides an overview of the HBV-specific sensitivities of currently

used NGS methodologies and their potential applicability for sequencing low VL clinical

samples. NGS approaches incorporating PCR amplification demonstrate sufficient

sensitivity for most HBV infections typically encountered in clinical practice, where VLs

exceed approximately 10 IU/ml. However, the presence of low-level contamination remains

a key limitation, restricting reliable genome identification at the lowest VLs. Future efforts

should focus on identifying and mitigating sources of background contamination to enhance

the specificity and reliability of NGS. Strengthening contamination control would enable

broader clinical implementation of NGS as a valuable tool for characterising HBV and other

viral pathogens in diverse diagnostic settings.

Acknowledgements: The authors thank Dr Belinda Singleton (NHS Blood and Transplant)

for helping develop the anellovirus PCR protocol. This study was primarily supported by the

National Institute for Health and Care Research [grant number NIHR203338], which

supported these authors under the Blood and Transplant Research Unit 'Genomics to

Enhance Microbiology Screening': MXF, KK, KR, RM, OETM, SB, SS, JBr, TG, PS, and HH.

The study was also supported by the Swedish Society for Medical Research (grant number

PG-23-0435) and the Gothenburg Society for Medicine (grant number GLS-1001038). SFL

is funded by a Wellcome Doctoral Training Fellowship (grant number 220549/Z/20/Z); this

research was funded in part by the Wellcome Trust. MFP and LH acknowledge funding from

Suomen Lääketieteen säätiö and Medicinska Understödsföreningen Liv & Hälsa. KH and

ZD acknowledge funding from the Jane and Aatos Erkko Foundation. TG is supported by

an Investigator Grant (GNT2025445) from the National Health and Medical Research

Council, Australia (NHMRC). PCM receives core funding from the Francis Crick Institute (ref

CC2223) and acknowledges funding support from the University College London NIHR

Biomedical Research Centre (BRC). MAA is supported by a Sir Henry Dale Fellowship

jointly funded by the Royal Society and Wellcome Trust (220171/Z/20/Z). The funding

bodies had no role in the study's design, data collection, analysis, or manuscript writing. For

the purpose of Open Access, the author has applied a CC BY public copyright licence to

any Author Accepted Manuscript version arising from this submission.

Author contributions: Conceptualisation: HH, PS, MXF. Formal analysis: MXF, HH, PS, WLI.

Investigation: MXF, MFP, SFL, JR, KK, KR, RM, OETM, LF, SB, JBo, JBS, ZD, CK, HC, MB,

LH, SS, GA, KH, MAA, EN, JBr, PM, TG. Writing – Original Draft: MXF. Writing – Review &

Editing: all authors. Supervision: HH, PS, MA. All authors approved the final version of the

submitted manuscript.

Conflicts of Interest: The authors declare no conflicts of interest that pertain to this work.

Data Availability Statement: The data supporting the findings of this study are included in

the manuscript. Raw sequencing data have been deposited in the NCBI BioProject

accession number PRJNA1364201, including BioSample accession numbers

SRR36116996 to SRR36117201. Consensus full-length genome sequences have been

deposited in GenBank under accession numbers PV786119 to PV786132.

References

Goodwin Sara, McPherson John D., McCombie W. Richard. Coming of age: ten years of next-generation sequencing technologies. *Nature Reviews Genetics 2016 17:6* 2016;**17**(6):333–51. Doi: 10.1038/nrg.2016.49.

Chiu Charles Y., Miller Steven A. Clinical metagenomics. *Nat Rev Genet* 2019;**20**(6):341–55. Doi: 10.1038/S41576-019-0113-7,.

Deng Xianding, Achari Asmeeta, Federman Scot, Yu Guixia, Somasekar Sneha, Bártolo Inês, et al. Metagenomic sequencing with spiked primer enrichment for viral diagnostics and genomic surveillance. *Nat Microbiol* 2020;**5**(3):443–54. Doi: 10.1038/S41564-019-0637-9,.

Thorburn Fiona, Bennett Susan, Modha Sejal, Murdoch David, Gunson Rory, Murcia Pablo R. The use of next generation sequencing in the diagnosis and typing of respiratory infections. *Journal of Clinical Virology* 2015;**69**:96. Doi: 10.1016/J.JCV.2015.06.082.

Matthews Philippa C., Downs Louise O., Vawda Sabeehah, Bester Phillip Armand, Lythgoe Katrina A., Wang Tingyan, et al. Bimodal distribution and set point HBV DNA viral loads in chronic infection: Retrospective analysis of cohorts from the UK and South Africa. *Wellcome Open Res* 2020;**5**. Doi: 10.12688/WELLCOMEOPENRES.15941.2.

Lumley Sheila F., Kent Chris, Jennings Daisy, Chai Haiting, Airey George, Waddilove Elizabeth, et al. Whole genome sequencing of hepatitis B virus using tiled amplicon (HEPTILE) and probe based enrichment on Illumina and Nanopore platforms. *Sci Rep* 2025;**15**(1):5795. Doi: 10.1038/S41598-025-87721-1;SUBJMETA=208,234,2483,2513,255,325,631,692,699;KWRD=VIRAL+GENETICS, VIRAL+HEPATITIS.

World Health Organization. *World health statistics 2024: monitoring health for the SDGs, Sustainable Development Goals*. Geneva; 2024.

McNaughton Anna L., D'Arienzo Valentina, Ansari M. Azim, Lumley Sheila F., Littlejohn Margaret, Revill Peter, et al. Insights From Deep Sequencing of the HBV Genome—Unique, Tiny, and Misunderstood. *Gastroenterology* 2019;**156**(2):384–99. Doi: 10.1053/j.gastro.2018.07.058.

Rajoriya Neil, Combet Christophe, Zoulim Fabien, Janssen Harry L.A. How viral genetic variants and genotypes influence disease and treatment outcome of chronic hepatitis B. Time for an individualised approach? *J Hepatol* 2017;**67**(6):1281–97. Doi: 10.1016/J.JHEP.2017.07.011.

Buddle Sarah, Forrest Leysa, Akinsuyi Naomi, Martin Bernal Luz Marina, Brooks Tony, Venturini Cristina, et al. Evaluating metagenomics and targeted approaches for diagnosis and surveillance of viruses. *Genome Medicine 2024 16:1* 2024;**16**(1):1–20. Doi: 10.1186/S13073-024-01380-X.

Lopez-Labrador F. Xavier, Huber Michael, Sidorov Igor A., Brown Julianne R., Cuypers Lize, Laenen Lies, et al. Multicenter benchmarking of short and long read wet lab protocols for clinical viral metagenomics. *Journal of Clinical Virology* 2024;**173**. Doi: 10.1016/J.JCV.2024.105695,.

Thomson Emma, Ip Camilla L.C., Badhan Anjna, Christiansen Mette T., Adamson Walt, Ansari M. Azim, et al. Comparison of Next-Generation Sequencing Technologies for Comprehensive Assessment of Full-Length Hepatitis C Viral Genomes. *J Clin Microbiol* 2016;**54**(10):2470. Doi: 10.1128/JCM.00330-16.

Stenbäck Joakim B., Schmidt Daniel, Noborg Ulrika, Gustafsson Joel, Norberg Peter, Andersson Maria E., et al. Accurate and Cost-Efficient Whole Genome Sequencing of

Hepatitis B Virus Using Nanopore. *J Med Virol* 2025;**97**(10):e70630. Doi: 10.1002/JMV.70630.

Fu Michael X., Simmonds Peter, Andreani Julien, Baklan Hatice, Webster Mhairi, Asadi Romisa, et al. Ultrasensitive PCR system for HBV DNA detection: Risk stratification for occult hepatitis B virus infection in English blood donors. *J Med Virol* 2023;**95**(10). Doi: 10.1002/JMV.29144.

Ono Yoshitaka, Onda Haruo, Sasada Reiko, Igarashi Koichi, Sugino Yukdo, Nishioka Kusuya. The complete nucleotide sequences of the cloned hepatitis B virus DNA; subtype adr and adw. *Nucleic Acids Res* 1983;**11**(6):1747. Doi: 10.1093/NAR/11.6.1747.

Simmonds Peter. SSE: A nucleotide and amino acid sequence analysis platform. *BMC Res Notes* 2012;**5**. Doi: 10.1186/1756-0500-5-50,.

GitHub - psimmond/SSE: SSE Sequence editor. Available at https://github.com/psimmond/SSE. Accessed November 14, 2025, n.d.

Vilsker Michael, Moosa Yumna, Nooij Sam, Fonseca Vagner, Ghysens Yoika, Dumon Korneel, et al. Genome Detective: an automated system for virus identification from high-throughput sequencing data. *Bioinformatics* 2019;**35**(5):871–3. Doi: 10.1093/BIOINFORMATICS/BTY695.

Fu Michael X, Larralde Osmany, Mayne Richard, Kean Kai, Reid Kaitlin, Andersson Monique, et al. Use of polyethylene glycol precipitation and ultracentrifugation to enhance the sensitivity of hepatitis B virus DNA detection. *Journal of Clinical Virology* 2025:105802. Doi: 10.1016/J.JCV.2025.105802.

Kapoor Vishal, Briese Thomas, Ranjan Amit, Donovan William M., Mansukhani Mahesh M., Chowdhary Rashmi, et al. Validation of the VirCapSeq-VERT system for

differential diagnosis, detection, and surveillance of viral infections. *J Clin Microbiol* 2023;**62**(1):e00612-23. Doi: 10.1128/JCM.00612-23.

Mourik Kees, Sidorov Igor, Carbo Ellen C., van der Meer David, Boot Arnoud, Kroes Aloysius C.M., et al. Comparison of the performance of two targeted metagenomic virus capture probe-based methods using reference control materials and clinical samples. *J Clin Microbiol* 2024;**62**(6). Doi: 10.1128/JCM.00345-24,.

Lewandowska Dagmara W., Zagordi Osvaldo, Geissberger Fabienne Desirée, Kufner Verena, Schmutz Stefan, Böni Jürg, et al. Optimization and validation of sample preparation for metagenomic sequencing of viruses in clinical samples. *Microbiome* 2017;**5**(1):94. Doi: 10.1186/S40168-017-0317-Z,.

Lemoine Maud, Eholié Serge, Lacombe Karine. Reducing the neglected burden of viral hepatitis in Africa: Strategies for a global approach. *J Hepatol* 2015;**62**(2):469–76. Doi: 10.1016/J.JHEP.2014.10.008.

López-Labrador F. Xavier, Brown Julianne R., Fischer Nicole, Harvala Heli, Van Boheemen Sander, Cinek Ondrej, et al. Recommendations for the introduction of metagenomic high-throughput sequencing in clinical virology, part I: Wet lab procedure. *Journal of Clinical Virology* 2021;**134**:104691. Doi: 10.1016/J.JCV.2020.104691.

Ye Fuqiang, Zhu Juanjuan, Zhang Xiaomin, Zhang Jiarong, Xie Zihan, Yang Tingting, et al. Characteristics and filtering of low-frequency artificial short deletion variations based on nanopore sequencing. *Gigascience* 2025;**14**:1–15. Doi: 10.1093/GIGASCIENCE/GIAF018.

Zhang Tianyuan, Li Hanzhou, Ma Silin, Cao Jian, Liao Hao, Huang Qiaoyun, et al. The newest Oxford Nanopore R10.4.1 full-length 16S rRNA sequencing enables the accurate resolution of species-level microbial community profiling. *Appl Environ*

Microbiol 2023;**89**(10). Doi: 10.1128/AEM.00605-23/SUPPL_FILE/AEM.00605-23-S0008.EPS.

Liu-Wei Wang, van der Toorn Wiep, Bohn Patrick, Hölzer Martin, Smyth Redmond P., von Kleist Max. Sequencing accuracy and systematic errors of nanopore direct RNA sequencing. *BMC Genomics* 2024;**25**(1):1–15. Doi: 10.1186/S12864-024-10440-W/FIGURES/6.

Spandole Sonia, Cimponeriu Dănuţ, Berca Lavinia Mariana, Mihăescu Grigore. Human anelloviruses: an update of molecular, epidemiological and clinical aspects. *Arch Virol* 2015;**160**(4):893–908. Doi: 10.1007/S00705-015-2363-9/FIGURES/4.

Pyöriä Lari, Pratas Diogo, Toppinen Mari, Hedman Klaus, Sajantila Antti, Perdomo Maria F. Unmasking the tissue-resident eukaryotic DNA virome in humans. *Nucleic Acids Res* 2023;**51**(7):3223–39. Doi: 10.1093/NAR/GKAD199.

Kamminga Sergio, van der Meijden Els, de Brouwer Caroline, Feltkamp Mariet, Zaaijer Hans. Prevalence of DNA of fourteen human polyomaviruses determined in blood donors. *Transfusion (Paris)* 2019;**59**(12):3689–97. Doi: 10.1111/TRF.15557,.

Edelman Daniel C. Human herpesvirus 8 - A novel human pathogen. *Virol J* 2005;**2**(1):1–32. Doi: 10.1186/1743-422X-2-78/TABLES/1.

Delbue Serena, Tremolada Sara, Elia Francesca, Carloni Camilla, Amico Serena, Tavazzi Eleonora, et al. Lymphotropic Polyomavirus is detected in peripheral blood from immunocompromised and healthy subjects. *Journal of Clinical Virology* 2010;**47**(2):156–60. Doi: 10.1016/j.jcv.2009.11.029.

Figure 1. Relationship between viral loads and HBV-specific bases read for each group of

methods on a common log scale. Pink indicates contaminating sequences, including where

HBV sequences were obtained for the negative HBV10 sample on the y-axis. Samples not

detected are found on the x-axis. Spearman's correlation test with Benjamini-Hochberg

adjustment deduced the significance of the association between linear viral loads and base

counts (ρ value and *p*-values listed in boxes). HBV: hepatitis B virus; MTG: untargeted

metagenomics; TAC: target capture; ONT: Oxford Nanopore Technologies.

Figure 2. Relationship between viral loads and breadth coverage of HBV assembled

sequences for each group of methods on a common axes scale. Pink indicates

contaminating sequences, including where HBV sequences were obtained for the negative

HBV10 sample on the y-axis. Samples not detected are found on the x-axis. Spearman's

correlation test with Benjamini-Hochberg adjustment deduced the significance of the

association between linear viral loads and genome coverage (ρ value and *p*-values listed in

boxes). HBV: hepatitis B virus; MTG: untargeted metagenomics; TAC: target capture; ONT:

Oxford Nanopore Technologies.

Figure 3. Normalised mean read depth of the HBV genome for each of the sequencing

methods. The normalised mean read depth was calculated as the number of bases at each

genome position as a proportion of the total reads of the sequence, then multiplied by the

total number of sites, where the expected mean value was one base per site. The eleven

highest viral load samples were included when detected, and only true-positive sequences

with more than 1000 total HBV-specific bases read were included; the number of samples

from each protocol that were included is indicated in each plot's key. Genome positions

were based on the D00330 reference sequence. A genome diagram of HBV, drawn to the x-

axis scale, shows gene positions in shaded boxes and regulatory regions in unshaded

boxes. TAC: target capture; MTG: untargeted metagenomics.

Figure 4. Relationship between viral loads and mean Shannon entropy values for

polymorphic sites for each group of methods on a common axes scale. Pink indicates

contaminating sequences, including where HBV sequences were obtained for the negative

HBV10 sample on the y-axis. Samples not detected are found on the x-axis. Spearman's

correlation test with Benjamini-Hochberg adjustment deduced the significance of the

association between linear viral loads and entropy values (ρ value and *p*-values listed in

boxes). HBV: hepatitis B virus; MTG: untargeted metagenomics; TAC: target capture; ONT:

Oxford Nanopore Technologies.

Figure 5. Comparison of a) time taken, and b) cost for each protocol based on the total

time and costs for a protocol-specific representative number of samples included in a

typical run. For Figure 5a, coloured bars to the left of the plot show per-person labour time

per sample, whereas the same-coloured bars with lighter-coloured diagonal lines to the

right show the waiting time per sample, which may include machine running times. For

visual comparison, sequencing time and costs per sample were standardised to the

shortest and cheapest Illumina or Nanopore protocol; the overall times reported include the

full sequencing durations as recorded by each laboratory. For clinical purposes, PCR-1 is

normally run overnight on the MinIon flow cell (average of 16 hours), but was run for 72

hours for this study. Costs were standardised using a 1:0.8959:0.7531 USD:EUR:GBP

exchange rate, as of May 18, 2025, and total costs were rounded to the nearest ten. USD:

United States Dollar; EUR: Euro; GBP: Great British Pound; QC: quality control; TAC: target

capture; MTG: untargeted metagenomics.

Table 1. Overview of the detection of HBV in sample sets. The percentage genome completeness is given in each cell. The left-hand side of the table with blue headers shows the sequences obtained from the NGS methods, whilst the right-hand side (calamine colour) shows the sequences obtained by Sanger sequencing (length of S amplicon = 395 nucleotides; SP = 973 nucleotides; XC = 698 nucleotides). Within-sample consensus sequences were assembled for each sample from the individual sequences of the NGS methods that had around 100% genome coverage and were genetically similar, which were assigned to HBV genotypes. The percentage genetic relatedness of the sequences from each method was compared to the within-sample consensus sequence, where assembled sequences similar to the within-sample consensus were shaded on a grayscale, and different strains (differing in >5% nucleotide sequence from the within-sample consensus) were shaded in a pink scale. The absence of sequences was shaded in black. Where within-sample consensus sequences could not be assembled (i.e. no two complete sequences were similar), genotypes were defined from Sanger sequences, and the NGS sequences were instead compared to one another. Samples are ranked from highest to lowest geometric mean viral load. For TAC-B: library preparation failed for sample HBV13; samples HBV4, HBV9, and HBV19 had reads of poor quality for the original pipeline to process, so the mean base quality was dropped below 1.0. #: For PCR-1, a negative water control was tested instead of the negative plasma control. Alphabetical superscripts (a, b, c) represent identical sequences. Numerical superscripts next to method identifiers indicate where the same laboratory performed the protocols.

Sample	Viral load (IU/ml)	PCR (Nanopore)		PCR (Illumina)		Target capture			Metagenomics		Within-sample consensus	Genotype assigned	Sanger sequencing		
		PCR-1	PCR-2[1]	PCR-A	PCR-B[1]	TAC-A[2]	TAC-B[2]	TAC-C	MTG-A	MTG-B[2]			S	SP	XC
HBV15	6207.0	100%	100%	91%	100%	11%	100%	90%	25%	72%	Y	A2	12%	30%	22%
HBV8	3154.0	100%	99%	98%	100%	91%[c]	97%	99%	50%	14%	Y	D1	12%		
HBV2	3108.0	100%	90%		100%	90%[c]	81%	100%		25%	Y	A2	12%		22%
HBV7	1058.0	100%	100%	98%	99%	17%	83%	93%	11%	25%	Y	C1	12%	29%	22%
HBV19	859.8	100%	49%		100%	22%	4%	88%	10%		Y	D3	12%	31%	22%
HBV16	599.2	100%	85%	59%	100%	9%		75%			Y	A1	12%		
HBV1	551.6	100%	50%	76%	100%	6%	38%	88%	4%		Y	D2	12%		
HBV14	522.3	100%	85%	92%	100%	9%	95%	80%		49%	Y	E	12%	30%	
HBV23	337.1	100%	57%	99%	98%	64%	71%	93%		16%	Y	D4	12%	31%	22%
HBV17	299.9	100%	100%	77%	60%		14%	50%			Y	E	12%	30%	
HBV9	138.1	100%	100%	33%	99%	9%	8%	35%	2%		Y	A1	12%	30%	
HBV5	128.4	100%			57%			66%			N	E	12%	30%	
HBV22	73.6	100%	54%		77%			54%			Y	D2	12%	31%	
HBV4	60.5	100%	85%	59%	100%	35%	7%	100%	12%		Y	B4	12%	30%	22%
HBV13	60.4	100%	85%		94%			91%	5%		Y	D1	12%	31%	22%
HBV21	41.0	100%	37%		73%[a]			7%	43%		N	A	12%	30%	
HBV6	11.3	100%			57%		3%	28%		3%	N	D3	12%	31%	
HBV18	8.5		37%*		33%*						N	A	12%*		
HBV20	3.3	100%	17%		43%[a]			26%	<1%		N	D	12%		
HBV12	1.8		17%		56%[a]			25%			N	A	12%		
HBV24	1.1				71%[b]						N	E	12%		
HBV11	0.3	99%	4%		29%[b]						N	D	12%		
HBV3	0.2		52%		74%[a]			5%			N	A3	5%		
Negative (HBV10)	-	100%[#]	17%	48%	45%						N	-			

Within-sample consensus	Shading key					
Y (% sequence identity to within-sample consensus)	No sequence	≥85%	≥90%	≥95%	≥98%	≥99.5%
N (similarity of sequences)	No sequence	Different sequence to others	Similar sequence			

Table 2. Detection of other viruses by the targeted capture and untargeted metagenomics methods, showing deduplicated read counts (genome coverage %). Results reported for TAC-C are for viruses where at least 10 reads were obtained. More than one TTV species may have been identified for MTG-A, but the most abundant was included. The furthest right column shows the PCR results, where PCR was performed for all samples when two or more methods detected a virus in the same sample. These PCRs were specific for alpha, beta and gamma-torquevirus, Merkel cell polyomavirus, and herpesviruses. '-' denotes where the method did not detect the virus, '=' denotes where the PCR was not performed. A numerical superscript indicates where the same laboratory performed the protocols. HHV: human herpesvirus; MCPyV: Merkel cell polyomavirus; AV: anellovirus; TTV: torquetenovirus; EBV: Epstein-Barr virus; KSHV: Kaposi's sarcoma-associated herpesvirus; JCPyV: John Cunningham virus.

Sample	Virus	MTG-A	MTG-B[1]	TAC-A[1]	TAC-B[1]	TAC-C	PCR
HBV1	AV	43 (54%)	-	6 (14%)	3 (3%)	37 (22%)	**Ct 13.1 (αTTV)**
	HHV7	-	-	-	-	885 (10%)	**Negative**
HBV2	MCPyV	-	-	-	-	14 (20%)	**Negative**
	AV	-	-	-	-	35 (21%)	**Ct 9.6 (αTTV), Ct 8.0 (βTTV), Ct 21.1 (γTTV)**
	HHV7	-	-	-	-	649 (10%)	**Negative**
HBV3	AV	9 (20%)	2 (1%)	7 (20%)	19 (12%)	167 (49%)	**Ct 6.3 (αTTV), Ct 5.8 (βTTV), Ct 8.5 (γTTV)**
	EBV	-	-	-	-	321 (6%)	**Negative**
	HHV7	-	-	-	-	1699 (9%)	**Negative**
HBV4	AV	26 (49%)	-	11 (20%)	14 (16%)	117 (45%)	**Ct 10.9 (αTTV), Ct 4.7 (βTTV), Ct 13.0 (γTTV)**
	Simian adenovirus 25	7 (0.74%)	-	-	-	-	**=**

HBV5	AV	-	5 (8%)	-	20 (16%)	15 (15%)	**Ct 22.2 (αTTV), Ct 11.0 (βTTV)**
	JCPyV	-	-	-	-	63 (52%)	**=**
HBV6	AV	6 (21%)	44 (40%)	7 (12%)	310 (75%)	84 (36%)	**Ct 7.2 (βTTV)**
	MCPyV	-	-	-	-	126 (65%)	**152 copies/mL**
	HPyV6	-	-	-	4 (4%)	26 (30%)	**Negative**
	HHV7	-	-	-	-	565 (10%)	**Negative**
	JCPyV	-	-	-	-	70 (56%)	**=**
HBV7	AV	-	2 (2%)	13 (6%)	11 (24%)	14 (17%)	**Ct 7.2 (γTTV)**
	HPV-2	-	-	-	-	10 (8%)	**=**
	HHV7	-	-	-	-	1112 (9%)	**Negative**
	JCPyV	-	-	-	-	34 (41%)	**=**
HBV8	EBV	12 (0.2%)	-	42 (14%)	3 (2%)	2020 (39%)	**Ct 35.9**
	AV	-	1 (1%)	1 (5%)	-	-	**Ct 12.5 (αTTV), Ct 8.3 (βTTV), Ct 21.4 (γTTV)**
	MCPyV	-	-	-	-	35 (31%)	**Negative**
HBV9	AV	-	-	3 (4%)	1 (1%)	-	**Ct 15.0 (αTTV), Ct 8.9 (βTTV), Ct 19.2 (γTTV)**
	HHV6	-	-	-	1 (1%)	1626 (15%)	**Ct 38.2**
HBV10	AV	-	3 (1%)	3 (4%)	1 (9%)	-	**Ct 12.6 (αTTV), Ct 10.7 (βTTV)**
	HHV7	-	-	-	-	730 (7%)	**Negative**
HBV11	AV	-	-	1 (6%)	2 (13%)	22 (10%)	**Ct 6.1 (αTTV), Ct 5.9 (βTTV), Ct 22.5 (γTTV)**
	EBV	-	-	-	-	774 (7%)	**Negative**
	HHV7	-	-	-	-	805 (7%)	**Negative**
HBV12	KSHV	18 (0.63%)	-	1 (5%)	-	7353 (93%)	**Ct 36.6**
	EBV	-	-	-	1 (1%)	-	**Negative**
	HHV7	-	-	-	-	1324 (9%)	**Negative**
	AV	-	2 (1%)	-	-	-	**Ct 26.1 (αTTV), Ct 13.3 (βTTV)**
HBV13	MCPyV	-	-	-	-	16 (15%)	**Negative**
	TTV	-	3 (2%)	-	-	20 (24%)	**Ct 11.9 (αTTV), Ct 5.1 (βTTV), Ct 13.0 (γTTV)**

	HHV7	-	-	-	-	825 (11%)	**Negative**
HBV14	AV	-	8 (11%)	-	88 (33%)	-	**Ct 7.0 (αTTV), Ct 10.1 (βTTV)**
	JCPyV	-	-	-	-	10 (10%)	**=**
HBV15	JCPyV	-	-	-	-	69 (59%)	**=**
	AV	7 (9.6%)	7 (1%)	-	16 (35%)	11 (8%)	**Ct 12.4 (αTTV), Ct 11.4 (βTTV), Ct 15.2 (γTTV)**
	EBV	-	-	-	1 (1%)	-	**Negative**
HBV16	AV	-	1 (1%)	-	1 (12%)	-	**Ct 13.8 (αTTV), Ct 9.1 (βTTV)**
HBV17	HHV7	-	-	-	-	413 (9%)	**Negative**
	AV	-	-	-	-	-	**Ct 13.3 (βTTV), Ct 15.3 (γTTV)**
HBV18	AV	-	-	1 (3%)	1 (1%)	-	**Ct 12.9 (αTTV), Ct 11.4 (βTTV), Ct 21.0 (γTTV)**
HBV19	AV	-	-	-	-	-	**Ct 19.1 (αTTV), Ct 16.2 (βTTV), Ct 9.5 (γTTV)**
HBV20	AV	27 (41%)	-	8 (13%)	2 (8%)	25 (18%)	**Ct 12.2 (αTTV), Ct 11.5 (βTTV), Ct 6.9 (γTTV)**
HBV21	TTV	-	-	1 (4%)	3 (9%)	-	**Ct 11.9 (βTTV), Ct 9.4 (γTTV)**
	HPV	-	-	-	-	17 (15%)	**=**
	HHV7	-	-	-	-	833 (9%)	**Negative**
	MCPyV	-	-	-	-	-	**219 copies/mL**
HBV22	AV	-	-	-	8 (11%)	-	**Ct 11.4 (αTTV), Ct 12.1 (βTTV), Ct 21.4 (γTTV)**
	HHV6	-	-	-	-	1070 (7%)	**Negative**
HBV23	AV	-	2 (1%)	11 (35%)	22 (26%)	56 (30%)	**Ct 5.8 (αTTV), Ct 6.4 (βTTV), Ct 19.8 (γTTV)**
	EBV	-	-	-	-	212 (6%)	**Negative**
	HHV6	-	-	-	-	1535 (7%)	**Negative**
	HHV7	-	-	-	-	761 (5%)	**Negative**
	MCPyV	-	-	-	-	10 (8%)	**Negative**

	AV	-	2 (1%)	22 (28%)	4 (6%)	31 (21%)	**Ct 7.8 (αTTV), Ct 7.0 (βTTV), Ct 13.5 (γTTV)**
HBV24	HHV6	-	-	-	-	616 (11%)	**Negative**
	HPyV7	-	-	-	-	13 (12%)	**=**
	HPyV6	-	-	-	-	27 (24%)	**Negative**
	KSHV	-	-	-	-	226 (5%)	**Negative**
	HHV7	-	-	-	-	326 (5%)	**Negative**