Type and developmental timing of childhood adversity predicts psychopathology symptoms

in a South African birth cohort

Sherief Y. Eldeeb[,1,2] – seldeeb@purdue.edu

Brooke G. McKenna[1,2,3] – mckenn34@purdue.edu

Marilyn T. Lake[7] – marilyn.lake@uct.ac.za

Nadia Hoffman[7] – nadia.hoffman@uct.ac.za

Alexandre A. Lussier[2,3] – alussier@mgh.harvard.edu

Esther Walton[4] – E.Walton@bath.ac.uk

Andrew J. Simpkin[5] – Andrew.simpkin@universityofgalway.ie

Andrew D.A.C. Smith[6] – andrew18.smith@uwe.ac.uk

Heather J Zar[8] – heather.zar@uct.ac.za

Dan J. Stein[7] – dan.stein@uct.ac.za

Erin C. Dunn[1,2,3] – erindunn@purdue.edu

[1]Department of Sociology, College of Liberal Arts, Purdue University, West Lafayette, IN

[2]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA

[3]Department of Psychiatry, Harvard Medical School, Boston, MA

[4] Department of Psychology, University of Bath, Bath, United Kingdom

[5]School of Mathematical and Statistical Sciences, University of Galway, Galway, Ireland

[6]Mathematics and Statistics Research Group, University of the West of England, Bristol, UK

[7] South African Medical Research Council (SAMRC) Unit on Risk and Resilience in Mental

Disorders, Department of Psychiatry and Mental Health & Neuroscience Institute, Faculty of

Health Sciences, University of Cape Town, South Africa

[8] South African Medical Research Council (SAMRC) Unit on Child & Adolescent Health, Cape

Town, South Africa Department of Paediatrics and Child Health, South African Medical

Research Council (SAMRC) Unit on Child and Adolescent Health, University of Cape Town,

South Africa

45   **Abstract**

Background: Childhood adversity is widespread globally and is one of the strongest predictors

of later psychopathology. However, the differential effects of type and timing of childhood

adversities on childhood psychopathology remain unclear, highlighting the need to explore

which life-course hypotheses (sensitive periods, accumulation of exposure, and/or recency of

exposure) best explain these associations. Of particular importance, there is a lack of research in

low- and middle-income countries (LMIC), where children experience higher rates of adversity

relative to children in high-income countries (HIC).

Methods: Participants included 787 children and their mothers from a South African birth

cohort, the Drakenstein Child Health Study. Mothers reported child exposure to adversity from

birth to 8 years of age across six adversity categories. We used the two-stage Structured Life

Course Modeling Approach (SLCMA) to examine life-course associations between childhood

adversity exposures and internalizing/externalizing symptoms measured using the Child

Behavior Checklist at age 8 years.

Results: Maternal psychopathology, maternal adverse events, child food insecurity, and child

exposure to community/domestic violence had the strongest associations with child

psychopathology symptoms, with varying life-course models selected. The accumulation

hypothesis best explained associations of maternal adverse events (partial R^2=2.3%) and child

exposure to community/domestic violence (partial R^2=1.6%) with internalizing symptoms. The

combined middle childhood sensitive period (age 5≤8 years) and recency hypotheses model best

explained associations between maternal psychopathology and internalizing (partial R^2=7.0%) or

externalizing (partial R^2=5.1) symptoms.

Conclusions: We identified that different types and timing of childhood adversity confer

differential risk for childhood psychopathology symptoms in this LMIC sample. Our work has

implications for strategically-timed intervention and prevention strategies to improve mental

health, which may need to be specifically designed for children in LMIC.

Keywords: childhood adversity, life-course hypotheses, sensitive period, low- and middle-

income countries (LMIC), SLCMA

Abbreviations: low- and middle-income countries (LMIC); Drakenstein Child Health Study

(DCHS); Structured Life-course Modeling Approach

Word count: 4927

90   **Introduction**

Mental health disorders are a leading cause of disability globally, with prevalence

estimates continuing to rise (Network, 2022; World Health Organization, 2017). Childhood

adversity remains one of the strongest predictors of later psychopathology, with lifetime risk

doubling for those who experience childhood adversity (Green et al., 2010; McLaughlin et al.,

2012). In a global study spanning 21 countries, approximately 38% of people reported at least

one form of adversity during childhood; the eradication of childhood adversity was estimated to

reduce the prevalence of mental health disorders by up to 42% globally (Kessler et al., 2010).

However, optimal targets and timing for intervention remain unclear, particularly for children in

low- and middle-income countries (LMIC). Gaining a clearer understanding of how the type and

timing of childhood adversity impacts risk for mental health disorders is an important step

towards developing effective and cost-efficient prevention and intervention efforts (Schaefer et

al., 2022).

Historically, childhood adversity research has disproportionally focused on WEIRD

(Western, Educated, Industrialized, Rich, and Democratic) samples, despite 88% of the world's

children residing in low- and middle-income settings (Ceccarelli et al., 2022; Henrich et al.,

2010; Network, 2022). Understanding the relationships between childhood adversities and risk

for mental health disorders in LMIC is crucial to expanding generalizability of results and

informing the adaptation of interventions to differing cultural and socioeconomic contexts.

Children in LMIC countries, such as South Africa, are exposed to higher rates of adversity

including intimate partner violence, parental substance abuse, and traumatic events (Ceccarelli et

al., 2022; Stein et al., 2015). One study comparing longitudinal birth cohorts found 91% of

participants from a South African cohort experienced at least one childhood adversity by age 10

years, compared to 64% and 67% in British and Dutch cohorts, respectively (Bigler et al., 2025).

Further, studies examining adversities generally focus on individual-level adversities, such as

interpersonal trauma, rather than community- and systemic-level factors, such as exposure to

community violence and chronic food insecurity (Stirling et al., 2015). Children in LMIC are

disproportionately exposed to these macro-level adversities, and increasing evidence points to

their negative impact on psychopathology risk (Hyde et al., 2022).

Beyond examining the *types* of childhood adversity that influence risk for

psychopathology, it is also vital to understand how *timing* of exposure may alter risk. Evidence

from a recent systematic review (Schaefer et al., 2022) and empirical studies suggest both type

and timing of childhood adversity influences the development of childhood psychopathology

(i.e., internalizing and externalizing) symptoms, which are early indicators of later

psychopathology risk (Dunn et al., 2018; Farooq et al., 2024; Juen et al., 2024). Life-course

theory proposes the timing of adversity exposure may differentially impact outcomes, based on:

exposure during a *sensitive period* of development, *accumulation* of exposure across childhood,

and the *recency* of an exposure. Sensitive period hypotheses posit there are specific

developmental periods when an exposure disproportionately impacts outcomes (Ben-Shlomo &

Kuh, 2002; Frankenhuis & Fraley, 2017; Schaefer et al., 2022). Accumulation hypotheses

contend that the number of exposures linearly increases risk, regardless of timing (Evans et al.,

2013; Guidi et al., 2020; Slopen et al., 2018). Recency hypotheses argue adversities occurring

closer to the timing of the outcome have stronger effects (Shanahan et al., 2011). Research

conducted in HIC contexts, like the United Kingdom and the Netherlands, provides mixed

evidence for developmental timing (Bagner et al., 2010; Yoon, 2022), accumulation (Buss et al.,

2011; Hogye et al., 2022; Mäntymaa et al., 2012; Rogosch et al., 2011), and recency of exposure

(Dunn et al., 2018) predicting childhood psychopathology symptoms. Very few studies have

examined which life-course hypotheses best explain child psychopathology symptoms in LMIC

contexts, with a single cross-sectional study conducted in Tanzania finding the recency of

exposure as the strongest predictor of childhood psychopathology compared to factors such as

accumulation, exposure at certain ages, and severity of exposure (Juen et al., 2024).

In the current study, we investigated which of three life-course hypotheses (sensitive

period, accumulation of risk, and/or recency of exposure) best explains the associations between

early adversity and childhood internalizing/externalizing symptoms. We examined these

adversities in a LMIC birth cohort of South African children who are disproportionately

impacted by poverty and other childhood adversities relative to children represented in HIC life-

course studies (Dunn et al., 2018; Farooq et al., 2024; Moss et al., 2023). We explored a broad

range of adversity types relative to previous work, with exposures spanning household-level –

maternal psychopathology, substance abuse, intimate partner violence (IPV), and maternal

adverse events – and the community-level – child food insecurity and exposure to community

and domestic violence.

151   **Methods**

Participants and Procedure

Data were obtained from the Drakenstein Child Health Study (DCHS), a

multidisciplinary longitudinal birth cohort from the Western Cape of South Africa (Stein et al.,

2015; Zar et al., 2015). Pregnant mothers were enrolled at 20-28 weeks' gestation from two

clinics, Mbekweni and TC Newman, during routine antenatal care appointments between March

2012 and March 2015. The Mbekweni clinic predominantly serves an isiXhosa-speaking Black

African population, whereas the TC Newman clinic predominantly serves an Afrikaans-speaking

mixed-race population of individuals self-identifying as 'Coloured'[1]. Both clinics are located in

the peri-urban Drakenstein subdistrict. To be eligible for the study, women needed to: (1) be

18☐years of age or older, (2) intend to attend antenatal care at one of the two clinics, and (3)

intend to remain in the area for at least a year.

Measures

Mothers answered in-person questionnaires on child exposure to adversity,

psychopathology symptoms, and demographic information. Trained study staff administered

measures in the preferred language of parents using the Research Electronic Data Capture

(REDCap) platform (Harris et al., 2019; Harris et al., 2009). All study measures were collected

within a target range of one month prior and three months following the child's birthdate, except

---

[1] In South Africa, the apartheid system divided individuals into different racial groups (e.g., "Black African", "Coloured", "White"). Under democracy these categories continue to be employed, partly for reasons of redress and partly due to South Africans' cultural embrace and celebration of these identities Dooms, T., & Chutel, L. E. (2023). *Coloured: How Classification Became Culture*. Jonathan Ball Publishers. In DCHS, participants self-identify as "Black African" or "Coloured". We are aware that these categories are not widely used internationally and may have painful racist connotations in other cultures. At the same time, we think it is important to honor South African participants' identities and culture by using the language and terminology they self-report. By using these categories, we are not aiming to reify social categories but instead hope to contribute to the study of ongoing health disparities stemming from systemic racism.

during the COVID-19 pandemic during which the range was expanded to six months following

the child's birthdate for most measures (**Figure S1**).

Predictors: Childhood Adversity

We assessed six types of childhood adversity: (1) maternal psychopathology, (2) maternal

adverse events, (3) child food insecurity, (4) child exposure to community and domestic

violence, (5) intimate partner violence between mother and a partner, and (6) maternal substance

abuse. Exposure to childhood adversity was measured between three to 12 timepoints, depending

on category, from birth to 8 years of age (see **Table S1** for a list of all measures and administered

timepoints). Measures assessing adversity exposure at birth or during the prenatal period were

excluded from the analyses. Participants were dichotomously coded as exposed or unexposed to

each category of adversity. Adversity measures are introduced below, with more detailed

descriptions and psychometrics provided in **Appendix S1**. Each adversity was analyzed

independently.

1. **Maternal Psychopathology.** Maternal psychopathology was measured using two

questionnaires. The first, the *Beck Depression Inventory (BDI-II)*, is an established and validated

21-item measure assessing the severity of major depressive disorder symptoms, including

sadness, pessimism, anhedonia, mood changes, and difficulty sleeping over the last two weeks

(Beck et al., 1996).

The second questionnaire, the *Self-Reporting Questionnaire (SRQ-20)*, is a 20-item tool

endorsed by the World Health Organization (WHO) to capture non-psychotic psychiatric

symptoms, including symptoms of depressive and anxiety disorders (Harding et al., 1980).

2. **Maternal Adverse Life Events.** Maternal adverse life events were measured using the

17-item *Life Events Questionnaire (LEQ)* (Myer et al., 2008). Items are scored according to

whether the event occurred in the last 12 months (0 = no, 1 = yes).

3. **Child Food Insecurity.** An adapted version of the US Department of Agriculture Short

Form Household Food Security Scale (USDA-HFSS) was used to assess perceived household

food insecurity resulting from financial difficulties in the last 6 months (Bickel et al., 2000).

4. **Child Exposure to Community and Domestic Violence.** Child exposure to

community and domestic violence were measured using the 35-item *Child Exposure to Violence*

Checklist (CECV), a validated measure of child exposure to violence in South Africa (Bruwer et

al., 2008; Fincham et al., 2009; Martin et al., 2013). Lifetime exposure was measured until 4.5

years, and past-year exposure was measured at later timepoints.

5. **Maternal Intimate Partner Violence (IPV).** Maternal intimate partner violence (IPV)

was measured using the 12-item *Intimate Partner Violence (IPV) Questionnaire*, a measure

adapted from the WHO country study on women's health and domestic violence against women

(Jewkes, 2002) and the Women's Health Study in Zimbabwe (Shamu et al., 2011).

6. **Maternal Substance Abuse.** Maternal substance abuse was measured using the

Alcohol, Smoking, and Substance Involvement Screening Test (ASSIST). The ASSIST is a 8-item

questionnaire developed by the WHO to measure substance use across 10 categories: tobacco,

alcohol, cannabis, cocaine, amphetamine-type stimulants, inhalants, sedatives, hallucinogens,

opioids, and other drug use over the past three months (Group, 2002). In the current study,

maternal substance abuse was dichotomized such that participants with scores in the low-risk

range for all substances, excluding tobacco, were considered unexposed (0) and participants with

scores in the moderate- and high-risk ranges for any substance, excluding tobacco, were

considered exposed (1). Maternal tobacco use was not considered an adversity exposure, but

rather was included as a covariate at birth.

Outcomes: Internalizing/Externalizing Problems

Parents reported on child psychopathology symptoms when children were 8 years old

using two widely used measures: the Child Behavior Checklist for Ages 6-18 (CBCL/6-18) and

the Strengths and Difficulties Questionnaire (SDQ). Our primary analyses focused on the

CBCL/6-18, as it more comprehensively assesses symptomatology (via 113 items, versus 25

items in the SDQ), demonstrates stronger internal and interrater reliability metrics (Achenbach et

al., 2008; Goodman, 2001), and offers greater clinical interpretability given its use of normed T-

scores and defined risk ranges relative to the raw continuous scores provided by the SDQ.

Nevertheless, many life-course studies solely use the SDQ. To facilitate greater comparability

with these studies, we repeated all analyses using the SDQ as our secondary outcome

(**Supplemental Materials**).

Covariates

Sociodemographic Questionnaire

A questionnaire adapted from the SASH study was administered to collect

sociodemographic information such as age, marital status, and income (Herman et al., 2009;

Myer et al., 2008). All analyses adjusted for the following covariates assessed at birth: child

race/ethnicity, child sex, maternal age, maternal marital status, number of mother's previous

pregnancies, highest level of maternal education, owned assets, household income, maternal

smoking, and maternal HIV status. Covariates were selected for their theoretical role as potential

confounders. Maternal smoking and HIV status at birth were included due to the high relative

prevalence in this population, as in previous studies using the DCHS cohort (Feil et al., 2022;

Moyakhe et al., 2023; Stein et al., 2015).

Data Analysis

Multiple Imputation

Among participants with complete outcome data (n=787), some were missing data on

covariates (up to 12.8%) and/or exposure variables (0.1%-23.4%) (see comparisons of total and

analytic samples in **Table S2** and **Figure S2**). To maximize our sample size, and in turn improve

statistical power and reduce biased conclusions from these data, we conducted multiple

imputation using the MICE package in R version 4.3.1 (van Buuren & Groothuis-Oudshoorn,

2011) within and across timepoints (**Appendix S2**).

Quantifying Life-course Hypotheses

We defined three developmental periods (very early childhood: $\leq$ 3 years old, early

childhood: 3$\leq$5, middle childhood: 5$\leq$8) for each adversity category. These age ranges are

consistent with previous work and were chosen to facilitate comparisons between studies (Dunn

et al., 2017; Dunn et al., 2018; Dunn et al., 2019). We considered a child as exposed to a specific

adversity if they were exposed at any timepoint within each developmental period.

Sensitive Period. We considered three sensitive period hypotheses corresponding to the

three developmental periods for each adversity, with one exception. For child exposure to

community and domestic violence only two developmental periods – the first five years ($\leq$5

years old) and middle childhood (5$\leq$8) – were tested due to the absence of past-year exposure

data prior to age 5.

Accumulation. Accumulation of exposure was defined as the sum of exposed

developmental periods for each adversity category. Values ranged from 0 (not exposed in any

developmental period) to 3 (exposed during all developmental periods).

Recency. Recency of exposure was defined as the weighted sum of exposed

developmental periods for each adversity category, weighted by the most recent timepoint

represented in the developmental period (i.e., weights of 3 for very early childhood, 5 for early

childhood, and 8 for middle childhood). Thus, children exposed in more recent developmental

periods had higher recency scores relative to children exposed earlier in development. For

example, if a participant was exposed in the very early childhood (3) and middle childhood (8)

developmental periods, their value was 11. Values ranged from 0 (not exposed in any

developmental periods) to 16 (exposed during all developmental periods).

Preliminary Analyses

Preliminary univariate descriptive statistics were used to examine the frequency of

demographics, adversity exposures, and covariates. Bivariate tetrachoric correlations were

conducted to examine the average correlations within each adversity category across

developmental periods, and the average correlations within each developmental period across

adversity categories. Multiple regression analyses were conducted to examine the "ever-

exposed" relationship for each adversity category on psychopathology outcome variables.

Primary Analyses: Structured Life-course Modeling Approach (SLCMA)

The two-stage Structured Life-course Modeling Approach (SLCMA; pronounced slick-

mah; Smith et al., 2016; Smith et al., 2015; Smith et al., 2021) was utilized to determine which

life-course hypotheses, individually or in combination, best explained the associations between

each type of adversity exposure and symptoms of psychopathology. A full description of the two-

stage process of SLCMA is described in the **Supplemental Materials**. Significance was set at a

p-value of .05. We present all results, regardless of p-value, as LARS enables the selection of the

best-fitting hypotheses through examination of elbow plots rather than relying on significance

thresholds (Smith et al., 2021) (**Appendix S2**).

Sensitivity Analyses

We conducted two sensitivity analyses to assess whether our findings varied by

differences in how life-course models were operationalized or covariates included. First, we

examined, for all adversities other than maternal psychopathology, whether results remained

consistent when controlling for maternal postpartum depression, as measured via the Edinburgh

Postpartum Depression Scale (EPDS; Cox et al., 1989). As in previous life-course hypothesis

work (Dunn et al., 2018), maternal postpartum depression was included as a covariate to control

for the potential bias maternal mood may have on reporting of child adversity exposure and

psychopathology (Chilcoat & Breslau, 1997; Holt et al., 2008; Ringoot et al., 2015). Second, we

tested whether averaging or summing data from individual timepoints influenced the life-course

hypotheses selected, to assess whether combining measurement timepoints into developmental

periods may have unduly influenced results (results reported in **Supplemental Material**).

<p align="center">**Results**</p>

Sample Demographics

A total of 1225 mothers were originally enrolled in the study, with 1137 giving live birth

to 1143 children. Data from 787 child-mother pairs were available after removing those with

non-singleton births (n=11 children) and without complete outcome data (n=339).

The analytic sample largely comprised families from disadvantaged backgrounds. Most

mothers were single (60%), completed less than secondary education (62.9%), and earned less

than half the national median household income at the time of data collection (87.6%;

GlobalData, 2022). Children were primarily of Black African (60.6%) descent, followed by those

of Coloured backgrounds (39.2%). Male and female children were equally represented (50.3%

female) (**Table 1**).

Adversity Characteristics

Adversity exposure prevalence ranged from 30.9% (child food insecurity) to 76.9%

(maternal adverse events; **Table 2**). Exposure prevalence tended to be highest in the earliest

developmental period measured (very early childhood), except for child food insecurity and child

exposure to community and domestic violence. Adversities were moderately correlated both

within adversity type across time (r=.22 to .53; **Figure 1**), as well as between adversities (r=.23

to .38; **Figure 1**).

Overall exposure, irrespective of timing, was associated with increased internalizing and

externalizing symptoms for all forms of childhood adversity. After controlling for covariates,

maternal psychopathology explained the greatest variance in both internalizing (partial R^2=3.5%)

and externalizing (partial R^2=2.4%) symptoms. After that, child exposure to domestic and

community violence (partial R^2=2.1%), child food insecurity (partial R^2=1.5%), and maternal

IPV (partial R^2=0.9%) explained the most variance for externalizing symptoms. Maternal

adverse events (partial R^2=1.8%), child exposure to domestic and community violence (partial

R^2=1.6%), and child food insecurity (partial R^2=1.5%) explained the most variance for

internalizing symptoms. Unstandardized effect estimates and partial R^2 are shown in **Figure 2**.

Life-course model selection

Results from the two-stage SLCMA are presented in **Table 3** for each adversity. Exposure

to maternal psychopathology in middle childhood increased internalizing (β=6.27, p=0.001, 95%

CI [2.69, 11.98]) and externalizing (β=6.38, p=0.004, 95% CI [1.79, 13.12]) symptoms.

Exposure to maternal adverse events in all developmental periods also increased internalizing

and externalizing symptoms, following an accumulation model for internalizing symptoms

(β=1.51, p<0.001, 95% CI [0.84, 2.15]) and a recency model for externalizing symptoms

(β=0.37, p<0.001, 95% CI [0.22, 0.51]). Finally, child exposure to community and domestic

violence increased internalizing symptoms (β=1.51, p=0.001, 95% CI [0.69, 2.44]) and child

food insecurity in early childhood increased externalizing symptoms (β=4.30, p=0.002, 95% CI

[1.58, 6.60]). No other adversities were associated with internalizing or externalizing symptoms

at the 5% significance threshold. Of note, the same hypotheses were selected when the SDQ was

used to assess child symptomatology (**Table S3**; **Figure S3**).

Figure 3 displays effect estimates across developmental periods for life-course models

demonstrating the strongest effects between childhood adversities and child psychopathology

symptoms. We present all other elbow plots in **Figure S4** and life-course models in **Figure S5.**

Sensitivity Analyses

Results were largely similar when controlling for maternal postpartum depression (**Table**

S4) and when utilizing alternative life-course theory operationalizations (**Table S5**).

<div align="center">**Discussion**</div>

Our findings suggest both the type and timing of childhood adversity influence risk for

childhood psychopathology symptoms among children in a LMIC context. These differential

timing and adversity effects align with previous evidence from HIC contexts (Dunn et al., 2020;

Dunn et al., 2018; Farooq et al., 2024) and highlight the importance of parsing type and timing of

childhood adversity exposure to inform risk evaluation, prevention, and targeted intervention

approaches.

Timing was particularly important in the case of maternal psychopathology, with later

exposure increasing risk. A combination of recency and the middle childhood sensitive period

hypotheses were selected, indicating that risk for childhood psychopathology symptoms

increased with later age of exposure and additional risk was conferred if the child was exposed

during middle childhood (ages 5 to 8). The selection of recency and the middle childhood

sensitive period hypotheses are somewhat inconsistent with previous evidence that earlier

exposure to maternal psychopathology exerts greater risk for later psychopathology symptoms

(Goodman et al., 2011; Luby et al., 2020) or that timing does not matter (Farooq et al., 2024).

However, Farooq and colleagues (2024) found cumulative exposure to parental mental health

problems best predicted later depression across two UK cohorts, with exposure during middle

childhood conferring the greatest risk across the developmental periods measured. Together,

these findings suggest middle childhood may warrant attention as a developmental period when

maternal psychopathology exposure may be especially influential in shaping child

psychopathology symptoms. Indeed, middle childhood is a time of rapid emotional development

in children, especially in the areas of empathy and emotion regulation (Berk, 2022; De

Raeymaecker & Dhar, 2022). Developmental advances in this period may therefore be more

sensitive to symptoms associated with maternal psychopathology such as negative emotions,

withdrawal, and low attachment security (Lovejoy et al., 2000; Suveg et al., 2011; Whittenburg

et al., 2023). Future research is needed to disentangle the potential mechanisms through which

maternal psychopathology during middle childhood may confer elevated risk for

psychopathology symptoms.

Overall it is also worth noting that, irrespective of timing, maternal psychopathology

demonstrated the strongest associations with internalizing and externalizing symptoms relative to

all other childhood adversities. Maternal psychopathology may be uniquely potent compared to

other adversity types due to the compound influence of both genetics and environment.

Heritability studies highlight the role of genetics in the transmission of psychopathology

symptoms from parent to child, and adoption studies support the additional role of environmental

exposure to maternal psychopathology symptoms (Cioffi et al., 2021; Jami et al., 2020).

Maternal psychopathology may also have a particularly profound influence on child symptoms

due to their bidirectional, or transactional, nature whereby parent and child symptoms influence

one another across time (Buchanan-Pascall et al., 2017). This finding highlights maternal

psychopathology as an important intervention target to reduce child risk for psychopathology,

especially in populations such as within the DCHS cohort where mothers are predominantly the

sole caretakers of children. Indeed, prior studies examining parent interventions aimed at

reducing maternal psychopathology indicate small to moderate effects in improving child

symptoms (Cuijpers et al., 2015; Gunlicks & Weissman, 2008), including in LMIC contexts

(Hoffmann et al., 2022). Of note, many of these interventions target mothers during their child's

first years of life (Cuijpers et al., 2015). Our work suggests interventions attempting to reduce

maternal psychopathology and improve child symptoms may have greater success if

implemented when children are between ages 5 and 8.

By contrast, food insecurity exposure during the early childhood sensitive period (ages 3

to 5) best predicted childhood psychopathology symptoms. Our results in this LMIC study differ

from those conducted in HIC, which have not pointed to a sensitive period when exposure is

disproportionately associated with impaired behavioral and emotional development (Shankar et

al., 2017). Our findings raise important questions about why the early childhood sensitive period

in particular emerged in this LMIC context. One explanation may be that food insecurity serves

as a proxy for malnutrition. Indeed, proper nutrition plays a particularly important role for neural

plasticity prior to age 5 (Georgieff et al., 2015) and the pre-frontal cortex, an area crucial for

emotion regulation, displays rapid development from ages 3 to 5 (Hodel, 2018). Given that

measures of deprivation, such as poverty and food insecurity, have been linked with slowed

development of the pre-frontal cortex (Hodel, 2018), it is possible that food insecurity between

ages 3 to 5 disproportionately impacts child pre-frontal cortex development, which, in turn,

influences later development of psychopathology symptoms. Taken together, our findings

suggest future research would benefit from better understanding how timing differences

influence the impact of food insecurity on child psychopathology symptoms.

Specificity in the timing of exposure mattered less for maternal adverse events and child

exposure to domestic/community violence, where the accumulation of exposures was most often

predictive of child symptoms. These findings align with previous research on the importance of

cumulative life stressors for both parents and children in predicting childhood psychopathology

symptoms (Benito-Gomez et al., 2019; Evans et al., 2013; Hogye et al., 2022). Research

conducted in the US in both cross-sectional and prospective longitudinal samples support our

findings, with additional exposure of community violence or maternal stressful life events

conferring corresponding increased risk for child internalizing and/or externalizing symptoms

(Fleckman et al., 2016; Rudd et al., 2022). By contrast, recency of child abuse was the best

predictor of child internalizing and externalizing symptoms in a LMIC sample from Southern

Tanzania, although the study was cross-sectional and retrospective which may account for the

difference with our results (Juen et al., 2024). The selection of the accumulation hypothesis for

maternal adverse events and child exposure to domestic/community violence suggests a more

general prevention strategy may be more advantageous than targeting specific developmental

periods. Overall, our study highlights the benefits of a secondary screening prevention strategy

for both parents and children experiencing cumulative life stressors where additional events may

indicate additional need for intervention. A risk strategy is especially critical in the context of

LMIC settings where stressors are more prevalent (Bigler et al., 2025).

Strengths and Limitations

422   Our study has several notable strengths. We separately examined six forms of childhood

adversity in a LMIC where children are disproportionately exposed to early life adversity,

focusing not only on individual-level factors, but also community- and systemic-level factors,

such as food insecurity and child exposure to domestic/community violence. Despite increasing

evidence that community- and systemic-level factors are negatively associated with child mental

health (Fleckman et al., 2016; Gard et al., 2021), these factors remain understudied in life-course

research examining risk for child psychopathology symptoms. In particular, literature examining

food insecurity in LMIC has predominantly focused on cognitive and motor skill development

rather than on indicators of child psychopathology symptoms (De Oliveira et al., 2021).

Moreover, by examining adversities separately rather than grouping them into a total exposure

score, we could identify unique timing effects for each category of adversity. The prospective

longitudinal cohort design allowed us to study timing effects with additional rigor and draw

wider conclusions than cross-sectional or retrospective designs.

435   Our study also had limitations. Our study had a smaller sample size compared to previous

life-course work in HIC samples. Thus, we grouped exposures into sensitive periods (e.g., early

childhood, ages 1-3) rather than examining each individual timepoint. While this strategy

improved statistical power, it also reduced the granularity of our findings related to timing of

exposure. During the COVID-19 pandemic, the study team expanded target windows for data

collection between June 10, 2021 and November 3, 2022. Although this approach increased

participation, it also increased variability in child age at collection, which may have impacted the

specificity of our timing findings.

Conclusion

To our knowledge, we have conducted the first prospective study in a LMIC sample

examining which life-course hypotheses best explain the associations between childhood

adversities and psychopathology symptoms in middle childhood. Our findings reinforce the

importance of considering the type and timing of adversity when assessing risk for child

psychopathology in highly adversity-exposed populations. If replicated, our findings can inform

the administration of strategically-timed interventions, as children exposed to adversity during

specific developmental periods may also be more sensitive to related interventions. The WHO

estimates a roughly four-fold return of global investment in anxiety and depression treatments

(Chisholm et al., 2016). By incorporating optimally-timed interventions that buffer the

deleterious impacts of adversity, there is potential to further maximize these benefits, particularly

among children in LMIC who are disproportionately impacted by childhood adversity.

> Key Points
> - Childhood adversity is widespread and is a strong predictor of later psychopathology.
> - Much of the research in childhood adversity and life-course hypotheses occurs in high-income countries.
> - Our work, conducted in the low-and middle-income country (LMIC) of South Africa, found that middle childhood and recency life-course hypotheses had the strongest associations with later childhood internalizing and externalizing symptoms.
> - Future research should continue exploring sensitive periods and mechanisms of childhood adversity in LMIC to inform prevention and intervention efforts.

Data Availability Statement: The Drakenstein Child Health Study is committed to the principle

of data sharing. De-identified data will be made available to requesting researchers as

appropriate. Requests for collaborations are welcome. More information can be found on our

website [http://www.paediatrics.uct.ac.za/scah/dclhs].

Ethical Information: The DCHS was approved by the Faculty of Health Sciences, Human

Research Ethics Committee, University of Cape Town (401/2009) and by the Western Cape

Provincial Health Research committee. Mothers provided informed consent at enrolment and

were re-consented annually. Consent was done in mother's preferred language: English,

Afrikaans or isiXhosa. Assent was obtained annually from children from 7 years of age

depending on their neurocognitive ability.

Acknowledgments: We greatly thank the parents and children who participated in this study. We

would also like to thank the study staff in Paarl, the study data and laboratory teams, and the

clinical and administrative staff of the Western Cape Government Health Department at Paarl

Hospital and at the clinics for support of the study. Additionally, we would like to thank Alison

Hoffnagle, Samantha Stoll, Isabel Schuurmans, Anke Hüls, and Anna Ruehlmann for their

assistance in preparing this manuscript. This manuscript reflects the views of the authors and

may not reflect the opinions or views of the NIH.

Funding: The Drakenstein Child Health Study (DCHS) was funded by the Bill and Melinda

Gates Foundation (OPP1017641 and OPP1017579), the National Institute of Mental

Health (1R21MH098662–01), the NIH/H3Africa (1U01AI110466-01A1), the National Research

Foundation, the South African Medical Research Council, and the Wellcome Trust

(221372/Z/20/Z). Research reported in this publication was supported by the National Institute of

Mental Health of the National Institutes of Health (R01 MH113930, PI: ECD). The NIMH had

no further role in study design; in the collection, analysis and interpretation of data; in the

writing of the report; and in the decision to submit the paper for publication. The content is

solely the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. EW received funding from UK Research and Innovation (UKRI)

under the UK government's Horizon Europe / ERC Frontier Research Guarantee [BrainHealth,

grant number EP/Y015037/1]. AAL is supported by an MQ Fellows Award from the MQ

Foundation (MQF22-9).

Conflict of Interest: The authors declare they have no competing interests.

Correspondence: Address correspondence to: Erin C. Dunn, ScD, MPH, Professor of Sociology,

College of Liberal Arts, Purdue University, Bindley Bioscience Center (BIND), 1203 W. State

Street, West Lafayette, IN 47906 erindunn@purdue.edu

Phone: 765-496-8206; Website: www.thedunnlab.com

508 References
509
Achenbach, T. M., Becker, A., Dopfner, M., Heiervang, E., Roessner, V., Steinhausen, H. C., &

Rothenberger, A. (2008). Multicultural assessment of child and adolescent

psychopathology with ASEBA and SDQ instruments: research findings, applications, and

future directions. *J Child Psychol Psychiatry*, *49*(3), 251-275.

https://doi.org/10.1111/j.1469-7610.2007.01867.x

Bagner, D. M., Pettit, J. W., Lewinsohn, P. M., & Seeley, J. R. (2010). Effect of maternal

depression on child behavior: A sensitive period? *Journal of the American Academy of*

Child & Adolescent Psychiatry, *49*(7), 699-707. https://doi.org/10.1097/00004583-

201007000-00010

Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory-II*.

Psychological Corporation.

Ben-Shlomo, Y., & Kuh, D. (2002). A life course approach to chronic disease epidemiology:

conceptual models, empirical challenges, and interdisciplinary perspectives. *Int J*

Epidemiol, *31*(2), 285-293.

Benito-Gomez, M., Fletcher, A. C., & Buehler, C. (2019). Youth parasympathetic functioning

moderates relations between cumulative family risk and internalizing behaviors. *Journal*

of Youth and Adolescence, *48*(11), 2307-2322.

Berk, L. E. (2022). *Development through the lifespan*. Sage Publications.

Bickel, G., Nord, M., Price, C., Hamilton, W., & Cook, J. (2000). *United States Department of*

Agriculture (USDA) Guide to Measuring Household Food Security. https://fns-

prod.azureedge.net/sites/default/files/FSGuide.pdf

Bigler, M. E., Schuurmans, I. K., Zhu, Y., Siddique, N., Hoffman, N., Koen, N., Zar, H. J., Stein,

D. J., Cecil, C. A. M., & Dunn, E. C. (2025). Are different types of childhood adversities

correlated within and across time? A comparison of data on adversity co-occurrence from

three longitudinal birth cohort studies. *Child Abuse Negl*, 107315.

https://doi.org/10.1016/j.chiabu.2025.107315

Bruwer, B., Emsley, R., Kidd, M., Lochner, C., & Seedat, S. (2008). Psychometric properties of

the Multidimensional Scale of Perceived Social Support in youth. *Comprehensive*

Psychiatry, *49*(2), 195-201.

https://doi.org/https://doi.org/10.1016/j.comppsych.2007.09.002

Buss, K. A., Davis, E. L., & Kiel, E. J. (2011). Allostatic and environmental load in toddlers

predicts anxiety in preschool and kindergarten. *Dev Psychopathol*, *23*(4), 1069-1087.

https://doi.org/10.1017/s0954579411000502

Ceccarelli, C., Prina, E., Muneghina, O., Jordans, M., Barker, E., Miller, K., Singh, R., Acarturk,

C., Sorsdhal, K., Cuijpers, P., Lund, C., Barbui, C., & Purgato, M. (2022). Adverse

childhood experiences and global mental health: avenues to reduce the burden of child

and adolescent mental disorders. *Epidemiology and psychiatric sciences*, *31*, e75, Article

e75. https://doi.org/10.1017/S2045796022000580

Chilcoat, H. D., & Breslau, N. (1997). Does psychiatric history bias mothers' reports? An

application of a new analytic approach. *Journal of the American Academy of Child and*

Adolescent Psychiatry, *36*(7), 971-979. https://doi.org/10.1097/00004583-199707000-

00020

Chisholm, D., Sweeny, K., Sheehan, P., Rasmussen, B., Smit, F., Cuijpers, P., & Saxena, S.

(2016). Scaling-up treatment of depression and anxiety: a global return on investment

analysis. *The Lancet Psychiatry*, *3*(5), 415-424. https://doi.org/10.1016/S2215-

0366(16)30024-4

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression.

Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal*

of Psychiatry, *150*, 782-786. http://www.ncbi.nlm.nih.gov/pubmed/3651732

De Oliveira, K. H., Buccini, G., Hernandez, D. C., Pérez-Escamilla, R., & Gubert, M. B. (2021).

Household food insecurity and early childhood development in Brazil: an analysis of

children under 2 years of age. *Public Health Nutr*, *24*(11), 3286-3293.

De Raeymaecker, K., & Dhar, M. (2022). The Influence of Parents on Emotion Regulation in

Middle Childhood: A Systematic Review. *Children (Basel)*, *9*(8).

https://doi.org/10.3390/children9081200

Dooms, T., & Chutel, L. E. (2023). *Coloured: How Classification Became Culture*. Jonathan Ball

Publishers.

Dunn, E. C., Nishimi, K., Neumann, A., Renaud, A., Cecil, C. A. M., Susser, E. S., & Tiemeier,

H. (2020). Time-dependent effects of exposure to physical and sexual violence on

psychopathology symptoms in late childhood: in search of sensitive periods in

development. *Journal of the American Academy of Child and Adolescent Psychiatry*,

59(2), 283-295. https://doi.org/10.1016/j.jaac.2019.02.022

Dunn, E. C., Nishimi, K., Powers, A., & Bradley, B. (2017). Is developmental timing of trauma

exposure associated with depressive and post-traumatic stress disorder symptoms in

adulthood? *Journal of Psychiatric Research*, *84*, 119-127.

https://doi.org/10.1016/j.jpsychires.2016.09.004

Dunn, E. C., Soare, T. W., Raffeld, M. R., Busso, D. S., Crawford, K. M., Davis, K. A., Fisher, V.

A., Slopen, N., Smith, A., Tiemeier, H., & Susser, E. S. (2018). What life course

theoretical models best explain the relationship between exposure to childhood adversity

and psychopathology symptoms: recency, accumulation, or sensitive periods?

Psychological Medicine, *48*(15), 2562-2572. https://doi.org/10.1017/s0033291718000181

Dunn, E. C., Soare, T. W., Zhu, Y., Simpkin, A. J., Suderman, M. J., Klengel, T., Smith, A.,

Ressler, K. J., & Relton, C. L. (2019). Sensitive periods for the effect of childhood

adversity on DNA methylation: results from a prospective, longitudinal study. *Biological*

Psychiatry, *85*(10), 838-849. https://doi.org/10.1016/j.biopsych.2018.12.023

Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development.

Psychological Bulletin, *139*(6), 1342.

Farooq, B., Russell, A. E., Howe, L. D., Herbert, A., Smith, A. D. A. C., Fisher, H. L., Baldwin,

J. R., Arseneault, L., Danese, A., & Mars, B. (2024). The relationship between type,

timing and duration of exposure to adverse childhood experiences and adolescent self-

harm and depression: findings from three UK prospective population-based cohorts.

Journal of Child Psychology and Psychiatry, *65*(10), 1369-1387.

https://doi.org/https://doi.org/10.1111/jcpp.13986

Feil, D., Abrishamcar, S., Christensen, G., Vanker, A., Koen, N., Kilanowski, A., Hoffman, N.,

Wedderburn, C., Donald, K., Kobor, M., Zar, H., Stein, D., & Hüls, A. (2022). DNA

methylation as a potential mediator of the association between indoor air pollution and

neurodevelopmental delay in a South African birth cohort. In: Research Square.

Fincham, D. S., Altes, L. K., Stein, D. J., & Seedat, S. (2009). Posttraumatic stress disorder

symptoms in adolescents: risk factors versus resilience moderation. *Compr Psychiatry*,

50(3), 193-199. https://doi.org/10.1016/j.comppsych.2008.09.001

Fleckman, J. M., Drury, S. S., Taylor, C. A., & Theall, K. P. (2016). Role of Direct and Indirect

Violence Exposure on Externalizing Behavior in Children. *Journal of Urban Health*,

93(3), 479-492. https://doi.org/10.1007/s11524-016-0052-y

Frankenhuis, W. E., & Fraley, R. C. (2017). What Do Evolutionary Models Teach Us About

Sensitive Periods in Psychological Development? *European Psychologist*, *22*(3), 141-

150. https://doi.org/10.1027/1016-9040/a000265

Gard, A. M., Maxwell, A. M., Shaw, D. S., Mitchell, C., Brooks☐Gunn, J., McLanahan, S. S.,

Forbes, E. E., Monk, C. S., & Hyde, L. W. (2021). Beyond family☐level adversities:

Exploring the developmental timing of neighborhood disadvantage effects on the brain.

Dev Sci, *24*(1), e12985.

Georgieff, M. K., Brunette, K. E., & Tran, P. V. (2015). Early life nutrition and neural plasticity.

Dev Psychopathol, *27*(2), 411-423. https://doi.org/10.1017/s0954579415000061

GlobalData. (2022). *Median Household Income in South Africa (2015 - 2021, Purchasing Power*

Parity in Current International Dollars). Retrieved March 10 from

https://www.globaldata.com/data-insights/macroeconomic/median-household-income-in-

south-africa-2091331/

Goodman, R. (2001). Psychometric properties of the strengths and difficulties questionnaire. *J*

Am Acad Child Adolesc Psychiatry, *40*(11), 1337-1345.

https://doi.org/10.1097/00004583-200111000-00015

Goodman, S. H., Rouse, M. H., Connell, A. M., Broth, M. R., Hall, C. M., & Heyward, D.

(2011). Maternal depression and child psychopathology: A meta-analytic review. *Clinical*

Child and Family Psychology Review, *14*, 1-27.

Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslvasky, A.

M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the

national comorbidity survey replication I: associations with first onset of DSM-IV

disorders. *Archives of General Psychiatry*, *67*(2), 113-123.

Group, W. A. W. (2002). The Alcohol, Smoking and Substance Involvement Screening Test

(ASSIST): development, reliability and feasibility. *Addiction*, *97*(9), 1183-1194.

https://doi.org/https://doi.org/10.1046/j.1360-0443.2002.00185.x

Guidi, J., Lucente, M., Sonino, N., & Fava, G. A. (2020). Allostatic load and its impact on health:

a systematic review. *Psychotherapy and Psychosomatics*, *90*(1), 11-27.

Harding, T. W., de Arango, M. V., Baltazar, J., Climent, C. E., Ibrahim, H. H., Ladrido-Ignacio,

L., Murthy, R. S., & Wig, N. N. (1980). Mental disorders in primary health care: a study

of their frequency and diagnosis in four developing countries. *Psychol Med*, *10*(2), 231-

241. https://doi.org/10.1017/s0033291700043993

Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., McLeod, L.,

Delacqua, G., Delacqua, F., Kirby, J., & Duda, S. N. (2019). The REDCap consortium:

Building an international community of software platform partners. *J Biomed Inform*, *95*,

103208. https://doi.org/10.1016/j.jbi.2019.103208

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research

electronic data capture (REDCap)--a metadata-driven methodology and workflow

process for providing translational research informatics support. *J Biomed Inform*, *42*(2),

377-381. https://doi.org/10.1016/j.jbi.2008.08.010

Henrich, J., Heine, S. J., & Norenzayan, A. (2010). The weirdest people in the world? *Behavioral

and Brain Sciences*, *33*(2-3), 61-83. https://doi.org/10.1017/S0140525X0999152X

Herman, A. A., Stein, D. J., Seedat, S., Heeringa, S. G., Moomal, H., & Williams, D. R. (2009).

The South African Stress and Health (SASH) study: 12-month and lifetime prevalence of

common mental disorders. *S Afr Med J*, *99*(5 Pt 2), 339-344.

Hodel, A. S. (2018). Rapid Infant Prefrontal Cortex Development and Sensitivity to Early

Environmental Experience. *Dev Rev*, *48*, 113-144.

https://doi.org/10.1016/j.dr.2018.02.003

Hoffmann, E. V., Duarte, C. S., Matsuzaka, C. T., Milani, A. C. C., Fossaluza, V., Mello, A. F., &

Mello, M. F. (2022). The positive impact of an intervention for maternal depression on

child emotional and behavioral symptoms in a low-resource setting. *Brazilian Journal of*

Psychiatry, *44*(6), 590-601.

Hogye, S. I., Lucassen, N., Jansen, P. W., Schuurmans, I. K., & Keizer, R. (2022). Cumulative

Risk and Internalizing and Externalizing Problems in Early Childhood: Compensatory

and Buffering Roles of Family Functioning and Family Regularity. *Adversity and*

Resilience Science, *3*(2), 149-167. https://doi.org/10.1007/s42844-022-00056-y

Holt, S., Buckley, H., & Whelan, S. (2008). The impact of exposure to domestic violence on

children and young people: a review of the literature. *Child Abuse & Neglect*, *32*(8), 797-

810. https://doi.org/10.1016/j.chiabu.2008.02.004

Hyde, L. W., Gard, A. M., Tomlinson, R. C., Suarez, G. L., & Westerman, H. B. (2022). Parents,

neighborhoods, and the developing brain. *Child Development Perspectives*, *16*(3), 148-

156.

Jewkes, R. (2002). Intimate partner violence: causes and prevention. *Lancet*, *359*(9315), 1423-

1429. https://doi.org/10.1016/S0140-6736(02)08357-5

Juen, F., Hecker, T., Hermenau, K., Teicher, M. H., Mikinga, G., Nkuba, M., Masath, F. B., &

Schalinski, I. (2024). Child maltreatment in a high adversity context: Associations of age,

type and timing of exposure with psychopathology in middle childhood. *Child Abuse*

Negl, *157*, 107060. https://doi.org/10.1016/j.chiabu.2024.107060

Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A.

M., Aguilar-Gaxiola, S., Alhamzawi, A. O., Alonso, J., Angermeyer, M., Benjet, C.,

Bromet, E., Chatterji, S., de Girolamo, G., Demyttenaere, K., Fayyad, J., Florescu, S.,

Gal, G., Gureje, O., . . . Williams, D. R. (2010). Childhood adversities and adult

psychopathology in the WHO World Mental Health Surveys. *The British Journal of*

Psychiatry, *197*(5), 378-385. https://doi.org/10.1192/bjp.bp.110.080499

Lovejoy, M. C., Graczyk, P. A., O'Hare, E., & Neuman, G. (2000). Maternal depression and

parenting behavior: A meta-analytic review. *Clinical Psychology Review*, *20*(5), 561-592.

Luby, J. L., Baram, T. Z., Rogers, C. E., & Barch, D. M. (2020). Neurodevelopmental

Optimization after Early-Life Adversity: Cross-Species Studies to Elucidate Sensitive

Periods and Brain Mechanisms to Inform Early Intervention. *Trends Neurosci*, *43*(10),

744-751. https://doi.org/10.1016/j.tins.2020.08.001

Mäntymaa, M., Puura, K., Luoma, I., Latva, R., Salmelin, R. K., & Tamminen, T. (2012).

Predicting Internalizing and Externalizing Problems at Five Years by Child and Parental

Factors in Infancy and Toddlerhood. *Child Psychiatry & Human Development*, *43*(2),

153-170. https://doi.org/10.1007/s10578-011-0255-0

Martin, L., Revington, N., & Seedat, S. (2013). The 39-item Child Exposure to Community

Violence (CECV) scale: exploratory factor analysis and relationship to PTSD

symptomatology in trauma-exposed children and adolescents. *Int J Behav Med*, *20*(4),

599-608. https://doi.org/10.1007/s12529-012-9269-7

McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R.

C. (2012). Childhood adversities and first onset of psychiatric disorders in a national

sample of US adolescents. *Archives of General Psychiatry*, *69*(11), 1151-1160.

https://jamanetwork.com/journals/jamapsychiatry/articlepdf/1389368/yoa110082_1151_1

160.pdf

Moss, K. M., Loxton, D., & Mishra, G. D. (2023). Does Timing Matter? Associations Between

Intimate Partner Violence Across the Early Life Course and Internalizing and

Externalizing Behavior in Children. *J Interpers Violence*, *38*(19-20), 10566-10587.

https://doi.org/10.1177/08862605231174505

Moyakhe, L., Dalvie, S., Lake, M., Huels, A., Donald, K., Hoffman, N., Zar, H., Stein, D., &

Koen, N. (2023). T64. GESTATIONAL EPIGENETIC AGE DEVIATION AND CHILD

DEVELOPMENTAL AND MENTAL HEALTH OUTCOMES IN A SOUTH AFRICAN

BIRTH COHORT STUDY. *European Neuropsychopharmacology*, *75*, S196.

https://doi.org/https://doi.org/10.1016/j.euroneuro.2023.08.349

Myer, L., Stein, D. J., Grimsrud, A., Seedat, S., & Williams, D. R. (2008). Social determinants of

psychological distress in a nationally-representative sample of South African adults. *Soc*

Sci Med, *66*(8), 1828-1840. https://doi.org/10.1016/j.socscimed.2008.01.025

Network, G. B. o. D. C. (2022). Global, regional, and national burden of 12 mental disorders in

204 countries and territories, 1990–2019: a systematic analysis for the Global

Burden of Disease Study 2019. *The Lancet Psychiatry*, *9*(2), 137-150.

https://doi.org/10.1016/S2215-0366(21)00395-3

Ringoot, A. P., Tiemeier, H., Jaddoe, V. W., So, P., Hofman, A., Verhulst, F. C., & Jansen, P. W.

(2015). Parental depression and child well-being: young children's self-reports helped

addressing biases in parent reports. *Journal of Clinical Epidemiology*, *68*(8), 928-938.

https://doi.org/10.1016/j.jclinepi.2015.03.009

Rogosch, F. A., Dackis, M. N., & Cicchetti, D. (2011). Child maltreatment and allostatic load:

consequences for physical and mental health in children from low-income families. *Dev

Psychopathol*, *23*(4), 1107-1124. https://doi.org/10.1017/s0954579411000587

Rudd, K. L., Cheng, S. S., Cordeiro, A., Coccia, M., Karr, C. J., LeWinn, K. Z., Mason, W. A.,

Trasande, L., Nguyen, R. H. N., Sathyanarayana, S., Swan, S. H., Barrett, E. S., & Bush,

N. R. (2022). Associations Between Maternal Stressful Life Events and Perceived

Distress during Pregnancy and Child Mental Health at Age 4. *Res Child Adolesc

Psychopathol*, *50*(8), 977-986. https://doi.org/10.1007/s10802-022-00911-7

Schaefer, J. D., Cheng, T. W., & Dunn, E. C. (2022). Sensitive periods in development and risk

for psychiatric disorders and related endpoints: A systematic review of child maltreatment

findings. *Lancet Psychiatry*, *9*(12), 978-991.

Shamu, S., Abrahams, N., Temmerman, M., Musekiwa, A., & Zarowsky, C. (2011). A systematic

review of African studies on intimate partner violence against pregnant women:

prevalence and risk factors. *PLoS One*, *6*(3), e17591.

https://doi.org/10.1371/journal.pone.0017591

Shanahan, L., Copeland, W. E., Costello, E. J., & Angold, A. (2011). Child-, adolescent- and

young adult-onset depressions: differential risk factors in development? *Psychological

Medicine*, *41*(11), 2265-2274. https://doi.org/10.1017/s0033291711000675

Shankar, P., Chung, R., & Frank, D. A. (2017). Association of Food Insecurity with Children's

Behavioral, Emotional, and Academic Outcomes: A Systematic Review. *Journal of*

Developmental & Behavioral Pediatrics, *38*(2), 135-150.

https://doi.org/10.1097/dbp.0000000000000383

Slopen, N., Meyer, C., & Williams, D. R. (2018). 75Cumulative Stress and Health. In C. D. Ryff

& R. F. Krueger (Eds.), *The Oxford Handbook of Integrative Health Science* (pp. 0).

Oxford University Press. https://doi.org/10.1093/oxfordhb/9780190676384.013.5

Smith, A. D., Hardy, R., Heron, J., Joinson, C. J., Lawlor, D. A., Macdonald-Wallis, C., &

Tilling, K. (2016). A structured approach to hypotheses involving continuous exposures

over the life course. *International Journal of Epidemiology*, *45*(4), 1271-1279.

https://doi.org/10.1093/ije/dyw164

Smith, A. D., Heron, J., Mishra, G., Gilthorpe, M. S., Ben-Shlomo, Y., & Tilling, K. (2015).

Model Selection of the Effect of Binary Exposures over the Life Course. *Epidemiology*,

26(5), 719-726. https://doi.org/10.1097/EDE.0000000000000348

Smith, B. J., Smith, A. D. A. C., & Dunn, E. C. (2021). Statistical Modeling of Sensitive Period

Effects Using the Structured Life Course Modeling Approach (SLCMA). In S. L.

Andersen (Ed.), *Sensitive Periods of Brain Development and Preventive Interventions*

(pp. 215–234). Springer. https://doi.org/https://doi.org/10.1007/7854_2021_280

Stein, D. J., Koen, N., Donald, K. A., Adnams, C. M., Koopowitz, S., Lund, C., Marais, A.,

Myers, B., Roos, A., Sorsdahl, K., Stern, M., Tomlinson, M., van der Westhuizen, C.,

Vythilingum, B., Myer, L., Barnett, W., Brittain, K., & Zar, H. J. (2015). Investigating the

psychosocial determinants of child health in Africa: The Drakenstein Child Health Study.

J Neurosci Methods, *252*, 27-35. https://doi.org/10.1016/j.jneumeth.2015.03.016

Stirling, K. J., Toumbourou, J. W., & Rowland, B. C. (2015). Community factors influencing

child and adolescent depression: A systematic review and meta-analysis. *Australian &*

New Zealand Journal of Psychiatry, *49*, 869 - 886.

Suveg, C., Shaffer, A., Morelen, D., & Thomassin, K. (2011). Links between maternal and child

psychopathology symptoms: mediation through child emotion regulation and moderation

through maternal behavior. *Child Psychiatry Hum Dev*, *42*(5), 507-520.

https://doi.org/10.1007/s10578-011-0223-8

van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate Imputation by Chained

Equations in R. *Journal of Statistical Software*, *45*, urn:issn:1548-7660.

Whittenburg, P. N., Stern, J. A., Brett, B. E., Straske, M. D., & Cassidy, J. (2023). Maternal

depressive symptoms and child behavior problems: Attachment security as a protective

factor. *Dev Psychopathol*, *35*(2), 678-688. https://doi.org/10.1017/s0954579421001802

World Health Organization. (2017). *Depression and Other Common Mental Disorders: Global*

Health Estimates.

Yoon, Y. (2022). Time Dimension of Childhood Adversities and Externalizing Behavior Among

Children of Teen Mothers: Sensitive Period Hypothesis vs. Accumulation Hypothesis.

Child Maltreatment, *27*(3), 389-399. https://doi.org/10.1177/1077559520984249

Zar, H. J., Barnett, W., Myer, L., Stein, D. J., & Nicol, M. P. (2015). Investigating the early-life

determinants of illness in Africa: the Drakenstein Child Health Study. *Thorax*, *70*(6), 592-

594. https://doi.org/10.1136/thoraxjnl-2014-206242

Table 1 Sample demographics

	Mbekweni (N=479)	TC Newman (N=308)	Overall (N=787)
Child Ethnicity[1]			
Black African	474 (99.0%)	4 (1.3%)	478 (60.6%)
Coloured	4 (0.8%)	304 (98.7%)	308 (39.2%)
Missing	1 (0.2%)	0 (0%)	1 (0.1%)
Sex			
Female	252 (52.6%)	144 (46.8%)	396 (50.3%)
Male	227 (47.4%)	164 (53.2%)	391 (49.7%)
Marital Status			
Single	302 (63.0%)	170 (55.2%)	472 (60.0%)
Married/Cohabiting	176 (36.7%)	138 (44.8%)	314 (39.9%)
Missing	1 (0.2%)	0 (0%)	1 (0.1%)
Highest Maternal Education at Birth			
Lower than Secondary	302 (63.0%)	193 (62.7%)	495 (62.9%)
Secondary or Higher	177 (37.0%)	115 (37.3%)	292 (37.1%)
Maternal HIV at Birth			
Negative	298 (62.2%)	298 (96.8%)	596 (75.7%)
Positive	181 (37.8%)	10 (3.2%)	191 (24.3%)
Number of Previous Pregnancies			
0	155 (32.4%)	114 (37.0%)	269 (34.2%)
1	185 (38.6%)	107 (34.7%)	292 (37.1%)
2	92 (19.2%)	55 (17.9%)	147 (18.7%)
3+	45 (9.4%)	30 (9.7%)	75 (9.5%)
Missing	2 (0.4%)	2 (0.6%)	4 (0.5%)
Maternal Age at Birth (in years)			
Mean (SD)	27.8 (5.75)	26.2 (5.53)	27.2 (5.71)
Median [Min, Max]	27.0 [18.0, 45.0]	25.0 [18.0, 44.0]	26.0 [18.0, 45.0]
ASSIST Tobacco Score[2]			

	Mbekweni (N=479)	TC Newman (N=308)	Overall (N=787)
Mean (SD)	1.13 (5.16)	11.9 (11.4)	5.44 (9.79)
Median [Min, Max]	0 [0, 31.0]	12.0 [0, 31.0]	0 [0, 31.0]
Missing	66 (13.8%)	33 (10.7%)	99 (12.6%)
Asset Sum[3]			
Mean (SD)	6.45 (1.99)	7.24 (1.71)	6.76 (1.92)
Median [Min, Max]	7.00 [2.00, 11.0]	7.00 [2.00, 10.0]	7.00 [2.00, 11.0]
Household Income[4]			
<R1000/m	184 (38.4%)	93 (30.2%)	277 (35.2%)
R1000-5000/m	249 (52.0%)	163 (52.9%)	412 (52.4%)
>R5000/m	46 (9.6%)	52 (16.9%)	98 (12.5%)

[1] In South Africa, the apartheid system divided individuals into different racial groups (e.g., "Black African", "Coloured", "White"). Under democracy these categories continue to be employed, partly for reasons of redress and partly due to South Africans' cultural embrace and celebration of these identities (Dooms & Chutel, 2023). In DCHS, participants self-identify as "Black African" or "Coloured". We are aware that these categories are not widely used internationally and may have painful racist connotations in other cultures. At the same time, we think it is important to honor South African participants' identities and culture by using the language and terminology they self-report. By using these categories, we are not aiming to reify social categories but instead hope to contribute to the study of ongoing health disparities stemming from systemic racism.

[2] A higher ASSIST Tobacco Score indicates greater tobacco usage during the antenatal visit.

[3] Asset sum is determined by the endorsement of 13 items, such as access to electricity, running water, or a motor vehicle, where a higher score indicates higher assets.

[4] The median household income in South Africa (2015) was R11,294 per month.

Table 2. Adversity exposure prevalence across developmental period

Adversity Category	Very Early Childhood (≤ 3)	Early Childhood ($3 \leq 5$)	Middle Childhood ($5 \leq 8$)	Ever Exposed
Maternal Psychopathology	196 (24.9%)	110 (14.0%)	61 (7.8%)	259 (32.9%)
Maternal Adverse Events	461 (58.6%)	313 (39.8%)	184 (23.4%)	605 (76.9%)
Child Food Insecurity	118 (15.0%)	93 (11.8%)	126 (16.0%)	243 (30.9%)
Child Exposure to Community and Domestic Violence*	192 (24.4%)		346 (44.0%)	415 (52.7%)
Maternal Intimate Partner Violence	303 (38.5%)	116 (14.7%)	61 (7.8%)	363 (46.1)
Maternal Substance Abuse	262 (33.3%)	193 (24.5%)	124 (15.8%)	362 (46.0%)

*Note. Child Exposure to Community and Domestic Violence developmental periods are first five years (age ≤ 5 years) and middle childhood (age $5 \leq 8$ years)

Table 3. Life-course hypotheses selected by SLCMA for each childhood adversity predicting childhood psychopathology symptoms measured via the CBCL

Internalizing Symptoms					
Adversity Category	**Partial R-squared %**	**Model Selected**	**Coefficient**	**P-value**	**CI (95%)**
Maternal Psychopathology	7.0	Middle Childhood	6.27	**.001**	2.69 - 11.98
		Recency	.20	.2	-.30 - .45
Maternal Adverse Event	2.3	Accumulation	1.51	**<.001**	.84 - 2.15
Child Food Insecurity	2.0	Early Childhood	2.8	.06	-.85 - 8.07
		Recency	.16	.36	-.52 - .41
Child Exposure to Community and Domestic Violence	1.6	Accumulation	1.61	**.001**	.69 - 2.44
Maternal IPV	.6	Middle Childhood	1.22	.61	-16.3 - 7.58
		Recency	.13	.30	-.57 - 1.07
Maternal Substance Abuse	.6	Middle Childhood	.95	.48	-8.34 - 6.94
		Recency	.09	.39	-.56 - .70
Externalizing Symptoms					
Adversity Category	**Partial R-squared %**	**Model Selected**	**Coefficient**	**P-value**	**CI (95%)**
Maternal Psychopathology	5.1	Middle Childhood	6.38	**.004**	1.79 - 13.12
		Recency	.21	.23	-.38 - .51
Maternal Adverse Event	2.6	Recency	.37	**<.001**	.22 - .51
Child Food Insecurity	1.4	Early Childhood	4.30	**.002**	1.58 - 6.60
Child Exposure to Community and Domestic Violence	1.5	Middle Childhood	.93	.53	-7.17 - 3.80
		Accumulation	1.28	.12	-1.18 - 5.65
Maternal IPV	.4	Accumulation	1.11	.12	-.87 - 2.1
Maternal Substance Abuse	.4	Accumulation	.96	.13	-.08 - 1.77

Note. The table presents results from 12 separate regression models, one for each adversity category and symptom outcome. All regression models adjusted for covariates. Coefficients represent unstandardized effect estimates. Bolded values indicate p < .05.

Figure 1. Average tetrachoric correlations between periods and adversities

Figure 1. Average tetrachoric correlations between periods and adversities. **A)** average tetrachoric correlation within each adversity category across developmental periods. **B)** depicts the average tetrachoric correlation of each adversity with all other adversity categories. IPV = intimate partner violence.

Figure 2. Associations of exposure to each type of adversity with CBCL-measured psychopathology symptoms. Bars display variance explained (R^2) by each adversity. Values displayed above each bar depict unstandardized effect estimates. Results were obtained from ever-exposed – i.e., irrespective of life-course hypothesis selection – multiple regression analyses, adjusted for covariates. IPV = intimate partner violence. CBCL = Child Behavior Checklist.

Internalizing Symptoms

A)
Maternal Psychopathology
Middle Childhood + Recency

B)
Maternal Adverse Event
Accumulation

C)
Child Exposure to Violence
Accumulation

Externalizing Symptoms

D)
Maternal Psychopathology
Middle Childhood + Recency

E)
Maternal Adverse Event
Recency

F)
Child Food Insecurity
Early Childhood

Figure 3. Effects of adversity exposure during each developmental period, based on the selected SLCMA life-course hypotheses via the CBCL. Figures illustrate the weighted effect of exposure during each developmental period based on the life-course hypothesis/es selected for each adversity. The y-axis was kept consistent across all figures, including negative values. *Internalizing Symptoms.* Panel A: For maternal psychopathology, the effect of exposure increases with each developmental period and there is an additional increased effect for exposures in middle childhood (age 5 ≤ 8 years). Panel B: For maternal adverse events, the effect of exposure is the same in each developmental period. Panel C: For child exposure to domestic and community violence, the effect of exposure is the same in each developmental period. *Externalizing Symptoms.* Panel D: For maternal psychopathology, the effect of exposure increases with each developmental period and there is an additional increased effect for exposures in middle childhood (age 5 ≤ 8 years). Panel E: For maternal adverse events, the effect of exposure increases with each developmental period. Panel F: For child food insecurity, exposure primarily had an effect if it occurred during the early childhood developmental period (age 3 ≤ 5 years). SLCMA = Structure Life-course Modeling Approach. CBCL = Child Behavior Checklist.