Efficacy of a parent-child program for 3- to 6-year-old children with stuttering: a

retrospective controlled wait-list group pilot study

Nicole E. Neef[a], Imke Niemann[b], Anna Merkel [b], Kristina Anders[b], Katja Hente[b], Johanna

Margarete Joisten[a], Alexander Wolff von Gudenberg[b], Christian H. Riedel[a]

[a]Department of Diagnostic and Interventional Neuroradiology, University Medical Center Göttingen, Georg August

University, Göttingen, Robert-Koch-Straße 40, 37075 Göttingen, Germany; [b]KST Institut GmbH, Feriendorfstraße 1,

34308 Bad Emstal, Germany; Corresponding author: Nicole Neef, nicole.neef@med.uni-goettingen.de

10
11
12
13

Abstract
We evaluated the efficacy of Frankini, a 12-month early parent-child intervention that combines
online parent counseling with hybrid speech restructuring to reduce stuttering severity and
promote fluency-supportive interaction.
This retrospective, nonrandomized pilot trial included cases enrolled between September 2019
and November 2023. For analysis, only participants who completed Module 1 (indirect parental
training) and Module 2 (the first hybrid speech restructuring module) were included. A total of 51
cases met all inclusion criteria, and 30 of these completed all three modules.
To simulate a wait-list-controlled design, eligible participants were divided into early and delayed
groups using median split. The early group completed Module 2 by month 9, while the delayed
group completed it by month 12. Groups were matched on key characteristics and differed only
in the timing of the first direct intervention. Blinded raters assessed stuttering severity.
Primary outcomes included the Stuttering Severity Index, parental severity rating, and a 10-item
parent report. At 9 months, the early group showed reduced stuttering severity, while the
delayed group showed no change (mean difference = -8.33 95%CI [-12.98, -3.68], p < 0.001,
with d = -1.14). By 12 months, both groups improved, and group difference were no longer
significant (mean difference = -3.37 95%CI [-8.23, 1.50], p = 0.168 and d = -0.48). Parental
ratings mirrored these outcomes showing consistent improvement after each module.
Early intervention significantly improved fluency and parents' confidence, supporting the value
of initiating treatment before age 6; however, follow-up is needed to assess long-term effects.
Trial Registration: DRKS00034731.

Introduction

Approximately 5% of children experience stuttering (Bloodstein et al., 2021). Overt symptoms include sound and syllable repetitions, sound prolongations, and speech blocks — involuntary disruptions that the speaker perceives as a loss of control — often resulting in struggle, compensatory behaviors, and a negative impact on life (Tichenor & Yaruss, 2021). Stuttering typically begins in early childhood, with an onset ranging from 2 to 5 years and a mean age of 36 months (Yairi & Ambrose, 2013). Spontaneous remission occurs in 75% of cases within the first 18 months (Reilly et al., 2009; Yairi & Ambrose, 2013), however no marker currently predicts whether stuttering will persist. Despite a high remission rate, 1% of the population continues to stutter throughout life (Yairi & Ambrose, 2013). A recent Cochrane review identified early treatment as the most effective treatment (Sjøstrand et al., 2021), and because of the high neuroplasticity in young children (Neef & Chang, 2024), early interventions are highly desirable. Treatment approaches for preschoolers who stutter are recommended in Germany (Katrin Neumann et al., 2017), however, referrals for therapy peak late, at 6 years of age (Sommer et al., 2021), and efficacy studies are rare (Brignell et al., 2021; Sjøstrand et al., 2021), with only one conducted in German preschoolers (Lattermann et al., 2008).

Worldwide, there are four treatment approaches for preschoolers who stutter that have been shown to reduce stuttering and improve communication attitudes in young children: The Lidcombe Program (Harris et al., 2002; Jones et al., 2005; Lattermann et al., 2008; Lewis et al., 2008; Onslow et al., 2003), the Palin Parent-Child Interaction therapy (Millard et al., 2009, 2018; Preston et al., 2022), the RESTART Demands and Capacities Model (Sonneville-Koedoot et al., 2015), and the combined fluency rules program (FRP) with parent-child interaction training (Bafrooei et al., 2025). The four approaches are broadly comparable in effectiveness but differ in methodology and focus, offering flexibility for tailoring treatment to the needs of the child and family. The Lidcombe Program is a direct therapy that explicitly focuses on reducing stuttering by reinforcing fluent speech and correcting stuttered speech through verbal contingencies.

Parents deliver therapy in everyday speaking situations guided by a trained clinician (Onslow et

al., 1994). Palin PCI and RESTART-DCM are mainly indirect therapies. Palin-PCI (Kelman &

Nicholas, 2019) focuses on enhancing parent-child interaction and reducing environmental

demands to support fluency, using video analysis, tailored feedback, and collaborative goal-

setting with parents. If the child continues to show concern about their stuttering, therapy may

be extended to include more direct strategies, such as increasing the child's understanding of

speech and communication and helping them develop personalized cognitive strategies for

managing stuttering (Franken et al., 2022). RESTART-DCM (Franken & Laroes, 2021) aims to

balance speaking demands with a child's speech capacities by reducing environmental pressure

with active parental involvement. The combined FRP with parent-child interaction training is a

flexible stuttering intervention that promotes speech control through self-monitoring,

individualized feedback, and non-verbal cues targeting both disfluencies and associated

behaviors together with parent education and coaching to support fluency-enhancing strategies

(Bafrooei et al., 2025). All four approaches require active parental involvement, but the intensity

and nature of the role vary. Lidcombe involves structured verbal feedback from parents; and

Palin PCI and RESTART-DCM focus on modifying interaction styles and environments rather

than targeting stuttering directly. The most recent trial combined both modifying stuttering and

educating parents about stuttering and parent-child interactions.

Thus, therapy for preschool children who stutter generally follows one of two

complementary approaches: parent-focused or child-focused. Parent-focused (indirect) therapy

aims to equip caregivers with strategies to foster a supportive, low-pressure communication

environment. Child-focused (direct) therapy, on the other hand, involves teaching children

techniques to manage and modify their stuttering directly. Each approach offers unique benefits

and presents certain considerations. While both depend on active parental involvement, tailoring

the level of engagement to individual family circumstances can enhance feasibility and success.

Direct approaches introduce children to practical fluency-enhancing tools, such as easy onsets

and slowed speech, and can empower them to develop awareness and resilience around their

speech patterns. However, it's important to match the intervention to a child's developmental

readiness, as younger children may need more support in integrating these strategies. For

some, focusing explicitly on speech may increase self-monitoring; with careful guidance, this

can be managed in a way that builds confidence rather than anxiety. Crucially, the field

continues to refine its understanding of what works best for whom. While promising outcomes

have been observed with both models (Brignell et al., 2021; K. Neumann et al., 2016; Katrin

Neumann et al., 2017; Sjøstrand et al., 2021), further research is needed to explore the long-

term impact of these strategies, individually and in combination. The diversity among children,

differences in temperament, stuttering severity, family dynamics, and cultural background,

underscores the value of flexible, individualized therapy planning (Millard et al., 2018). Finally,

the evolving definitions of therapeutic success, ranging from fluency gains to increased

communicative confidence and quality of life, reflect the broader, more holistic view emerging in

the field (Brundage et al., 2021). This diversity of goals enriches clinical practice and

encourages ongoing dialogue about best practices in early stuttering intervention.

Here, we assessed the efficacy of Frankini, an early intervention program designed for

parents and preschoolers, that combines online parent counseling and training with speech

restructuring techniques delivered both in-person and via telehealth. The program aimed to

reduce stuttering severity while fostering interaction styles that promote fluent communication.

By integrating both indirect and direct therapeutic elements, Frankini seeks to address the

multifaceted nature of early childhood stuttering; that is supporting the child's speech

development while empowering caregivers to create a communicatively supportive home

environment. The hybrid delivery format enhances accessibility for families who may face

logistical or geographical barriers to in-person services, and allows for continuity of care across

settings. This approach reflects a growing recognition of the importance of early, comprehensive

intervention that aligns with both the child's speech capacities and the family's resources and

dynamics. The program is an adaptation of the Kasseler Stuttering Therapy, a computer-

assisted fluency-shaping program developed for adults who stutter (Euler et al., 2009; Wolff-

von-Gudenberg & Euler, 2000). Previously, we introduced an adoption of this program for

school-aged children (FRANKA, (Euler et al., 2021). The new program, Frankini, combines

indirect and direct treatment components across three modules spread over the course of 12

months (Figure 1A). Since spontaneous remission is still possible between the ages of three

and six, and the course of stuttering often fluctuates, parents and therapists decide together

after each module whether to continue, pause, or end treatment. After Module 1, key indicators

for initiating intensive in-person therapy included the level of distress experienced by the child

and parents, the severity of stuttering, and the prognosis for its progression.

Randomized controlled trials of behavioral interventions in neurodevelopmental

disorders are notoriously difficult to conduct. Recruiting and retaining families is difficult because

of the time-intensive nature of participation, especially for those already burdened by the

disorder. The required long-term commitment often leads to high dropout rates, affecting study

validity. Withholding treatment from the control groups raises ethical concerns, particularly when

early intervention is crucial. Additionally, blinding is challenging because parents and clinicians

often know if an intervention is active. We present an alternative approach by simulating a

waitlist group design through the retrospective allocation of participants into early and delayed

groups. Waiting times after registration varied, as parents were assigned to courses based on

availability and compatibility with the family's schedule. At nine months post-baseline, the early

group had completed a direct treatment module, while the delayed group had not yet started the

same module, and an interaction between group and time would indicate an effect of the

intervention. Six recordings of speech samples throughout the program allowed for the

comparison of objectively quantified stuttering severity at this relative point in time. This study

design accounted for maturation processes and spontaneous remission, ensuring that the

likelihood of these influences was equally distributed across both groups. Additionally, raters of

the stuttering severity assessment were blinded to the time point of recording, reducing bias

related to expectations of progress over time.

To our knowledge, this is the first study to apply a retrospective simulated wait-list design.

In contrast to a prospective wait-list control group, where participants are knowingly delayed in

receiving treatment, this design uses pre-existing clinical data to simulate a control group, with

all outcome measures collected before group assignment. This approach offers distinct ethical

and practical advantages, particularly in early childhood stuttering intervention, where timely

access to treatment is critical and delays may contribute to emotional strain and frustration for

both children and their families. By avoiding intentional withholding of therapy, the design

reduces the risk of demotivation, dropout, and dissatisfaction that can occur in families placed

on a traditional wait list. At the same time, the retrospective nature of the design entails certain

methodological limitations, such as the absence of randomization and the potential for selection

bias. Nonetheless, it provides a feasible and ethically responsible alternative for evaluating

intervention effects in real-world clinical settings where randomized controlled trials may be

difficult to implement.

To summarize, the present study evaluated the efficacy of a novel hybrid intervention

program for preschool children who stutter. The program integrates telehealth coaching of

parents, in-person online and in-person individual and group sessions to support both children

and their families. Primary outcomes included objectively measured stuttering severity, parent-

reported severity ratings, and coping measures assessed through online surveys. These

surveys captured dimensions such as confidence in communication, emotional regulation, and

personal feelings and concerns related to stuttering.

Methods

This retrospective study was approved by the Ethical Review Board of the University

Medical Center Göttingen and registered at the UMG study center (ID 2024-03288). The study

protocol was registered in the German Clinical Trials Register

https://www.drks.de/DRKS00034731.

Participants Flow

The trial was a retrospective, two-armed, controlled, monocentric trial with a prospective

simulated wait-list design. In *the retrospective phase,* eligible cases were identified from all 79

families who entered the program between September 2019 and November 2023. Families

contacted us through various channels, including referrals from pediatricians and speech-

language therapists, or through independent online searches. No formal exclusion criteria were

applied at intake, other than age (3–6 years) and clinical indicators consistent with

developmental stuttering. Eligibility required parental consent for repeated clinical and research

assessments, video calls, daily video recordings, and storage and processing of

pseudonymized data for treatment evaluation. No systematic sampling strategy was applied.

Inclusion criteria required participation in Module 1 (indirect parent training) and Module 2 (first

speech restructuring module). Families who paused after Module 1 or began directly with

Module 2 were excluded. Additionally, four families discontinued without mutual agreement, one

family transitioned to another therapy, and one child was withdrawn due to illness. As shown in

the participant flow chart (Figure 2), 51 cases met the eligibility criteria.

In *the prospective phase*, the 51 eligible cases identified during the retrospective phase

were assigned to early or delayed intervention groups based on naturally occurring variation in

the time between their initial clinical assessment (T0) and the completion of Module 2 (T2). This

variation in treatment initiation allowed for a simulation of a waitlist-controlled design.

Participants were divided into two groups using a median split of the time interval between T0

and T2 (Mdn = 6.3 months). The cutoff of 6.3 months was chosen because it represented the

median waiting time among the 42 participating families. This approach ensured an even split

between the shorter and longer waiting time groups and allowed for a balanced comparison The

early group completed Module 2 on average within nine months, while the delayed group

reached the same point after approximately twelve months (see Table 1, Figure 1B).

To reduce potential confounds, group matching was performed on key variables: sex,

age, and age of stuttering onset. During this process, three participants had to be excluded due

to matching limitations, resulting in a final sample of 48 children, 24 in each group, who were

included in the study analyses. SSI-4 ratings were then conducted by three blinded raters for

the final matched sample. During this phase, some speech samples were found to be ineligible

for analysis due to non-compliance. At the same time, reasons for missing samples were

investigated for the first time, a documentation oversight on our part that could not be corrected

retrospectively. As a result, valid SSI-4 data were obtained for 42 children (see Participant Flow

Chart for details).

Participants

Of the 42 children included in the analysis, 23 were assigned to the early group and 19

to the delayed group. Participants ranged in age from 3 to 5 years and included 29 boys and 13

girls. The mother tongue of 39 children was German. Among the 10 bilingual children, three had

a different primary language, Albanian, English, or Russian, with German as their second L1

language. The remaining seven were native German speakers with a second L1 language,

including English, Polish, Persian, or Russian.

A family history of stuttering was reported for 26 children, while 14 had no known family

history; for 2 children, this information was unavailable. Twenty children had not received prior

treatment for stuttering, while 22 had previously undergone some form of therapy. Information

on family history, bilingualism, and prior treatment is summarized in Supplementary Table 1.

Individual demographic data are presented in Supplementary Table 8.

At baseline (T0), stuttering severity as assessed by the SSI-4 was categorized as very

mild in 10 children, mild in 15, moderate in 16, and severe in one case.

Intervention

The Frankini program is a structured, year-long parent-child intervention for preschool-

aged children who stutter (ages 3 - 6). It comprises three consecutive modules combining

indirect and direct therapeutic elements, with parents actively involved throughout (Figure 1A). A

description of the program has been published in a German clinical journal (Anders et al., 2023).

Below we provide an overview of the intervention.

All therapists participating in this study were experienced members of the Kasseler

Stottertherapie team (Euler et al., 2009, 2014, 2021; Wolff-von-Gudenberg & Euler, 2000) a

specialized center for intensive stuttering treatment. Group sessions are co-led by two

therapists to support consistency and collaborative delivery. As this was a pilot study, formal

fidelity monitoring procedures (e.g., checklists, session recordings, or inter-rater reliability) were

not yet implemented, and establishing fidelity was not a primary focus. However, informal fidelity

was supported through internal supervision, shared structured materials, and the use of a

common online therapy platform. These elements helped promote coherence across sessions

and therapists. A formal treatment manual is currently in development to guide future studies,

including a planned randomized controlled trial with more rigorous fidelity monitoring.

The program aims to promote speech confidence, reduce stuttering severity, and foster

acceptance and openness around stuttering. Rather than enforcing fluency, the focus is on

building communicative agency and early use of supportive strategies.

Module 1 - Parent-Only Online Training: Module 1 was delivered entirely online and

focused on parent counseling and training, without direct child involvement. Through individual

and small-group video sessions, parents received education on stuttering, developed a more

confident and accepting stance, and learned to communicate openly and age-appropriately

about stuttering. Practical strategies included structuring speaking situations, adjusting linguistic

demands, and practicing soft voice onsets, an early fluency-shaping technique, through home-

based exercises. These were practiced by parents alone before being introduced to children in

later modules. Parents also submitted video recordings of interactions with their child, which

were reviewed with therapists.

An integral part of this phase was the flunatic.mini software (Euler et al., 2009), which

provides visual feedback on vocal onset to support the learning of soft voice onsets.

Flunatic.mini is part of an online platform, that also offers written guidance, video

demonstrations, fluency tracking tools, and reflection prompts.

Modules 2 and 3 - In-Person Parent-Child Intervention: Modules 2 and 3 each began

with a four-day in-person phase involving group and individual parent-child sessions. In Module

2, many children encountered peers who stutter for the first time, helping to reduce stigma and

encourage open discussion about stuttering. Children learned to identify and name stuttering

symptoms using visual symbols in a supportive, pressure-free environment. Parents were

present throughout and could step in during exercises if needed.

The core fluency technique, soft voice onset ("feather words"), was introduced through

structured, play-based exercises that increased in linguistic complexity. Children practiced the

technique in both group and individual settings, with reinforcement through flunatic.mini. Post-

intensive phases included biweekly online sessions for individualized feedback.

Therapy was not aimed at achieving consistent use of soft onsets or prompting children

to use them continuously. Instead, parents were guided to select appropriate home activities

and adjust linguistic demands to match their child's capabilities. Most children began practicing

at the word or sentence level in Module 2, with Module 3 focusing on longer utterances and

simple transfer tasks (e.g., ordering ice cream or sending a voice message). Maintaining

speaking enjoyment remained central. Group participation also fostered learning by modeling,

both from therapists and from peers, and helped desensitize children and parents to stuttering

and fluency techniques.

<u>Goals and Decision Points:</u> Therapy goals across modules were tailored to each child's

developmental level and family context. These included reducing stuttering severity, enhancing

communication confidence, increasing self-efficacy in both child and parent, and promoting a

non-stigmatizing, open approach to stuttering. At the end of each module, therapists and

parents jointly decided whether to continue, pause, or conclude therapy, based on observed

progress, current symptom severity, family distress, and the potential for spontaneous recovery.

(Euler et al., 2009)The program courses between September 2019 and November 2023

involved five to eight families per course, with a total of 13 completed courses. The 13 courses

were conducted by eight therapists, including the coauthors I.N., K.A., A.M., and K.H.

Outcomes

The objective assessment of stuttering severity was conducted using the Stuttering

Severity Index (SSI-4, (Riley, 2009), which was conducted at six time points: at the initial

assessment before Module 1, at the end of Module 1, and at the beginning and end of the two

hybrid modules (T0 - T5, Figure 1A). At each time point, a speech sample was recorded during

a video call in which the therapist engaged the child in a conversation. Additionally, at T0, T3,

and T5, parents provided videos of the children in everyday situations. Because speech

samples were not available for every child at each time point, all available samples for a child

were analyzed at each time point. The number of analyzed syllables varied among the children,

with a median of 348 syllables in the early group and 316 syllables in the delayed group at T0

(Table 1). Parent ratings of stuttering severity (PR) and the online questionnaire on parents'

experiences with the child's stuttering (OQ) were conducted before Module 1 and at the end of

each module (T0, T1, T3, and T5; Figure 1A). Parent-reported measures were scored on a

Likert scale.

The surveys assessed parents' subjective perception of their child's stuttering severity as

well as their attitudes and feelings about stuttering. The severity rating involved evaluating the

child's current symptoms on a scale of one (no stuttering) to 10 (very severe). Additionally, five

questions gauged parents' confidence in handling and discussing stuttering (e.g., "I can handle

my child's stuttering calmly."), while another five questions measured their emotional responses

and concerns (e.g., "My child's stuttering worries me."). Responses were recorded on a scale

ranging from one (do not agree at all) to five (completely agree).

Blinded assessment of stuttering severity

K.A. pseudonymized the speech samples and assigned them to three raters, including

the coauthor A.M., all of whom are clinical speech scientists. Initially, T0 samples were assessed

to test for baseline differences between the groups, with none found (Table 1). Subsequently,

the remaining samples (T1–T5) were randomized according to the time point for analysis. Each

rater analyzed all samples from a given child. No rater was involved in the intervention for that

child.

The objective assessment of stuttering severity was conducted using the SSI-4 (Riley,

2009), which involves analyzing the frequency of stuttered syllables, measuring the duration of

the three longest stuttering events, and evaluating physical concomitants such as grimacing and

head and limb movements. SSI-4 provides a total score that encompasses three subscales:

frequency, duration, and physical concomitants.

Statistical methods

To test the efficacy of the parent-child program, we fitted the SSI-4 total scores with a

linear mixed-effects model using maximum likelihood estimation (Supplementary Table 2). The

baseline model included sex and age as fixed-effects variables, and subject and time as

random-effects variables. We tested whether the model fit improved by including (i) group (early

versus delayed treatment group), (ii) time (pre-, 9 months post-T0, 12 months post-T0), and (iii)

group-by-time interaction. Additionally, we set the following orthogonal and non-orthogonal

contrasts: The contrast for the group tested whether the early treatment group had a greater

reduction than the delayed group; contrasts for time tested (i) whether the 9-month condition

showed smaller SSI values than the pre-condition, and (ii) whether the 12-month condition

showed smaller SSI values than the pre-condition. Post hoc unpaired *t*-tests were conducted to

assess group differences at 9 and 12 months after baseline. To assess the effect size, we

calculated Cohen's *d*. Additionally, we conducted one-sample *t*-tests to determine whether the

reduction in stuttering, as reflected by the subjective parent rating score, was significant. These

tests were performed after treatment Module 2 and Module 3.

To test whether the parents' experience with the child's stuttering improved, we

calculated the Wilcoxon signed rank test with continuity correction between baseline scores and

scores at T1, T3, and T5. Additionally, we plotted the difference in scores between these time

points and the baseline to visualize individual changes.

Interrater agreement

We tested interrater agreement among all three raters. Therefore, each rater analyzed

one randomly chosen speech sample from each child. This resulted in a total of 42 speech

samples per rater (19%). Krippendorff's alpha (Hughes, 2021), along with the macro developed

by Hayes for ordinal and interval data (Hayes & Krippendorff, 2007), was used to calculate

interrater agreement for both frequency (interval scale) and SSI total score (ordinal scale).

Additionally, we calculated the intraclass correlation (ICC) with a two-way model, type

agreement, to provide the 95% confidence interval for the ICC population values. Statistical

analysis was performed using R version 4.4.0 using *irr, multcomp, nlme, pastecs, stats,* and*,*

tidyr.

Results

Objective measure

Our analysis showed that treatment led to a significant reduction in stuttering severity, as

reflected in the effect of time, $\chi^2(9) = 22.0$, $p < 0.001$. This reduction differed between the early

and delayed treatment groups, as reflected in a significant group effect, $\chi^2(7) = 5.1$, $p < 0.024$

(Figure 3A). Most importantly, the treatment × group interaction was also significant, $\chi^2(11) =$

10.6, $p = 0.005$, indicating that the extent of improvement in stuttering severity depended on the

timing of treatment.

By 9 months, children in the early treatment group showed a marked reduction in

stuttering with a mean SSI-4 score of 7.6 (95% CI [4.6, 10.5]), a mean change of -8.7 points (95%

CI [-12.2, -5.2]) from baseline. In contrast, the delayed group showed virtually no change during

the same period, with a mean SSI-4 score of 15.9 (95% CI [12.1, 19.6]) and a negligible mean

change of 0.3 (95% CI [-3.1, 3.6]). A post hoc unpaired *t*-test confirmed that the difference in

SSI-4 total scores between the groups at 9 months post-baseline was statistically significant,

with a mean difference of -8.33 (95% CI [-12.98, -3.68]), t(36.3) = -3.63, p < 0.001. This

corresponds to a large effect size of d = -1.14 (95% CI [-1.82, -0.47]).

At the 12 month mark, both groups showed a reduction in stuttering severity. The early

group manifested their improvement, with a mean score of 7.2 (95% CI [3.2, 11.1]), while the

delayed group also improved to a mean score of 10.5 (95% CI [7.4, 13.7]). However, the

difference between the groups at this point in time was no longer statistically significant (t(30.1)

= -1.44, p = 0.168), although the effect size still suggested a moderate advantage for the early

group (d = 0.49, 95% CI [-1.18, 0.21]). The mean change from baseline at 12 months was -10.6

(95% CI [-18.2, -3.3]) for the early group and -5.1 (95% CI [-8.0, -2.2]) for the delayed group.

Supplementary Table 2 lists the details of the statistical analysis of the SSI total scores.

Additional analyses of percent stuttered syllables and SSI-4-subscores for frequency

and duration confirmed these findings (Supplementary Tables 3 to 5). Group-wise mean

changes in SSI-4 total scores from baseline (T0) to the end of Module 2 (T3) and Module 3 (T5)

are reported in Table 2. Figure 3B shows the trajectory of stuttering severity over the course of

the program for all 42 participants, regardless of the group assignment. Importantly, the most

substantial improvements were observed after completion of Module 2, with gains largely

maintained following Module 3 in children who continued the program.

Stuttering status after Module 2 and after Module 3

According to the SSI-4 severity assessment conducted after Module 2, at T3, 6 children

were classified as having no stuttering, 23 as very mild, 5 as mild, and 8 as moderate.

Compared to baseline, at T0, stuttering severity decreased by three grades in 1 case, by two

grades in 10 cases, and by one grade in 15 cases. In 14 cases, severity remained unchanged,

of these, 6 were already classified as very mild. In 2 cases, severity increased by one grade.

Following Module 3, 2 children were classified as having no stuttering, 18 as very mild, 6

as mild, and 2 as moderate. Compared to the end of Module 2 (T3), severity decreased by two

grades in 3 cases and by one grade in 4 cases. In 15 cases, severity remained unchanged, of

these, 11 were already very mild and 1 had been classified as no stuttering. Severity increased

by one grade in 6 cases and by two grades in 1 case.

Finally, 29% of the cases were excluded from Module 3 analysis due to attrition, with 19%

of families, that is 8 cases, choosing not to continue treatment. While some level of attrition bias

is unavoidable given the structure of the program, it is important to note that most of these

children had already demonstrated meaningful progress by the end of Module 2. As shown in

Figure 3B (red lines), stuttering severity improved in all but one of the children excluded at this

stage. Among the eight families who chose to pause after Module 2, 1 child was classified as

having no stuttering, 5 as very mild, 1 as mild, and 1 as moderate. Stuttering severity was

reduced in five children by 2 grades, and in 1 child by 1 grad. It remained at a very mild level in

2 children, and was moderate in 1 child (see Supplementary Table 7). Notably, the two children

who remained in the very mild range still demonstrated improvement, with SSI-4 total scores

decreasing from 9 to 6 and from 4 to 2, respectively. The child with moderate stuttering also

showed measurable progress, with the SSI-4 total score decreasing from 25 to 20 and percent

stuttered syllables reducing from 12% to 8%. Taken together, these data suggest that the

majority of children who discontinued after Module 2, had experienced clinically meaningful

improvements in speech fluency.

Parents' reports

Parents' subjective ratings confirmed the positive impact of treatment on their children's

stuttering. After completing Module 2, parents reported a significant reduction in perceived

stuttering severity, with an average change score of -2.4 (95% CI [-3.2, -1.6], $T(31) = -5.82$, p

< .001). This improvement continued following Module 3, with a mean change of -2.6 (95% CI [-

3.7, -1.5], $T(24)$ = -5.10, p < .001).

Among the eight families who discontinued after Module 2, five provided severity ratings

via the online survey. Of these, three parents rated their child's stuttering severity as 1, and two

rated it as 2, suggesting continued low perceived severity even without completing the final

module.

Online questionary about coping strategies

Parents' were also asked to rate their experience across a set of items assessing how

they coped with and managed their child's stuttering. This questionnaire was administered as

part of a cohort design and was not controlled, as data were collected at only four time points:

baseline (T0), and after each intervention module (T1, T3, T5; see Figure 1A). Figure 3C

presents a heatmap illustrating changes in parental ratings from baseline (T0) to the end of

Module 2 (T3). Each column represents an individual case, with the rightmost column displaying

the average change across all participants.

For the first five items, where higher scores indicate improved coping, positive changes

are visualized using shades of red, with darker reds representing greater improvement. This

visual approach summarizes the pattern of responses across families. For instance, the Item 1,

"*I can handle my child's stuttering calmly,*" showed an increase in scores for 27 families, no

change for 3 families, and a decrease for 3 families. A one sample *t*-test revealed a significant

improvement, $T(32)$= 6.18, p > 0.001, as indicated by three asterisks in the rightmost cell of this

row (also see Supplementary Table 6).

For the remaining items, where lower scores reflect better outcomes, improvements are

represented in shades of blue, with darker blues indicating larger reductions in score. For

example, Item 8, "*I feel uncomfortable talking to others about stuttering,*" showed a decrease in

scores for 12 families, no change for 14 families, and an increase for 6 families. This change

was not statistically significant, $T(32)= -1.38$, $p = 1.0$, as denoted by the absence of asterisks.

Overall, most items demonstrated clear improvement across participants. Statistically

significant changes were observed in all items except for Item 8, as reflected in the asterisk

markers in the average column on the right side of Figure 3C and detailed in Supplementary

Table 6.

At the end of Module 1, significant improvements were observed for Items 1 to 3 and

Item 5 among the positively worded items, and for Items 6 and 10 among the negatively worded

items. These results suggest that early in the intervention, parents began to feel more capable

and experienced some relief in their emotional responses related to their child's stuttering.

Following completion of Module 3, the remaining families reported improvements across nearly

all items. However, no significant changes were observed for three items: Item 4 ("*I feel

confident talking about stuttering with people I trust.*"), Item 8 ("*I feel uncomfortable talking to

others about stuttering*"), and Item 9 ("*When my child reacts emotionally to his/her stuttering or

talks about it, I often feel overwhelmed*"). These results suggest that while the intervention had a

broad impact on parental coping, some aspects, particularly those related to social support and

public disclosure, may require additional focus.

Interrater-reliability

Interrater agreement was found to be almost-perfect, following the interpretation of

Hayes and Krippendorff (Hayes & Krippendorff, 2007). Specifically, Krippendorff's alpha was

0.88 for the total SSI score and 0.92 for stuttering frequency, indicating strong consistency

among raters. These findings were further supported by intraclass correlation coefficients (ICCs),

which also reflected high reliability. The ICC for the total SSI score was 0.88, with a 95% CI

[0.80, 0.93], while the ICC for stuttering frequency was even higher at 0.92 (95% CI [0.86, 0.96]).

Discussion

The newly developed parent-child program significantly reduced stuttering, yielding a large effect size within 9 months post-baseline. Our findings provide the first evidence that child-friendly fluency shaping can be feasibly integrated into a parent-child therapeutic framework and effectively delivered to preschoolers. The program's efficacy appears comparable to outcomes reported in previous randomized controlled trials of stuttering interventions for children aged 3 to 6 years (Harris et al., 2002; Jones et al., 2005; Lattermann et al., 2008; Lewis et al., 2008). The use of a simulated wait-list control helps to mitigate the influence of spontaneous remission, strengthening confidence in treatment effect. Although attrition was relatively high, it is noteworthy that nearly all children whose families paused after Module 2 still demonstrated measurable reductions in stuttering. These key aspects will be discussed in more detail in the sections that follow.

Fluency shaping is a well-established method for treating adults who stutter and seek increased fluency (Euler et al., 2014; Webster, 1980; Wolff-von-Gudenberg & Euler, 2000), and has also shown effectiveness in school-age children (Euler et al., 2021). Given the young age of the participants, the use of fluency shaping may be seen as controversial, underscoring the importance of monitoring its impact on children's attitudes toward their speech and overall well-being. Although we did not directly assess the children's experiences with the speech technique, we gathered parental perceptions at the end of Module 3. Parents were asked to rate, on a scale from 1 to 5, how comfortable they believed their child felt with relaxed phonation and soft speech. Of the families who continued treatment through Module 3, 25 responded via the online survey. One family reported that the child felt very comfortable, 11 reported feeling comfortable, 9 reported partial comfort, 3 reported less comfort, and 1 reported that the child did not feel comfortable at all. While this is a qualitative and self-reported measure, and responses are

limited to a subset of participants who remained in treatment, the responses suggest general

acceptance of the technique among participating families, with some variability in individual

comfort levels. These findings underscore the importance of further research into children's lived

experiences with fluency-shaping methods in early intervention contexts.

The evaluation of a therapeutic approach can be designed in several ways. In our study,

we choose the simulated wait-list design, which allowed us to control for the influence of

spontaneous remission, a particularly relevant factor in early childhood stuttering (Yairi &

Ambrose, 2013). Another common design is to compare a novel or untested intervention with a

treatment that has already demonstrated efficacy (Sonneville-Koedoot et al., 2015). Such

comparisons can reveal whether the new approach is inferior, equivalent, or superior to the

established one. A third design strategy involves comparing two untested interventions, offering

the potential to identify which yields better outcomes (Bafrooei et al., 2025). Given that young

children exhibit high neuroplasticity and that stuttering follows dynamic developmental courses

at this age (Chow et al., 2023; Chow & Chang, 2017), it is especially important to account for

spontaneous remission. Therefore, designs that incorporate a control for natural recovery are

particularly valuable when evaluating early interventions.

The timing and type of intervention are critical considerations in the treatment of

developmental stuttering, particularly with regard to fluency-shaping approaches in preschool-

age and school-age children. Our findings suggest that early intervention may yield more

favorable outcomes than interventions initiated later in childhood. This advantage may stem

from the brain's heightened neuroplasticity during early childhood, which supports more efficient

speech motor learning and adaptation. This neural flexibility, combined with a generally lower

emotional burden for both children and their parents at this age, may contribute to the more

robust outcomes observed. While these interpretations align with developmental neuroscience

(Chang et al., 2025; Chow et al., 2023) , they warrant further investigation using neuroimaging

methods to examine how early intervention interacts with neural mechanisms of speech

production and self-regulation.

In contrast, a recent randomized controlled trial involving German school-age children

who stutter found no significant effect on the SSI-4 total score after 3 months (d = 0.092) and

only a modest improvement after 12 months (Kohmäscher 2023). These results reflect a

broader trend in school-age intervention studies, where the focus shifts from fluency

enhancement to reducing the negative impact of stuttering on quality of life by addressing their

attitude toward stuttering, with moderate improvements d = 0.62 (Kohmäscher 2023).

Given these findings, there is a pressing need for well-controlled randomized trials to

evaluate the efficacy of fluency-shaping techniques in children. Such studies could help clarify

whether the limited effects observed to date are due to reduced neuroplasticity, suboptimal

timing, or mismatches between the intervention approach and developmental stage. Our own

findings, while preliminary, emphasize the potential benefits of early fluency-based treatment

and underscore the importance of optimizing both timing and method. Similar benefits observed

in early indirect approaches, such as those reported in the RESTART-DCM (Sonneville-Koedoot

et al., 2015) further support the case for early, well-targeted interventions. However, the

complex relationship between intervention timing, treatment type, and long-term psychosocial

outcomes remains an open question and requires further study (Leclercq et al., 2024).

A key challenge in evaluating interventions for preschoolers who stutter is the high

probability of spontaneous remission (Reilly et al., 2009; Yairi & Ambrose, 2013). As noted

earlier, previous prospective controlled trials with both school-age (Craig et al., 1996; Euler et al.,

2021; Kohmäscher et al., 2023) and preschool-age children (Harris et al., 2002; Jones et al.,

2005; Lattermann et al., 2008; Lewis et al., 2008) have addressed this confound using wait-list

control designs to help distinguishing treatment effects from spontaneous recovery. Following

this approach, our study simulated a parallel wait-list control group that did not receive direct

intervention until approximately six months after baseline. At study entry, both groups had

comparable stuttering durations since stuttering onset (*Mdn* = 20 months). While the early

intervention group showed a significant reduction by 9 months post-baseline, after completing

Module 2, the wait-list group showed no comparable improvement during this same period (See

Figure 3A). This strengthens the interpretation that the observed improvements were primarily

attributable to the intervention. However, we acknowledge that individual recovery potential may

interact with treatment outcomes, and we do not suggest that intervention alone accounts for all

observed improvements.

Our data offer only a limited snapshot of the developmental trajectory of assisted

stuttering remission, limited to a three-month intervention window. During this period, children

who continued beyond Module 1 showed notable improvements in fluency. Still, the fluctuating

nature of early stuttering is evident, with some children showing both improvement and later

regression in SSI-4 scores across T0, T3, and T5. Although the intervention extended over 12

months, this does not constitute long-term follow-up, and we are therefore cautious in making

definitive classifications of remission or persistence at T5. In some cases, stuttering severity

increased between T3 and T5 despite earlier gains. To avoid overinterpreting these short-term

fluctuations, we have chosen not to stratify the data by remission status, as such distinctions

would require longer-term follow-up data to be reliable.

Having said this, we consider the findings of the current study preliminary as the

retrospective design did not allow for random assignment to early treatment or wait-list-control

groups, introducing potential selection bias. We controlled for variables, such as sex, age,

stuttering onset, time since onset, and baseline severity to minimize such biases. Observer bias

is also a concern, as SSI-4 raters knew that all participants received treatment, possibly leading

to favorable interpretations of ambiguous cases. To address this, we simulated a waitlist group

and blinded raters for the timing of the speech samples. The observed improvement in the early

group after 9 months, with no change in the delayed group, supports the efficacy of the

treatment. Additionally, since participants and parents knew they were receiving treatment,

placebo effects or response bias could have affected subjective ratings, although the objective

SSI-4 measure is less prone to these biases.

A key strength of this study lies in its use of a retrospective simulated wait-list design,

which, to our knowledge, has not previously been applied in early childhood stuttering research.

This approach offers the ethical and practical advantages of a wait-list comparison without

requiring families to knowingly delay access to treatment. By leveraging pre-existing clinical

data and collecting outcome measures prior to group assignment, the design avoids potential

demotivation, dropout, and dissatisfaction often associated with prospective wait-list controls.

Importantly, the study was conducted within a real-world clinical setting, reflecting how the

intervention is delivered in everyday practice. This enhances the ecological validity of the

findings and underscores the feasibility of applying the method in typical service contexts. In

clinical scenarios where timely intervention is critical, such as early childhood stuttering, this

methodology offers a promising framework for evaluating treatment efficacy while minimizing

ethical concerns. We believe it holds potential for broader use in developmental and pediatric

intervention research.

Limitations

A key limitation of this study is the absence of direct outcome measures assessing the

emotional and psychological effects of the intervention in children. We cannot draw conclusions

about the long-term emotional impact of the therapy, particularly for those children who

continued to stutter after treatment. Although the primary focus was on speech fluency and

parent-reported coping, this leaves important questions unanswered regarding how children

themselves experience and internalize fluency-shaping interventions.

Given concerns that such approaches may inadvertently influence children's attitudes or

self-perception, especially in those for whom stuttering persists, future research should

incorporate validated measures of emotional, social, and cognitive outcomes such as the

Communication Attitude Test for Preschoolers and Kindergartners (KiddyCat, (Vanryckeghem et

al., 2005) and the Overall Assessment of the Speaker's Experience of Stuttering for School-age

children (OASES-S, (Yaruss et al., 2012; Yaruss & Quesal, 2014) to more fully assess the

broader impact of intervention.

Furthermore, the study design did not allow for systematic long-term follow-up of

stuttering outcomes at later time points such as T5. As such, we are unable to report on the

persistence or relapse of stuttering beyond the initial post-treatment phase. This is an important

limitation in assessing the sustained efficacy of Frankini. Future studies, particularly prospective

randomized controlled trials, should include longitudinal follow-ups and measures of

maintenance or change in stuttering severity over time. These should be combined with child-

centered assessments to ensure that both objective and subjective outcomes are adequately

captured, ultimately providing a more comprehensive evaluation of treatment effectiveness.

Another limitation of the present study is the potential for selection bias related to

differences between children who began therapy earlier versus those in the delayed treatment

group. As the study was retrospective, we could not systematically capture the reasons behind

delayed therapy initiation. Importantly, the intervention was delivered in a group format with

limited capacity, and logistical constraints likely influenced enrollment timing. For example,

some families may have experienced delays because courses were already fully booked, while

in other cases, the institution had to postpone starting a group until a minimum number of

participants had registered. These factors suggest that delayed therapy was not necessarily a

reflection of lower parental motivation or child-related variables. Nevertheless, we cannot rule

out the possibility that differences in family engagement, help-seeking behavior, or other

unmeasured factors contributed to treatment timing and outcomes. Future prospective studies

should aim to control for such potential confounders or use randomization to ensure more

balanced group comparisons.

While some degree of attrition bias was unavoidable due to the program's flexible design,

stuttering severity improved for all but one of the children who discontinued after Module 2.

These improvements are consistent with parental ratings available for five of these children,

suggesting that attrition likely did not systematically bias the results toward more favorable

outcomes. As this study is based on retrospectively recorded clinical data, outcome measures

are inherently vulnerable to missing data due to participant dropout. In future prospective

studies, this limitation could be addressed by informing families from the outset about the

importance of completing follow-up assessments at multiple time points, even if treatment is

paused or ended early. This approach would help ensure more complete and systematic

outcome data.

Clinical implications and recommendations for practitioners

The findings of this study have several important implications for clinical practice. First,

the observed benefits of early intervention support current guidelines that emphasize timely

referral and treatment initiation in preschool-aged children who stutter (Brignell et al., 2021;

Keilmann et al., 2018; K. Neumann et al., 2016; Sjøstrand et al., 2021). The results also suggest

that even brief delays in therapy access may impact outcomes, reinforcing the need for efficient

resource allocation and reduced waiting times. Additionally, the use of parent-reported coping

outcomes highlights the importance of involving families in the therapeutic process

(Bergþórsdóttir et al., 2022) and considering both objective and subjective indicators of progress

(Onslow et al., 2018). Clinicians may also consider how structured group formats can be

effectively implemented within clinical settings while balancing logistical constraints.

Conclusion

Early intervention is crucial as it targets stuttering before it becomes deeply ingrained,

leading to improved long-term outcomes and reducing its impact on the child's overall quality of

life. Our novel parent-child program effectively reduced stuttering severity in preschoolers, while

enhancing parents' attitudes and coping strategies. Future research should include follow-up

assessments to determine the long-term sustainability of the intervention's benefits along with

measures of children's self-perception of their speech and communication abilities.

Acknowledgments

This project was supported by the University Medical Center Göttingen, Germany. IN,

AM, KA, and KA have received institutional support from KST GmbH for the development of the

parent-child program.

Data Availability Statement

Owing to ethical concerns, supporting data cannot be made openly available. Further

information regarding the data and conditions for access is available from the corresponding

author.

References

Anders, K., Hente, K. & Merkel, A. (2023). Frankini Therapie für 3- bis 6-jährige stotternde
Kinder und deren Eltern Ein intensivtherapeutischer Behandlungsansatz in der Gruppe.
Sprachtherapie Aktuell: Praxis – Beruf – Verband, 12–17.

Bafrooei, E. B., Farazi, M., Maroufizadeh, S. & Darouie, A. (2025). Development and evaluation
of a combined fluency rules program (FRP) with parent-child interaction training for
preschool children who stutter: A randomized controlled trial. *International Journal of*
Pediatric Otorhinolaryngology, *188*, 112189. https://doi.org/10.1016/j.ijporl.2024.112189

Bergþórsdóttir, Í. Ö., Crowe, K. & Einarsdóttir, J. T. (2022). Implementation fidelity in parent-
implemented interventions for stuttering. *Clinical Linguistics & Phonetics*, *36*(10), 904–927.
https://doi.org/10.1080/02699206.2021.1965659

Bloodstein, O., Ratner, N. B. & Brundage, S. (2021). *A Handbook on Stuttering* (7th ed.). Plural
Publishing, Inc.

Brignell, A., Krahe, M., Downes, M., Kefalianos, E., Reilly, S. & Morgan, A. (2021).
Interventions for children and adolescence who stutter: A systematic review, meta-analysis,
and evidence map. *Journal of Fluency Disorders*, *70*, 105843.
https://doi.org/10.1016/j.jfludis.2021.105843

Brundage, S. B., Ratner, N. B., Boyle, M. P., Eggers, K., Everard, R., Franken, M.-C.,
Kefalianos, E., Marcotte, A. K., Millard, S., Packman, A., Vanryckeghem, M. & Yaruss, J. S.
(2021). Consensus Guidelines for the Assessments of Individuals Who Stutter Across the
Lifespan. *American Journal of Speech-Language Pathology*, *30*(6), 2379–2393.
https://doi.org/10.1044/2021_ajslp-21-00107

Chang, S.-E., Below, J. E., Chow, H. M., Guenther, F. H., Wray, A. M. H., Jackson, E. S., Max,
L., Neef, N. E., SheikhBahaei, S., Shekim, L., Tichenor, S. E., Walsh, B., Watkins, K. E.,
Yaruss, J. S. & Ratner, N. B. (2025). Stuttering: Our Current Knowledge, Research
Opportunities, and Ways to Address Critical Gaps. *Neurobiology of Language*, *6*,
nol_a_00162. https://doi.org/10.1162/nol_a_00162

Chow, H. M. & Chang, S. (2017). White matter developmental trajectories associated with
persistence and recovery of childhood stuttering. *Human Brain Mapping*, *38*(7), 3345–3359.
https://doi.org/10.1002/hbm.23590

Chow, H. M., Garnett, E. O., Koenraads, S. P. C. & Chang, S.-E. (2023). Brain developmental
trajectories associated with childhood stuttering persistence and recovery. *Developmental*
Cognitive Neuroscience, *60*, 101224. https://doi.org/10.1016/j.dcn.2023.101224

Craig, A., Hancock, K., Chang, E., McCready, C., Shepley, A., McCaul, A., Costello, D.,
Harding, S., Kehren, R., Masel, C. & Reilly, K. (1996). A Controlled Clinical Trial for
Stuttering in Persons Aged 9 to 14 Years. *Journal of Speech, Language, and Hearing*
Research, *39*(4), 808–826. https://doi.org/10.1044/jshr.3904.808

Euler, H. A., Gudenberg, A. W. von, Jung, K. & Neumann, K. (2009). Computergestützte
Therapie bei Redeflussstörungen: Die langfristige Wirksamkeit der Kasseler Stottertherapie
(KST). *Sprache, Stimme, Gehör*, *33*(04), 193–202. https://doi.org/10.1055/s-0029-1242747

693 Euler, H. A., Lange, B. P., Schroeder, S. & Neumann, K. (2014). The effectiveness of stuttering
694  treatments in Germany. *Journal of Fluency Disorders*, *39*, 1–11.
695  https://doi.org/10.1016/j.jfludis.2014.01.002

696 Euler, H. A., Merkel, A., Hente, K., Neef, N., Gudenberg, A. W. von & Neumann, K. (2021).
697  Speech restructuring group treatment for 6-to-9-year-old children who stutter: A therapeutic
698  trial. *Journal of Communication Disorders*, *89*, 106073.
699  https://doi.org/10.1016/j.jcomdis.2020.106073

700 Franken, M.-C. & Laroes, E. (2021). *RESTART-DCM Method*. https://restartdcm.nl/wp-
701  content/uploads/2021/05/RestartDCM-Method-2021_online.pdf

702 Franken, M.-C., Millard, S. & Hearne, A. (2022). 10 Preschool-age children. In P. M.
703  Zebrowski, J. D. Anderson & E. G. Conture (Eds.), *Stuttering and related disorders of fluency*
704  (pp. 153–173). Thieme.

705 Harris, V., Onslow, M., Packman, A., Harrison, E. & Menzies, R. (2002). An experimental
706  investigation of the impact of the Lidcombe Program on early stuttering. *Journal of Fluency
707  Disorders*, *27*(3), 203–214. https://doi.org/10.1016/s0094-730x(02)00127-4

708 Hayes, A. F. & Krippendorff, K. (2007). Answering the Call for a Standard Reliability Measure
709  for Coding Data. *Communication Methods and Measures*, *1*(1), 77–89.
710  https://doi.org/10.1080/19312450709336664

711 Hughes, J. (2021). krippendorffsalpha: An R Package for Measuring Agreement Using
712  Krippendorff's Alpha Coefficient. *The R Journal*, *13*(1), 413. https://doi.org/10.32614/rj-
713  2021-046

714 Jones, M., Onslow, M., Packman, A., Williams, S., Ormond, T., Schwarz, I. & Gebski, V.
715  (2005). Randomised controlled trial of the Lidcombe programme of early stuttering
716  intervention. *BMJ*, *331*(7518), 659. https://doi.org/10.1136/bmj.38520.451840.e0

717 Keilmann, A., Neumann, K., Zöller, D. & Freude, C. (2018). Clinical trial of the D.E.L.P.H.I.N.
718  speech treatment for children and adolescents who stutter. *Logopedics Phoniatrics Vocology*,
719  *43*(4), 155–168. https://doi.org/10.1080/14015439.2018.1498917

720 Kelman, E. & Nicholas, A. (2019). *Practical Intervention for Early Childhood Stammering,
721  Palin PCI Approach*. https://doi.org/10.4324/9781315169200

722 Kohmäscher, A., Primaßin, A., Heiler, S., Avelar, P. D. C., Franken, M.-C. & Heim, S. (2023).
723  Effectiveness of Stuttering Modification Treatment in School-Age Children Who Stutter: A
724  Randomized Clinical Trial. *Journal of Speech, Language, and Hearing Research*, *66*(11),
725  4191–4205. https://doi.org/10.1044/2023_jslhr-23-00224

Lattermann, C., Euler, H. A. & Neumann, K. (2008). A randomized control trial to investigate the impact of the Lidcombe Program on early stuttering in German-speaking preschoolers. *Journal of Fluency Disorders*, *33*(1), 52–65. https://doi.org/10.1016/j.jfludis.2007.12.002

Leclercq, A.-L., Waelkens, V., Roelant, E., Allegaert, M., Verhaegen, I., Claes, K., Dauvister, E., Snijders, S., Eggers, K., Moyse, A. & Eerdenbrugh, S. V. (2024). Treatment for preschool age children who stutter: Protocol of a randomised, non-inferiority parallel group pragmatic trial with Mini-KIDS, social cognitive behaviour treatment and the Lidcombe Program— TreatPaCS. *PLOS ONE*, *19*(7), e0304212. https://doi.org/10.1371/journal.pone.0304212

Lewis, C., Packman, A., Onslow, M., Simpson, J. M. & Jones, M. (2008). A Phase II Trial of Telehealth Delivery of the Lidcombe Program of Early Stuttering Intervention. *American Journal of Speech-Language Pathology*, *17*(2), 139–149. https://doi.org/10.1044/1058-0360(2008/014)

Millard, S. K., Edwards, S. & Cook, F. M. (2009). Parent-child interaction therapy: Adding to the evidence. *International Journal of Speech-Language Pathology*, *11*(1), 61–76. https://doi.org/10.1080/17549500802603895

Millard, S. K., Zebrowski, P. & Kelman, E. (2018). Palin Parent–Child Interaction Therapy: The Bigger Picture. *American Journal of Speech-Language Pathology*, *27*(3S), 1211–1223. https://doi.org/10.1044/2018_ajslp-odc11-17-0199

Neef, N. E. & Chang, S.-E. (2024). Knowns and unknowns about the neurobiology of stuttering. *PLOS Biology*, *22*(2), e3002492. https://doi.org/10.1371/journal.pbio.3002492

Neumann, K., Euler, H. A., Bosshardt, H. G., Cook, S., Sandrieser, P., Schneider, P., Sommer, M. & Thum, G. (2016). *Pathogenese, Diagnostik und Behandlung von Redeflussstörungen. Evidenz- und konsensbasierte S3- Leitlinie* (D. G. für P. und Pädaudiologie, Ed.).

Neumann, Katrin, Euler, H. A., Bosshardt, H.-G., Cook, S., Sandrieser, P. & Sommer, M. (2017). The Pathogenesis, Assessment and Treatment of Speech Fluency Disorders. *Deutsches Aerzteblatt Online*, *114*(22–23), 383–390. https://doi.org/10.3238/arztebl.2017.0383

Onslow, M., Andrews, C. & Lincoln, M. (1994). A Control/Experimental Trial of an Operant Treatment for Early Stuttering. *Journal of Speech, Language, and Hearing Research*, *37*(6), 1244–1259. https://doi.org/10.1044/jshr.3706.1244

Onslow, M., Jones, M., O'Brian, S., Packman, A., Menzies, R., Lowe, R., Arnott, S., Bridgman, K., Sonneville, C. de & Franken, M.-C. (2018). Comparison of Percentage of Syllables Stuttered With Parent-Reported Severity Ratings as a Primary Outcome Measure in Clinical Trials of Early Stuttering Treatment. *Journal of Speech, Language, and Hearing Research*, *61*(4), 811–819. https://doi.org/10.1044/2017_jslhr-s-16-0448

Onslow, M., Packman, A. & Harrison, R. E. (2003). *The Lidcombe Program of early stuttering*
Intervention: A clinician's guide. Pro-Ed.

Preston, R., Halpin, M., Clarke, G. & Millard, S. (2022). Palin parent-child interaction therapy
with children with autism spectrum disorder and stuttering. *Journal of Communication*
Disorders, *97*, 106217. https://doi.org/10.1016/j.jcomdis.2022.106217

Reilly, S., Onslow, M., Packman, A., Wake, M., Bavin, E. L., Prior, M., Eadie, P., Cini, E.,
Bolzonello, C. & Ukoumunne, O. C. (2009). Predicting Stuttering Onset by the Age of 3
Years: A Prospective, Community Cohort Study. *Pediatrics*, *123*(1), 270–277.
https://doi.org/10.1542/peds.2007-3219

Riley, G. D. (2009). *SSI-4: Stuttering Severity Instrument* (Fourth Edition).

Sjøstrand, Å., Kefalianos, E., Hofslundsengen, H., Guttormsen, L. S., Kirmess, M., Lervåg, A.,
Hulme, C. & Næss, K.-A. B. (2021). Non□pharmacological interventions for stuttering in
children six years and younger. *Cochrane Database of Systematic Reviews*, *2021*(9),
CD013489. https://doi.org/10.1002/14651858.cd013489.pub2

Sommer, M., Waltersbacher, A., Schlotmann, A., Schröder, H. & Strzelczyk, A. (2021).
Prevalence and Therapy Rates for Stuttering, Cluttering, and Developmental Disorders of
Speech and Language: Evaluation of German Health Insurance Data. *Frontiers in Human*
Neuroscience, *15*, 645292. https://doi.org/10.3389/fnhum.2021.645292

Sonneville-Koedoot, C. de, Stolk, E., Rietveld, T. & Franken, M.-C. (2015). Direct versus
Indirect Treatment for Preschool Children who Stutter: The RESTART Randomized Trial.
PLoS ONE, *10*(7), e0133758. https://doi.org/10.1371/journal.pone.0133758

Tichenor, S. E. & Yaruss, J. S. (2021). Variability of Stuttering: Behavior and Impact. *American*
Journal of Speech-Language Pathology, *30*(1), 75–88. https://doi.org/10.1044/2020_ajslp-20-
00112

Vanryckeghem, M., Brutten, G. J. & Hernandez, L. M. (2005). A comparative investigation of
the speech-associated attitude of preschool and kindergarten children who do and do not
stutter. *Journal of Fluency Disorders*, *30*(4), 307–318.
https://doi.org/10.1016/j.jfludis.2005.09.003

Webster, R. L. (1980). *The Precision Fluency Shaping Program: Speech Reconstruction for*
Stutterers.

Wolff-von-Gudenberg, A. & Euler, H. A. (2000). Evaluation study of the Kasseler Stuttering
Therapy (KST). *Journal of Fluency Disorders*, *25*(3), 195. https://doi.org/10.1016/s0094-
730x(00)80227-2

Yairi, E. & Ambrose, N. (2013). Epidemiology of stuttering: 21st century advances. *Journal of*
Fluency Disorders, *38*(2), 66–87. https://doi.org/10.1016/j.jfludis.2012.11.002

Yaruss, J. S., Coleman, C. E. & Quesal, R. W. (2012). Stuttering in School-Age Children: A
Comprehensive Approach to Treatment. *Language, Speech, and Hearing Services in Schools*,
43(4), 536–548. https://doi.org/10.1044/0161-1461(2012/11-0044)

Yaruss, J. S. & Quesal, R. W. (2014). *OASES: Overall Assessment of the Speaker's Experience*
of Stuttering ((Deutsche Übersetzung: H. A. Euler & & A. Alpermann), Trans.).

Tables

Table 1. Baseline demographic and clinical characteristics

Characteristic	Early-treatment $n = 23$	Delayed-treatment $n = 19$	Difference $T(df)$ p
Sex, No. (%)			
Male	16 (70)	13 (68)	[a]0.007 (1.0)
Female	7 (30)	6 (32)	.936
Age stuttering onset, mo			
M (*SD*)	33.7 (8.7)	32.1 (7.0)	0.66 (40.0)
Mdn (*IQR*)	33 (8.5)	30 (10)	.515
Min-Max	18-54	24-48	
Age start Module 1, mo			
M (*SD*)	53.6 (8.8)	58.0 (8.0)	-0.40 (39.6)
Mdn (*IQR*)	55.5 (11.2)	58.0 (11.7)	.690
Min-Max	36.0 - 69.7	38.0 - 67.6	
Δ start Module 1 - age onset, mo			
M (*SD*)	19.9 (9.3)	22.6 (9.0)	-0.93 (39.1)
Mdn (*IQR*)	19.5 (12.9)	20.9 (12.8)	.358
Min-Max	4.3 - 43.3	7.0 - 37.0	
Δ T2 - T0, mo			
M (*SD*)	5.3 (0.5)	8.1 (1.0)	-10.8 (26.0)
Mdn (*IQR*)	5.3 (0.7)	8.5 (1.5)	**< .001**
Min-Max	3.9 - 6.1	6.3 - 9.4	
N Syllables of T0 sample			
M (*SD*)	377 (193)	389 (164)	[c]78.5 (.079)
Mdn (*IQR*)	348 (348)	316 (246)	
Min-Max	23 - 614	69 - 615	
SSI-4 total score at T0			
M (*SD*)	16.2 (7.3)	15.6 (5.4)	0.34 (39.5)
Mdn (*IQR*)	13.5 (13)	15.9 (6.5)	.734
Min-Max	6.0 - 31.5	4.0 - 26.0	
Stuttering frequency at T0 (%)			
M (*SD*)	7.6 (5.1)	7.0 (4.6)	0.44 (39.7)
Mdn (*IQR*)	6.5 (7.5)	5.7 (6.5)	.656
Min-Max	1.2 - 18.6	0.3 - 16.9	
Parent rating at T0			
M (*SD*)	6.0 (2.1)	[b]5.3 (2.2)	1.06 (33.5)
Mdn (*IQR*)	6.0 (2.0)	5.0 (2.0)	.297
Min-Max	2 - 10	2 – 10	
Online Questionnaire at T0			
…Q1 *Mdn* (*IQR*)	3.0 (1.0)	3.0 (0.5)	[c]198 (.954)
…Q2 *Mdn* (*IQR*)	2.8 (2.0)	3.0 (1.0)	[c]151 (.295)
…Q3 *Mdn* (*IQR*)	3.0 (2.1)	3.0 (2.0)	[c]157 (.395)
…Q4 *Mdn* (*IQR*)	5.0 (1.8)	4.0 (1.0)	[c]204 (.610)
…Q5 *Mdn* (*IQR*)	3.0 (2.0)	3.0 (1.5)	[c]112 (.556)

…Q6 *Mdn* (*IQR*)	2.0 (2.0)	2.0 (2.0)	[c]188 (.844)
…Q7 *Mdn* (*IQR*)	4.0 (2.0)	5.0 (1.0)	[c]167 (.414)
…Q8 *Mdn* (*IQR*)	2.0 (2.0)	2.0 (2.9)	[c]167 (.553)
…Q9 *Mdn* (*IQR*)	3.0 (2.0)	2.0 (1.3)	[c]130 (.468)
…Q10 *Mdn* (*IQR*)	2.5 (2.0)	2.0 (1.5)	[c]250 (.069)

805   [a]χ^2-test, [b]n = 17, [c]*W* (*p*-value) after the Wilcoxon rank sum test with continuity correction

808 **Table 2 Summary of mean changes from baseline per group**

		Early-treatment		Delayed-treatment	
		at 9 months $n = 23$	at 12 months $n = 16$	at 9 months $n = 19$	at 12 months $n = 19$
Stuttered syllables %	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-5.3 [-7.3 to -3.2] -4.0 (-16.7 to 1.4)	-6.0 [-9.0 to -3.1] -5.1 (-15.4 to 3.5)	-0.8 [-3.6 to 1.9] -0.8 (-11.7 to 7.6)	-3.8 [-5.4 to -2.1] -2.9 (-10.2 to 0.6)
SSI-4 total	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-8.7 [-12.2 to -5.2] -7.5 (-23.0 to 8.5)	-10.6 [-18.2 to -3.3] 17.0 (-31.5 to 18)	0.3 [-3.1 to 3.6] 0.5 (-12.0 to 13.0)	-5.1 [-8.0 to -2.2] -6.0 (-14.5 to 4)
Frequency	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-5.4 [-7.4 to -3.3] -5.0 (-15.0 to 5.0)	-5.6 [-8.5 to -2.6] -4.5 (-15 to 5)	0.2 [-2.3 to 2.7] 0.0 (-7.0 to 9.0)	-2.7 [-4.6 to -0.8] -3.0 (-9.0 to 4.0)
Duration	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-2.9 [-4.1 to -1.6] -2.0 (-8.0 to 2.0)	-2.6 [-4.1 to -1.1] -3.0 (-8.0 to 4.0)	-0.1 [-1.0 to 0.8] 0.0 (-4.0 to 3.0)	-1.6 [-2.4 to -0.7] -1.0 (-4.0 to 1.0)
Physical concomitants	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-0.5 [-1.0 to 0.1] 0.0 (-4.5 to 1.5)	-0.8 [-1.8 to 0.2] -0.5 (-6.5 to 2.5)	0.3 [-0.7 to 1.4] 0.0 (-2.0 to 7.0)	-0.7 [-1.4 to -0.01] -0.5 (-4.5 to 2.0)
Subjective parent rating	$M\,[CI_{95\%}]$ $Mdn\,(Min$ to $Max)$	-2.75 [-3.9 to -1.6] -3 (-8 to 3)	-3.1 [-4.6 to -1.5] -3 (-7 to 2)	nan	-1.8 [-3.2 to -0.5] -2 (-5 to 2)

Figure Legends

Figure 1. Procedure of the parent child program and participant's group allocation. A) T0

– T5 are assessment time points. Assessments included Stuttering Severity Index (SSI),

subjective parent rating (PR), online questionnaire on parents' experience with the child's

stuttering (OQ). Modules included online sessions (blue) therapy in presence (red), individual

therapy (IT) and group therapy (GT). B) Individual time schedules and group assignments of the

participants.

Figure 2. Participant flow diagram illustrates the sequence of identifying participants for

inclusion in the analysis.

Figure 3. Changes in stuttering severity and parents' experiences of their child's

stuttering over time. A) Stuttering severity (SSI) reduced after 9 months in the early treatment

group, with no change in the delayed group. Plotted are group averages and standard errors.

The dotted line represents the SSI cut-off value for stuttering. B) Box plots and individual curves

for SSI total scores, separately plotted for the three treatment modules. Red lines indicate cases

not participating in Module 3. Blue circles indicate cases with missing values. C) Heat map

showing changes in parental experience with their child's stuttering between baseline and T3

(after Module 2). White cells indicate missing data. The last column indicates the grand mean

for a given item with $*p < .05$, $**p < .01$, and $***p < .001$ (Wilcoxon signed rank test with

continuity correction).

A

SSI total score

delayed
early

pre   9   12   16 months

B

Module 1
n = 42

Module 2
n = 42

Module 3
n = 30

SSI total score

T0   T1   T2   T3   T4   T5

C

Parental coping with child stuttering [ΔT3-T0]

Deal calmly with my child's stuttering — ***
Confident talking to my child — ***
Confident talking to others — ***
Confident talking to close contacts — **
I can handle my childs emotions — ***
My child's stuttering embarrasses me — **
My child's stuttering worries me — ***
Talking about stuttering is uncomfortable —
My child's reactions overwhelm me — *
Stuttering disrupts family communication — *

Nr. participant   Group average

Δ Score