1 **Title**
2 Functional Variant Discovery identifies a novel genetic link between SPRY2, wood smoke, and
3 asthma
4

5 **Authors**
6 Arnav Gupta[1,7], Amber Dahlin[2], Alejandra Macario[1], Fabienne Gally[1,3], Michael R. Weaver[1],
7 Samuel Guarino[1], Louisa Kahn[1], Lynn Sanford[4], Margaret A. Gruca[4], Michael H. Cho[2], Robin D.
8 Dowell[4,5,6], Scott T. Weiss[2], Sarah K. Sasse[1], Anthony N. Gerber[1,3,7]
9
1. Department of Medicine, National Jewish Health
2. Channing Division of Network Medicine, Harvard Medical School
3. Department of Immunology and Genomic Medicine, National Jewish Health
4. BioFrontiers Institute, University of Colorado Boulder
5. Molecular, Cell, and Developmental Biology, University of Colorado Boulder
6. Computer Science, University of Colorado Boulder
7. Department of Medicine, University of Colorado Anschutz Medical Campus
17
18 Corresponding Author: Arnav Gupta
19 guptaa@njhealth.org
20 214-649-9559

27
28

Abstract
As a consequence of climate change and land use policies, there has been a historic rise in
wildfire smoke across the United States and the world. While the deleterious effects of wildfire
smoke and associated air pollution on asthma outcomes are established epidemiologically,
genetic risks and molecular mechanisms of how wildfire smoke affects asthma are unknown.
This knowledge gap hinders the identification of high-risk individuals and the creation of
targeted therapies or recommendations to protect these individuals. We identified 52 genetic
risk variants that colocalized with genomic responses to wood smoke particles (WSP), a model
of wildfire particulate matter, and associated with asthma in the Genetic Epidemiology Research
on Aging (GERA) cohort. We used additional filters to prioritize variants for direct testing of
allele-dependent transcriptional regulatory function in plasmid reporters. We found that the
rs3861144 variant (Odds Ratio$_{asthma}$ = 1.036) changes *SPRY2* responses to WSP in airway
epithelial cells, which are involved in Interleukin-8 secretion, Extracellular Signal-related Kinase
(ERK) activation, and mechanical scratch repair in cell culture. These findings provide insights
into the molecular pathways through which WSP may influence asthma risk and propose
genetic candidates that warrant further study for their potential as clinical tools for asthma.

Introduction
Climate change is contributing to a historic rise in wildfire smoke and reversing decades of
improvements in air quality made by environmental policy (1–3). The scope of this problem is
extending across North America and the world, increasing the population exposed to wildfire air
pollution (4). Wildfire smoke is difficult to regulate or prevent. Climate policy and wildland
management strategies will take decades to reverse the trend of expanding wildfires, and
deteriorating air quality is expected over the next 30 years (5). It is therefore critical to
understand the health consequences of wildfire air pollution and implement detailed strategies
to manage and mitigate this developing healthcare crisis.

The profound health impacts of wildfire smoke and air pollution are well established at the
population scale (6,7) and the impact on respiratory diseases, such as asthma, is a major
concern. Wildfire smoke increases levels of ambient air pollutants, such as particulate matter
less than 2.5 um (PM2.5), and causes morbid asthma outcomes, such as loss of asthma
control, emergency room visits and hospitalizations (8–10). Data suggest ambient PM2.5 may
also contribute to reduced lung function and the development of asthma (11–13). These major
concerns highlight critical knowledge gaps in our current understanding of how PM2.5 from
wildfire smoke impacts respiratory biology and asthma. First, the molecular genetic mechanisms
that connect wildfire PM2.5 with asthma have not been identified, preventing the development of
targeted treatments that mitigate asthma development or exacerbation. Second, there is
significant variation in how individuals respond to airborne exposures, such as PM2.5 or wildfire
smoke (14), and public health guidelines, which only advise exposure avoidance, have limited
efficacy, as restricting large population segments from outdoor activity for weeks during wildfire
events is impractical. A better understanding of how PM2.5 and wildfire air pollution contributes
to asthma risk and identification of high-risk individuals are critical to address this growing
problem.

Genetic analysis is a powerful tool to inform disease mechanisms, define individual risk, and
advance our understanding of the relationships between PM2.5, wildfire smoke, and asthma.
However, there is a significant gap between the effect sizes of established genetic variants and
asthma heritability, which is estimated at up to 70% of causation in certain studies (15). This
suggests that additional asthma risk factors that are not captured in genome-wide association
studies (GWAS), such as environmental exposures, can influence genetic effects. This concept
is supported by discoveries of gene mechanisms that connect air pollution exposure with

asthma, such as mutations regulating *NQO1*, *GSTM1* and *GSTP1* (16–18). Identification of
additional conditional genetic risk variants, which primarily contribute to asthma risk in the
presence of exposure, has the potential to define novel asthma mechanisms associated with
wildfire PM2.5 and individuals with high risk.
84
Several challenges limit the identification of exposure-conditional risk variants, despite the major
advances achieved by asthma genetic studies. First, integrating environmental variables in
GWAS is highly challenging due to a myriad of reasons, including exposure variability, duration
of exposure effects, and types of exposures, among others. Genetic-environment interaction
studies have approached this challenge by quantifying environmental variables through
geolocalization and retrospective measurements of local air quality (19), but these studies are
limited by the precision of geolocalization, resolution of air pollutant measurements, and
difficulty with compounding exposures based on where study subjects have lived over decades.
Second, linking specific genetic variants with cell and molecular functions can be challenging.
GWAS lead variants often reside in non-coding regions (20) and represent statistical but not
functional association priorities. Tools, such as functional scoring and expression quantitative
trait locus (eQTL) analysis, can improve associations between GWAS variants and gene
regulation but fail to leverage the full power of GWAS results, which may contain additional
functional variants that do not meet statistical thresholds. These tools also do not account for
the effects of relevant environmental exposures. Therefore, novel approaches that build on top
of existing GWAS data, incorporate environmental exposures, and directly connect variants with
function are necessary to realize the full potential of genetic analysis.
102
In this study, we applied our novel approach to variant discovery (21) to identify functional
genetic variants and molecular mechanisms that connect a wildfire PM2.5 model with asthma
risk. Using wood smoke particles (WSP) (22,23) to model wildfire PM2.5, we identified and
biologically validated a set of exposure-conditional genetic variants using population data of
lung function, transcription factor binding motifs, and epigenetic features of live airway epithelia
obtained from bronchoscopy. We directly tested the potential for high-priority variants to change
transcriptional function, connected variants with gene targets, and tested the function of these
target genes in airway epithelial cell biology. Our results were highlighted by the discovery of
rs3861144, a functional variant that changes *SPRY2* responses to WSP exposure and
potentially contributes to asthma through cytokine secretion and injury-repair programs.
113
114   **Methods and Materials**
115   Additional methods are provided in the Online Data Supplement due to word limit constraints.
116
Genetic discovery data set and Analysis
Genome-wide and phenotype data were utilized from the Genetic Epidemiologic Research on
Aging (GERA) study, a large, multi-ethnic asthma cohort including over 110,000 subjects
(68,623 total asthma cases across four ethnic groups) with extensive electronic medical records
and genome-wide SNP genotypes (>8 million typed and imputed markers). We previously
reported multiple novel associations with asthma in four major ethnic groups in GERA (24). The
current analysis was limited to the ethnic subset of the cohort with the highest percentage of
asthma cases and therefore the greatest statistical power; the final data set included information
from 16,274 non-Hispanic white asthmatic cases and 38,269 controls (54,543 total samples).
We applied methods previously developed by our group (21) to regulatory elements identified in
WSP-exposed Beas-2B cells by Precision Run-on sequencing (PRO-seq) (n=18,552) (25).
SNPs that overlapped regions within 50 bp (+/-) of regulatory element centers were selected for
association analysis. Tests for association of additive genotype with asthma status
(dichotomized outcome variable) were conducted using multivariate logistic regression models,

adjusted for covariates, and the significance of associations was determined using a max(T)
permutation-based approach with 5000 permutations and a false-discovery rate (FDR) threshold
of 0.05. All analyses were performed using R statistical computing software and PLINK 1.9.
134
CRISPR-Cas9 RNP transfection and nucleofection
We performed genomic deletion of the transcriptional regulatory element (TRE) containing
rs3861144 and *SPRY2* using the Alt-R CRISPR-Cas9 system supplied and guided by IDT.
Custom-designed crRNAs, which target two sites within the TRE or the *SPRY2* gene body, and
a negative control crRNA (IDT) were duplexed (5 min at 95C) with tracrRNAs and then
complexed with Cas9 enzyme (1:1 molar ratio of duplex to Cas9) to a final concentration of 1
uM. This ribonucleic protein complex was transfected into Beas-2B cells using Lipofectamine
CRISPRmax (Life Technologies) or electroporated into primary small airway epithelial cells
(SAECs) using the Lonza nucleofector IIb system and the Basic Kit for Epithelial Cells.
Nucleofection protocol was adapted from a previous study (26) for the I/IIb system using the T-
020 program. Genome editing was confirmed by PCR amplification of genomic DNA (DNeasy
Blood and Tissue kit – Qiagen) or Western Blot. Guide RNA sequences are provided in
Supplemental Table E2.
148
Results
Identification of functional variant candidates associated with asthma risk and wood smoke
particle exposure. To connect airway epithelial wood smoke particle (WSP) responses with
mechanisms of asthma, we re-analyzed nascent transcription profiles from an airway epithelial
cell line (Beas-2B) exposed to WSP for 30 or 120 minutes (25). Within 18,552 dynamic
transcriptional regulatory element (TRE, transcriptional start sites excluded) responses to WSP
(100 bp regions, increased or decreased transcription between vehicle/control and WSP at 30
or 120 minutes), we tested 4373 single nucleotide polymorphisms (SNPs, minor allele frequency
> 0.05) and identified 52 candidates associated with asthma phenotype in the non-Hispanic-
white subpopulation of the GERA cohort (False Discovery Rate, FDR < 0.05, max(T)
permutation), which are designated "WSP-SNPs" in the remainder of this manuscript (Figure
1A). These candidates predominantly reside in non-coding genomic regions (Figure 1B). A
representative example of a WSP-SNP, rs28362965, is depicted in Figure 1C. Here, WSP
exposure increases transcription of both the TRE containing rs28362965 and *HK2.* A full list of
the 52 WSP-SNPs is provided in Table 1.
164
WSP-SNPs associated with measures of lung function in an independent genetic data set. As
an initial validation approach, we analyzed the 52 WSP-SNPs for associations with lung function
in previously published GWAS data. We used the Open Targets Genetics platform (27,28) and
the Lung Disease Knowledge Portal (29) to navigate GWAS data of lung function
measurements, such as forced vital capacity (FVC), forced expiratory volume at 1 second
(FEV1), or peak expiratory flow (PEF) (30,31). We identified 6 WSP-SNPs significantly
associated (p_{adj} < 0.05) with measures of lung function (Table 2). Three WSP-SNPs (rs3861144,
rs3129975 and rs28517751) achieved a statistical threshold for genome-wide association (p <
$5x10^{-8}$). Associations between WSP-SNPs and measurements of lung function, which are
objective features of asthma, supported the validity of our discovery data set and motivated
further biological study of the 52 WSP-SNPs.
176
Aggregate analysis of the 52 WSP-SNPs suggested regulation of diverse cellular processes. To
understand biologic pathways regulated by the WSP-SNPs, we imputed their gene targets using
two methods. First, we used expression quantitative trait locus (eQTL) data from the Open
Targets Genetics database to assign gene targets to SNPs. This approach was not
comprehensive, as multiple WSP-SNPs had no eQTL associations. However, we identified 154

genes associated with at least one WSP-SNP. Functional annotation using DAVID (32) identified
no enriched categories for these 154 genes. Second, we applied a permissive approach that
identified all genes that were located within 500 kb and shared nascent transcriptional kinetics
with the WSP-SNPs' TRE. This approach identified 376 genes, and one significantly enriched
(p_{adj} = 2.1e-7) annotation term – "keratin filament." A complete list of the putative gene targets
from each approach is provided in Supplemental Tables E4 and E5. Based on the limited
aggregate clustering, we concluded that the WSP-SNPs regulate diverse mechanisms, and
focused analysis was necessary to identify functional SNPs.
190
Focused biologic analysis using chromatin accessibility and bioinformatic transcription factor
binding predictions identified WSP-SNP candidates for functional analysis. We next applied
cellular and molecular features relevant to asthma and wildfire PM2.5 exposures to evaluate
regulatory regions containing WSP-SNPs. We found that 23 WSP-SNPs overlapped with
chromatin accessibility in live airway epithelial cells obtained from bronchoscopy from three
healthy non-smoking adult donors (Figure 2A), supporting the biologic validity of these WSP-
SNPs in airway epithelial function. A representative example demonstrates chromatin
accessibility in live airway epithelium at the rs9903086 SNP and its neighboring gene *JUP*
(Figure 2B). We next scanned genome sequences of TREs containing WSP-SNPs for binding
motifs from two classes of transcription factors that are relevant to WSP and PM2.5 exposures –
aryl hydrocarbon receptor (AHR) and nuclear factor kappa B (NFkB) (33,34). Using a 500 bp
radius to define the regulatory region containing each WSP-SNP, we encountered these motifs
at 38 WSP-SNP loci, and 23 WSP-SNP loci contained motifs for both AHR and NFkB factors
(Figure 2C), supporting our previous study that demonstrated cross talk between these factors
as a mechanism mediating inflammatory responses to WSP exposure (25). The TRE containing
rs11883931, which neighbors *LRRFIP1*, is one example that contains both AHR nuclear
transporter (ARNT2) and NFkB (REL) transcription factor binding motifs (Figure 2D). These
genomic features biologically validated and prioritized a subset of WSP-SNPs for further
functional analysis.
210
Functional WSP-SNP genotype changed transcriptional regulatory potential in plasmid reporter
systems. We selected 15 prioritized WSP-SNPs to test transcriptional function in plasmid
systems (Figure 3A) and cloned 700-800 bp regions surrounding these SNPs into luciferase
reporters. Within this reporter set, 5 WSP-SNP constructs exhibited enhancer activity at
baseline or in response to WSP exposure, and single base allele change of 2 WSP-SNPs
changed the regulatory activity of the entire construct (Figure 3B). Based on this, we defined
rs3861144 and rs11832287 as functional genetic variants that alter the transcriptional regulatory
activity of their resident TREs based on haplotype.
219
WSP-SNPs identified functional variants at statistically prioritized GWAS loci. The functional
genetic variants that we identified were not previously reported but had statistical associations
with lung function, such as rs3861144 with FEV1 (Table 2). We next identified the leading
statistical candidate SNPs that colocalize with WSP-SNPs to compare regulatory functions
using the luciferase reporter system. We performed Bayesian fine mapping (35) for the
continuous FVC trait (36) at the rs3861144 and rs9903086 loci. No SNPs were identified in the
credible set at rs3861144, but rs4885681 was identified as the local lead variant based on the
lowest p value for association (p = 1.3×10^{-11}). The rs3861144 plasmid reporter had greater
transcriptional regulatory activity in the luciferase assay when compared with a plasmid reporter
similarly constructed for rs4885681, which resides in a non-annotated and non-functional
genomic region (Figure 3C-D). We selected three SNPs within the credible set (posterior
probability > 0.95) of association with FVC at the rs9903086 locus – rs9913846, rs9911655, and
rs33914436 (Supplemental Table E6). Again, the transcriptional regulatory activity of the

rs9903086-containing plasmid reporter at baseline and in response to WSP exposure was
greater than reporters containing these local fine-mapped SNPs (Figure 3E), which reside in
genomic regions without functional annotations (Supplemental Figure E1). These results
demonstrate that within a genomic locus associated with lung function, the genetic variants
identified by our approach have greater transcriptional regulatory function than the lead variants
reported by prior GWAS.
239
Shared nascent transcriptional kinetics between gene and TRE identified SPRY2 as the target
of the rs3861144 WSP-SNP. We applied PRO-seq data, three-dimensional chromatin analysis,
and genome editing to propose and confirm the gene target of one selected WSP-SNP,
rs3861144. Prior studies have connected TREs with target genes based on shared activity and
genomic proximity (37). Using this concept, the regulatory element containing rs3861144, a
WSP-SNP independently associated with lung function with allele-dependent transcriptional
regulatory activity, regulates *SPRY2* in response to WSP exposure (Figure 4A). Three-
dimensional chromatin analysis using micrococcal nuclease chromosome conformation capture
(Micro-C) in unstimulated Beas-2B cells (38) demonstrated that rs3861144 and *SPRY2* reside in
the same topologically associated domain and may interact indirectly (Supplemental Figure E2).
To definitively link the TRE harboring rs3861144 with control of *SPRY2*, we deleted the
rs3861144 TRE using CRISPR-Cas9 and analyzed *SPRY2* expression using qPCR in Beas-2B
cells. PCR amplification of genomic DNA from cells transfected with negative control or
rs3861144-targeting guide RNA demonstrated efficient deletion of this TRE (Figure 4B). Cells
with rs3861144 TRE deletion showed a reduction in *SPRY2* expression response to WSP
(Figure 4C). These results connected the rs3861144 TRE with *SPRY2* expression in response
to WSP and demonstrated the facility of functional SNPs to connect exposure, gene
mechanism, and phenotype.
258
Cytokine kinetics are regulated by SPRY2 knockdown in Beas-2B cells exposed to WSP. We
next studied how *SPRY2* knockdown interacts with WSP exposure by applying short-interfering
RNA targeted to *SPRY2* (si*SPRY2*) in Beas-2B cells and WSP exposure at multiple time points.
We first performed RNA sequencing in si*SPRY2* and control cells exposed to WSP for 2 and 4
hours. A principal component plot of sample clustering is presented in Supplemental Figure E4.
Within gene expression differences between WSP-exposed si*SPRY2* and control cells at 4
hours of exposure, we observed 180 differentially expressed genes enriched in si*SPRY2* cells
(p_{adj} < 0.05, log_2FoldChange > 0.5), and using DAVID (32) for functional annotation, we
identified "Growth Factor" (FDR = 0.028) and "TNF signaling" (FDR = 0.023) terms enriched in
this gene set. Supplemental Table E7 contains a complete list of significant functional annotation
terms (FDR < 0.05). These results were supported by changes in ERK1/2 phosphorylation
kinetics in si*SPRY2* cells compared with controls under acute WSP exposure (Supplemental
figure E5). Next, independent comparisons in si*SPRY2* and control cells demonstrated 103
genes with increased expression between 2- and 4-hour WSP exposures in control cells and
180 genes in si*SPRY2* cells (Figure 5B-C). Functional annotation terms from these gene sets
had notable differences in cytokine enrichment, which suggested an interaction between
si*SPRY2* and cytokine gene expression kinetics (Table 3, Supplemental Tables E8-E9).
However, when we tested Interleukin-8 as a representative cytokine in si*SPRY2* and control
cells exposed to WSP for 24 hours, we observed reduced *CXCL8* gene expression and reduced
IL8 cytokine secretion by Enzyme Linked Immunosorbent Assay in WSP-exposed si*SPRY2* cells
(Figure 5D, Supplemental Figure E6). This change was consistent with a reduction in ERK1/2
phosphorylation in si*SPRY2* cells at later time points (Figure 5E), and no significant cytotoxicity
was observed at this time point (Supplemental Figure E7). These results indicated that *SPRY2*
knockdown exhibits variable interactions with cytokine gene expression in Beas-2B cells

exposed to WSP, and sustained WSP exposure results in reduced IL8 secretion and ERK1/2
phosphorylation in cells with *SPRY2* knockdown.
285
SPRY2 knockdown resulted in impaired mechanical scratch repair in Beas-2B cells. Given the
association between *SPRY2* knockdown and reduced IL8 responses to WSP exposure and prior
reports associating *SPRY2* with growth factors and cell proliferation (39–41), we next studied
the interaction between *SPRY2* knockdown and epithelial monolayer injury by mechanical
scratch. In the absence of WSP exposure, we measured the closure of a mechanical scratch
defect in submerged culture of si*SPRY2* and control Beas-2B cells. Scratch area in si*SPRY2*
cells separated from control cells at later time points, with the largest difference at 16 hours
following injury (Figure 6A), consistent with later time points that demonstrated reduced IL8
secretion following WSP exposure. Representative images from one experiment are presented
in Figure 6B. We again observed reduced ERK1/2 phosphorylation in unstimulated proliferating
si*SPRY2* cells compared with controls (Figure 6C). These results established the functional
significance of *SPRY2* knockdown and rs3861144 in airway epithelial cell physiology.
298
SPRY2 is regulated in primary human airway cells. To determine whether *SPRY2* is regulated in
primary human airway cells, we next examined responses in relevant primary human airway
models. We demonstrated that *SPRY2* expression increases in primary SAECs cultured and
exposed to WSP in submerged/basal conditions (Figure 7A). The gene expression kinetics of
SPRY2 were similar to kinetics in Beas-2B cells, peaking at 2 hours and decreasing by 4 hours
of WSP exposure. We also demonstrated that *SPRY2* expression was regulated in primary
human airway smooth muscle (HASM) cells in response to TNF-alpha, a secondary mediator of
WSP released by airway epithelial and innate immune cells, and primary HASM cells from one
asthma donor exhibited reduced *SPRY2* expression (Figure 7B). Finally, we performed *SPRY2*
knockdown using CRISPR-Cas9 in SAECs from one donor (Sm36) and demonstrated a
reduced ratio of phosphorylated to dephosphorylated ERK1/2 in *SPRY2*-knockdown basal
SAECs. (Supplemental Figure E9). Therefore, *SPRY2* is regulated in primary human airway
cells and controls MAPK signaling by regulating ERK1/2 expression and phosphorylation.
312
Discussion
Wildfire related PM2.5 air pollution is increasing and has a direct impact on asthma morbidity
(10,42,43). However, the genetic and molecular mediators of these malign effects are poorly
understood. To address these gaps, we developed a genetic approach to discover novel
relationships between asthma, SNPs, and gene expression responses to WSP exposure in an
airway epithelial cell culture model. Through this approach, we identified 52 common variant
SNPs within WSP-responsive TREs that were nominally associated with an asthma diagnosis in
the GERA cohort. We prioritized SNPs based on primary human airway chromatin accessibility
profiles and bioinformatic predictions of transcription factor binding, and we identified two
functional asthma variants, rs3861144 and rs11832287. Transcriptional regulatory activity of the
regions harboring rs3861144 and rs11832287 was controlled by haplotype, and based on
congruent TRE and gene transcriptional responses, we identified the putative gene targets of
rs3861144 and rs11832287 as *SPRY2* and *CD9*. We confirmed the regulatory association
between rs3861144 and *SPRY2* using a CRISPR-based deletion strategy, and we demonstrated
that *SPRY2* is a plausible candidate for asthma pathobiology through cellular functions that
include regulating ERK1/2 phosphorylation, cytokine secretion, and monolayer scratch repair.
Our findings establish a powerful pipeline for genetic discovery of associations between asthma
and woodsmoke and highlight a novel link between *SPRY2* regulation by WSP and risk of an
asthma diagnosis.
332

Our pipeline for genetic discovery builds on genome-wide association studies by re-analyzing
existing data to narrow the focus to functional variants, which can be more directly connected
with gene targets than statistically prioritized results. We first used PRO-seq-annotated WSP
responses to pre-select SNPs to test for asthma association that reside in the functional centers
of TREs, which are enriched for transcriptional machinery (44,45) and regulatory SNPs (46,47).
This constraint enriched our discovery for functional SNPs and reduced the genomic sequences
for discovery to a total of ~1 Mb. The reduced number of total SNPs tested allowed us to apply
permissive permutation-based statistics and identify variants that would not meet genome-wide
statistical thresholds. Previous studies that have applied genomic annotations to prioritize
functional variants from GWAS results (48–51) are limited in only analyzing variants that meet
these rigorous statistical thresholds and may miss functional variants with weaker statistical
associations. Second, we pre-selected SNPs that reside in WSP-responsive TREs to enrich
genetic discovery for SNPs whose transcriptional regulatory activity depends on WSP exposure
in cell culture models. We further prioritized SNPs based on proximity to regulatory motifs that
are well-established to mediate responses to combusted particles, such as wood smoke, *in vivo*.
While this method does not directly integrate the complexity of environmental exposures with
the genetic risk of asthma, it does provide insight into asthma risk in the context of genomic
responses to relevant exposures.
351
The use of PRO-seq annotations in our pipeline facilitated connecting functional variants with
target genes that respond to pertinent exposures based on transcriptional kinetics. The location
of most genetic risk variants in non-coding and non-annotated genomic regions (52) presents a
general challenge for defining relevant target genes and associated mechanisms from GWAS
results. Expression quantitative trait locus (eQTL) analysis can address this problem (53) but
requires knowledge of the functional cell type and access to large tissue gene expression data
sets. Direct exposure measurements in relationship to gene expression are also challenging for
eQTL studies, and unaccounted variable exposures can confound eQTL results. We connected
functional genetic variants with gene targets by applying PRO-seq data from our cell exposure
model to an established paradigm for regulatory element-gene association (37). The
advantages of our pipeline relative to other methods led to the discovery of rs3861144 and
SPRY2 as novel mechanisms that connect WSP with asthma. Future studies that integrate our
method with existing tools may facilitate additional discovery of variants and target genes that
influence gene-by-environment risk in asthma.
366
We demonstrated the utility of our genetic approach by prioritizing rs3861144 as the functional
variant within its locus and connecting the disease variant with a reduced *SPRY2* gene
expression response to WSP exposure. *SPRY2* is a kinase modulator of the mitogen-activated
protein kinase (MAPK) pathway that plays diverse cellular roles in regulating cell growth,
survival, and proliferation (54–56). Prior reports have connected *SPRY2* gene regulation with
functional consequences (41,57). Previous studies have also reported that genetic signals in the
SPRY2 locus control lung function. Multiple linked SNPs, including rs3861144, were statistically
associated ($p < 5\times10^{-9}$) with measurements of lung function, such as FEV1 (36). However, this
study prioritized rs4885681 as the lead SNP based on the strongest association statistics and
connected rs4885681 with the LINC00382 long non-coding RNA, based on its genomic location.
We demonstrated that rs4885681 did not exhibit features of a functional genetic variant, such as
genomic annotations or transcriptional regulatory function in luciferase reporters. Therefore, the
mechanism derived from rs4885681 was unlikely to explain the connection between genetic
signals at this locus and lung function. In contrast, we prioritized rs3861144 as a functional
variant at this locus, and using nascent transcript annotations and genome editing, we identified
SPRY2 as its gene target and the plausible mechanism that connected the genetic signals in
this locus with lung function. These data demonstrate that defining the functional variants within

a statistically associated genetic locus provides additional utility to GWAS in connecting gene
mechanisms with phenotypes.
386
We also demonstrated changes in airway epithelial cell physiology associated with *SPRY2*
knockdown that suggest rs3861144 relates to asthma mechanisms. In our model system,
sustained WSP exposure resulted in increased *SPRY2* protein expression at latent time points
(8-24 hours), and at these latent time points, *SPRY2* knockdown resulted in reduced IL8
secretion, reduced ERK1/2 phosphorylation, and an impaired ability to close a mechanical
scratch in submerged cell culture. These results were consistent with prior reports in airway
epithelial cells (58) and rhabdomyosarcoma cell lines (59) and suggest that rs3861144 and
SPRY2 have functional roles in the airway epithelium that relate to asthma pathobiology. We
also observed a distinct context associated with acute WSP exposure, where *SPRY2*
knockdown increased ERK1/2 phosphorylation and cytokine gene expression kinetics between
2-6 hours of WSP exposure. Some of these changes, such as increased *CXCL8* gene
expression in *SPRY2* knockdown cells, were not recapitulated in functional signals, such as IL8
secretion, suggesting that the latent program was more relevant to *SPRY2* function. However,
our observations were consistent with prior studies that have reported that *SPRY2* function is
highly dependent on cell context, such as growth factor presence (41), and further study of the
acute context is necessary to resolve the complex roles of *SPRY2* in the airway epithelium.
403
Our study had several limitations. The positive predictive value of our primary discovery in
relationship to our secondary analyses was modest. Only 6/52 SNPs had independent
associations with lung function and 2/52 SNPs were rigorously defined as functional variants.
Low thresholds and limitations in accurately defining TREs bioinformatically and coding the
asthma diagnosis may have contributed to these results, as well as our application of lung
function genetics as a surrogate validation tool for our primary findings. We also performed our
initial genetic analyses in mostly non-Hispanic white cohorts and used a single reference for
linkage disequilibrium analysis, limiting the generalizability of the results and increasing potential
bias. Future validation in an independent multi-ancestry asthma cohort with a whole genome
sequencing data set will improve the rigor and confidence of the results.
414
Our WSP and cell culture exposure system also had limitations. We used aged, desiccated, and
UV-exposed WSP from a single combustion source, and the WSP dose that we selected (1
mg/ml or 0.26 mg/cm^2) lacks a specific *in-vivo* correlation. We cited prior studies of *in-vivo*
exposure (22), comparison with established toxicological doses of TCDD (60,61), and the
induction of relevant pathways, such as aryl hydrocarbon receptor and inflammasome signaling
(62–64), as evidence to support our use of this dose. However, this concentration of WSP
promotes a bias towards cell stress pathways and away from potentially important metabolic
responses, which have also been associated with asthma (65), and limits in the interpretation of
our results overall. Inconsistencies between the Beas-2B immortalized airway epithelial cell line
and normal human airway epithelia, such as supraphysiologic concentrations of growth factors
(66), lack of active metabolic pathways (67), cilia, and mucus functions (68) also resulted in a
bias towards transcriptional programs involved in cell proliferation in our model exposure
system. Lacking from our model were important airway epithelial programs, such as mucus
dysregulation, metabolism, and cell junction integrity, which also contribute to asthma
development (69–72). The acute 30- and 120-minute exposure time points, while potentially
enriched for direct exposure mediators, may fail to capture important sustained or secondary
responses, such as oxidative stress or epithelial-to-mesenchymal transformation, which have
been suggested as mechanisms that connect exposures with asthma previously (68,73). Finally,
our understanding of *SPRY2* was limited to the effects of *SPRY2* knockdown, including its
effects on scratch repair. While *SPRY2*-knockdown recapitulated the effect of the rs3861144 risk

variant, a better understanding of *SPRY2* biology would require additional experiments, such as
SPRY2 overexpression and a detailed analysis of additional factors that interact with *SPRY2*
and ERK1/2 signaling. Our study proposes several avenues for further investigation into this
complex but relevant pathway in airway cell physiology and asthma.
439
Despite these limitations, our study expands the understanding of the genetic factors that
influence asthma risk in the context of wildfire PM2.5 exposure, a problem of growing
importance due to global climate change. We identified common SNPs that may contribute to
the personal risk of asthma associated with wildfire PM2.5 and cellular mechanisms that
mediate this risk. We characterized one important mechanism by demonstrating that *SPRY2*
expression in response to WSP exposure is mediated by rs3861144, which modifies asthma risk
and lung function. This connection between WSP and *SPRY2* may alter asthma risk by
impacting airway epithelial cell proliferation and injury repair through the MAPK pathway or
increasing pro-inflammatory responses to exposures. Irrespective of the precise mechanism,
our results implicate *SPRY2* as a mediator of the adverse effects of wildfire PM2.5 exposure on
asthma risk and define *SPRY2* as a potential therapeutic target for asthma.

Table 1 – WSP-SNPs

SNP	CHR	BP (HG38)	MAF	OR	FDR
rs16860727	1	183272646	0.12620	1.049	0.043200
rs11590328	1	209399237	0.3555	1.045	0.007899
rs75459144	10	75294962	0.15020	0.941	0.006199
rs6584144	10	97700730	0.19740	0.9563	0.028900
rs138744334	11	94643137	0.0694	0.938	0.045100
rs12321808	12	6167792	0.37620	1.037	0.024000
rs11832287	12	6182141	00.46510	0.9602	0.011800
rs10747612	12	51394035	0.37200	0.9623	0.020300
rs9509353	13	20722128	0.4838	0.9683	0.042700
rs17065695	13	44310244	0.0598	0.9236	0.015800
rs3861144	13	79936471	0.26700	1.036	0.042800
rs1275823	14	69726964	0.13050	0.9479	0.021300
rs9939094	16	29268980	0.35850	0.9569	0.008099
rs253140	16	83963743	0.11640	1.068	0.008099
rs3968348	17	2732850	0.4839	0.9575	0.007999
rs60701308	17	4858995	0.0946	0.9418	0.025700
rs9896022	17	16405309	0.12300	1.052	0.037300
rs10491130	17	41377775	0.10090	0.9437	0.029200
rs9903086	17	41793264	0.0625	0.9322	0.034900
rs3848425	17	66467803	0.39440	1.035	0.033900
rs644821	18	11954652	0.10280	0.9441	0.027400
rs4806835	19	2266880	0.13350	1.054	0.031300
rs10424200	19	38792918	0.27260	1.039	0.036800
rs10418662	19	46426768	0.34810	1.04	0.020100
rs7246311	19	680378	0.4121	1.05	0.024000
rs770665	2	197729533	0.3457	1.055	0.001400
rs7591472	2	201025028	0.17090	1.044	0.044000
rs11883931	2	237315599	0.21540	1.046	0.022500
rs10208406	2	10057488	0.19520	0.9596	0.041000
rs2139379	2	12574801	0.17300	0.9573	0.038000
rs28362965	2	74845567	0.0516	1.078	0.031600
rs6023060	20	38109598	0.17150	1.047	0.028100
rs764025	22	33853729	0.22390	0.9545	0.019000
rs2173620	3	177538135	0.23200	0.959	0.026500
rs7428132	3	177745986	0.0541	0.9083	0.007299
rs6784916	3	189937753	0.35420	0.9612	0.017000
rs12630368	3	5013455	0.11000	1.052	0.045400
rs2633783	3	9560123	0.1346	0.9438	0.012100
rs56984996	4	168138760	0.36030	1.041	0.016100
rs72705761	4	185847661	0.29980	1.043	0.016800
rs7669195	4	187421022	0.24990	0.9633	0.039100

rs6448484	4	26857636	0.4736	1.051	0.001900
rs1371799	4	74112120	0.38900	0.9682	0.045900
rs28517751	5	120624434	0.08031	0.9169	0.003900
rs162481	5	123719195	0.19460	0.9445	0.004400
rs6871945	5	56278753	0.27250	0.9601	0.023600
rs1873333	6	168008153	0.2205	1.049	0.012900
rs3129975	6	30764277	0.21440	1.075	0.000400
rs979701	6	77034107	0.24350	1.038	0.044100
rs524953	9	110844289	0.2588	0.9593	0.025500
rs7038732	9	127764264	0.41900	0.967	0.036200
rs3739819	9	127965830	0.0989	1.067	0.016700

453 Table 2 – WSP-SNPs with significant associations in lung function GWAS

		Open Targets Genetics	Lung Disease Knowledge Portal
SNP	**Association**	**P value**	**P value**
rs3861144	FEV1	6.80E-10	8.79E-10
rs28517751	FEV1/FVC	4.20E-08	4.12E-08
rs3129975	PEF	1.30E-26	1.74E-25
rs11883931	FEV1/FVC	3.90E-05	3.36E-05
rs16860727	FEV1/FVC	7.60E-06	3.80E-06
rs1873333	PEF	6.50E-04	8.53E-04

455 Table 3 – Functional annotation categories for genes upregulated between 2 and 4 hours of
456 WSP exposure in control (siCTRL) and *SPRY2* knockdown (si*SPRY2*) cells

siCTRL WSP4h > WSP2h		
Term	**Enrichment score**	**FDR**
Secreted	2.3	0.031
Extracellular Matrix	2	0.025
Benzodiazepine Receptor Binding	1.5	0.031
si*SPRY2* WSP4h > WSP2h		
Term	**Enrichment score**	**FDR**
IL-17 signaling	4.2	2.00E-05
Cytokine	3.7	0.00098
NFkB	3.5	0.024
Secreted	3.4	0.00075
Cell Adhesion	2.3	0.006
Small Cell Lung Cancer	2.3	0.0058
Positive regulation of angiogenesis	2.3	0.024
RAGE signaling	2.1	0.0081
PI3K-AKT signaling	2	0.036
NOD-like receptor signaling	2	0.034
RHD DNA binding domain	2	0.0063

Inflammatory Bowel Disease	2	0.038
Extracellular Matrix	1.7	0.044
Amoebiasis	1.7	0.0014
Neutrophil Chemotaxis	1.5	0.049

Figure legends
Figure 1 – A permutation statistical approach identifies 52 common single nucleotide
polymorphisms (SNPs) that reside within transcriptional regulatory elements (TREs) that
respond to wood smoke particle (WSP) exposure and associate with asthma diagnosis. A.
Schematic of the approach that applies WSP-dependent TREs identified by Precision Run-on
sequencing (PRO-seq) in Beas-2B airway epithelial cells to asthma variant discovery. B.
Genomic features of the 52 WSP-SNPs. C. Representative example of WSP-SNP which resides
within an intronic TRE in the *HK2* gene. PRO-seq data is visualized in the Integrative Genomics
Viewer (IGV) genome browser based on counts per million mapped reads (vertical scales).
Positive (blue) signal indicates reads annotated to the sense strand while negative (red) signal
reflects reads annotated to the antisense strand. The Transcription Start Site (TSS) and
direction of transcription are indicated by the arrow at the top of the panel.
471
Figure 2 – Subsets of the 52 WSP-SNPs exhibit biologic functions associated with asthma. A.
Venn diagram demonstrates 23 genomic regions that overlap (> 1 bp), based on genomic
coordinates, between WSP-SNPs and chromatin accessibility features in live proximal airway
epithelial tissue obtained from healthy non-smoking subjects. B. The representative example
rs9903086 WSP-SNP resides in an intergenic region near the *JUP* TSS and overlaps with
bidirectional PRO-seq signal in Beas-2B cells and chromatin accessibility signal in live airway
epithelium. PRO-seq (blue and red) and ATAC-seq (green and teal) data visualized in the IGV
genome browser based on counts per million mapped reads (vertical scales). The TSS is
indicated by the arrow at the top of the panel. C. Venn diagram demonstrating sites of genomic
overlap between a 500 bp radius surrounding each WSP-SNP and consensus transcription
factor binding motifs for aryl hydrocarbon receptor (AHR) and nuclear factor kappa b (NFkB)
family transcription factors. Twenty-three WSP-SNP regions contain motifs from both families.
D. A representative example at the rs11883931 locus, which contains both ARNT2 (AHR nuclear
transporter) and REL (NFkB family) transcription factor binding motifs.
486
Figure 3 – Luciferase activity in plasmid reporter systems demonstrates that WSP-SNPs reside
in active TREs in comparison with local lead variants predicted by GWAS. A. Schematic for
selection of WSP-SNP candidates prioritized by biologic variables of live airway chromatin
accessibility and colocalization with both AHR and NFkB family transcription factor binding
motifs. B. Normalized luciferase activity of plasmid reporters transfected into Beas-2B cells and
treated with vehicle (veh) or WSP (1 mg/ml) for 8 hrs. A 700 – 800 bp genomic region
surrounding each WSP-SNP was cloned into the plasmid system and mutated for allele change,
with "WT" indicating the allele native to Beas-2B cells and genotype letter change indicating the
mutated reporter. Bars indicate mean reporter activity (+/- standard deviation, n = 4) relative to
an internal control Renilla luciferase that was co-transfected into cells; * $p < 0.01$. C. Normalized
luciferase activity of plasmid reporters containing the WSP-SNP rs3861144 and rs4885681, the
lead variant at this locus prioritized by GWAS statistics. A 350 – 400 bp radius genomic region
surrounding each SNP was cloned into the plasmid system, and luciferase activity was
measured after transfection into Beas-2B cells and treatment with vehicle (veh) or WSP (1
mg/ml) for 8 hrs. Data are presented as described for (B). D. Beas-2B PRO-seq and live airway
epithelium ATAC-seq signal at the WSP-SNP rs3861144 and rs4885681 loci, located
approximately 380 kb 3' to the *SPRY2* gene. E. Normalized luciferase activity of plasmid
reporters containing a 350-400 bp radius around the WSP-SNP rs9903086 and three SNPs
(rs9913846, rs991655 and rs39914436) prioritized by statistical fine mapping (FINEMAP (35))
for association with Forced Vital Capacity using summary statistics from an independent data
set (30). Bars indicate mean reporter activity (+/- standard deviation, n = 4) relative to Renilla; *
$p < 0.01$.
509

Figure 4 – The rs3861144 functional variant resides in a TRE that regulates *SPRY2* in response
to WSP exposure in Beas-2B cells. A. Beas-2B PRO-seq signal visualized in IGV at the
rs3861144 and *SPRY2* gene locus demonstrates shared transcriptional kinetics between these
regions at 30 and 120 minutes of WSP exposure. Below – green arrows indicate two target
guide RNA (gRNA) sites designed for Cas9 genome editing of the rs3861144 TRE; purple
arrows indicate target sites for polymerase chain reaction (PCR) primers used to amplify the
region following Cas9 editing and confirm genomic deletion. B. Beas-2B cells were transfected
with two on-target Cas9-gRNA RNP complexes indicated by green arrow sites in the rs3861144
TRE in (A) (rs3861144-Cas9) or negative control Cas9-gRNA RNP (NC). Gel electrophoresis
confirmed genomic deletion at the rs3861144 target region in cells transfected with targeting
gRNAs. C. CRISPR-Cas9 transfected cells were exposed to WSP at 1 mg/ml (0.26 mg/cm^2) for
2 hours. Gene expression of the canonical WSP target *CYP1A1* and *SPRY2* were measured
using quantitative PCR (qPCR). Bars represent mean normalized C_T values on a $\log_2$ scale (+/-
standard deviation, n = 4) relative to vehicle-treated controls; * $p < 0.01$.

Figure 5 – *SPRY2* knockdown by siRNA results in acute and latent responses to WSP
exposure. A. Volcano plot from RNA-sequencing data demonstrates differentially expressed
genes between WSP-exposed (4 hours) non-targeting control (siCTRL) or *SPRY2*-targeting
siRNA (si*SPRY2*) transfected Beas-2B cells. Each point represents a single gene, with the x-
axis showing $\log_2$ fold change and the y-axis showing the log transformed p_{adj} statistic. Dashed
lines indicate $\log_2$ fold change of 0.5, which was the threshold used for this analysis. Number of
differentially expressed genes ($p_{adj} < 0.05$ and $\log_2$ fold change > 0.5) are listed below the plot.
B. Volcano plot demonstrates differentially expressed genes between 2- and 4-hour WSP
exposure in siCTRL cells. C. Volcano plot demonstrates differentially expressed genes between
2- and 4-hour WSP exposure in si*SPRY2* cells. D. ELISA analysis of IL-8 concentration in the
supernatant of siCTRL and si*SPRY2* Beas-2B cells exposed to WSP (1 mg/ml of cell culture
media or 0.26 mg/cm^2 cell culture surface) at latent time points. Bars indicate mean IL-8
concentration (+/- SD); n = 4/group, *$p < 1\times10^{-4}$ for indicated comparisons. E. Western blot
demonstrates knockdown in *SPRY2* expression with *SPRY2*-siRNA transfection (si*SPRY2*) is
associated with reduced phosphorylated ERK1/2 in vehicle- and WSP-exposed (24 hour) Beas-
2B cells compared with control-transfected cells (siCTRL). GAPDH served as the loading
control.

Figure 6 – *SPRY2* knockdown by siRNA delays scratch repair in a Beas-2B monolayer. A.
Scratch assay in siCTRL or si*SPRY2* Beas-2B cells demonstrated a larger remaining defect at
8, 12, and 16 hours in cells with *SPRY2* knockdown. Solid line depicts the interpolated slope
between the means of measurements at 2, 6, 8, 10, 12, 16 and 25 hours. Individual
measurements are plotted as blue (non-targeting control "siCTRL") or red (*SPRY2*-siRNA)
points (n=6-7/condition). * $p < 0.01$, ** $p < 0.05$. B. Representative images from three wells of
siCTRL and si*SPRY2* scratch assay at three time points. C. Western blot demonstrates
knockdown in *SPRY2* expression in si*SPRY2* cells is associated with a reduction in
phosphorylated ERK1/2 signal in siCTRL cells. GAPDH served as the loading control.

Figure 7 – *SPRY2* is expressed and regulated in primary human airway cells. A. Primary small
airway epithelial cells obtained from four healthy, non-smoking donors were exposed to WSP (1
mg/ml, 0.26 mg/cm^2) in submerged culture for 2 or 4 hours. Gene expression for *SPRY2* was
measured using qPCR. Bars represent mean normalized C_T values on a $\log_2$ scale (+/- standard
deviation, n = 4) relative to vehicle-treated controls. B. *SPRY2* expression in normal HASM cells
is increased by TNF-alpha stimulation but reduced in cells from an asthmatic donor. GAPDH
was the loading control.
560

References

1. Westerling ALR. Increasing western US forest wildfire activity: Sensitivity to changes in the timing of spring. Philos Trans R Soc B Biol Sci [Internet]. 2016 Jun 5 [cited 2020 Oct 13];371(1696). Available from: /pmc/articles/PMC4874415/?report=abstract

2. Burke M, Childs ML, de la Cuesta B, Qiu M, Li J, Gould CF, et al. The contribution of wildfire to PM2.5 trends in the USA. Nature [Internet]. 2023 Oct 26 [cited 2024 Nov 26];622(7984):761–6. Available from: https://pubmed.ncbi.nlm.nih.gov/37730996/

3. Childs ML, Li J, Wen J, Heft-Neal S, Driscoll A, Wang S, et al. Daily Local-Level Estimates of Ambient Wildfire Smoke PM2.5for the Contiguous US. Environ Sci Technol [Internet]. 2022 Oct 4 [cited 2024 Nov 26];56(19):13607–21. Available from: https://pubs.acs.org/doi/abs/10.1021/acs.est.2c02934

4. Cunningham CX, Williamson GJ, Bowman DMJS. Increasing frequency and intensity of the most extreme wildfires on Earth. Nat Ecol Evol 2024 88 [Internet]. 2024 Jun 24 [cited 2024 Nov 26];8(8):1420–5. Available from: https://www.nature.com/articles/s41559-024-02452-2

5. Foundation FS. First Street Foundation Aggregated Wildfire Risk Summary Statistics Version 1.0 [Internet]. Zenodo; 2022. Available from: https://doi.org/10.5281/zenodo.6564731

6. Gould CF, Heft-Neal S, Johnson M, Aguilera J, Burke M, Nadeau K. Health Effects of Wildfire Smoke Exposure. Annu Rev Med [Internet]. 2024 Jan 29 [cited 2024 Nov 26];75(Volume 75, 2024):277–92. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-med-052422-020909

7. Gao Y, Huang W, Yu P, Xu R, Yang Z, Gasevic D, et al. Long-term impacts of non-occupational wildfire exposure on human health: A systematic review. Environ Pollut. 2023 Mar 1;320:121041.

8. Skinner R, Luther M, Hertelendy AJ, Khorram-Manesh A, Sorensen J, Goniewicz K, et al. A Literature Review on the Impact of Wildfires on Emergency Departments: Enhancing Disaster Preparedness. Prehosp Disaster Med [Internet]. 2022 Oct 25 [cited 2024 Nov 26];37(5):657. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC9470518/

9. Beyene T, Harvey ES, Van Buskirk J, McDonald VM, Jensen ME, Horvat JC, et al. 'Breathing Fire': Impact of Prolonged Bushfire Smoke Exposure in People with Severe Asthma. Int J Environ Res Public Health [Internet]. 2022 Jun 1 [cited 2023 Oct 2];19(12):7419. Available from: /pmc/articles/PMC9224478/

10. Noah TL, Worden CP, Rebuli ME, Jaspers I. The Effects of Wildfire Smoke on Asthma and Allergy. Curr Allergy Asthma Rep [Internet]. 2023 Jul 1 [cited 2024 Nov 26];23(7):1. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10176314/

11. Gehring U, Wijga AH, Koppelman GH, Vonk JM, Smit HA, Brunekreef B. Air pollution and the development of asthma from birth until young adulthood. 2020 Jul 1 [cited 2023 Jan 29];56(1). Available from: https://pubmed.ncbi.nlm.nih.gov/32299858/

12. McConnell R, Berhane K, Yao L, Jerrett M, Lurmann F, Gilliland F, et al. Traffic, susceptibility, and childhood asthma. Environ Health Perspect [Internet]. 2006 May [cited 2024 Nov 26];114(5):766–72. Available from: https://ehp.niehs.nih.gov/doi/10.1289/ehp.8594

13. Gauderman WJ, McConnell R, Gilliland F, London S, Thomas D, Avol E, et al. Association between Air Pollution and Lung Function Growth in Southern California Children. https://doi.org/101164/ajrccm16249909096 [Internet]. 2012 Dec 14 [cited 2025 May 9];162(4 I):1383–90. Available from: /doi/pdf/10.1164/ajrccm.162.4.9909096?download=true

14. Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical Review of Health Impacts of Wildfire Smoke Exposure. Environ Health Perspect [Internet]. 2016 Sep [cited 2019 Jun 25];124(9):1334–43. Available from:

612   http://www.ncbi.nlm.nih.gov/pubmed/27082891

15. Thomsen SF, Ulrik CS, Kyvik KO, Hjelmborg J von B, Skadhauge LR, Steffensen I, et al. Genetic and environmental contributions to hay fever among young adult twins. Respir Med [Internet]. 2006 Dec [cited 2024 Nov 26];100(12):2177–82. Available from: https://pubmed.ncbi.nlm.nih.gov/16650971/

16. FRYER AA, BIANCO A, HEPPLE M, JONES PW, STRANGE RC, SPITERI MA. Polymorphism at the Glutathione S-transferase GSTP1 Locus. Am J Respir Crit Care Med [Internet]. 2000;161(5):1437–42. Available from: https://doi.org/10.1164/ajrccm.161.5.9903006

17. Goodrich GG, Goodman PH, Budhecha SK, Pritsos CA. Functional polymorphism of detoxification gene NQO1 predicts intensity of empirical treatment of childhood asthma. Mutat Res Toxicol Environ Mutagen. 2009 Mar 31;674(1–2):55–61.

18. Strange RC, Spiteri MA, Ramachandran S, Fryer AA. Glutathione-S-transferase family of enzymes. Mutat Res Mol Mech Mutagen. 2001 Oct 1;482(1–2):21–6.

19. Melbourne CA, Mesut Erzurumluoglu A, Shrine N, Chen J, Tobin MD, Hansell AL, et al. Genome-wide gene-air pollution interaction analysis of lung function in 300,000 individuals. Environ Int [Internet]. 2022 Jan 15 [cited 2024 Nov 26];159:107041. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC8739564/

20. Zhu Y, Tazearslan C, Suh Y. Challenges and progress in interpretation of non-coding genetic variants associated with human disease. Exp Biol Med [Internet]. 2017 Jul 1 [cited 2024 Nov 26];242(13):1325. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5529005/

21. Sasse SK, Dahlin A, Sanford L, Gruca MA, Gupta A, Gally F, et al. Enhancer RNA transcription pinpoints functional genetic variants linked to asthma. Nat Commun 2025 161 [Internet]. 2025 Mar 31 [cited 2025 Apr 29];16(1):1–14. Available from: https://www.nature.com/articles/s41467-025-57693-x

22. Ghio AJ, Soukup JM, Case M, Dailey LA, Richards J, Berntsen J, et al. Exposure to wood smoke particles produces inflammation in healthy volunteers. Occup Environ Med [Internet]. 2012 Mar [cited 2019 Jun 18];69(3):170–5. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21719562

23. Ghio AJ, Soukup JM, Dailey LA, Tong H, Kesic MJ, Budinger GRS, et al. Wood Smoke Particle Sequesters Cell Iron to Impact a Biological Effect. Chem Res Toxicol [Internet]. 2015 Nov 16 [cited 2019 Jul 3];28(11):2104–11. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26462088

24. Dahlin A, Sordillo JE, Ziniti J, Iribarren C, Lu M, Weiss ST, et al. Large-scale, multi-ethnic genome-wide association study identifies novel loci contributing to asthma susceptibility in adults. J Allergy Clin Immunol [Internet]. 2019 Apr 1 [cited 2021 Jul 13];143(4):1633. Available from: /pmc/articles/PMC6451881/

25. Gupta A, Sasse SK, Gruca MA, Sanford L, Dowell RD, Gerber AN. Deconvolution of multiplexed transcriptional responses to wood smoke particles defines rapid aryl hydrocarbon receptor signaling dynamics. J Biol Chem [Internet]. 2021 Oct 1;297(4). Available from: https://doi.org/10.1016/j.jbc.2021.101147

26. Koh KD, Siddiqui S, Cheng D, Bonser LR, Sun DI, Zlock LT, et al. Efficient RNP-directed human gene targeting reveals SPDEF is required for IL-13-induced mucostasis. Am J Respir Cell Mol Biol. 2020 Mar 1;62(3):373–81.

27. Ghoussaini M, Mountjoy E, Carmona M, Peat G, Schmidt EM, Hercules A, et al. Open Targets Genetics: systematic identification of trait-associated genes using large-scale genetics and functional genomics. Nucleic Acids Res [Internet]. 2021 Jan 8 [cited 2024 Feb 21];49(D1):D1311–20. Available from: https://pubmed.ncbi.nlm.nih.gov/33045747/

28. Mountjoy E, Schmidt EM, Carmona M, Schwartzentruber J, Peat G, Miranda A, et al. An open approach to systematically prioritize causal variants and genes at all published

human GWAS trait-associated loci. Nat Genet [Internet]. 2021 Nov 1 [cited 2024 Feb 21];53(11):1527–33. Available from: https://pubmed.ncbi.nlm.nih.gov/34711957/

29. Lung Disease Knowledge Portal (lungdiseasegenetics.org) [Internet]. [cited 2024 Nov 12]. Available from: https://lung.hugeamp.org/

30. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, et al. New genetic signals for lung function highlight pathways and chronic obstructive pulmonary disease associations across multiple ancestries. Nat Genet [Internet]. 2019 Feb 25 [cited 2022 Dec 27];51(3):481–93. Available from: https://europepmc.org/articles/PMC6397078

31. Shrine N, Izquierdo AG, Chen J, Packer R, Hall RJ, Guyatt AL, et al. Multi-ancestry genome-wide association analyses improve resolution of genes and pathways influencing lung function and chronic obstructive pulmonary disease risk. Nat Genet 2023 553 [Internet]. 2023 Mar 13 [cited 2024 Nov 12];55(3):410–22. Available from: https://www.nature.com/articles/s41588-023-01314-0

32. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc [Internet]. 2009 [cited 2021 Jan 21];4(1):44–57. Available from: https://pubmed.ncbi.nlm.nih.gov/19131956/

33. Vogel CFA, Van Winkle LS, Esser C, Haarmann-Stemmann T. The aryl hydrocarbon receptor as a target of environmental stressors – Implications for pollution mediated stress and inflammatory responses [Internet]. Vol. 34, Redox Biology. Elsevier B.V.; 2020 [cited 2020 Sep 29]. Available from: /pmc/articles/PMC7327980/?report=abstract

34. Cui C, Yang R, Chen H, Li D, Sun X, Wang Y, et al. Air toxins disorder the NF-kB Pathway leads to immune disorders and immune diseases in the human health. Ecotoxicol Environ Saf [Internet]. 2025 Sep 1 [cited 2025 Jul 18];302:118474. Available from: https://www.sciencedirect.com/science/article/pii/S0147651325008140

35. Benner C, Spencer CCA, Havulinna AS, Salomaa V, Ripatti S, Pirinen M. FINEMAP: efficient variable selection using summary data from genome-wide association studies. Bioinformatics [Internet]. 2016 May 5 [cited 2024 Feb 21];32(10):1493. Available from: /pmc/articles/PMC4866522/

36. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, et al. New genetic signals for lung function highlight pathways and chronic obstructive pulmonary disease associations across multiple ancestries. Nat Genet [Internet]. 2019 Mar 1 [cited 2024 Nov 26];51(3):481–93. Available from: https://pubmed.ncbi.nlm.nih.gov/30804560/

37. Fulco CP, Nasser J, Jones TR, Munson G, Bergman DT, Subramanian V, et al. Activity-by-Contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. Nat Genet [Internet]. 2019 Dec 1 [cited 2024 Feb 7];51(12):1664. Available from: /pmc/articles/PMC6886585/

38. El Kharbili M, Sasse SK, Sanford L, Jacobson S, Aviszus K, Gupta A, et al. Noncoding SNPs decrease expression of FABP5 during COPD exacerbations. J Clin Invest. 2024 Feb 1;134(3).

39. Chandramouli S, Chye YY, Yusoff P, Lao DH, Hwei FL, Mizuno K, et al. Tesk1 Interacts with Spry2 to Abrogate Its Inhibition of ERK Phosphorylation Downstream of Receptor Tyrosine Kinase Signaling. J Biol Chem. 2008 Jan 18;283(3):1679–91.

40. Ding W, Warburton D. Down-regulation of Sprouty2 via p38 MAPK plays a key role in the induction of cellular apoptosis by tumor necrosis factor-α. Biochem Biophys Res Commun [Internet]. 2008 Oct 24 [cited 2025 Apr 21];375(3):460. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2575769/

41. Masoumi-Moghaddam S, Amini A, Morris DL. The developing story of Sprouty and cancer. Cancer Metastasis Rev [Internet]. 2014 Apr 18 [cited 2025 Apr 24];33(2–3):695–720. Available from: https://link.springer.com/article/10.1007/s10555-014-9497-1

42. Cascio WE. Wildland fire smoke and human health. 2018 May 15 [cited 2019 Jun 25];624:586–95. Available from: /pmc/articles/PMC6697173/?report=abstract

43. McArdle CE, Dowling TC, Carey K, DeVies J, Johns D, Gates AL, et al. Asthma-Associated Emergency Department Visits During the Canadian Wildfire Smoke Episodes — United States, April– August 2023. MMWR Morb Mortal Wkly Rep [Internet]. 2023 Aug 25 [cited 2023 Oct 2];72(34):926–32. Available from: https://www.cdc.gov/mmwr/volumes/72/wr/mm7234a5.htm

44. Panigrahi A, O'Malley BW. Mechanisms of enhancer action: the known and the unknown. Genome Biol 2021 221 [Internet]. 2021 Apr 15 [cited 2024 Nov 26];22(1):1–30. Available from: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-021-02322-1

45. Gasperini M, Tome JM, Shendure J. Towards a comprehensive catalogue of validated and target-linked human enhancers. Nat Rev Genet [Internet]. 2020 May 1 [cited 2024 Nov 26];21(5):292. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC7845138/

46. Boix CA, James BT, Park YP, Meuleman W, Kellis M. Regulatory genomic circuitry of human disease loci by integrative epigenomics. Nat 2021 5907845 [Internet]. 2021 Feb 3 [cited 2024 Nov 26];590(7845):300–7. Available from: https://www.nature.com/articles/s41586-020-03145-z

47. Gusev A, Lee SH, Trynka G, Finucane H, Vilhjálmsson BJ, Xu H, et al. Partitioning heritability of regulatory and cell-type-specific variants across 11 common diseases. Am J Hum Genet [Internet]. 2014 [cited 2024 Nov 26];95(5):535–52. Available from: https://pubmed.ncbi.nlm.nih.gov/25439723/

48. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. Nature [Internet]. 2015 Feb 2 [cited 2024 Feb 25];518(7539):317. Available from: /pmc/articles/PMC4530010/

49. Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, et al. An integrated encyclopedia of DNA elements in the human genome. Nat 2012 4897414 [Internet]. 2012 Sep 5 [cited 2024 Feb 25];489(7414):57–74. Available from: https://www.nature.com/articles/nature11247

50. Kircher M, Witten DM, Jain P, O'roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. Nat Genet 2014 463 [Internet]. 2014 Feb 2 [cited 2024 Feb 25];46(3):310–5. Available from: https://www.nature.com/articles/ng.2892

51. Ionita-Laza I, Mccallum K, Xu B, Buxbaum JD. A spectral approach integrating functional genomic annotations for coding and noncoding variants. Nat Genet 2015 482 [Internet]. 2016 Jan 4 [cited 2024 Feb 25];48(2):214–20. Available from: https://www.nature.com/articles/ng.3477

52. Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, et al. Transcriptional super-enhancers connected to cell identity and disease. Cell [Internet]. 2013 Nov 7 [cited 2025 May 7];155(4):10.1016/j.cell.2013.09.053. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC3841062/

53. Gibson G, Powell JE, Marigorta UM. Expression quantitative trait locus analysis for translational medicine. Genome Med 2015 71 [Internet]. 2015 Jun 24 [cited 2024 Nov 26];7(1):1–14. Available from: https://genomemedicine.biomedcentral.com/articles/10.1186/s13073-015-0186-7

54. Edwin F, Anderson K, Ying C, Patel TB. Intermolecular Interactions of Sprouty Proteins and Their Implications in Development and Disease. Mol Pharmacol [Internet]. 2009 Oct [cited 2025 Apr 24];76(4):679. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2769046/

55. Hacohen N, Kramer S, Sutherland D, Hiromi Y, Krasnow MA. sprouty Encodes a Novel Antagonist of FGF Signaling that Patterns Apical Branching of the Drosophila Airways. Cell. 1998 Jan 23;92(2):253–63.

765   56.   Walker DJ, Land SC. Regulation of vascular signalling by nuclear Sprouty2 in fetal lung
766   epithelial cells: Implications for co-ordinated airway and vascular branching in lung
767   development. Comp Biochem Physiol Part B Biochem Mol Biol. 2018 Oct 1;224:105–14.

768   57.   Clarke M, Volpe G, Sheriff L, Walton D, Ward C, Wei W, et al. Transcriptional regulation
769   of SPROUTY 2 by MYB influences myeloid cell proliferation and stem cell properties by
770   enhancing responsiveness to IL-3. Leukemia [Internet]. 2016 Apr 1 [cited 2025 May
771   7];31(4):957. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5302168/

772   58.   Liu W, Tundwal K, Liang Q, Goplen N, Rozario S, Quayum N, et al. Establishment of
773   Extracellular Signal-Regulated Kinase 1/2 Bistability and Sustained Activation through
774   Sprouty 2 and Its Relevance for Epithelial Function. Mol Cell Biol [Internet]. 2010 Apr 1
775   [cited 2024 Mar 26];30(7):1783. Available from: /pmc/articles/PMC2838067/

776   59.   Saini M, Verma A, Mathew SJ. SPRY2 is a novel MET interactor that regulates metastatic
777   potential and differentiation in rhabdomyosarcoma. Cell Death Dis 2018 92 [Internet].
778   2018 Feb 14 [cited 2025 Apr 24];9(2):1–15. Available from:
779   https://www.nature.com/articles/s41419-018-0261-2

780   60.   Abraham K, Krowke R, Neubert D. Pharmacokinetics and biological activity of 2,3,7,8-
781   tetrachlorodibenzo-p-dioxin - 1. Dose-dependent tissue distribution and induction of
782   hepatic ethoxyresorufin O-deethylase in rats following a single injection. Arch Toxicol
783   [Internet]. 1988 Dec [cited 2025 Jul 18];62(5):359–68. Available from:
784   https://pubmed.ncbi.nlm.nih.gov/3242446/

785   61.   Harvey WA, Jurgensen K, Pu X, Lamb CL, Cornell KA, Clark RJ, et al. Exposure to
786   2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) increases human hepatic stellate cell
787   activation. Toxicology [Internet]. 2016 Feb 17 [cited 2025 Jul 18];344–346:26. Available
788   from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4792734/

789   62.   Guerrina N, Traboulsi H, Eidelman DH, Baglole CJ. The Aryl Hydrocarbon Receptor and
790   the Maintenance of Lung Health. Int J Mol Sci [Internet]. 2018 Dec 1 [cited 2022 Nov
791   8];19(12). Available from: http://www.ncbi.nlm.nih.gov/pubmed/30563036

792   63.   Iu M, Zago M, Rico de Souza A, Bouttier M, Pareek S, White JH, et al. RelB attenuates
793   cigarette smoke extract-induced apoptosis in association with transcriptional regulation of
794   the aryl hydrocarbon receptor. Free Radic Biol Med [Internet]. 2017 Jul 1 [cited 2020 Feb
795   20];108:19–31. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28254546

796   64.   Wegesser TC, Pinkerton KE, Last JA. California wildfires of 2008: Coarse and fine
797   particulate matter toxicity. Environ Health Perspect [Internet]. 2009 [cited 2020 Sep
798   30];117(6):893–7. Available from: /pmc/articles/PMC2702402/?report=abstract

799   65.   Líbalová H, Závodná T, Vrbová K, Sikorová J, Vojtíšek-Lom M, Beránek V, et al.
800   Transcription profiles in BEAS-2B cells exposed to organic extracts from particulate
801   emissions produced by a port-fuel injection vehicle, fueled with conventional fossil
802   gasoline and gasoline-ethanol blend. Mutat Res Toxicol Environ Mutagen [Internet]. 2021
803   Dec 1 [cited 2025 Apr 25];872:503414. Available from:
804   https://www.sciencedirect.com/science/article/pii/S1383571821001054?via%3Dihub

805   66.   Zhao F, Klimecki WT. CULTURE CONDITIONS PROFOUNDLY IMPACT PHENOTYPE
806   IN BEAS-2B, A HUMAN PULMONARY EPITHELIAL MODEL. J Appl Toxicol [Internet].
807   2014 Aug 1 [cited 2025 May 7];35(8):945. Available from:
808   https://pmc.ncbi.nlm.nih.gov/articles/PMC4474793/

809   67.   Garcia-Canton C, Minet E, Anadon A, Meredith C. Metabolic characterization of cell
810   systems used in in vitro toxicology testing: Lung cell system BEAS-2B as a working
811   example. Toxicol Vitr [Internet]. 2013 Sep 1 [cited 2025 Apr 25];27(6):1719–27. Available
812   from: https://www.sciencedirect.com/science/article/pii/S0887233313001239

813   68.   Han X, Na T, Wu T, Yuan BZ. Human lung epithelial BEAS-2B cells exhibit
814   characteristics of mesenchymal stem cells. PLoS One [Internet]. 2020 Jan 1 [cited 2025
815   Apr 25];15(1):e0227174. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC6941928/

69.   Georas SN, Rezaee F. Epithelial barrier function: at the frontline of asthma immunology
and allergic airway inflammation. J Allergy Clin Immunol [Internet]. 2014 Sep 1 [cited
2025 May 7];134(3):509. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC4170838/

70.   Gref A, Merid SK, Gruzieva O, Ballereau S, Becker A, Bellander T, et al. Genome-wide
interaction analysis of air pollution exposure and childhood asthma with functional follow-
up. Am J Respir Crit Care Med [Internet]. 2017 May 15 [cited 2024 Feb
25];195(10):1373–83. Available from: /pmc/articles/PMC5443897/

71.   Melén E, Kho AT, Sharma S, Gaedigk R, Leeder JS, Mariani TJ, et al. Expression
analysis of asthma candidate genes during human and murine lung development. Respir
Res 2011 121 [Internet]. 2011 Dec 1 [cited 2025 May 7];12(1):1–10. Available from:
https://respiratory-research.biomedcentral.com/articles/10.1186/1465-9921-12-86

72.   Evans CM, Kim K, Tuvim MJ, Dickey BF. Mucus hypersecretion in asthma: causes and
effects. Curr Opin Pulm Med [Internet]. 2009 Jan [cited 2025 May 7];15(1):4. Available
from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2709596/

73.   Grilli A, Bengalli R, Longhin E, Capasso L, Proverbio MC, Forcato M, et al.
Transcriptional profiling of human bronchial epithelial cell BEAS-2B exposed to diesel and
biomass ultrafine particles. BMC Genomics [Internet]. 2018 Apr 27 [cited 2025 Apr
25];19(1):302. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5923024/

A

Dynamically regulated WSP-TREs (Tfit)
N = 18,552

Common SNPs (MAF > 0.05)
(+/-) 50 bp WSP-TRE centers
N = 4,373

Associated with Asthma
phenotype = WSP-SNPs
N = 52

B

- Exon - coding
- 3' UTR
- 5' UTR
- Intergenic
- Intron

C

Beas-2B
PRO-seq

HK2

VEH

WSP30

WSP120

500 bp

0.6
-0.3
0.6
-0.3
0.6
-0.3

rs28362965