1   **Title**
2   Functional Variant Discovery Identifies SPRY2 as a Genetic Mechanism Linking Wood Smoke
3   with Asthma
4
5   **Authors**

6   Arnav Gupta[1,7], Amber Dahlin[2], Alejandra Macario[1], Fabienne Gally[1,3], Michael R. Weaver[1], Lynn
7   Sanford[4], Margaret A. Gruca[4], Michael H. Cho[2], Robin D. Dowell[4,5,6], Scott T. Weiss[2], Sarah K.
8   Sasse[1], Anthony N. Gerber[1,3,7]
9
1. Department of Medicine, National Jewish Health
2. Channing Division of Network Medicine, Harvard Medical School
3. Department of Immunology and Genomic Medicine, National Jewish Health
4. BioFrontiers Institute, University of Colorado Boulder
5. Molecular, Cell, and Developmental Biology, University of Colorado Boulder
6. Computer Science, University of Colorado Boulder
7. Department of Medicine, University of Colorado Anschutz Medical Campus
17
18   Corresponding Author: Arnav Gupta
19   guptaa@njhealth.org
20   214-649-9559

27
28

Abstract

As a consequence of climate change and land use policies, there has been a historic rise in wildfire smoke across the United States and the world. While the deleterious effects of wildfire smoke and associated air pollution on asthma outcomes are established epidemiologically, genetic risks and molecular mechanisms of how wildfire smoke affects asthma are unknown. This knowledge gap hinders the identification of high-risk individuals and the creation of targeted therapies or recommendations to protect these individuals. We identified 52 genetic risk variants that colocalized with genomic responses to wood smoke particles (WSP), a model of wildfire particulate matter, and associated with asthma in the Genetic Epidemiology Research on Aging (GERA) cohort. We used additional filters to prioritize variants for direct testing of allele-dependent transcriptional regulatory function in plasmid reporters. We found that the rs3861144 variant (Odds Ratio$_{asthma}$ = 1.036) changes *SPRY2* responses to WSP airway epithelial cells, which we showed is involved in mechanical scratch repair in cell culture by regulating Extracellular Signal-related Kinase (ERK) phosphorylation. These findings provide insights into the molecular pathways through which WSP may influence asthma risk and propose genetic candidates that warrant further study for their potential as clinical tools for asthma.

Introduction

Climate change is contributing to a historic rise in wildfire smoke and reversing decades of improvements in air quality made by environmental policy (1–3). The scope of this problem is extending across North America and the world, increasing the population exposed to wildfire air pollution (4). Wildfire smoke is difficult to regulate or prevent. Climate policy and wildland management strategies will take decades to reverse the trend of expanding wildfires, and deteriorating air quality is expected over the next 30 years at least (5). It is therefore critical to understand the health consequences of wildfire air pollution and implement detailed strategies to manage and mitigate this developing healthcare crisis.

The profound health impacts of wildfire smoke and air pollution are well established at the population scale (6,7) and the impact on respiratory diseases, such as asthma, is a major concern. Wildfire smoke increases levels of ambient air pollutants, such as particulate matter less than 2.5 um (PM2.5), and causes morbid asthma outcomes, such as loss of asthma control, emergency room visits and hospitalizations (8–10). Data suggest ambient PM2.5 may also contribute to reduced lung function and the development of asthma (11–13). These major concerns highlight critical knowledge gaps in our current understanding of how PM2.5 from wildfire smoke impacts respiratory biology and asthma. First, the molecular genetic mechanisms that connect wildfire PM2.5 with asthma have not been identified, preventing the development of targeted treatments that mitigate asthma development or exacerbation. Second, there is significant variation in how individuals respond to airborne exposures, such as PM2.5 or wildfire smoke (14), and there are no biomarkers that predict individual risk of asthma complications resulting from exposure. As a result, public health guidelines, which only advise exposure avoidance, have limited efficacy, as restricting large population segments from outdoor activity for weeks during wildfire events is impractical, and methods to guide high-risk individuals do not exist. A better understanding of how PM2.5 and wildfire air pollution contributes to asthma risk and who is affected is critical to advise large populations in the face of this problem.

Genetic analysis is a powerful tool to inform disease mechanisms, define individual risk, and advance our understanding of the relationships between PM2.5, wildfire smoke, and asthma. However, there is a significant gap between the effect sizes of established genetic variants and asthma heritability, which is estimated at up to 70% of causation in certain studies (15). This suggests that additional asthma risk factors that are not captured in genome-wide association

studies (GWAS), such as environmental exposures, can influence genetic effects. This concept
is supported by discoveries of gene mechanisms that connect air pollution exposure with
asthma, such as mutations regulating *NQO1*, *GSTM1* and *GSTP1* (16–18). Identification of
additional conditional genetic risk variants, which primarily contribute to asthma risk in the
presence of exposure, has the potential to define novel asthma mechanisms associated with
wildfire PM2.5 and individuals with high risk.
86
Several challenges limit the identification of exposure-conditional risk variants, despite the major
advances achieved by asthma genetic studies. First, integrating environmental variables in
GWAS is highly challenging due to a myriad of reasons, including exposure variability, duration
of exposure effects, and types of exposures, among others. Genome-wide interaction studies
(GWIS) have approached this challenge by quantifying environmental variables through
geolocalization and retrospective measurements of local air quality (19), but these studies are
limited by the precision of geolocalization, resolution of air pollutant measurements, and
difficulty with compounding exposures based on where study subjects have lived over decades.
Second, linking specific genetic variants with cell and molecular functions can be challenging.
GWAS lead variants often reside in non-coding regions (20) and represent statistical but not
functional association priorities. Tools, such as functional scoring and expression quantitative
trait locus (eQTL) analysis, can improve associations between GWAS variants and gene
regulation but fail to leverage the full power of GWAS results, which may contain additional
functional variants that do not meet statistical thresholds. These tools also do not account for
the effects of relevant environmental exposures. Therefore, novel approaches that build on top
of existing GWAS data, incorporate environmental exposures, and directly connect variants with
function are necessary to realize the full potential of genetic analysis.
104
In this study, we applied our novel approach to variant discovery (21) to identify functional
genetic variants and molecular mechanisms that connect a wildfire PM2.5 model with asthma
risk. Using wood smoke particles (WSP) (22,23) to model wildfire PM2.5, we identified and
biologically validated a set of exposure-conditional genetic variants using population data of
lung function, transcription factor binding motifs, and epigenetic features of live airway epithelia
obtained from bronchoscopy. We directly tested the potential for high-priority variants to change
transcriptional function, connected variants with gene targets, and tested the function of these
target genes in airway epithelial cell biology. Our results were highlighted by the discovery of
rs3861144, a functional variant that changes *SPRY2* responses to WSP exposure and
potentially contributes to asthma development by affecting inflammatory gene expression and
airway epithelial repair following mechanical injury.
116
117 **Methods and Materials**
118 Additional methods are provided in the Online Data Supplement due to word limit constraints.
119
Genetic discovery data set and Analysis
For this analysis, large-scale genome-wide and phenotype data were utilized from the Genetic
Epidemiologic Research on Aging (GERA) study, a large, multi-ethnic asthma cohort. GERA
includes over 110,000 subjects (68,623 total asthma cases across four ethnic groups) with
extensive electronic medical records (EMR) information and genome-wide SNP genotypes (>8
million typed and imputed markers). We previously reported multiple novel associations with
asthma in four major ethnic groups in GERA (24). The current analysis was limited to the ethnic
subset of the cohort with the highest percentage of asthma cases and therefore the greatest
statistical power; the final data set included information from 16,274 non-Hispanic white
asthmatic cases and 38,269 controls (54,543 total samples). We applied methods previously
developed by our group (21) to regulatory elements identified in WSP-exposed Beas-2B cells by

Precision Run-on sequencing (PRO-seq) (n=18,552) (25). SNPs that overlapped regions within
50 bp (+/-) of regulatory element centers were selected for association analysis. Tests for
association of additive genotype with asthma status (dichotomized outcome variable) were
conducted using multivariate logistic regression models, adjusted for covariates, and the
significance of associations was determined using a max(T) permutation-based approach with
5000 permutations and a false-discovery rate (FDR) threshold of 0.05. All analyses were
performed using R statistical computing software and PLINK 1.9.
138
CRISPR-Cas9 RNP transfection and nucleofection
We performed genomic deletion of the transcriptional regulatory element (TRE) containing
rs3861144 and *SPRY2* using the Alt-R CRISPR-Cas9 system supplied and guided by IDT.
Custom-designed crRNAs, which target two sites within the TRE or the *SPRY2* gene body, and
a negative control crRNA (IDT) were duplexed (5 min at 95C) with tracrRNAs and then
complexed with Cas9 enzyme (1:1 molar ratio of duplex to Cas9) to a final concentration of 1
uM. This ribonucleic protein complex was transfected into Beas-2B cells using Lipofectamine
CRISPRmax (Life Technologies) based on manufacturer recommendations or electroporated
into primary small airway epithelial cells (SAECs) using the Lonza nucleofector IIb system and
the Basic Kit for Epithelial Cells. Nucleofection protocol was adapted from a previous study (26)
for the I/IIb system using the T-020 program. Genome editing was confirmed by PCR
amplification of genomic DNA (DNeasy Blood and Tissue kit – Qiagen) or Western Blot. Guide
RNA sequences are provided in Supplemental Table E2.
152
Results
Identification of functional variant candidates associated with asthma risk and wood smoke
particle exposure. To connect airway epithelial wood smoke particle (WSP) responses with
mechanisms of asthma, we re-analyzed nascent transcription profiles from an airway epithelial
cell line (Beas-2B) exposed to WSP for 30 or 120 minutes (25). 18,552 transcriptional regulatory
elements (TREs), excluding transcriptional start sites, responded dynamically to WSP in 6
kinetic programs (increased or decreased transcription between vehicle/control, WSP 30
minutes and WSP 120 minutes). Within a 50 bp radius of bioinformatically-defined center points
of these TREs, we tested 4373 common variant single nucleotide polymorphisms (SNPs, minor
allele frequency > 0.05) and identified 52 candidates associated with asthma phenotype using a
permutation-based statistical method applied to data from the non-Hispanic-white subpopulation
of the GERA cohort (False Discovery Rate, FDR < 0.05), which are designated "WSP-SNPs" in
the remainder of this manuscript (Figure 1A). These candidates predominantly reside in non-
coding genomic regions (Figure 1B). A representative example of a WSP-SNP, rs28362965, is
depicted in Figure 1C. Here, WSP exposure increases transcription of both the TRE containing
rs28362965 and *HK2.* A full list of the 52 variants is provided in Table 1.
169
WSP-SNPs associated with measures of lung function in an independent genetic data set. As
an initial validation approach, we analyzed the 52 WSP-SNPs for associations with lung function
in previously published GWAS data. We used the Open Targets Genetics platform (27,28) and
the Lung Disease Knowledge Portal (29) to navigate GWAS data of lung function
measurements, such as forced vital capacity (FVC), forced expiratory volume at 1 second
(FEV1), or peak expiratory flow (PEF) (30,31). We identified 6 WSP-SNPs significantly
associated ($p_{adj} < 0.05$) with measures of lung function (Table 2). Three WSP-SNPs (rs3861144,
rs3129975 and rs28517751) achieved a statistical threshold for genome-wide association (p <
$5\text{x}10^{-8}$). Associations between WSP-SNPs and measurements of lung function, which are
objective features of asthma, supported the validity of our discovery data set and motivated
further biological study of the 52 WSP-SNPs.
181

Aggregate analysis of the 52 WSP-SNPs suggested regulation of diverse cellular processes. To
understand biologic pathways regulated by the WSP-SNPs, we imputed their gene targets using
two methods. First, we used expression quantitative trait locus (eQTL) data from a public
genetic repository (Open Targets Genetics) to assign gene targets to SNPs. This approach was
not comprehensive, as multiple WSP-SNPs had no eQTL associations. However, we identified
154 genes associated with at least one WSP-SNP. Functional annotation using DAVID (32)
identified no enriched categories for these 154 genes. Second, we applied a permissive
approach that identified all genes that were located within 500 kb and shared nascent
transcriptional kinetics with the WSP-SNP's TRE. This approach identified 376 genes, and one
significantly enriched (p_{adj} = 2.1e-7) annotation term – "keratin filament." A complete list of the
putative gene targets from each approach is provided in Supplemental Tables E4 and E5.
Based on the limited functional clustering, we concluded that the WSP-SNPs regulate diverse
genes and cellular processes, and further filtering steps were necessary to identify functional
SNPs and associated mechanisms.
196
Biologic filtering using chromatin accessibility and bioinformatic transcription factor binding
predictions identified a subset of WSP-SNP candidates for functional analysis. We next filtered
the set of WSP-SNPs using cellular and molecular features relevant to asthma and wildfire
PM2.5 exposures. We found that 23 WSP-SNPs overlapped with chromatin accessibility in live
airway epithelial cells obtained from bronchoscopy from three healthy non-smoking adult donors
(Figure 2A), supporting the biologic validity of these WSP-SNPs in airway epithelial function. A
representative example demonstrates chromatin accessibility in live airway epithelium at the
rs9903086 SNP and its neighboring gene *JUP* (Figure 2B). We next scanned genome
sequences of TREs containing WSP-SNPs for binding motifs from two classes of transcription
factors that are relevant to WSP and PM2.5 exposures – aryl hydrocarbon receptor (AHR) and
nuclear factor kappa B (NFkB). Using a 500 bp radius to define the regulatory region containing
each WSP-SNP, we encountered these motifs at 38 WSP-SNP loci, and 23 WSP-SNP loci
contained motifs for both AHR and NFkB factors (Figure 2C), supporting our previous study that
demonstrated cross talk between these factors as a mechanism mediating inflammatory
responses to WSP exposure (25). The TRE containing rs11883931, which neighbors *LRRFIP1*,
is one example that contains both AHR nuclear transporter (ARNT2) and NFkB (REL)
transcription factor binding motifs (Figure 2D). We used these filters to select high-priority WSP-
SNPs for further functional testing.
215
Functional WSP-SNP genotype changed transcriptional regulatory potential in plasmid reporter
systems. We selected 15 WSP-SNPs prioritized by these filters to test transcriptional function in
plasmid reporters (Figure 3A) and cloned 700-800 bp regions surrounding these SNPs into
luciferase reporters. Within this reporter set, 5 WSP-SNP constructs exhibited enhancer activity
at baseline or in response to WSP exposure, and single base allele change of 2 WSP-SNPs
changed the regulatory activity of the entire construct (Figure 3B). We defined rs3861144 and
rs11832287 as functional genetic variants that alter the transcriptional regulatory activity of their
resident TREs based on haplotype.
224
WSP-SNPs identified functional variants at statistically prioritized GWAS loci. The functional
genetic variants that we identified were not previously reported but had statistical associations
with lung function, such as rs3861144 with FEV1 (Table 2). We next identified the statistically
prioritized variants at the WSP-SNP loci to compare the regulatory function of these lead or
causal variants with WSP-SNPs using the luciferase reporter system. We performed Bayesian
fine mapping (33) for the continuous FVC trait using summary statistics from a previously
published GWAS (34) at rs3861144 and rs9903086 loci. No SNPs were identified in the credible
set at rs3861144, but rs4885681 was identified as the local lead variant based on the lowest p

value for association ($p = 1.3 \times 10^{-11}$). The rs3861144 plasmid reporter had greater transcriptional
regulatory activity in the luciferase assay when compared with a plasmid reporter similarly
constructed for rs4885681, which resides in a non-annotated and non-functional genomic region
(Figure 3C-D). We selected three SNPs within the credible set of association with FVC at the
rs9903086 locus – rs9913846, rs9911655, and rs33914436 (Supplemental Table E6). Again, the
transcriptional regulatory activity of the rs9903086-containing plasmid reporter at baseline and
in response to WSP exposure was greater than reporters containing these local fine-mapped
SNPs (Figure 3E), which reside in genomic regions without functional annotations
(Supplemental Figure E1). While statistical approaches support the association between WSP-
SNP loci and lung function, our discovery approach improves the precision of defining functional
variants, which facilitate the connections between genetic signals, gene mechanisms, and
disease.
245
Shared nascent transcriptional kinetics between genes and regulatory elements identified
SPRY2 as the target of the rs3861144 WSP-SNP. We next leveraged one functional genetic
variant to define a gene mechanism that connects WSP exposure with asthma. Prior studies
have connected TREs with target genes based on shared activity and genomic proximity (35).
We applied this concept to identify genes regulated by WSP-SNPs based on nascent
transcription. Using this assumption, the regulatory element containing rs3861144, a WSP-SNP
independently associated with lung function with allele-dependent transcriptional regulatory
activity, regulates *SPRY2* in response to WSP exposure (Figure 4A). Three-dimensional
chromatin analysis using micrococcal nuclease chromosome conformation capture (Micro-C) in
unstimulated Beas-2B cells (36) demonstrated that rs3861144 and *SPRY2* reside in the same
topologically associated domain and may interact indirectly (Supplemental Figure E2). To
definitively link the TRE harboring rs3861144 with control of *SPRY2*, we deleted the rs3861144
TRE using CRISPR-Cas9 and analyzed *SPRY2* expression using qPCR in Beas-2B cells. PCR
amplification of genomic DNA from cells transfected with rs3861144 TRE-targeting guide RNAs
or negative control guide RNA demonstrated efficient deletion of this TRE (Figure 4B). Cells that
had undergone bulk editing showed a reduction in *SPRY2* expression response to WSP (Figure
4C). We also demonstrated that the *SPRY2* gene expression response to WSP correlated with
an increase in *SPRY2* protein (Supplemental Figure E3). These results connected the
rs3861144 TRE with *SPRY2* expression in response to WSP and demonstrated the facility of
functional SNPs to connect exposure, gene mechanism, and phenotype.
266
SPRY2 knockdown resulted in impaired mechanical scratch repair in Beas-2B cells by reducing
ERK1/2 phosphorylation. Based on previous studies that have reported *SPRY2* as a regulator of
growth factor responses, cell proliferation, and ERK phosphorylation (37–39), we studied the
function of *SPRY2* in Beas-2B cells by performing siRNA (small-interfering RNA) knockdown
and measuring the closure of a mechanical scratch defect in submerged culture. Cells with
SPRY2 knockdown had a greater scratch area at 16 hours (Figure 5A). Representative images
from one experiment are presented in Figure 5B. We demonstrated that *SPRY2* knockdown was
associated with reduced ERK1/2 phosphorylation in unstimulated Beas-2B cells, but following
WSP exposure, *SPRY2* knockdown resulted in prolonged ERK1/2 phosphorylation (Figure 5C).
These results demonstrated paradoxical functions of *SPRY2* in unstimulated and WSP-
stimulated conditions.
278
SPRY2 knockdown prolonged inflammatory gene expression responses to acute WSP
exposure. We applied RNA sequencing to study the role of *SPRY2* in regulating gene
expression responses to WSP exposure in Beas-2B cells. A principal component plot
demonstrating sample clustering is presented in Supplemental Figure E4. Consistent with our
prior studies (25), we observed an early-peaking response in differential gene expression with

591 upregulated genes at 2 hours and 408 upregulated genes at 4 hours of WSP exposure ($\log_2$
fold change > 0.5 and p_{adj} < 0.05). Transcriptional programs previously associated with WSP
exposure, such as canonical aryl hydrocarbon receptor targets AHRR and CYP1A1 or Nuclear
Factor Kappa-B targets CXCL8 and ICAM1, were upregulated at 2 hours of WSP exposure. We
identified 117 upregulated and 229 downregulated genes in *SPRY2* knockdown cells compared
with control cells in unstimulated conditions (Figure 6A). Functional annotation using DAVID (32)
identified "Growth Factor" (FDR = 0.0088) as the top annotation term in the upregulated gene
set. Next, we identified 164 or 122 genes upregulated by 2- or 4-hour WSP exposure in control
cells but not *SPRY2* knockdown cells (Figure 6B, Supplemental Table E6). The significant (FDR
< 0.05) functional annotation terms for these gene sets are provided in Table 3. We next directly
compared genes up- or downregulated by WSP exposure between *SPRY2* knockdown and
control cells and identified 132 upregulated and 320 downregulated genes (2-hour exposure)
and 152 upregulated and 320 downregulated genes (4-hour exposure) (Figure 6C). Significant
functional annotation terms for each of these gene sets are listed in Table 4. Notably, in *SPRY2*
knockdown cells after 4-hour WSP exposure, in addition to the annotation term "Growth Factor"
(FDR = 0.0081), which is consistent with prolonged ERK1/2 phosphorylation, we also
encountered "TNF signaling" as an upregulated functional cluster (FDR = 0.023). Finally, we
identified 86 genes in control cells and 157 genes in *SPRY2* knockdown cells that were
upregulated between the 2- and 4-hour WSP exposure time points (Supplemental Table E7 and
E8). Functional annotations of these gene sets identified enrichment of inflammatory pathways
in *SPRY2* knockdown cells but not control cells (Table 5). These data suggest that *SPRY2*
knockdown increases inflammatory gene expression at longer durations of WSP exposure and
SPRY2 may play an important role as a negative feedback regulator.
307
SPRY2 is regulated in primary human airway cells. We next demonstrated that *SPRY2*
expression increases in primary SAECs cultured and exposed to WSP in submerged/basal
conditions (Figure 7A). The gene expression kinetics of *SPRY2* were similar to kinetics in Beas-
2B cells, peaking at 2 hours and decreasing by 4 hours of WSP exposure. No significant
cytotoxicity was observed at 4 or 24 hours of WSP exposure in these cells in this or prior studies
(25,40). We also demonstrated that *SPRY2* expression was regulated in primary human airway
smooth muscle (HASM) cells in response to TNF-alpha, a secondary mediator of WSP released
by airway epithelial and innate immune cells, and primary HASM cells from one asthma donor
exhibited reduced *SPRY2* expression (Figure 7B). Finally, we performed *SPRY2* knockdown
using CRISPR-Cas9 in SAECs from one donor (Sm36) and demonstrated a reduced ratio of
phosphorylated to dephosphorylated ERK1/2 in *SPRY2*-knockdown basal SAECs.
(Supplemental Figure E7). Therefore, *SPRY2* is regulated in primary human airway cells and
controls MAPK signaling by regulating ERK1/2 expression and phosphorylation.
321
Discussion
Wildfire related PM2.5 air pollution is increasing and has a direct impact on asthma morbidity
(10,41,42). However, the genetic and molecular mediators of these malign effects are poorly
understood. To address these gaps, we developed a genetic approach to discover novel
relationships between asthma, SNPs, and transcriptional regulatory responses to WSP
exposure in a cell culture model. Through this approach, we identified 52 common variant SNPs
within WSP-responsive TREs that were nominally associated with an asthma diagnosis in the
GERA cohort. We prioritized SNPs based on primary human airway chromatin accessibility
profiles and bioinformatic predictions of transcription factor binding, and we identified two
functional asthma variants, rs3861144 and rs11832287. Transcriptional regulatory activity of the
regions harboring rs3861144 and rs11832287 was controlled by haplotype, and based on
congruent TRE and gene transcriptional responses, we identified the putative gene targets of
rs3861144 and rs11832287 as *SPRY2* and CD9. We confirmed the regulatory association

between rs3861144 and *SPRY2* using a CRISPR-based deletion strategy. Further, we
connected *SPRY2* to ERK1/2 phosphorylation, epithelial monolayer scratch repair, and pro-
inflammatory gene expression. Our findings establish a powerful pipeline for genetic discovery
of associations between asthma and woodsmoke and highlight a novel link between *SPRY2*
induction by WSP and risk of an asthma diagnosis.
340
SPRY2 is a kinase modulator of the mitogen-activated protein kinase (MAPK) pathway that
plays diverse cellular roles in regulating cell growth, survival, and proliferation (43–45). We
demonstrated that *SPRY2* expression increases following WSP exposure, which aligns with
prior reports that connect *SPRY2* regulation and function (39,46). Previous studies have
implicated the *SPRY2* locus in controlling lung function based on genetic signals. Multiple linked
SNPs, including rs3861144, were statistically associated ($p < 5 \times 10^{-9}$) with measurements of lung
function, such as FEV1 (34). This study prioritized rs4885681 as the lead SNP based on the
strongest association statistics and connected rs4885681 with the LINC00382 long non-coding
RNA, based on its genomic location. However, we demonstrated that rs4885681 did not exhibit
features of a functional genetic variant, such as genomic annotations or transcriptional
regulatory function. Therefore, the mechanism derived from rs4885681 was unlikely to explain
the connection between genetic signals at this locus and lung function. In contrast, we identified
rs3861144 as a functional variant at this locus, and using nascent transcript annotations, we
identified *SPRY2* as its gene target and the plausible mechanism that connected the genetic
signals in this locus with FEV1. These data demonstrate that defining the functional variants
within a statistically associated genetic locus provides additional utility to GWAS by narrowing in
on gene mechanisms that contribute to the phenotype.
358
Although the precise role of WSP-dependent *SPRY2* expression in asthma development
remains to be defined, we observed *SPRY2*-dependent cellular effects that align with previous
reports of *SPRY2* function and identify plausible mechanisms linking *SPRY2* and asthma. In our
model, *SPRY2* regulates MAPK signaling and phosphorylation of mediators, such as ERK1/2
(37,38,47). In unstimulated Beas-2B cells, we observed a direct correlation between *SPRY2*
knockdown, reduced ERK1/2 phosphorylation, and delayed ability of *SPRY2* knockdown cells to
close a mechanical scratch in a submerged airway epithelial monolayer, consistent with prior
studies in airway epithelial cells (48) and rhabdomyosarcoma cell lines (47). However, in
response to WSP exposure, *SPRY2* knockdown increased ERK1/2 phosphorylation, suggesting
significant differences in *SPRY2* function between unstimulated conditions and WSP exposure.
These observations are supported by previous studies that reported cell context and growth
factors as variables that determine the positive or negative correlation between *SPRY2* and
ERK1/2 phosphorylation (39). We also observed changes in pro-inflammatory gene expression
kinetics in *SPRY2* knockdown cells exposed to WSP. In our model, the pro-inflammatory
response to WSP is under negative feedback and peaks in magnitude at 2 hours of exposure
(25). However, in *SPRY2* knockdown cells, pro-inflammatory genes, such as CXCL8 and IL1B,
increased in magnitude between 2 and 4 hours of exposure. These results support a negative
feedback mechanism for *SPRY2* in regulating inflammatory responses to WSP by regulating
MAPK signaling. Further investigations of WSP exposure, ERK1/2 phosphorylation and de-
phosphorylation, and mechanistic partners of the *SPRY2*-inflammation relationship are
necessary to clarify the role of *SPRY2* in WSP exposure and the development of asthma.
380
In addition to identifying the connection between *SPRY2*, WSP exposure, and asthma, our
approach to linking WSP exposure with asthma had several advantages. First, by focusing on
PRO-seq-annotated WSP responses, we pre-selected SNPs to test for asthma association that
reside in the functional centers of TREs, which are enriched for transcriptional machinery
(49,50) and regulatory SNPs (51,52). This constraint enriched our discovery for functional SNPs

and reduced the genomic sequences for discovery to a total of ~1 Mb. The reduced number of
total SNPs tested allowed us to apply permissive permutation-based statistics and identify
variants that would not enrich in GWAS results. While previous studies have applied genomic
annotations to prioritize functional variants from GWAS results (53–56), these methods are
limited to testing variants that meet the rigorous statistical thresholds of GWAS first, which may
miss functional variants with weaker statistical associations. Second, we pre-selected SNPs that
reside in WSP-responsive TREs to enrich genetic discovery for SNPs whose transcriptional
regulatory activity depends on WSP exposure. While this method does not directly integrate the
complexity of environmental exposures with the genetic risk of asthma, it does provide insight
into asthma risk in the context of genomic responses to relevant exposures. Finally, PRO-seq
annotations facilitated connecting functional variants with target genes that respond to pertinent
exposures based on transcriptional kinetics. The location of most genetic risk variants in non-
coding and non-annotated genomic regions (57) presents a general challenge for defining
relevant target genes and associated mechanisms from GWAS results. Expression quantitative
trait locus (eQTL) analysis can address this problem (58) but requires knowledge of the
functional cell type and access to large tissue gene expression data sets. Direct exposure
measurements in relationship to gene expression are also challenging for eQTL studies, and
unaccounted variable exposures can confound eQTL results. We connected functional genetic
variants with gene targets by applying PRO-seq data from our cell exposure model to an
established paradigm for regulatory element-gene association (35). The advantages of our
pipeline relative to other methods led to the discovery of rs3861144 and *SPRY2* as novel
mechanisms that connect wildfire PM2.5 exposure, modeled by WSP, with asthma. Future
studies that integrate our method with existing tools may facilitate additional discovery of
variants and target genes that influence gene-by-environment risk in asthma.
410
Our study had several limitations. The positive predictive value of our primary discovery in
relationship to our secondary analyses was modest. Only 6/52 SNPs had independent
associations with lung function and 2/52 SNPs were rigorously defined as functional variants.
Low thresholds and limitations in accurately defining TREs bioinformatically and coding the
asthma diagnosis may have contributed to these results, as well as our application of lung
function genetics as a surrogate validation tool for our primary findings We also performed our
initial genetic analyses in mostly non-Hispanic white cohorts and used a single reference for
linkage disequilibrium analysis, limiting the generalizability of the results and increasing potential
bias. Future validation in an independent multi-ancestry asthma cohort with a whole genome
sequencing data set will improve the rigor and confidence of the results.
421
Our WSP and cell culture exposure system also had limitations. We used aged, desiccated, and
UV-exposed WSP from a single combustion source. Although this model induces relevant
pathways, such as aryl hydrocarbon receptor and inflammasome signaling (59–61), it also
comprises a narrow representation of long-distance PM2.5 air pollution from wildfire smoke.
Inconsistencies between the Beas-2B immortalized airway epithelial cell line and normal human
airway epithelia, such as supraphysiologic concentrations of growth factors (62), lack of active
metabolic pathways (63), cilia, and mucus functions (64) resulted in a bias towards
transcriptional programs involved in cell proliferation in our model exposure system. Lacking
from our model were important airway epithelial programs, such as mucus dysregulation,
metabolism, and cell junction integrity, which also contribute to asthma development (65–68).
Finally, the dose (1 mg/ml or 0.26 mg/cm^2) and duration of WSP exposure introduced bias of
transcriptional responses towards cell stress pathways and away from potentially important
metabolic responses, which have also been associated with asthma (69). The acute 30- and
120-minute exposure time points, while potentially enriched for direct exposure mediators, may
fail to capture important sustained or secondary responses, such as oxidative stress or

epithelial-to-mesenchymal transformation, which have been suggested as mechanisms that
connect exposures with asthma previously (64,70).
439
Despite these limitations, our study expands the understanding of the genetic factors that
influence asthma risk in the context of wildfire PM2.5 exposure, a problem of growing
importance due to global climate change. We identified common SNPs that may contribute to
the personal risk of asthma associated with wildfire PM2.5 and cellular mechanisms that
mediate this risk. We characterized one important mechanism by demonstrating that *SPRY2*
expression in response to WSP exposure is mediated by rs3861144, which modifies asthma risk
and lung function. This connection between WSP and *SPRY2* may alter asthma risk by
impacting airway epithelial cell proliferation and injury repair through the MAPK pathway or
increasing pro-inflammatory responses to exposures. Irrespective of the precise mechanism,
our results implicate *SPRY2* as a mediator of the adverse effects of wildfire PM2.5 exposure on
asthma risk and define *SPRY2* as a potential therapeutic target for asthma.
451

452   Table 1 – WSP-SNPs

SNP	CHR	BP (HG38)	MAF	OR	FDR
rs16860727	1	183272646	0.12620	1.049	0.043200
rs11590328	1	209399237	0.3555	1.045	0.007899
rs75459144	10	75294962	0.15020	0.941	0.006199
rs6584144	10	97700730	0.19740	0.9563	0.028900
rs138744334	11	94643137	0.0694	0.938	0.045100
rs12321808	12	6167792	0.37620	1.037	0.024000
rs11832287	12	6182141	00.46510	0.9602	0.011800
rs10747612	12	51394035	0.37200	0.9623	0.020300
rs9509353	13	20722128	0.4838	0.9683	0.042700
rs17065695	13	44310244	0.0598	0.9236	0.015800
rs3861144	13	79936471	0.26700	1.036	0.042800
rs1275823	14	69726964	0.13050	0.9479	0.021300
rs9939094	16	29268980	0.35850	0.9569	0.008099
rs253140	16	83963743	0.11640	1.068	0.008099
rs3968348	17	2732850	0.4839	0.9575	0.007999
rs60701308	17	4858995	0.0946	0.9418	0.025700
rs9896022	17	16405309	0.12300	1.052	0.037300
rs10491130	17	41377775	0.10090	0.9437	0.029200
rs9903086	17	41793264	0.0625	0.9322	0.034900
rs3848425	17	66467803	0.39440	1.035	0.033900
rs644821	18	11954652	0.10280	0.9441	0.027400
rs4806835	19	2266880	0.13350	1.054	0.031300
rs10424200	19	38792918	0.27260	1.039	0.036800
rs10418662	19	46426768	0.34810	1.04	0.020100
rs7246311	19	680378	0.4121	1.05	0.024000
rs770665	2	197729533	0.3457	1.055	0.001400
rs7591472	2	201025028	0.17090	1.044	0.044000
rs11883931	2	237315599	0.21540	1.046	0.022500
rs10208406	2	10057488	0.19520	0.9596	0.041000
rs2139379	2	12574801	0.17300	0.9573	0.038000
rs28362965	2	74845567	0.0516	1.078	0.031600
rs6023060	20	38109598	0.17150	1.047	0.028100
rs764025	22	33853729	0.22390	0.9545	0.019000
rs2173620	3	177538135	0.23200	0.959	0.026500
rs7428132	3	177745986	0.0541	0.9083	0.007299
rs6784916	3	189937753	0.35420	0.9612	0.017000
rs12630368	3	5013455	0.11000	1.052	0.045400
rs2633783	3	9560123	0.1346	0.9438	0.012100
rs56984996	4	168138760	0.36030	1.041	0.016100
rs72705761	4	185847661	0.29980	1.043	0.016800
rs7669195	4	187421022	0.24990	0.9633	0.039100

rs6448484	4	26857636	0.4736	1.051	0.001900
rs1371799	4	74112120	0.38900	0.9682	0.045900
rs28517751	5	120624434	0.08031	0.9169	0.003900
rs162481	5	123719195	0.19460	0.9445	0.004400
rs6871945	5	56278753	0.27250	0.9601	0.023600
rs1873333	6	168008153	0.2205	1.049	0.012900
rs3129975	6	30764277	0.21440	1.075	0.000400
rs979701	6	77034107	0.24350	1.038	0.044100
rs524953	9	110844289	0.2588	0.9593	0.025500
rs7038732	9	127764264	0.41900	0.967	0.036200
rs3739819	9	127965830	0.0989	1.067	0.016700

454 Table 2 – WSP-SNPs with significant associations in lung function GWAS

		Open Targets Genetics	Lung Disease Knowledge Portal
SNP	**Association**	**P value**	**P value**
rs3861144	FEV1	6.80E-10	8.79E-10
rs28517751	FEV1/FVC	4.20E-08	4.12E-08
rs3129975	PEF	1.30E-26	1.74E-25
rs11883931	FEV1/FVC	3.90E-05	3.36E-05
rs16860727	FEV1/FVC	7.60E-06	3.80E-06
rs1873333	PEF	6.50E-04	8.53E-04

456 Table 3 – Functional Annotation Categories for Genes upregulated by WSP exposure in control
457 (siCTRL) cells but not *SPRY2* knockdown (si*SPRY2*) cells

siCTRL-WSP2h > veh NOT si*SPRY2*-WSP2h > veh		
Term	**Enrichment score**	**FDR**
Phosphoprotein	2.8	0.019
p53-like DNA binding domain	2.4	7.00E-05
Endocytosis	1.9	0.035
siCTRL-WSP4h > veh NOT si*SPRY2*-WSP4h > veh		
Term	**Enrichment score**	**FDR**
DNA binding Transcription Factor Activity	1.5	0.013

459 Table 4 – Functional Annotation Categories for Genes upregulated or downregulated in *SPRY2*
460 knockdown-WSP exposed (si*SPRY2*-WSP) cells compared with control-WSP exposed (siCTRL-
461 WSP) cells.

si*SPRY2*-WSP2h > siCTRL-WSP2h		
Term	**Enrichment score**	**FDR**
Endoplasmic Reticulum	2.7	0.012
Synapse	1.9	0.022
TGF-beta	1.9	0.017
Glycoprotein	1.8	0.022

siSPRY2-WSP2h < siCTRL-WSP2h		
Term	**Enrichment score**	**FDR**
Metalloprotease	2.1	0.037
Kinase	1.9	0.0062
Transmembrane	1.8	0.003
siSPRY2-WSP4h > siCTRL-WSP4h		
Term	**Enrichment score**	**FDR**
Growth Factor	2.5	0.008
Secreted	2.1	0.027
TNF signaling pathway	1.8	0.023
siSPRY2-WSP4h < siCTRL-WSP4h		
Term	**Enrichment score**	**FDR**
None		

463 Table 5 – Functional annotation categories for genes upregulated between 2 and 4 hours of
464 WSP exposure in control (siCTRL) and *SPRY2* knockdown (si*SPRY2*) cells

siCTRL WSP4h > WSP2h		
Term	**Enrichment score**	**FDR**
Secreted	2.3	0.031
Extracellular Matrix	2	0.025
Benzodiazepine Receptor Binding	1.5	0.031
si*SPRY2* WSP4h > WSP2h		
Term	**Enrichment score**	**FDR**
IL-17 signaling	4.2	2.00E-05
Cytokine	3.7	0.00098
NFkB	3.5	0.024
Secreted	3.4	0.00075
Cell Adhesion	2.3	0.006
Small Cell Lung Cancer	2.3	0.0058
Positive regulation of angiogenesis	2.3	0.024
RAGE signaling	2.1	0.0081
PI3K-AKT signaling	2	0.036
NOD-like receptor signaling	2	0.034
RHD DNA binding domain	2	0.0063
Inflammatory Bowel Disease	2	0.038
Extracellular Matrix	1.7	0.044
Amoebiasis	1.7	0.0014
Neutrophil Chemotaxis	1.5	0.049

465
466

Figure legends

Figure 1 – A permutation statistical approach identifies 52 common single nucleotide polymorphisms (SNPs) that reside within transcriptional regulatory elements (TREs) that respond to wood smoke particle (WSP) exposure and associate with asthma diagnosis. A. Schematic of the approach that applies WSP-dependent TREs identified by Precision Run-on sequencing (PRO-seq) in Beas-2B airway epithelial cells to asthma variant discovery. B. Genomic features of the 52 WSP-SNPs. C. Representative example of WSP-SNP which resides within an intronic TRE in the *HK2* gene. PRO-seq data is visualized in the Integrative Genomics Viewer (IGV) genome browser based on counts per million mapped reads (vertical scales). Positive (blue) signal indicates reads annotated to the sense strand while negative (red) signal reflects reads annotated to the antisense strand. The Transcription Start Site (TSS) and direction of transcription are indicated by the arrow at the top of the panel.

Figure 2 – Subsets of the 52 WSP-SNPs exhibit biologic functions associated with asthma. A. Venn diagram demonstrates 23 genomic regions that overlap (> 1 bp), based on genomic coordinates, between WSP-SNPs and chromatin accessibility features in live proximal airway epithelial tissue obtained from healthy non-smoking subjects. B. The representative example rs9903086 WSP-SNP resides in an intergenic region near the *JUP* TSS and overlaps with bidirectional PRO-seq signal in Beas-2B cells and chromatin accessibility signal in live airway epithelium. PRO-seq (blue and red) and ATAC-seq (green and teal) data visualized in the IGV genome browser based on counts per million mapped reads (vertical scales). The TSS is indicated by the arrow at the top of the panel. C. Venn diagram demonstrating sites of genomic overlap between a 500 bp radius surrounding each WSP-SNP and consensus transcription factor binding motifs for aryl hydrocarbon receptor (AHR) and nuclear factor kappa b (NFkB) family transcription factors. Twenty-three WSP-SNP regions contain motifs from both families. D. A representative example at the rs11883931 locus, which contains both ARNT2 (AHR nuclear transporter) and REL (NFkB family) transcription factor binding motifs.

Figure 3 – Luciferase activity in plasmid reporter systems demonstrates that WSP-SNPs reside in active TREs in comparison with local lead variants predicted by GWAS. A. Schematic for selection of WSP-SNP candidates prioritized by biologic variables of live airway chromatin accessibility and colocalization with both AHR and NFkB family transcription factor binding motifs. B. Normalized luciferase activity of plasmid reporters transfected into Beas-2B cells and treated with vehicle (veh) or WSP (1 mg/ml) for 8 hrs. A 700 – 800 bp genomic region surrounding each WSP-SNP was cloned into the plasmid system and mutated for allele change, with "WT" indicating the allele native to Beas-2B cells and genotype letter change indicating the mutated reporter. Bars indicate mean reporter activity (+/- standard deviation, n = 4) relative to an internal control Renilla luciferase that was co-transfected into cells; * $p < 0.01$. C. Normalized luciferase activity of plasmid reporters containing the WSP-SNP rs3861144 and rs4885681, the lead variant at this locus prioritized by GWAS statistics. A 350 – 400 bp radius genomic region surrounding each SNP was cloned into the plasmid system, and luciferase activity was measured after transfection into Beas-2B cells and treatment with vehicle (veh) or WSP (1 mg/ml) for 8 hrs. Data are presented as described for (B). D. Beas-2B PRO-seq and live airway epithelium ATAC-seq signal at the WSP-SNP rs3861144 and rs4885681 loci, located approximately 380 kb 3' to the *SPRY2* gene. E. Normalized luciferase activity of plasmid reporters containing a 350-400 bp radius around the WSP-SNP rs9903086 and three SNPs (rs9913846, rs991655 and rs39914436) prioritized by statistical fine mapping (FINEMAP (33)) for association with Forced Vital Capacity using summary statistics from an independent data set (30). Bars indicate mean reporter activity (+/- standard deviation, n = 4) relative to Renilla; * $p < 0.01$.

Figure 4 – The rs3861144 functional variant resides in a TRE that regulates *SPRY2* in response
to WSP exposure in Beas-2B cells. A. Beas-2B PRO-seq signal visualized in IGV at the
rs3861144 and *SPRY2* gene locus demonstrates shared transcriptional kinetics between these
regions at 30 and 120 minutes of WSP exposure. Below – green arrows indicate two target
guide RNA (gRNA) sites designed for Cas9 genome editing of the rs3861144 TRE; purple
arrows indicate target sites for polymerase chain reaction (PCR) primers used to amplify the
region following Cas9 editing and confirm genomic deletion. B. Beas-2B cells were transfected
with two on-target Cas9-gRNA RNP complexes indicated by green arrow sites in the rs3861144
TRE in (A) (rs3861144-Cas9) or negative control Cas9-gRNA RNP (NC). Gel electrophoresis
confirmed genomic deletion at the rs3861144 target region in cells transfected with targeting
gRNAs. C. CRISPR-Cas9 transfected cells were exposed to WSP at 1 mg/ml (0.26 mg/cm^2) for
2 hours. Gene expression of the canonical WSP target *CYP1A1* and *SPRY2* were measured
using quantitative PCR (qPCR). Bars represent mean normalized C_T values on a log_2 scale (+/-
standard deviation, n = 4) relative to vehicle-treated controls; * $p < 0.01$.

Figure 5 – *SPRY2* knockdown by siRNA delays scratch repair in a Beas-2B monolayer and
controls ERK1/2 phosphorylation. A. Scratch assay in Beas-2B cells transfected with short
interfering RNA (siRNA) targeting *SPRY2* or non-targeting control demonstrated a larger
remaining defect at 16 hours in cells with *SPRY2* knockdown. Solid line depicts the interpolated
slope between the means of measurements at 2, 6, 8, 10, 12, 16 and 25 hours. Individual
measurements are plotted as blue (non-targeting control "siCTRL") or red (*SPRY2*-siRNA)
points (n=6-7/condition). * $p_{adj} < 0.05$. B. Representative images from three wells of control-
transfected (siCTRL) and *SPRY2* knockdown (si*SPRY2*) scratch assay at three time points. C.
Western blot demonstrates knockdown in *SPRY2* expression with *SPRY2*-siRNA transfection
(si*SPRY2*) is associated with reduced phosphorylated ERK1/2 in unstimulated Beas-2B cells
compared with control-transfected cells (siCTRL). The increase in ERK1/2 phosphorylation
following WSP exposure is increased at 6 hours of WSP exposure in *SPRY2* knockdown
(si*SPRY2*) cells. GAPDH served as the loading control.

Figure 6 – RNA sequencing data of *SPRY2* knockdown Beas-2B cells exposed to WSP for 2
and 4 hours. A. Volcano plot demonstrates up- and downregulated genes in unstimulated
control vs *SPRY2*-knockdown cells. Each point represents a single gene, with the x-axis
showing log_2 fold change and the y-axis showing log transformed p_{adj} statistic. Dashed lines
indicate log_2 fold change of 0.5, which was the threshold used for this analysis. Horizontal axis
line indicates $p_{adj} < 0.05$. B. Venn diagrams demonstrate overlapping and non-overlapping
genes between two comparisons. Top – significant genes identified in the vehicle vs WSP-2hr
exposure comparison for each transfection condition – control (siCTRL) or *SPRY2*-targeting
(siSPRY2) siRNA. Bottom – similar comparison for vehicle vs WSP-4hr exposure. C. Volcano
plots showing differentially expressed genes at 2 (top) and 4 (bottom) hours of WSP exposure
between siCTRL and siSPRY2 cells. D. Volcano plot showing differentially expressed genes
between 2 and 4 hours of WSP exposure in siCTRL (top) and siSPRY2 (bottom) cells.

Figure 7 – *SPRY2* is expressed and regulated in primary human airway cells. A. Primary small
airway epithelial cells obtained from four healthy, non-smoking donors were exposed to WSP (1
mg/ml, 0.26 mg/cm^2) in submerged culture for 2 or 4 hours. Gene expression for *SPRY2* was
measured using qPCR. Bars represent mean normalized C_T values on a log_2 scale (+/- standard
deviation, n = 4) relative to vehicle-treated controls. B. *SPRY2* expression in normal HASM cells
is increased by TNF-alpha stimulation but reduced in cells from an asthmatic donor. GAPDH
was the loading control.

References

1.   Westerling ALR. Increasing western US forest wildfire activity: Sensitivity to changes in
the timing of spring. Philos Trans R Soc B Biol Sci [Internet]. 2016 Jun 5 [cited 2020 Oct
13];371(1696). Available from: /pmc/articles/PMC4874415/?report=abstract

2.   Burke M, Childs ML, de la Cuesta B, Qiu M, Li J, Gould CF, et al. The contribution of
wildfire to PM2.5 trends in the USA. Nature [Internet]. 2023 Oct 26 [cited 2024 Nov
26];622(7984):761–6. Available from: https://pubmed.ncbi.nlm.nih.gov/37730996/

3.   Childs ML, Li J, Wen J, Heft-Neal S, Driscoll A, Wang S, et al. Daily Local-Level
Estimates of Ambient Wildfire Smoke PM2.5for the Contiguous US. Environ Sci Technol
[Internet]. 2022 Oct 4 [cited 2024 Nov 26];56(19):13607–21. Available from:
https://pubs.acs.org/doi/abs/10.1021/acs.est.2c02934

4.   Cunningham CX, Williamson GJ, Bowman DMJS. Increasing frequency and intensity of
the most extreme wildfires on Earth. Nat Ecol Evol 2024 88 [Internet]. 2024 Jun 24 [cited
2024 Nov 26];8(8):1420–5. Available from: https://www.nature.com/articles/s41559-024-
02452-2

5.   Foundation FS. First Street Foundation Aggregated Wildfire Risk Summary Statistics
Version 1.0 [Internet]. Zenodo; 2022. Available from:
https://doi.org/10.5281/zenodo.6564731

6.   Gould CF, Heft-Neal S, Johnson M, Aguilera J, Burke M, Nadeau K. Health Effects of
Wildfire Smoke Exposure. Annu Rev Med [Internet]. 2024 Jan 29 [cited 2024 Nov
26];75(Volume 75, 2024):277–92. Available from:
https://www.annualreviews.org/content/journals/10.1146/annurev-med-052422-020909

7.   Gao Y, Huang W, Yu P, Xu R, Yang Z, Gasevic D, et al. Long-term impacts of non-
occupational wildfire exposure on human health: A systematic review. Environ Pollut.
2023 Mar 1;320:121041.

8.   Skinner R, Luther M, Hertelendy AJ, Khorram-Manesh A, Sorensen J, Goniewicz K, et al.
A Literature Review on the Impact of Wildfires on Emergency Departments: Enhancing
Disaster Preparedness. Prehosp Disaster Med [Internet]. 2022 Oct 25 [cited 2024 Nov
26];37(5):657. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC9470518/

9.   Beyene T, Harvey ES, Van Buskirk J, McDonald VM, Jensen ME, Horvat JC, et al.
'Breathing Fire': Impact of Prolonged Bushfire Smoke Exposure in People with Severe
Asthma. Int J Environ Res Public Health [Internet]. 2022 Jun 1 [cited 2023 Oct
2];19(12):7419. Available from: /pmc/articles/PMC9224478/

10.   Noah TL, Worden CP, Rebuli ME, Jaspers I. The Effects of Wildfire Smoke on Asthma
and Allergy. Curr Allergy Asthma Rep [Internet]. 2023 Jul 1 [cited 2024 Nov 26];23(7):1.
Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10176314/

11.   Gehring U, Wijga AH, Koppelman GH, Vonk JM, Smit HA, Brunekreef B. Air pollution and
the development of asthma from birth until young adulthood. 2020 Jul 1 [cited 2023 Jan
29];56(1). Available from: https://pubmed.ncbi.nlm.nih.gov/32299858/

12.   McConnell R, Berhane K, Yao L, Jerrett M, Lurmann F, Gilliland F, et al. Traffic,
susceptibility, and childhood asthma. Environ Health Perspect [Internet]. 2006 May [cited
2024 Nov 26];114(5):766–72. Available from:
https://ehp.niehs.nih.gov/doi/10.1289/ehp.8594

13.   Gauderman WJ, McConnell R, Gilliland F, London S, Thomas D, Avol E, et al.
Association between Air Pollution and Lung Function Growth in Southern California
Children. https://doi.org/101164/ajrccm16249909096 [Internet]. 2012 Dec 14 [cited 2025
May 9];162(4 I):1383–90. Available from:
/doi/pdf/10.1164/ajrccm.162.4.9909096?download=true

14.   Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical Review of
Health Impacts of Wildfire Smoke Exposure. Environ Health Perspect [Internet]. 2016
Sep [cited 2019 Jun 25];124(9):1334–43. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/27082891

15.   Thomsen SF, Ulrik CS, Kyvik KO, Hjelmborg J von B, Skadhauge LR, Steffensen I, et al.
Genetic and environmental contributions to hay fever among young adult twins. Respir
Med [Internet]. 2006 Dec [cited 2024 Nov 26];100(12):2177–82. Available from:
https://pubmed.ncbi.nlm.nih.gov/16650971/

16.   FRYER AA, BIANCO A, HEPPLE M, JONES PW, STRANGE RC, SPITERI MA.
Polymorphism at the Glutathione S-transferase GSTP1 Locus. Am J Respir Crit Care
Med [Internet]. 2000;161(5):1437–42. Available from:
https://doi.org/10.1164/ajrccm.161.5.9903006

17.   Goodrich GG, Goodman PH, Budhecha SK, Pritsos CA. Functional polymorphism of
detoxification gene NQO1 predicts intensity of empirical treatment of childhood asthma.
Mutat Res Toxicol Environ Mutagen. 2009 Mar 31;674(1–2):55–61.

18.   Strange RC, Spiteri MA, Ramachandran S, Fryer AA. Glutathione-S-transferase family of
enzymes. Mutat Res Mol Mech Mutagen. 2001 Oct 1;482(1–2):21–6.

19.   Melbourne CA, Mesut Erzurumluoglu A, Shrine N, Chen J, Tobin MD, Hansell AL, et al.
Genome-wide gene-air pollution interaction analysis of lung function in 300,000
individuals. Environ Int [Internet]. 2022 Jan 15 [cited 2024 Nov 26];159:107041. Available
from: https://pmc.ncbi.nlm.nih.gov/articles/PMC8739564/

20.   Zhu Y, Tazearslan C, Suh Y. Challenges and progress in interpretation of non-coding
genetic variants associated with human disease. Exp Biol Med [Internet]. 2017 Jul 1
[cited 2024 Nov 26];242(13):1325. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC5529005/

21.   Sasse SK, Dahlin A, Sanford L, Gruca MA, Gupta A, Gally F, et al. Enhancer RNA
transcription pinpoints functional genetic variants linked to asthma. Nat Commun 2025
161 [Internet]. 2025 Mar 31 [cited 2025 Apr 29];16(1):1–14. Available from:
https://www.nature.com/articles/s41467-025-57693-x

22.   Ghio AJ, Soukup JM, Case M, Dailey LA, Richards J, Berntsen J, et al. Exposure to wood
smoke particles produces inflammation in healthy volunteers. Occup Environ Med
[Internet]. 2012 Mar [cited 2019 Jun 18];69(3):170–5. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/21719562

23.   Ghio AJ, Soukup JM, Dailey LA, Tong H, Kesic MJ, Budinger GRS, et al. Wood Smoke
Particle Sequesters Cell Iron to Impact a Biological Effect. Chem Res Toxicol [Internet].
2015 Nov 16 [cited 2019 Jul 3];28(11):2104–11. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/26462088

24.   Dahlin A, Sordillo JE, Ziniti J, Iribarren C, Lu M, Weiss ST, et al. Large-scale, multi-ethnic
genome-wide association study identifies novel loci contributing to asthma susceptibility
in adults. J Allergy Clin Immunol [Internet]. 2019 Apr 1 [cited 2021 Jul 13];143(4):1633.
Available from: /pmc/articles/PMC6451881/

25.   Gupta A, Sasse SK, Gruca MA, Sanford L, Dowell RD, Gerber AN. Deconvolution of
multiplexed transcriptional responses to wood smoke particles defines rapid aryl
hydrocarbon receptor signaling dynamics. J Biol Chem [Internet]. 2021 Oct 1;297(4).
Available from: https://doi.org/10.1016/j.jbc.2021.101147

26.   Koh KD, Siddiqui S, Cheng D, Bonser LR, Sun DI, Zlock LT, et al. Efficient RNP-directed
human gene targeting reveals SPDEF is required for IL-13-induced mucostasis. Am J
Respir Cell Mol Biol. 2020 Mar 1;62(3):373–81.

27.   Ghoussaini M, Mountjoy E, Carmona M, Peat G, Schmidt EM, Hercules A, et al. Open
Targets Genetics: systematic identification of trait-associated genes using large-scale
genetics and functional genomics. Nucleic Acids Res [Internet]. 2021 Jan 8 [cited 2024
Feb 21];49(D1):D1311–20. Available from: https://pubmed.ncbi.nlm.nih.gov/33045747/

28.   Mountjoy E, Schmidt EM, Carmona M, Schwartzentruber J, Peat G, Miranda A, et al. An
open approach to systematically prioritize causal variants and genes at all published

human GWAS trait-associated loci. Nat Genet [Internet]. 2021 Nov 1 [cited 2024 Feb
21];53(11):1527–33. Available from: https://pubmed.ncbi.nlm.nih.gov/34711957/

29. Lung Disease Knowledge Portal (lungdiseasegenetics.org) [Internet]. [cited 2024 Nov 12].
Available from: https://lung.hugeamp.org/

30. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, et al.
New genetic signals for lung function highlight pathways and chronic obstructive
pulmonary disease associations across multiple ancestries. Nat Genet [Internet]. 2019
Feb 25 [cited 2022 Dec 27];51(3):481–93. Available from:
https://europepmc.org/articles/PMC6397078

31. Shrine N, Izquierdo AG, Chen J, Packer R, Hall RJ, Guyatt AL, et al. Multi-ancestry
genome-wide association analyses improve resolution of genes and pathways influencing
lung function and chronic obstructive pulmonary disease risk. Nat Genet 2023 553
[Internet]. 2023 Mar 13 [cited 2024 Nov 12];55(3):410–22. Available from:
https://www.nature.com/articles/s41588-023-01314-0

32. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. Nat Protoc [Internet]. 2009 [cited 2021 Jan
21];4(1):44–57. Available from: https://pubmed.ncbi.nlm.nih.gov/19131956/

33. Benner C, Spencer CCA, Havulinna AS, Salomaa V, Ripatti S, Pirinen M. FINEMAP:
efficient variable selection using summary data from genome-wide association studies.
Bioinformatics [Internet]. 2016 May 5 [cited 2024 Feb 21];32(10):1493. Available from:
/pmc/articles/PMC4866522/

34. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, et al.
New genetic signals for lung function highlight pathways and chronic obstructive
pulmonary disease associations across multiple ancestries. Nat Genet [Internet]. 2019
Mar 1 [cited 2024 Nov 26];51(3):481–93. Available from:
https://pubmed.ncbi.nlm.nih.gov/30804560/

35. Fulco CP, Nasser J, Jones TR, Munson G, Bergman DT, Subramanian V, et al. Activity-
by-Contact model of enhancer-promoter regulation from thousands of CRISPR
perturbations. Nat Genet [Internet]. 2019 Dec 1 [cited 2024 Feb 7];51(12):1664. Available
from: /pmc/articles/PMC6886585/

36. El Kharbili M, Sasse SK, Sanford L, Jacobson S, Aviszus K, Gupta A, et al. Noncoding
SNPs decrease expression of FABP5 during COPD exacerbations. J Clin Invest. 2024
Feb 1;134(3).

37. Chandramouli S, Chye YY, Yusoff P, Lao DH, Hwei FL, Mizuno K, et al. Tesk1 Interacts
with Spry2 to Abrogate Its Inhibition of ERK Phosphorylation Downstream of Receptor
Tyrosine Kinase Signaling. J Biol Chem. 2008 Jan 18;283(3):1679–91.

38. Ding W, Warburton D. Down-regulation of Sprouty2 via p38 MAPK plays a key role in the
induction of cellular apoptosis by tumor necrosis factor-α. Biochem Biophys Res
Commun [Internet]. 2008 Oct 24 [cited 2025 Apr 21];375(3):460. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC2575769/

39. Masoumi-Moghaddam S, Amini A, Morris DL. The developing story of Sprouty and
cancer. Cancer Metastasis Rev [Internet]. 2014 Apr 18 [cited 2025 Apr 24];33(2–3):695–
720. Available from: https://link.springer.com/article/10.1007/s10555-014-9497-1

40. Gupta A, Sasse SK, Berman R, Gruca MA, Dowell RD, Wei H, et al. Integrated Genomics
Approaches Identify Transcriptional Mediators and Epigenetic 1 Responses to Afghan
Desert Particulate Matter in Small Airway Epithelial Cells 2 3. Available from:
https://doi.org/10.1101/2022.04.27.489761

41. Cascio WE. Wildland fire smoke and human health. 2018 May 15 [cited 2019 Jun
25];624:586–95. Available from: /pmc/articles/PMC6697173/?report=abstract

42. McArdle CE, Dowling TC, Carey K, DeVies J, Johns D, Gates AL, et al. Asthma-
Associated Emergency Department Visits During the Canadian Wildfire Smoke Episodes

— United States, April– August 2023. MMWR Morb Mortal Wkly Rep [Internet]. 2023 Aug 25 [cited 2023 Oct 2];72(34):926–32. Available from: https://www.cdc.gov/mmwr/volumes/72/wr/mm7234a5.htm

43. Edwin F, Anderson K, Ying C, Patel TB. Intermolecular Interactions of Sprouty Proteins and Their Implications in Development and Disease. Mol Pharmacol [Internet]. 2009 Oct [cited 2025 Apr 24];76(4):679. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2769046/

44. Hacohen N, Kramer S, Sutherland D, Hiromi Y, Krasnow MA. sprouty Encodes a Novel Antagonist of FGF Signaling that Patterns Apical Branching of the Drosophila Airways. Cell. 1998 Jan 23;92(2):253–63.

45. Walker DJ, Land SC. Regulation of vascular signalling by nuclear Sprouty2 in fetal lung epithelial cells: Implications for co-ordinated airway and vascular branching in lung development. Comp Biochem Physiol Part B Biochem Mol Biol. 2018 Oct 1;224:105–14.

46. Clarke M, Volpe G, Sheriff L, Walton D, Ward C, Wei W, et al. Transcriptional regulation of SPROUTY 2 by MYB influences myeloid cell proliferation and stem cell properties by enhancing responsiveness to IL-3. Leukemia [Internet]. 2016 Apr 1 [cited 2025 May 7];31(4):957. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5302168/

47. Saini M, Verma A, Mathew SJ. SPRY2 is a novel MET interactor that regulates metastatic potential and differentiation in rhabdomyosarcoma. Cell Death Dis 2018 92 [Internet]. 2018 Feb 14 [cited 2025 Apr 24];9(2):1–15. Available from: https://www.nature.com/articles/s41419-018-0261-2

48. Liu W, Tundwal K, Liang Q, Goplen N, Rozario S, Quayum N, et al. Establishment of Extracellular Signal-Regulated Kinase 1/2 Bistability and Sustained Activation through Sprouty 2 and Its Relevance for Epithelial Function. Mol Cell Biol [Internet]. 2010 Apr 1 [cited 2024 Mar 26];30(7):1783. Available from: /pmc/articles/PMC2838067/

49. Panigrahi A, O'Malley BW. Mechanisms of enhancer action: the known and the unknown. Genome Biol 2021 221 [Internet]. 2021 Apr 15 [cited 2024 Nov 26];22(1):1–30. Available from: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-021-02322-1

50. Gasperini M, Tome JM, Shendure J. Towards a comprehensive catalogue of validated and target-linked human enhancers. Nat Rev Genet [Internet]. 2020 May 1 [cited 2024 Nov 26];21(5):292. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC7845138/

51. Boix CA, James BT, Park YP, Meuleman W, Kellis M. Regulatory genomic circuitry of human disease loci by integrative epigenomics. Nat 2021 5907845 [Internet]. 2021 Feb 3 [cited 2024 Nov 26];590(7845):300–7. Available from: https://www.nature.com/articles/s41586-020-03145-z

52. Gusev A, Lee SH, Trynka G, Finucane H, Vilhjálmsson BJ, Xu H, et al. Partitioning heritability of regulatory and cell-type-specific variants across 11 common diseases. Am J Hum Genet [Internet]. 2014 [cited 2024 Nov 26];95(5):535–52. Available from: https://pubmed.ncbi.nlm.nih.gov/25439723/

53. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. Nature [Internet]. 2015 Feb 2 [cited 2024 Feb 25];518(7539):317. Available from: /pmc/articles/PMC4530010/

54. Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, et al. An integrated encyclopedia of DNA elements in the human genome. Nat 2012 4897414 [Internet]. 2012 Sep 5 [cited 2024 Feb 25];489(7414):57–74. Available from: https://www.nature.com/articles/nature11247

55. Kircher M, Witten DM, Jain P, O'roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. Nat Genet 2014 463 [Internet]. 2014 Feb 2 [cited 2024 Feb 25];46(3):310–5. Available from: https://www.nature.com/articles/ng.2892

56. Ionita-Laza I, Mccallum K, Xu B, Buxbaum JD. A spectral approach integrating functional

genomic annotations for coding and noncoding variants. Nat Genet 2015 482 [Internet]. 2016 Jan 4 [cited 2024 Feb 25];48(2):214–20. Available from: https://www.nature.com/articles/ng.3477

57. Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, et al. Transcriptional super-enhancers connected to cell identity and disease. Cell [Internet]. 2013 Nov 7 [cited 2025 May 7];155(4):10.1016/j.cell.2013.09.053. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC3841062/

58. Gibson G, Powell JE, Marigorta UM. Expression quantitative trait locus analysis for translational medicine. Genome Med 2015 71 [Internet]. 2015 Jun 24 [cited 2024 Nov 26];7(1):1–14. Available from: https://genomemedicine.biomedcentral.com/articles/10.1186/s13073-015-0186-7

59. Guerrina N, Traboulsi H, Eidelman DH, Baglole CJ. The Aryl Hydrocarbon Receptor and the Maintenance of Lung Health. Int J Mol Sci [Internet]. 2018 Dec 1 [cited 2022 Nov 8];19(12). Available from: http://www.ncbi.nlm.nih.gov/pubmed/30563036

60. Iu M, Zago M, Rico de Souza A, Bouttier M, Pareek S, White JH, et al. RelB attenuates cigarette smoke extract-induced apoptosis in association with transcriptional regulation of the aryl hydrocarbon receptor. Free Radic Biol Med [Internet]. 2017 Jul 1 [cited 2020 Feb 20];108:19–31. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28254546

61. Wegesser TC, Pinkerton KE, Last JA. California wildfires of 2008: Coarse and fine particulate matter toxicity. Environ Health Perspect [Internet]. 2009 [cited 2020 Sep 30];117(6):893–7. Available from: /pmc/articles/PMC2702402/?report=abstract

62. Zhao F, Klimecki WT. CULTURE CONDITIONS PROFOUNDLY IMPACT PHENOTYPE IN BEAS-2B, A HUMAN PULMONARY EPITHELIAL MODEL. J Appl Toxicol [Internet]. 2014 Aug 1 [cited 2025 May 7];35(8):945. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4474793/

63. Garcia-Canton C, Minet E, Anadon A, Meredith C. Metabolic characterization of cell systems used in in vitro toxicology testing: Lung cell system BEAS-2B as a working example. Toxicol Vitr [Internet]. 2013 Sep 1 [cited 2025 Apr 25];27(6):1719–27. Available from: https://www.sciencedirect.com/science/article/pii/S0887233313001239

64. Han X, Na T, Wu T, Yuan BZ. Human lung epithelial BEAS-2B cells exhibit characteristics of mesenchymal stem cells. PLoS One [Internet]. 2020 Jan 1 [cited 2025 Apr 25];15(1):e0227174. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6941928/

65. Georas SN, Rezaee F. Epithelial barrier function: at the frontline of asthma immunology and allergic airway inflammation. J Allergy Clin Immunol [Internet]. 2014 Sep 1 [cited 2025 May 7];134(3):509. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4170838/

66. Gref A, Merid SK, Gruzieva O, Ballereau S, Becker A, Bellander T, et al. Genome-wide interaction analysis of air pollution exposure and childhood asthma with functional follow-up. Am J Respir Crit Care Med [Internet]. 2017 May 15 [cited 2024 Feb 25];195(10):1373–83. Available from: /pmc/articles/PMC5443897/

67. Melén E, Kho AT, Sharma S, Gaedigk R, Leeder JS, Mariani TJ, et al. Expression analysis of asthma candidate genes during human and murine lung development. Respir Res 2011 121 [Internet]. 2011 Dec 1 [cited 2025 May 7];12(1):1–10. Available from: https://respiratory-research.biomedcentral.com/articles/10.1186/1465-9921-12-86

68. Evans CM, Kim K, Tuvim MJ, Dickey BF. Mucus hypersecretion in asthma: causes and effects. Curr Opin Pulm Med [Internet]. 2009 Jan [cited 2025 May 7];15(1):4. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2709596/

69. Líbalová H, Závodná T, Vrbová K, Sikorová J, Vojtíšek-Lom M, Beránek V, et al. Transcription profiles in BEAS-2B cells exposed to organic extracts from particulate emissions produced by a port-fuel injection vehicle, fueled with conventional fossil

gasoline and gasoline-ethanol blend. Mutat Res Toxicol Environ Mutagen [Internet]. 2021
Dec 1 [cited 2025 Apr 25];872:503414. Available from:
https://www.sciencedirect.com/science/article/pii/S1383571821001054?via%3Dihub
70.   Grilli A, Bengalli R, Longhin E, Capasso L, Proverbio MC, Forcato M, et al.
Transcriptional profiling of human bronchial epithelial cell BEAS-2B exposed to diesel and
biomass ultrafine particles. BMC Genomics [Internet]. 2018 Apr 27 [cited 2025 Apr
25];19(1):302. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5923024/
830

A

Dynamically regulated WSP-TREs (Tfit)
N = 18,553

Common SNPs (MAF > 0.05)
(+/-) 50 bp WSP-TRE centers

Associated with Asthma
phenotype = WSP-SNPs
N = 52

B

- Exon - coding
- 3' UTR
- 5' UTR
- Intergenic
- Intron

C

Beas-2B
PRO-seq

HK2

VEH

WSP30

WSP120

500 bp

0.6
-0.3
0.6
-0.3
0.6
-0.3

rs28362965

Table 3

Table 4

Table 5

A

WSP (1 mg/ml) 2h
WSP (1 mg/ml) 4h

B HASM
   Normal   Asthma
   VEH   TNF   VEH   TNF

SPRY2 ————————————————— 37 kDa

GAPDH ————————————————— 37 kDa