Resurgence of Large-Scale Influenza A (H1N1) Outbreaks: the Interplay of Transmission, Loss of Immunity, and Vaccination.

Chanaka Kottegoda[1], Claudia T. Codeço[2], Claudio J. Struchiner[3], and Lucas M. Stolerman[4,*]

[1]Department of Mathematics and Physics, Marshall University, Huntington, WV, USA

[2]Programa de Computação Científica, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

[3]School of Applied Mathematics (EMAp FGV), Brazil

[4]Department of Mathematics, Oklahoma State University, Stillwater, OK, USA

[*]Corresponding author. *E-mail address:* lucas.martins_stolerman@okstate.edu

Abstract

Influenza is a highly transmissible respiratory virus that can cause devastating pandemics. The last influenza pandemic, caused by a strain of A(H1N1) virus, resulted in millions of cases and hundreds of thousands of deaths worldwide. After the initial outbreaks, many regions experienced subsequent waves driven by environmental conditions, waning immunity, and other factors. More dramatically, several locations saw large resurgent outbreaks, yet the mechanisms behind these severe resurgences remain poorly understood. Here we investigate the dynamics of these large-scale H1N1 outbreaks using a mechanistic model with time-varying rates of infection, loss of immunity, and vaccination. In particular, we explore two simple

mathematical functions to model dynamic loss of immunity, capturing how viral evolution may reduce the duration of immune protection. Numerical simulations reveal regimes influencing resurgence timing, and the model reproduces key features of H1N1 flu case data. We also fit the model to data from nine locations, demonstrating its ability to capture resurgence patterns and track associated changes in infection and immunity-loss rates. Our findings highlight the complex interplay among environmental drivers, population immunity, and viral mutation, contributing to a better understanding of large-scale H1N1 dynamics in the post-pandemic era.

Keywords. A(H1N1) influenza; Resurgent outbreaks; Mechanistic modeling

Introduction

A new strain of influenza A(H1N1) virus emerged in the spring of 2009 in central Mexico and rapidly spread worldwide [1]. The Centers for Disease Control and Prevention (CDC) reported the first US cases in April 2009, and the World Health Organization (WHO) declared a pandemic on June 11, 2009 [2]. From April 2009 to April 2010, the CDC reported 60.8 million cases, 274,304 hospitalizations, and 12,469 deaths in the United States alone [3]. Globally, many countries – including Brazil, China, Australia, and the United Kingdom – experienced significant surges between 2009 and 2010 [4, 5, 6, 7, 8]. The 2009 H1N1 virus has continued to circulate, causing subsequent outbreaks [9].

The mechanisms underlying sustained H1N1 transmission in the post-pandemic period are understood to an extent. During the 2009 outbreaks, susceptible populations were not fully depleted, allowing for subsequent waves of infection. As the number of susceptible individuals declined below a critical threshold, outbreaks became less frequent, resulting in "skip-and-resurgence" patterns [10]. Complex oscillations in influenza epidemics are thought to arise from a combination of epidemiological, immunological (humoral), and seasonal drivers, many of which have been successfully incorporated into mathematical models. Immunity acquired through natural infection wanes

over time, while cellular immunity might last longer and play a role in the final dynamics [11]. Seasonal variations in transmissibility drive periodic increases in transmission [12, 13, 14, 15]. Poor vaccination coverage, suboptimal vaccine efficacy, and demographic changes further contribute to sustaining outbreaks [16, 17, 18]. Antigenic changes shape outbreak dynamics by rendering previously immune individuals susceptible to emerging strains [19, 20, 21].

Large H1N1 outbreaks have occurred worldwide in the post-pandemic era, often reaching incidence levels comparable to those of the initial 2009 pandemic. For instance, an estimated 30 million influenza illnesses, 13 million medical visits, 347,000 hospitalizations, and 38,000 deaths occurred in the United States during the 2013–2014 influenza season, when 2009 H1N1 viruses were the predominant circulating strains, primarily affecting young and middle-aged adults [22, 23, 24]. In 2016, Brazil experienced a severe outbreak with more than a thousand confirmed deaths [25]. Iran similarly faced a substantial resurgence during the 2015–2016 season, with elevated case counts and mortality [26, 27]. Similar spikes were reported in Turkey (2016), South Africa (2011), Croatia (2011), among other regions [28, 29, 30]. Although prior studies have proposed explanations for influenza epidemics, to our knowledge, no modeling efforts have explored how the dynamic landscape of transmission, waning immunity, and vaccination coverage contributes to large resurgent H1N1 epidemics.

In this study, we developed a mechanistic model to investigate large-scale H1N1 outbreaks in the post-pandemic era. Our framework incorporates time-varying rates of infection, loss of immunity – which may occur naturally or through viral evolution – and vaccination. In particular, we explore two simple functions to model dynamic immunity loss, capturing how viral evolution can shorten the duration of immunity in recovered individuals. Through numerical simulations, we identify parameter regimes that influence resurgence timing and show that our model reproduces key patterns observed in H1N1 time series data. By fitting the model to data from nine locations, we explore its ability to capture large resurgent outbreaks, uncovering temporal trends in infection and loss-of-immunity rates.

Materials and Methods

Description of epidemiological and demographic datasets

We identified locations that experienced large resurgent H1N1 outbreaks in the post-pandemic period by examining their weekly reported H1N1 case counts. Defining what constitutes a large resurgence is challenging and can be approached in multiple ways. Here, a large resurgent H1N1 outbreak is defined as one whose epidemic peak (maximum number of reported cases) exceeds 50% of the major peak observed during the 2009 pandemic. Major outbreaks in 2010 were classified as late-pandemic activity rather than resurgences. Fig. 1 A) displays the nine selected locations: São Paulo State in Brazil, Turkey, Iran, New Zealand, the U.S. Department of Health and Human Services (HHS) Regions 2 and 6, Colombia, South Africa, and Croatia. The HHS Region 2 includes the states of New Jersey and New York, as well as Puerto Rico and the U.S. Virgin Islands. The HHS Region 6 comprises the states of Arkansas, Louisiana, New Mexico, Oklahoma, and Texas.

Fig. 1 B) displays the time series of weekly reported H1N1 cases. Some epidemiological datasets exhibited gaps in the data. In particular, New Zealand had 103 missing data points out of 273 weeks, Iran had eight missing data points out of 351 weeks, and Turkey had 69 missing data points out of 364 weeks. In this work, we did not perform data imputation or estimation for the missing values; only the available data were used in the model-fitting analysis. In Fig. 1 B), the large resurgent outbreaks are marked with a red star for each location. Epidemiological data from HHS Regions 2 and 6 were obtained from the Fluview interactive system [31]. H1N1 case counts for São Paulo were collected from the Brazilian Information System for Influenza Epidemiological Surveillance (SIVEP). The rest of the epidemiological data for other locations were obtained from FluNet [32], the World Health Organization's (WHO) influenza surveillance system, which provides real-time, publicly accessible epidemiological data. For each location, we also collected population data from 2009 to the year of the large resurgence outbreak, sourced

from Worldbank.org [33]. For datasets and further details, we refer the reader to the supplementary material.

Figure 1: **Weekly reported A(H1N1) influenza cases in selected regions.** (A) Geographical locations of nine selected regions from eight countries that experienced large resurgent H1N1 outbreaks after the 2009 pandemic. (B) Weekly reported H1N1 cases in the selected regions, starting from the 2009 pandemic until the year of the large resurgent outbreak (marked by a red star). The visible gaps between the blue curves in New Zealand and Turkey indicate missing data.

Mathematical Model

We present our SVIRS model to investigate the dynamics of large resurgent H1N1 outbreaks, as illustrated in Fig. 2 A) i). To maintain mathematical simplicity, the model omits age-structured dynamics, heterogeneous contact patterns, and exposure periods. Instead, we focus on the sole interplay of time-varying rates of infection, loss of immunity, and vaccination to provide insights into the mechanisms driving large resurgent outbreaks. The governing equations are given by the following system:

$$
\begin{aligned}
\dot{S} &= \Lambda - \beta(t)S\frac{I}{N} - \phi S + \gamma(t)R - \epsilon v(t)S + \omega V, \\
\dot{V} &= \epsilon v(t)S - \omega V - \beta_v(t)V\frac{I}{N}, \\
\dot{I} &= \beta(t)S\frac{I}{N} + \phi S - rI + \beta_v(t)V\frac{I}{N}, \\
\dot{R} &= rI - \gamma(t)R,
\end{aligned}
\tag{1}
$$

Here, S, V, I, and R represent the susceptible, vaccinated, infectious, and recovered individuals, respectively. Overdots denote derivatives with respect to time. The total population is given by $N = S + V + I + R$. The time-varying rates are assumed to reflect dynamic changes driven by complex biological factors, which are discussed in detail in the next subsection. Here, $\beta(t)$ denotes the infection rate at which susceptible individuals become infectious. The parameter $\gamma(t)$ represents the loss-of-immunity rate, and $v(t)$ the vaccination rate. Additionally, $\beta_v(t) = \xi\beta(t)$ denotes the infection rate at which vaccinated individuals become infectious. Here ξ is a factor between 0 and 1 representing the vaccine efficacy ($\xi = 0$ corresponds to a vaccine that confers perfect protection). The recruitment rate Λ is constant and thus $N(t)$ satisfies the differential equation $\dot{N} = \Lambda$, leading to $N(t) = \Lambda t + N_{2009}$, where N_{2009} denotes the total population in 2009. This linear trend provides a suitable approximation for the complex demographic evolution observed in the

selected locations during the initial years following the 2009 H1N1 pandemic (see supplementary material). Natural recovery rates are represented by the parameter r, while ω describes the rate at which vaccine-induced immunity wanes. Finally, ϵ and ϕ represent the vaccine effectiveness and a small importation rate, respectively. We list all fixed (time-independent) parameters in Table 1 (see supplementary material for details).

Parameter	Description	Value
Λ	Recruitment rate into the susceptible class	Location-specific; estimated from demographic data (see supplementary material)
r	Natural recovery rate	1.17 week^{-1} [34]
ϕ	Importation rate	10^{-6} week^{-1} (one imported case per million susceptibles per week)
ϵ	Vaccine effectiveness	0.7 [35]
ω	Waning rate of vaccine-induced immunity	0.019 week^{-1} [36]

Table 1: Time-independent model parameters: description and values.

Model assumptions for time-varying rates.

Periodic infection rates

Influenza transmission depends on multiple environmental and social factors with seasonal components, including humidity [37, 38, 39], human mobility [40, 41, 42], and school terms [43, 44]. We adopt a time-varying periodic infection rate, $\beta(t)$, given by the following expression:

$$\beta(t) = A \sin\left(Bt + C\right) + D. \tag{2}$$

The parameters B and C correspond to the angular frequency and phase shift of the sine function, while A and D represent its amplitude and vertical shift. To parameterize $\beta(t)$ in terms of its

maximum and minimum values $\beta_{\max}$ and $\beta_{\max}$, we can simply write $A = \dfrac{\beta_{\max} - \beta_{\min}}{2}$ and $D = \dfrac{\beta_{\max} + \beta_{\min}}{2}$. Fig. 2 B) i) illustrates an example of a periodic $\beta(t)$, representing periodic changes in the infection rate.

Loss-of-immunity rates

The influenza virus evolves through two main processes: antigenic drift and antigenic shift [45]. While antigenic shift can lead to the abrupt emergence of novel subtypes – such as the 2009 H1N1 pandemic strain – our modeling approach focuses on antigenic drift, the gradual accumulation of mutations that help the virus evade host immune responses [46, 47, 48]. We represent drift using two formulations: a linear model, which assumes continuous immune escape over time, and a jump model, which describes a punctuated antigenic change. This latter mechanism is consistent with evidence that a single mutation at a key site, such as on the hemagglutinin (HA) protein, can render the virus antigenically distinct [49]. Mathematically, we consider the following functions:

$$\gamma(t) = \left(\frac{\gamma_{\max} - \gamma_0}{T} \right) t + \gamma_0 \tag{3}$$

and

$$\gamma(t) = \gamma_0 + (\gamma_{\max} - \gamma_0)U(t - \tau). \tag{4}$$

For the linear model (Eq. (3)), γ_0 and $\gamma_{\max}$ denote the baseline and maximum rates of loss of immunity, respectively, with T the interval over which γ rises from γ_0 to $\gamma_{\max}$. We also set T in weeks based on the time between the initial outbreak and the year of the large resurgence in each location. Throughout this paper, we fix $\gamma_0 = 3.2 \times 10^{-3}\,\text{week}^{-1}$ (see supplementary material for details). In the jump model (Eq. (4)), $\gamma_{\max}$ again defines the post-jump maximum loss rate, and τ specifies the time of the discrete increase. The term $U(t - \tau)$ represents a step function assuming the values 0 if $t < \tau$ and 1 if $t \geq \tau$. In what follows, we refer to the *linear γ model* whenever the SVIRS model (Eqs. (1)) is equipped with the linear function given by Eq. (3). Similarly, the *jump*

γ *model* refers to the SVIRS model with $\gamma(t)$ given by Eq. (4).

Vaccination rates

The frequent antigenic changes prompt re-evaluations of influenza vaccines, which are typically administered yearly before the flu season. For example, in Brazil, vaccines are offered to high-risk groups through annual campaigns between April and May, a period that generally precedes the major outbreaks [50]. In our model, we adopted a rectangular pulse vaccination rate $v(t)$ that achieves the maximum level v_{max} during the vaccination period for a given location (see supplementary material). Fig. 2 B) iii) illustrates the format of the $v(t)$ function.

Figure 2: **Large resurgent H1N1 outbreaks and our proposed SVIRS mathematical model.** A) i) The resurgence of large H1N1 influenza outbreak in year x is defined by the condition $r_x > 0.5r_{2009}$. Here r_{2009} represents the epidemic peak during the 2009 H1N1 pandemic, while r_x is the same quantity in year x. A) ii) A compartmental diagram of our SVIRS model. A population is divided into susceptible (S), vaccinated (V), infectious (I), and recovered individuals (R). The rates of Infection, loss of immunity, and vaccination are assumed to change over time. B) i) A sine function was used to approximate the periodic infection rate $\beta(t)$, with minimum and maximum values $\beta_{\min}$ and $\beta_{\max}$. B) ii). Linear and Jump functions represent different effects of antigenic drift on the loss-of-immunity rate $\gamma(t)$ that increases from γ_0 to $\gamma_{\max}$. B) iii). A rectangular pulse function represented the vaccination rate, which increases to $v_{\max}$ during vaccination periods.

Numerical Simulations

We performed numerical simulations to explore parametric regimes that influence the timing of large resurgent outbreaks and to evaluate our model's ability to capture the overall characteristics of reported H1N1 case time series. Fig. 3 displays four sets of simulations conducted under different

vaccine efficacy factors (ξ), considering the linear γ model (Eq. (3)). Results for $\xi = 0, 0.3, 0.6$, and 0.9 are shown in panels A)–D). The colormaps depict the time interval (in years) between large outbreak peaks as a function of the minimum immunity period $D_{\min} = \frac{1}{\gamma_{\max}}$ and the maximum infection rate $\beta_{\max}$. For all colormaps, the time interval between large outbreaks tends to increase as the immunity period lengthens. Lower vaccine efficacy (i.e., larger ξ values) resulted in shorter time intervals between large outbreaks. That can be observed as colormaps become darker as ξ increases. We also display sample model trajectories for selected points in the colormaps, which exhibit decreasing time intervals between large outbreaks as ξ increases. The leftmost time series of each colormap shows the modeled weekly H1N1 cases for low $\beta_{\max}$ and $D_{\min}$. The resulting time interval between the first and large resurgent outbreaks decreases from 4 years to 1 year, as ξ changes from 0 to 0.6. Within each colormap, longer immunity periods tend to delay the occurrence of large resurgent outbreaks. For instance, in Fig. 3 A), three sample trajectories are highlighted. At the lowest $D_{\min}$ (approximately 8 weeks [51]), a large resurgent outbreak occurs 4 years after the initial outbreak. In contrast, at the highest $D_{\min}$ value (approximately 30 weeks), no large resurgent outbreak is observed within ten years, as indicated by the dark blue region marked (NR).

Figure 3: **Years between large outbreak peaks along with sample trajectories (linear γ model).** Panels (A)–(D) show colormaps of the time interval (in years) between large outbreak peaks for increasing values of ξ (0, 0.3, 0.6, 0.9), corresponding to decreasing vaccine efficacy. Dark blue regions labeled "NR" indicate no large outbreaks within the past ten years. Beneath each colormap are sample trajectories of the normalized infectious population. Gray dashed lines indicate the threshold used to define large resurgent outbreaks (red colored). Parameter values were taken from Table 1. Initial population $N_{2009} = 42,075,716$ and recruitment rate $\Lambda = 137.9351$ week^{-1} were taken from São Paulo, Brazil. Other fixed parameters include $\beta_{\min} = 1$ week^{-1}, $v_{\max} = 0.02$ week^{-1}, $\gamma_0 = 3.2 \times 10^{-3}$ week^{-1}, and $T = 520$ weeks (10 years). For the periodic infection rate, we set $B = -0.1202$ week^{-1} and $C = -10.1360$ to produce annual cycles.

Fig. 4 shows numerical simulations of the jump γ model (Eq. (4)). We set $\tau = 130$ weeks (2.5 years), representing antigenic changes occurring 130 weeks after the initial outbreak, and $v_{\text{max}} = 0.01$ week^{-1} while all other parameters remain the same as in Fig. 3. The predominance of light blue colormap regions across panels A–D indicates immediate large resurgent outbreaks following the jump, while yellow regions highlight outbreaks delayed by several years. Sample trajectories below each colormap illustrate these varied dynamics; Notably, the middle time series in panels C and D exhibit resurgent outbreaks that slightly exceed the threshold (gray dashed line) approximately seven years after the initial wave.

Figure 4: **Years between large outbreak peaks along with sample trajectories (jump γ model).** Panels (A) - (D) exhibit four colormaps for increasing values of ξ, corresponding to decreasing vaccine efficacy levels. We considered four different ξ values ($\xi = 0, 0.3, 0.6,$ and 0.9). Here, $\xi = 0$ represents a vaccine that confers perfect protection (panel A). Dark blue regions indicate no large resurgent outbreaks within ten years (labeled "NR"). Here, we set $\tau = 2.5$ years and $v_{max} = 0.01$ week^{-1} while all other parameter values remain the same as in Fig. 3.

Fitting the model to H1N1 data

We evaluated our model's ability to fit H1N1 case data across the nine locations shown in Fig. 1 – São Paulo State in Brazil, Turkey, Iran, New Zealand, HHS Regions 2 and 6 in the US, Colombia, South Africa, and Croatia. These selected locations experienced a large resurgence of H1N1 outbreaks within a decade of the 2009 pandemic, a timeframe that captures the combined effects of waning immunity, cycles of higher transmission, and continued vaccination campaigns. For each location, we perform a least-squares optimization, fitting the simulated reported cases – modeled as a constant fraction (ρ) of the total infectious population at a given time and location – to the available case data. Here, ρ represents the reporting rate, that is, the proportion of actual infections that are detected and recorded in surveillance systems.

For the linear γ model, we estimated eight unknown parameters for each location: the angular frequency and phase shift (B, C) of the periodic infection rate (Eq. (2)), the maximum infection rate ($\beta_{\max}$), the maximum loss-of-immunity rate ($\gamma_{\max}$), the vaccination level ($v_{\max}$), the vaccine efficacy (ξ), the reporting rate (ρ), and the initial infectious population $I(0)$ (while fixing $V(0) = R(0) = 0$). For simulations involving the jump γ model (Eq. (4)), we estimated a total of nine parameters: the eight parameters listed above, plus the timing of the jump τ in the γ function (Eq. (4)). The remaining parameters were fixed at the values listed in Table 1, along with $\gamma_0 = 3.2 \times 10^{-3}$ week^{-1} (see supplementary material). We also fixed $\beta_{\min} = 1$ week^{-1} so that during off-peak seasons we have $\beta(t) < r$, ensuring natural epidemic decline in troughs, while allowing $\beta_{\max}$ to drive seasonal surges. To estimate the best-fit curves, each parameter was assigned four initial guesses, evenly spaced between reasonable lower and upper bounds (see data and codes for details). For the linear γ model, this resulted in $4^8 (65,536)$ total simulations. The system of ODEs was numerically integrated in MATLAB using the `ode45` function, which employs a fourth-order Runge–Kutta method. Optimization was performed with MATLAB's `lsqcurvefit` function. For each initial guess, a best-fit trajectory was obtained by running the least-squares procedure. Among all 4^8 resulting trajectories, the final best-fit curve was selected as the one that minimized

the sum of squared residuals (SSR). A similar procedure was applied to the jump γ model, where three equally spaced initial guesses were used per parameter, resulting in 3^9 $(19,683)$ simulations. As this was a sparser search compared to the 4^8 initial guesses used for the linear model, we conducted an additional 10,000 simulations using randomly sampled starting points to enhance the search for the global minimum.

The best-fit curves for the linear γ model are presented in Fig. 5. Uncertainty bounds, shown in purple shades, were obtained using a parametric bootstrapping approach [52]. Overall, the linear γ model accurately captured the timing of most initial and large resurgent outbreaks. The closest fits occurred in South Africa, Croatia, and HHS Region 2 (USA). The model fits also showed reasonable agreement for São Paulo, New Zealand, and Iran, although sometimes over- or underestimating outbreak magnitudes. Remarkably, despite substantial gaps in the New Zealand data, the model recovered the large outbreaks and even inferred a surge around 2012 during the unreported interval. In contrast, mid-sized surges were less accurately captured in other locations. Model performance was slightly weaker in HHS Region 6 (USA), Turkey, and Colombia; notably, Colombia was the only location where the model failed to capture a clear resurgent wave, likely reflecting higher noise in the reported data.

At the bottom of each best-fit panel in Fig. 5, the estimated rates of infection and loss of immunity illustrate the interplay between these two factors in shaping H1N1 dynamics up to the large resurgent outbreak. A flexible range of oscillations in the infection rate (dark blue) was observed, suggesting that complex transmission spikes arise from factors beyond seasonal influences. This implies that climate-driven oscillations alone may not fully explain the intricate temporal patterns of H1N1 outbreaks, particularly the resurgence events. The loss-of-immunity rate $\gamma(t)$ is plotted in green on a separate y-axis. Notably, Iran's panel shows a virtually flat $\gamma(t)$ – no appreciable increase in immunity loss – yet the model still achieves a good fit. This outcome demonstrates our model's ability to infer a possible absence of a significant antigenic drift over the considered period. All estimated parameters can be found in the supplementary material (Tables S3 and S4).

HHS Regions 2 and 6 in the United States experienced two pronounced H1N1 waves—in 2009 and again in 2010. Croatia followed a similar trajectory, with a modest preliminary wave occurring before its major 2009 outbreak. To account for these early peaks in 2009, we fitted the first outbreak an $SIRS$ model with constant rates of infection and loss of immunity. The resulting values of S, I, and R, together with $V = 0$, were then used as initial conditions for a second simulation using our SVIRS model, targeting the period between the 2009–2010 waves and the later large resurgence.

Figure 5: **Best-fit curves and optimal time-varying rates (linear γ model).** The top panels show observed (blue) and modeled (red) weekly H1N1 cases for nine locations, with shaded uncertainty bounds. Bottom panels display the fitted infection rate $\beta(t)$ (purple) and linearly increasing immunity-loss rate $\gamma(t)$ (green).

Fig. 6 presents a similar set of best-fit results for the jump γ model. The model captures the overall timing and magnitude of large resurgent outbreaks in most locations. The bottom panels display the fitted rates of infection and loss of immunity. Interestingly, the jump γ model enabled exploration of antigenic change timing. For instance, in South Africa, a jump was inferred between 2010 and 2011, reasonably close to the resurgent outbreak. In contrast, in São Paulo, the jump occurred closer to 2010, while the resurgent outbreak only took place in 2016. This finding suggests that large resurgent outbreaks can be preceded by antigenic changes occurring several years earlier. In other locations, such as Croatia and HHS Region 2 (USA), the best-fit solutions exhibited jumps at the beginning of the time series, suggesting that a constant γ model might better describe influenza dynamics in these regions. Oscillation periods of $\beta(t)$ varied by location, as in the linear γ model. Best-fit solutions with higher-frequency $\beta(t)$ oscillations were observed in South Africa, Croatia, and São Paulo, while lower-frequency oscillations were found in HHS Regions 2 and 6 (USA). As in the linear γ model, these variable oscillations likely reflect complex phenomena such as the interplay between seasonal factors and fluctuations in contact patterns.

Figure 6: **Best-fit curves and time-varying rates (jump γ model).** Top panels show observed (blue) and modeled (red) weekly H1N1 cases for nine locations, with shaded uncertainty bounds. Bottom panels display the fitted infection rate $\beta(t)$ (purple) and jump immunity-loss rate $\gamma(t)$ (green).

Table 2 summarizes the performances of the linear and jump γ models across different locations. Model performance was evaluated using both Root Mean Square Error (RMSE) and Mean Absolute Error (MAE), with the lowest errors highlighted in yellow. The linear γ model outperformed the jump γ model on both metrics for São Paulo, Croatia, New Zealand, South Africa, and HHS Region 2 (USA). The jump γ model provided the best performance only in Turkey, although neither model captured the large outbreaks there particularly well. In Colombia, Iran, and HHS Region 6 (USA), the preferred model differed depending on the error metric. Overall, RMSE and MAE values were relatively low across most locations, indicating good fitting accuracy. Turkey was a notable exception, with much larger errors for both models, suggesting greater difficulty in capturing outbreak dynamics. In locations where the preferred model differed depending on the metric (Colombia, Iran, and HHS Region 6), the differences in error values were small, indicating that both models provided comparable fits.

Location	γ Model	RMSE	MAE
São Paulo, Brazil	linear	43.5778	13.8190
	jump	47.2714	16.7395
Colombia	linear	17.5558	9.7925
	jump	17.6997	9.1485
Croatia	linear	22.3348	11.1401
	jump	26.3099	12.9522
Iran	linear	49.7471	18.77895
	jump	67.3555	15.4817
New Zealand	linear	13.8952	6.2309
	jump	15.7949	6.6599
South Africa	linear	14.3537	8.3187
	jump	15.9148	9.4717
Turkey	linear	178.1758	51.76
	jump	175.4567	48.2271
HHS Region 2, USA	linear	21.3360	10.9899
	jump	22.0727	11.9082
HHS Region 6, USA	linear	77.6263	29.6823
	jump	76.2881	30.0496

Table 2: **Model performance comparison (RMSE and MAE).** Root Mean Square Error (RMSE) and Mean Absolute Error (MAE) are shown for the linear and jump γ models across all locations. The lowest error for each location is highlighted in yellow.

Discussion

Large-scale resurgent H1N1 outbreaks have emerged worldwide in the post-pandemic era, often reaching incidence levels comparable to those observed during the 2009 pandemic. To investigate

these resurgences, we developed a mechanistic SVIRS model with assumptions grounded in biological and demographic drivers. Infection rates fluctuated periodically to reflect the impact of environmental and social factors on transmission. We modeled loss of immunity dynamically using linear and jump γ functions to represent different effects of antigenic drift. Vaccination rates were represented by a time-varying pulse to reflect targeted seasonal campaigns. We investigated how different parameter combinations influence outbreak dynamics and assessed model fits to historical H1N1 data across nine locations.

Using numerical simulations, we identified parameter regimes that influence the timing of large resurgent outbreaks. As shown in Fig. 3, longer immunity durations increased the interval between large peaks, while reduced vaccine efficacy shortened these intervals. We then fitted both linear and jump γ models to historical H1N1 data from nine locations. Both models generally captured the timing of initial and resurgent outbreaks, though accuracy varied across locations. The closest fits occurred for South Africa, Croatia, and HHS Region 2 (USA), where the models effectively captured outbreak peaks. In contrast, fits for São Paulo, New Zealand, Iran, Turkey, and Colombia showed some discrepancies, particularly in capturing outbreak magnitudes or precisely aligning with observed peaks. The fitted infection-rate oscillations varied considerably, suggesting that additional drivers beyond seasonal forcing may contribute to resurgent outbreaks.

Two time-varying functions were proposed to model how waning immunity replenishes the susceptible pool. Previous work by Yaari et al. employed a more sophisticated version of the jump γ model, incorporating multiple jumps corresponding to different influenza seasons to capture seasonal dynamics [53]. In contrast, our approach simplifies this idea by modeling immunity loss through a single jump or a linear increase. Biological evidence supports this simplification, as the H1N1pdm strain has remained antigenically stable in recent years [54]. To our knowledge, a linear function has not previously been used to model the effects of antigenic drift. It remains an open question whether using an evolution-informed approach—as in Du et al. [21]—would yield substantially different estimates for $\gamma(t)$. In some scenarios, such as regions that experienced a single

dominant antigenic change, we suspect that our simplified $\gamma(t)$ functions may closely approximate more complex evolution-informed structures.

While our model captured key features of large resurgent outbreaks, the modest fitting performance in some locations suggests that further refinement is needed to capture some epidemic dynamics. Our current framework does not account for multi-strain dynamics, which are known to influence influenza patterns and may contribute to discrepancies in outbreak timing and magnitude [55, 56]. Additional extensions could involve age-structured transmission, human mobility, and heterogeneous contact patterns [57, 58, 59]. Future studies can also explore hybrid models combining linear and jump components to reflect biologically plausible regimes in which variants with few mutations are followed by those with many more, as observed in chronic SARS-CoV-2 infections [60].

Our modeling framework and insights, though developed in the context of H1N1 outbreaks, are broadly applicable to other influenza viruses with similar antigenic and seasonal characteristics. The interplay between transmission variability, waning immunity, and vaccination coverage is not unique to H1N1 and could help explain resurgence dynamics across different influenza subtypes. In this sense, our approach contributes to a broader understanding of how time-dependent biological and behavioral factors shape the dynamics of respiratory infections. By exploring the effects of environmental variability, immune landscape evolution, and vaccination strategies within a simplified modeling framework, this work enhanced our understanding of large-scale influenza resurgence and may help inform future preparedness strategies.

Acknowledgments

We thank Priscila Born at IOC/FIOCRUZ, Leonardo Bastos, Marcelo Gomes, and the team at PROCC/FIOCRUZ for the insightful comments and support on this project. LMS acknowledges

support from the National Science Foundation (NSF DMS-2327844). Some of the computing for this project was performed at the High-Performance Computing Center at Oklahoma State University, supported in part by the National Science Foundation (NSF OAC-1531128).

References

[1] Walter N Harrington, Christina M Kackos, and Richard J Webby. The evolution and future of influenza pandemic preparedness. *Experimental & molecular medicine*, 53(5):737–749, 2021.

[2] Centers for Disease Control and Prevention. 2009 pandemic timeline, 2009. Available at https://archive.cdc.gov/www_cdc_gov/flu/pandemic-resources/2009-pandemic-timeline.html. Accessed: 2025-02-16.

[3] National Library of Medicine. The world's first public health emergency of international concern, 2020. Available at https://circulatingnow.nlm.nih.gov/2020/04/21/the-worlds-first-public-health-emergency-of-international-concern/. Accessed: 2025-02-16.

[4] Cláudia Torres Codeço, Josiane da Silva Cordeiro, Arthur Weiss da Silva Lima, Rodrigo Amarante Colpo, Oswaldo Gonçalves Cruz, Flavio Codeço Coelho, Paula Mendes Luz, Claudio José Struchiner, and Fernando Ribeiro de Barros. The epidemic wave of influenza a (h1n1) in brazil, 2009. *Cadernos de Saúde Pública*, 28:1325–1336, 2012.

[5] Wanderson Kleber de Oliveira, Eduardo Hage Carmo, Gerson Oliveira Penna, Ricardo de Souza Kuchenbecker, HB Santos, Wildo Navegantes de Araújo, Regina Malaguti, Bruce B Duncan, Maria Inês Schmidt, surveillance Team for the pandemic influenza A (H1N1) 2009 in the Ministry of Health, et al. Pandemic h1n1 influenza in brazil: analysis of

the first 34,506 notified cases of influenza-like illness with severe acute respiratory infection (sari). *Eurosurveillance*, 14(42):19362, 2009.

[6] W Liang, L Feng, C Xu, N Xiang, Y Zhang, Y Shu, H Wang, H Luo, H Yu, X Liang, et al. Response to the first wave of pandemic (h1n1) 2009: experiences and lessons learnt from china. *Public health*, 126(5):427–436, 2012.

[7] Tarun S Weeramanthri, Andrew G Robertson, Gary K Dowse, Paul V Effler, Muriel G Leclercq, Jeremy D Burtenshaw, Susan J Oldham, David W Smith, Kathryn J Gatti, and Helen M Gladstones. Response to pandemic (h1n1) 2009 influenza in australia–lessons from a state health department perspective. *Australian Health Review*, 34(4):477–486, 2010.

[8] RG Pebody, E McLean, Hongxin Zhao, Paul Cleary, S Bracebridge, Kirsty Foster, Andre Charlett, P Hardelid, P Waight, Joanna Ellis, et al. Pandemic influenza a (h1n1) 2009 and mortality in the united kingdom: risk factors for death, april 2009 to march 2010. *Eurosurveillance*, 15(20):19571, 2010.

[9] Daihai He, Roger Lui, Lin Wang, Chi Kong Tse, Lin Yang, and Lewi Stone. Global spatio-temporal patterns of influenza in the post-pandemic era. *Scientific reports*, 5(1):1–11, 2015.

[10] Lewi Stone, Ronen Olinky, and Amit Huppert. Seasonal dynamics of recurrent epidemics. *Nature*, 446(7135):533–536, 2007.

[11] Saranya Sridhar, Shaima Begom, Alison Bermingham, Katja Hoschler, Walt Adamson, William Carman, Thomas Bean, Wendy Barclay, Jonathan J Deeks, and Ajit Lalvani. Cellular immune correlates of protection against symptomatic pandemic influenza. *Nature medicine*, 19(10):1305–1312, 2013.

[12] Benjamin D Dalziel, Stephen Kissler, Julia R Gog, Cecile Viboud, Ottar N Bjørnstad, C Jessica E Metcalf, and Bryan T Grenfell. Urbanization and humidity shape the intensity of influenza epidemics in us cities. *Science*, 362(6410):75–79, 2018.

[13] Jonathan Dushoff, Joshua B Plotkin, Simon A Levin, and David JD Earn. Dynamical resonance can account for seasonality of influenza epidemics. *Proceedings of the National Academy of Sciences*, 101(48):16915–16916, 2004.

[14] James Tamerius, Martha I Nelson, Steven Z Zhou, Cécile Viboud, Mark A Miller, and Wladimir J Alonso. Global influenza seasonality: reconciling patterns across temperate and tropical regions. *Environmental health perspectives*, 119(4):439–445, 2011.

[15] Ana Carolina Carioca da Costa, Cláudia Torres Codeço, Elias Teixeira Krainski, Marcelo Ferreira da Costa Gomes, and Aline Araújo Nobre. Spatiotemporal diffusion of influenza a (h1n1): starting point and risk factors. *PLOS ONE*, 13(9):1–20, 2018.

[16] Arnold S Monto. Seasonal influenza and vaccination coverage. *Vaccine*, 28:D33–D44, 2010.

[17] Danuta M Skowronski, Catharine Chambers, Suzana Sabaiduc, Gaston De Serres, Anne-Luise Winter, James A Dickinson, Mel Krajden, Jonathan B Gubbay, Steven J Drews, Christine Martineau, et al. A perfect storm: impact of genomic variation and serial vaccination on low influenza vaccine effectiveness during the 2014–2015 season. *Clinical Infectious Diseases*, 63(1):21–32, 2016.

[18] Shweta Bansal, Babak Pourbohloul, Nathaniel Hupert, Bryan Grenfell, and Lauren Ancel Meyers. The shifting demographic landscape of pandemic influenza. *PloS one*, 5(2):e9360, 2010.

[19] Jacob Bock Axelsen, Rami Yaari, Bryan T Grenfell, and Lewi Stone. Multiannual forecasting of seasonal influenza dynamics reveals climatic and evolutionary drivers. *Proceedings of the National Academy of Sciences*, 111(26):9538–9542, 2014.

[20] David JD Earn, Jonathan Dushoff, and Simon A Levin. Ecology and evolution of the flu. *Trends in ecology & evolution*, 17(7):334–340, 2002.

[21] Xiangjun Du, Aaron A King, Robert J Woods, and Mercedes Pascual. Evolution-informed forecasting of seasonal influenza a (h3n2). *Science translational medicine*, 9(413):eaan5325, 2017.

[22] Centers for Disease Control and Prevention. 2013-2014 influenza season summary, 2014. Available at https://archive.cdc.gov/www_cdc_gov/flu/pastseasons/1314season.htm. Accessed: 2025-02-18.

[23] Centers for Disease Control and Prevention. 2013-2014 estimated influenza illnesses, medical visits, hospitalizations, and deaths averted by vaccination in the united states. https://archive.cdc.gov/#/details?url=https://www.cdc.gov/flu/about/burden/2013-2014.html, 2014. Accessed: 2025-05-20.

[24] S. L. Linderman, B. S. Chambers, S. J. Zost, K. Parkhouse, Y. Li, C. Herrmann, A. H. Ellebedy, D. M. Carter, S. F. Andrews, N. Y. Zheng, and M. Huang. Potential antigenic explanation for atypical h1n1 infections among middle-aged adults during the 2013–2014 influenza season. *Proceedings of the National Academy of Sciences*, 111(44):15798–15803, 2014.

[25] The Disease Daily. Over 1,000 deaths in h1n1 outbreak in brazil. https://www.diseasedaily.org/2016/07/12/over-1000-deaths-h1n1-outbreak-brazil-71216/, 2016. Accessed: 2025-04-29.

[26] Farhang Babamahmoodi, Mohammadreza Haghshenas, Fatemeh Ahangarkani, Alireza Davoudi, Elham A Ashkezari, and Lotfollah Davoodi. Survey of mortality due to influenza a in north of iran, 2015-2016. *Current Respiratory Medicine Reviews*, 13(2):110–114, 2017.

[27] Atieh Sedghian, Shahab MohammadEbrahimi, Benn Sartorius, and Behzad Kiani. National spatiotemporal patterns of influenza-like illness in iran: A pre-covid-19 study (2015–2019). *PloS one*, 20(4):e0320990, 2025.

[28] Dilek Guldemir, Fatma Filiz Coskun-Ari, Ayse Basak Altas, Zekiye Bakkaloglu, Ozlem Unaldi, Fatma Bayraktar, Gulay Korukluoglu, Ali Riza Aktas, and Riza Durmaz. Molecular characterization of the influenza a (h1n1) pdm09 isolates collected in the 2015-2016 season and comparison of ha mutations detected in turkey since 2009. *Journal of medical virology*, 91(12):2074–2082, 2019.

[29] European Centre for Disease Prevention and Control. Start of 2011 influenza season in the southern hemisphere – south africa. https://www.ecdc.europa.eu/en/news-events/start-2011-influenza-season-southern-hemisphere-south-africa, 2011. Accessed: 2025-04-29.

[30] Balkan Insight. Croatia: Swine flu cases rise to 150. https://balkaninsight.com/2011/01/05/croatia-swine-flu-cases-rise-to-150/, 2011. Accessed: 2025-04-29.

[31] Centers for Disease Control and Prevention. National, regional, and state level outpatient illness and viral surveillance, 2023. Available at https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html. Accessed: 2025-05-06.

[32] A. Flahault and et al. Flunet as a tool for global monitoring of influenza on the web. *JAMA*, 280:1330–1332, 1998.

[33] World bank. Available at https://www.worldbank.org/ext/en/home.

[34] Peter Bogner, Ilaria Capua, David J Lipman, and Nancy J Cox. A global initiative on sharing avian flu data. *Nature*, 442(7106):981–981, 2006.

[35] Belongia E Osterholm MT, Sommer NS. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *The Lancet Infectious Diseases*, 12(1):36–44, 2012.

[36] Blanca García-Barreno, Teresa Delgado, Sonia Benito, Inmaculada Casas, Francisco Pozo, María Teresa Cuevas, Vicente Mas, Alfonsina Trento, Ariel Rodriguez-Frandsen, Ana Falcon, et al. Characterization of an enhanced antigenic change in the pandemic 2009 h1n1 influenza virus haemagglutinin. *Journal of General Virology*, 95(5):1033–1042, 2014.

[37] Wan Yang, Marc Lipsitch, and Jeffrey Shaman. Inference of seasonal and pandemic influenza transmission dynamics. *Proceedings of the National Academy of Sciences*, 112(9):2723–2728, 2015.

[38] Jeffrey Shaman and Melvin Kohn. Absolute humidity modulates influenza survival, transmission, and seasonality. *Proceedings of the National Academy of Sciences*, 106(9):3243–3248, 2009.

[39] Daihai He, Jonathan Dushoff, Raluca Eftimie, and David JD Earn. Patterns of spread of influenza a in canada. *Proceedings of the Royal Society B: Biological Sciences*, 280(1770):20131174, 2013.

[40] Vivek Charu, Scott Zeger, Julia Gog, Ottar N Bjørnstad, Stephen Kissler, Lone Simonsen, Bryan T Grenfell, and Cécile Viboud. Human mobility and the spatial transmission of influenza in the united states. *PLoS computational biology*, 13(2):e1005382, 2017.

[41] Stefano Merler and Marco Ajelli. The role of population heterogeneity and human mobility in the spread of pandemic influenza. *Proceedings of the Royal Society B: Biological Sciences*, 277(1681):557–565, 2010.

[42] Eduardo Massad, Marcelo Nascimento Burattini, Francisco Antonio Coutinho, and Cláudio José Struchiner. The risk of acquiring the new influenza a (h1n1) for brazilian travelers to chile, argentina and the usa. *Memórias do Instituto Oswaldo Cruz*, 105:179–183, 2010.

[43] David JD Earn, Daihai He, Mark B Loeb, Kevin Fonseca, Bonita E Lee, and Jonathan Dushoff. Effects of school closure on incidence of pandemic influenza in alberta, canada. *Annals of internal medicine*, 156(3):173–181, 2012.

[44] Simon Cauchemez, Neil M Ferguson, Claude Wachtel, Anders Tegnell, Guillaume Saour, Ben Duncan, and Angus Nicoll. Closure of schools during an influenza pandemic. *The Lancet infectious diseases*, 9(8):473–481, 2009.

[45] Hyunsuh Kim, Robert G Webster, and Richard J Webby. Influenza virus: dealing with a drifting and shifting pathogen. *Viral immunology*, 31(2):174–183, 2018.

[46] Meghan O Altman, Davide Angeletti, and Jonathan W Yewdell. Antibody immunodominance: the key to understanding influenza virus antigenic drift. *Viral immunology*, 31(2):142–149, 2018.

[47] Carolien E Van de Sandt, Joost HCM Kreijtz, and Guus F Rimmelzwaan. Evasion of influenza a viruses from innate and adaptive immune responses. *Viruses*, 4(9):1438–1476, 2012.

[48] Joel O Wertheim. The re-emergence of h1n1 influenza virus in 1977: a cautionary tale for estimating divergence times using biologically unrealistic sampling dates. *PloS one*, 5(6):e11184, 2010.

[49] Centers for Disease Control and Prevention. How the flu virus can change: "drift" and "shift". https://www.cdc.gov/flu/php/viruses/change.html, 2022. Accessed: 2025-05-20.

[50] Michael T Osterholm, Nicholas S Kelley, Alfred Sommer, and Edward A Belongia. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *The Lancet infectious diseases*, 12(1):36–44, 2012.

[51] Yizhuo Sun, Chao Bian, Ke Xu, Weibin Hu, Tongyan Wang, Jun Cui, Hongqiang Wu, Zhiyang Ling, Yongyong Ji, Guomei Lin, et al. Immune protection induced on day 10 following administration of the 2009 a/h1n1 pandemic influenza vaccine. *PLoS One*, 5(12):e14270, 2010.

[52] Gerardo Chowell. Fitting dynamic models to epidemic outbreaks with quantified uncertainty: A primer for parameter uncertainty, identifiability, and forecasts. *Infectious Disease Modelling 2*, no. 3:379–398, 2017.

[53] Rami Yaari, Guy Katriel, Amit Huppert, JB Axelsen, and Lewi Stone. Modelling seasonal influenza: the role of weather and punctuated antigenic drift. *Journal of The Royal Society Interface*, 10(84):20130298, 2013.

[54] Teagan Guarnaccia, Louise A Carolan, Sebastian Maurer-Stroh, Raphael TC Lee, Emma Job, Patrick C Reading, Stephen Petrie, James M McCaw, Jodie McVernon, Aeron C Hurt, et al. Antigenic drift of the pandemic 2009 a (h1n1) influenza virus in a ferret model. *PLoS pathogens*, 9(5):e1003354, 2013.

[55] Julia R. Gog and Bryan T. Grenfell. Dynamics and selection of many-strain pathogens. *Proceedings of the National Academy of Sciences*, 99(26):17209–17214, 2002.

[56] Pavel Minayev and Neil M Ferguson. Improving the realism of deterministic multi-strain models: implications for modelling influenza a. *Journal of the Royal Society Interface*, 6(35):509–518, 2009.

[57] Marco Ajelli, Piero Poletti, Alessia Melegaro, and Stefano Merler. The role of different social contexts in shaping influenza transmission during the 2009 pandemic. *Scientific Reports*, 4:7218, 2014.

[58] Duygu Balcan, Vittoria Colizza, Bruno Gonçalves, Hao Hu, José J Ramasco, and Alessandro Vespignani. Multiscale mobility networks and the spatial spreading of infectious diseases. *Proceedings of the National Academy of Sciences*, 106(51):21484–21489, 2009.

[59] Shweta Bansal, Bryan T Grenfell, and Lauren Ancel Meyers. When individual behaviour matters: homogeneous and network models in epidemiology. *Journal of the Royal Society Interface*, 4(16):879–891, 2007.

[60] Jie Zhou, Thomas P. Peacock, Judith C. Brown, Daniel H. Goldhill, Rebecca Frise, Laura Baillon, Benjamin J. Micklem, Timothy J. Green, Tetyana I. Vasylyeva, Emma L. Wise, Ajit Lalvani, John Sillitoe, and Wendy S. Barclay. Sars-cov-2 evolution and immune escape in immunocompromised patients. *Nature Microbiology*, 8:764–777, 2023.

Supplementary Material

The supplementary material details the estimation of fixed (time-independent) model parameters, the parameters obtained from least-squares optimization, and a link to the data and code used in this study.

Time-independent model parameters

Population data

To estimate the recruitment rate Λ into the susceptible class, we fitted a linear model to population data from 2009 onward. For each location, the model was initialized with the recorded 2009 population (N_{2009}), and the slope was estimated using the method of least squares, minimizing the sum of squared errors between the model and the observed data up to the year of the large resurgence (see Fig. S1). Overall, population changes were gradual and well approximated by a linear trend across all locations. The estimated growth rates, expressed in week^{-1}, are summarized in Table S1.

Figure S1: Population data for each location with best fit linear curve

Location	Population in 2009	Growth rate Λ (week^{-1})
São Paulo, Brazil	42, 075, 716	137.9351
Colombia	44, 313, 917	186.0822
Croatia	4, 382, 130	-5.4751
Iran	74, 322, 685	439.9057
New Zealand	4, 302, 600	14.4896
HHS Region 2	28, 0626, 68	38.4428
HHS Region 6	37, 944, 626	194.0172
South Africa	51, 170, 779	356.6660
Turkey	72, 561, 312	343.8197

Table S1: **Population in 2009 and estimated growth rates.**

Baseline loss-of-immunity rate (γ_0)

The constant parameter γ_0 represents the baseline rate at which individuals in the recovered class lose immunity. Its reciprocal, $\dfrac{1}{\gamma_0}$, corresponds to the baseline immunity duration, which we use to estimate the loss-of-immunity rate in 2009. Based on influenza studies reporting immunity durations ranging from 4 to 8 years [13], we assumed an average duration of 6 years. Converting to a weekly scale yields

$$\gamma_0 = \frac{1}{6 \times 52} = 3.2 \times 10^{-3} \text{ week}^{-1}.$$

Waning rate of vaccine-induced immunity (ω) and vaccination periods

The reciprocal $\dfrac{1}{\omega}$ represents the average duration of vaccine-induced immunity for an individual. In this study, we assumed 52 weeks based on [36], leading to

$$\omega = \frac{1}{52} = 0.019 \text{ week}^{-1}.$$

Vaccination periods for each location were estimated based on the months preceding winter and are summarized in Table S2.

Location	Winter period	Estimated vaccination period
Brazil	June - August	April - May
Colombia	December - February*	October - November
Croatia	December - February	October - November
New Zealand	June - August	April - May
Iran	December - February	October - November
Turkey	December - February	October - November
South Africa	June - August	April - May
HHS Region 2	December - February	October - November
HHS Region 6	December - February	October - November

Table S2: **Estimated vaccination periods.** Winter periods were defined according to the typical winter months in each region, accounting for their respective hemispheres and climate zones. In Colombia, where a true winter season does not occur due to its tropical climate, the rainy season was considered the equivalent of winter.

Recovery rate (r)

The reciprocal $\dfrac{1}{r}$ represents the average infectious period, typically ranging from 5 to 7 days [34].

We assumed an average duration of 6 days, which corresponds to

$$r = \frac{1}{0.85} = 1.17 \text{ week}^{-1}.$$

Estimated parameters from least-squares optimization

Location	$\beta_{\max}$	B	C	$\gamma_{\max}$	$\upsilon_{\max}$	ξ	ρ
Brazil	2.0599	0.0198	6.2823	0.0032	0.005	0.5	2.5466×10^{-4}
Colombia	10	0	4.2556	0.02	0.0105	0	1.5227×10^{-4}
Croatia	7.0501	0.2398	5.1492	0.0257	0.4996	0.08	8.0991×10^{-4}
Iran	7.9904	0.0062	4.0604	0.0032	0.0783	0	1.9073×10^{-4}
NZ	3.2591	0.0459	0.4632	0.0587	0.005	0.5	2.9337×10^{-4}
HHS Region 2	9.9906	0.0196	1.8298	0.004	0.0541	0.5	1.8985×10^{-4}
HHS Region 6	6.9894	0.0059	4.1677	0.0480	0.0562	0.3330	1×10^{-4}
SA	10	0.0203	3.8920	0.0041	0.0562	0.0002	1.0240×10^{-4}
Turkey	3.8400	0.0180	6.2828	0.0129	0.005	0.5	1.0556×10^{-4}

Table S3: **Estimated parameters from best-fit curves (linear γ model)**

Location	$\beta_{\max}$	B	C	$\gamma_{\max}$	τ	$\upsilon_{\max}$	ξ	ρ
Brazil	2.5160	0.0295	2.5244	0.0086	48.8403	0.005	0	1.3675×10^{-4}
Colombia	1.9094	6.2768	0	0.0032	60.3514	0.01	0.5	5.5207×10^{-4}
Croatia	4.9944	0.1290	5.8164	0.01	1	0.01343	0.5	9.3802×10^{-4}
Iran	1.8452	0.0287	0.7030	0.01	176	0.08	0.0611	1.0876×10^{-4}
NZ	3.9148	0.0233	0.7030	0.01	1	0.005	0.5	3.3975×10^{-4}
HHS Region 2	12.1608	0.0222	1.2788	0.0032	1	0.08	0.4997	1.7207×10^{-4}
HHS Region 6	5.9994	0.0294	0	0.0032	239.0694	0.0163	0.0001	1×10^{-4}
SA	2	0.1280	2.0913	0.027	59	0.0122	0.4995	1×10^{-4}
Turkey	2.8783	0	0.0534	0.01	221.9301	0.08	0.5	1.5917×10^{-4}

Table S4: **Estimated parameters from best-fit curves (jump γ model)**

Data and Codes

All of the data and code are available at the following link.