Undetected SARS-CoV-2 transmission in a Malawian pregnancy cohort: A longitudinal

serological surveillance study

Louise M. Randall, PhD[1*], Nicholas Kiernan-Walker[1*], Ernest Moya, PhD[2], Glory Mzembe, MD[2],

Alistair R. D. McLean, PhD[1,3], Rebecca Harding, PhD[1], Ramin Mazhari, PhD[1], Gomezgani

Mhango, MSc[2], Katherine L. Fielding, MBBS[1], Ivo Mueller, PhD[1,4], Martin N. Mwangi, PhD[2,5],

Sabine Braat, MSc[1,3], Kamija Phiri, PhD[2], Sant-Rayn Pasricha, PhD[1,7,8], Emily M. Eriksson,

PhD[1,4*+], Ricardo Ataíde, PhD[1*+]

[1] Infection and Global Health, The Walter and Eliza Hall Institute, VIC, Australia

[2] Training and Research Unit of Excellence (TRUE), Chichiri, Blantyre, Malawi

[3] Methods and Implementation Support for Clinical and Health Sciences Research Hub, University

of Melbourne, Melbourne, Victoria, Australia

[4] Department of Medical Biology, University of Melbourne, Melbourne, VIC, Australia

[5] The Micronutrient Forum, Healthy Mothers Healthy Babies Consortium, Washington, DC, USA.

[6] Clinical Hematology at the Royal Melbourne Hospital and Peter MacCallum Cancer Centre,

Parkville, Australia

[7] Diagnostic Hematology, The Royal Melbourne Hospital, Parkville, Australia

[8] School of Population and Global Health, University of Melbourne, Melbourne, Australia

* These authors contributed equally to the work.

+ Corresponding authors: Ricardo Ataíde, ataide.a@wehi.edu.au, tel: +61 (03) 9345 2555;

Address: The Walter and Eliza Institute for Medical Research, 1G Royal Parade, Parkville, 3052

VIC, Australia, Emily M. Eriksson, eriksson@wehi.edu.au, tel: +61 (03) 9345 2555; Address: The

Walter and Eliza Institute for Medical Research, 1G Royal Parade, Parkville, 3052 VIC, Australia,

Abstract

Background: Despite COVID-19's global impact, surveillance remains challenging in resource-

limited settings. This study investigated IgG to SARS-CoV-2 antigens in Malawian pregnant

women participating in the REVAMP trial, where no COVID-19 clinical cases or positive SARS-

CoV-2 tests were reported during the pandemic (April 2020 to September 2021).

Methods: We analysed serum samples collected from November 2018 to September 2021 from

852 pregnant women at enrolment (second trimester) and delivery, measuring IgG against five

SARS-CoV-2 antigens using a multiplex bead assay. IgG to Tetanus toxoid served as a positive

control, and IgG to four seasonal coronavirus antigens and Influenza A were used for context.

Findings: Women recruited after the start of the pandemic were younger than those recruited before

the emergence of COVID-19 (median age 19 vs 21 years) and more likely primigravid (60.1% vs

51.1%). IgG levels to SARS-CoV-2 antigens showed sharp increases of 18.5% - 29.7% every 30

days during COVID waves 2 and 3. Overall seropositivity to SARS-CoV-2 antigens reached

39.3% during the pandemic; however, the 14.7% seropositivity detected pre-pandemic

demonstrates the presence of cross-reactive antibody responses against other antigens in Malawi.

Surprisingly, pregnancies within the pandemic showed improved outcomes, with longer gestations

(mean difference: 0.6 weeks [95% CI: 0.2 - 0.9]) and higher birth weights (mean difference:

169.3g, [65.9 - 272.6]) when compared to pregnancies occurring pre-pandemic. We found no

evidence that SARS-CoV-2 IgG levels were associated with pregnancy outcomes.

Interpretations: This study demonstrates that serological surveillance reveals undetected SARS-

CoV-2 exposure where traditional testing was limited. The improvement in pregnancy outcomes

during the pandemic suggests community-level exposure did not offset the social impacts of the

pandemic in this population. These findings highlight pregnancy cohorts as valuable sentinel

populations for infectious disease surveillance, emphasising the importance of serosurveillance in

resource-limited settings.

Introduction:

The COVID-19 pandemic presented unique surveillance challenges in resource-limited settings, where constrained testing capacity and limited healthcare access complicated efforts to track SARS-CoV-2 transmission.[1] In Malawi, despite early public health measures, significant challenges in disease surveillance and response coordination left the true extent of viral spread uncertain. While official statistics suggested low case numbers, compared to other global regions, limited molecular testing capacity and accessibility, particularly in rural areas, meant that actual infection rates remained difficult to ascertain. [2,3]

This surveillance challenge was strikingly illustrated in the REVAMP trial—a randomized controlled trial of iron supplementation in pregnancy conducted in southern Malawi.[4] Running from November 2018 through September 2021, the trial spanned pre-pandemic and pandemic periods, including three distinct COVID-19 waves that occurred in Malawi.[5] Despite regular participant monitoring through at least four study visits between second-trimester enrolment and delivery, no trial participant reported receiving a positive SARS-CoV-2 test or presented with respiratory symptoms indicative of COVID-19. A seroprevalence survey of blood donations at the Malawi Blood Transfusion Service, suggested that the seropositivity in Malawi could reach 65%.[6] As such, the absence of reported cases within REVAMP, a closely monitored cohort going through a pandemic, suggested that infections were going undetected.

Serological surveillance offers a powerful tool for addressing such detection challenges, particularly when supported by appropriate validation measures. By determining antibody responses rather than active infections, serosurveillance can reveal historical exposure patterns and

provide insights into population-level transmission dynamics that traditional surveillance methods

might miss.[7] The value of this approach has been well-demonstrated in various infectious disease

contexts.[8–11]

Pregnancy cohorts are particularly valuable for serosurveillance, as they provide year-round

contact points for sample collection and monitoring.[12,13] For example, in malaria-endemic areas of

Africa, serosurveillance of pregnant women is increasingly seen as a reliable and easy way to

determine trends in malaria transmission across the population.[10,12,13] Additionally, pregnancy-

specific interventions, such as vaccinations during governmental health programs, create built-in

immunological controls that can validate changes seen in incidental serological findings— e.g.

tetanus vaccination during pregnancy provides a reliable immunological stimulus that can verify

both healthcare delivery and antibody detection methods.[14]

The REVAMP trial presented an ideal platform for investigating potential undetected transmission

of SARS-CoV-2 through serosurveillance during pregnancy. Its timeline included reliable pre- and

post-pandemic samples from the same population—rare in COVID-19 serological studies—while

its regular monitoring schedule ensured consistent sample collection throughout the pandemic

period. The trial's comprehensive data collection on several demographics, clinical outcomes, and

healthcare delivery patterns offered multiple avenues for validating the serological findings. We

aimed to leverage this unique opportunity to assess evidence of undetected SARS-CoV-2 exposure

through validated serological approaches. In addition, we evaluated the impact of changes to

SARS-CoV-2 serology on pregnancy outcomes.

Methods

Study design

All samples are from the REVAMP trial, which took place in southern Malawi, in the Zomba and

Blantyre districts. The trial protocol, statistical analysis plan and primary results have been

published.[4,15]   Briefly, pregnant women with ultrasound-confirmed singleton second-trimester

pregnancy were individually randomized to receive either an intravenous iron infusion of Ferric

Carboxymaltose [Vifor Pharma] or standard of care oral iron if they met the eligibility criteria.

Eligibility criteria included haemoglobin of <10.0g/dL (by HemoCue 301+ [Angelholm,

Sweden]), and a negative malaria rapid diagnostic test, among others. The primary endpoint of the

trial was anaemia prevalence at 36 weeks' gestation. Venous blood samples were taken throughout

the pregnancy, including at enrolment (13-26 weeks' gestation), and delivery. Blood was

processed within two hours of collection and plasma was aliquoted and stored locally at -80 ºC

until being shipped to Melbourne, Australia. All women were eligible to receive pregnancy

immunisations from the Malawi Government, following the national Expanded Programme of

Immunisation strategy. These include a dose of Tetanus toxoid (Tt) at the first antenatal visit (often

the time of first contact with the REVAMP trial team), a second dose 4 weeks after the first dose

and a third dose 6 months after the second dose.[16]

The trial received ethics approvals from the College of Medicine, University of Malawi, Malawi

(P.02/18/2357), and The Walter and Eliza Hall Institute (WEHI), Australia (18/02) and was

prospectively registered (ACTRN12618001268235). An independent data and safety monitoring

board oversaw the trial. Secondary analyses of the trial samples were included in the study

protocol.

COVID-19 Setting

All dates related to the COVID-19 pandemic were obtained from publicly available Malawian Government sources.[5] The first official case of COVID-19 in Malawi was reported on April 2nd 2020, and marked the beginning of the first wave of infections (Figure 1).[5] Following the establishment of containment policies, and at the request of the Malawi Ministry of Health, REVAMP amended its protocol and adopted COVID measures in May of 2020 (i.e. limiting the number of contact visits between the team and the participants, as well as limiting the duration of those visits).[15] In January 2021, the Malawi government implemented a national lockdown. Vaccinations were introduced in Malawi on March 11th 2021, with the ChAdOx1-S COVID-19 vaccine being delivered.[5] Study participants were deliberately not asked whether they had received COVID-19 vaccines as there was a lot of misconception, and it was thought that this question could affect retention in subsequent visits. However, at each visit, participants were asked to report any concomitant medication they were on or any prior visits to a health centre, and we had no indication that any of the study participants received the vaccination. Malawi experienced three waves of COVID-19 during the lifespan of REVAMP.[5] During the entire trial, there were no participant self-reported nor trial team-confirmed COVID-19 cases within the cohort, and there were no observed or reported positive SARS-CoV-2 tests.

Serological multiplex bead-array

A multiplex serological assay was used to detect antigen-specific IgG antibodies targeting the SARS-CoV-2 Spike protein, Spike-derived antigens (S1, S2 and receptor binding domain (RBD)) and nucleoprotein (NP), as previously described.[17] Additionally, we assessed antibodies to

seasonal coronaviruses antigens (NL63, OC43, HKU1 and 229E) and Influenza virus A (H1N1).

Antibodies to Tt were also measured as an immunological positive control.[16] Briefly, individual

plasma samples were incubated with fluorescent magnetic beads coated with the relevant antigens.

Antigen-specific antibodies were then identified using a secondary anti-human IgG Fc antibody

conjugated to phycoerythrin (PE) and analyzed with the MAGPIX® system. All samples were run

with the analysts blinded to the participants' characteristics.

Statistical analysis

The sample size was determined by the available data of all randomised women with a plasma

sample at study enrolment or delivery. Baseline characteristics were summarised and compared

before and after COVID-19 emergence in Malawi using either two-sample t or Wilcoxon rank-

sum test for continuous data or chi-square or Fisher's exact test for categorical data. Antibody

levels were quantified as adjusted mean fluorescent intensity (aMFI) derived from standard curves.

Seropositivity for SARS-CoV-2 antibodies was determined as IgG levels exceeding the mean plus

two standard deviations from pre-pandemic pregnancies. Longitudinal $\log_e$-transformed IgG levels

were visualised using locally weighted (LOWESS) regression with restricted cubic spline curve,

and 10 smoothing points.[18] Linear regression models with two inflexion points (start of the

COVID-19 pandemic in Malawi - April 2nd 2020; beginning of the second COVID-19 wave -

January 1st 2021), resulted in three time segments. Models accounted for variables assumed to be

prognostic or predictive of the outcomes: maternal age, body mass index, primiparity, sex of the

newborn, HIV status, income source and treatment group. Estimates and confidence intervals (CI)

were back-transformed to reflect a per cent of change over 30 days.

Linear regression models were used to quantify associations between pregnancy outcomes of

interest (gestational age, birth weight and low birth weight) and serological data at delivery as

exposures with adjustment for confounding and prognostic variables - maternal age,

primigravidity, body mass index, HIV status, and treatment group. Spearman correlations assessed

the IgG relationships during the Pre-COVID or COVID periods. No adjustment for multiple testing

was done.

Small for gestational age was defined as a birthweight below the 10th percentile for gestational age

according to INTERGROWTH-21 standards;[19] Low birth weight is defined as a birth weight

<2500 g; Premature birth is a birth occurring at <37 weeks' gestation; Foetal loss represents a

composite of pregnancy loss and stillbirth. All statistical analyses were conducted in Stata v18.0

(StataCorp, College Station, TX) and GraphPad Prism v10 (GraphPad Software, San Diego, CA,

USA).

Results

Trial Population reflected COVID-19 impacts

The REVAMP trial recruited 862 participants between November 2018 and March 2021, with the

last delivery in the trial occurring in September 2021, thus pregnancies in REVAMP spanned pre-

pandemic and pandemic periods (Figure 1a). A total of 852/862 (98.8%) women had available

serum samples and were included in this nested study (Table 1), including 147 samples from

enrolment only, 30 samples from delivery only, and 675 samples for both enrolment and delivery.

The baseline characteristics of the women with a sample analysed at only one time point were

similar to women who had samples analysed at both time points (Table S1).

A total of 569 (66.8%) women were enrolled before and 283 (33.2%) women were enrolled after

April 2nd 2020, the date of the first case of COVID-19 registered in Malawi (Table 1) while 358

(42.0%) pregnancies were spent solely under the Pre-COVID period and 135 (15.8%) solely under

the COVID periods (Figure 1a).

We observed differences in demographic characteristics of the trial participants before and after

the emergence of COVID-19 in Malawi (Table 1). The impact of time of recruitment on

demographics revealed maternal age decreased across the period of the trial, from a median of 21

years (IQR [19-27]) in the pre-COVID period to 19 years (IQR [17-23]) ($P=0.0010$) during the

COVID period. Consistent with this, we observed an increase in the percentage of women in their

first pregnancy towards the end of the trial (51.1% (291/569) pre-COVID vs 60.1% (170/283)

COVID; $P=0.014$). Income source also changed significantly from the pre-COVID to the COVID

periods ($P<0.001$), with more women reporting practicing Subsistence farming (13.1% (74/567)

pre-COVID vs 25.9% (73/282) COVID) and less women reporting being Employed (21.2% pre-

COVID vs 11.4% COVID).

Serological outcomes validation – Tetanus toxoid

We measured antibodies to Tetanus toxoid (Tt) as a validation tool for our serological assays. As

was expected from Malawi's vaccination policy,[16] IgG toward Tt were higher at delivery versus

enrolment (Figure 1b), and in women with two or more pregnancies compared to women in their

first pregnancy – both at enrolment and, though to a lesser degree, at delivery (Figure 1c-d). The

overall levels of anti-Tt antibodies measured at delivery were unaffected in those pregnancies

carried solely within the COVID period vs those solely in the pre-COVID period (Figure 1e).

Serological outcomes – SARS-CoV-2

Antibodies to all five of SARS-CoV-2 antigens increased from the beginning of 2021, coinciding

with the emergence of COVID waves 2 and 3 in Malawi (Figure 2a and Supplementary figure 1).

A wide range of IgG levels to SARS-CoV-2 antigens were present even before the emergence of

the COVID-19 pandemic, indicating the occurrence of cross-reactive immune responses in this

population. In general, there was a subtle positive trend in antibody levels over time, with values

remaining relatively stable through to the end of COVID wave 1, followed by an increase through

COVID waves 2 and 3. Linear modelling of this relationship revealed a significant increase in

antibodies to SARS-CoV-2 antigens during COVID waves 2 and 3 (Figure 2b, Supplementary

Table 2). This increase was also noticeable when comparing mean antibody levels in those

pregnancies carried solely within the COVID period versus those occurring in the pre-COVID

(Figure 2c). In addition, the correlations between IgG levels to all SARS-CoV-2 antigens also

significantly increased from those pregnancies entirely in the pre-COVID period to those entirely

in the COVID period (Figure 2e-d).

A total of 102 samples (17.3%) were seropositive for at least one of the SARS-CoV-2 antigens

(ranging from 4.3% for RBD to 9.1% against SPIKE). Half of the 102 seropositive samples (49%)

were seropositive for 1 of the antigens and only 5 samples (4.1%) were seropositive for all five

SARS-CoV-2 antigens. The overall seroprevalence calculated for those pregnancies entirely

outside of the COVID-19 period in Malawi was 14.7% (48/327), revealing a significant number

of cross-reactive responses to other antigens. In pregnancies occurring entirely within the COVID

period, the overall seroprevalence was 39.3% (48/122).

Serological outcomes – Seasonal coronaviruses and Influenza A

Antibodies to seasonal viruses presented a distinct pattern across the timeline of the trial (Figure

3a-b and Supplementary figure 1). LOWESS visualization showed little variation across the

entirety of the trial (Figure 3a). Linear modelling over the three-segments of the trial revealed a

general stability over the pre-COVID period— maximum variation was for IgG towards

Coronavirus HKU1 (mean % change over 30 days: -4.1% [-5.8, -2.3]) (Supplementary Table 2).

This was followed by a small increase in IgG levels during the second period, and an overall

decrease during the last period (Figure 3b, Supplementary Table 2). However, the levels of

antibodies to these common human coronaviruses, including types NL63, OC43, and HKU1, and

to Influenza A (H1N1) remained relatively unchanged when analysing those pregnancies occurring

outside or inside the COVID period, except for antibodies to Coronavirus 229E that declined

significantly at delivery between solely pre-COVID and solely COVID pregnancies (Figure 3c).

There was no significant change to the correlations between antigens to the common coronaviruses

and Influenza A between COVID periods (Figure 3c-d).

Pregnancy outcomes of the trial according to COVID period and association with SARS-CoV-2

antibodies

Throughout the trial, and after accounting for confounding variables, gestation duration increased

significantly between pregnancies that delivered during the COVID period compared to pre-

COVID (adjusted mean difference 0.5 weeks [0.2-0.8]) (Table 2). This difference was still

observed when looking at women with pregnancies entirely in the COVID period versus entirely

in the pre-COVID period (mean difference 0.6 weeks [0.2, 0.9]). Accordingly, babies had higher

birth weights when delivered during the COVID period compared with those delivered during the

pre-COVID period (mean difference 69.4 g [-2.4-141.1]), although this increased birthweight was

only significant when comparing those pregnancies occurring solely within the COVID period

versus occurring solely in pre-COVID (mean difference 169.3 g [65.9, 272.6]). Low birth weight

decreased during the COVID period (Odds Ratio (OR) COVID vs pre-COVID: 0.57 [0.37-0.89]),

and in those pregnancies solely occurring during COVID versus in pre-COVID (OR Solely

COVID vs pre-COVID: 0.40 (0.20-0.77)) (Table 2).

In women whose pregnancies occurred within the COVID period, we found no evidence of a

statistically significant association between seropositivity to SARS-CoV-2 antigens and pregnancy

outcomes. However, the estimates around the associations had very large 95% CI, which is

compatible with large effects in either direction (Supplementary Table 3).

Discussion

In this study we used samples from the REVAMP clinical trial to examine serological patterns and

pregnancy outcomes before and during the COVID-19 pandemic in a well-characterized cohort of

Malawian pregnant women. We found that, despite a lack of clinical symptoms and laboratory-

confirmed positive tests, levels of antibodies to key antigens of SARS-CoV-2 significantly

increased in the population, as did the overall seropositivity prevalence, from a background

estimate of 14.7% seropositivity before COVID to an overall seropositivity of 39.3% for

pregnancies within the COVID period. Over the full span of the trial, we observed an increase in

the mean gestation duration and birth weight of the babies, accompanied by a decrease in the

overall prevalence of low birth weight. However, we found no evidence that exposure to SARS-

CoV2—as measured by IgG to 5 different antigens — was associated with pregnancy and birth

outcomes.

The REVAMP trial commenced in November 2018 and had its last delivery in September 2021.

This resulted in continuous sampling throughout pregnancy and up to delivery for women before

the emergence of the COVID-19 pandemic and up to the height of the 2021 COVID-19 waves in

Malawi. We observed significant demographic shifts throughout the trial's duration. Women

enrolled during the pandemic were significantly younger than those enrolled in the pre-COVID

period. This change was accompanied by significant changes to the parity, education and income

source measures of the trial cohort—reflecting the younger population. This shift in population

age observed during our trial timeline may reflect the broader socio-economic changes that

occurred during the pandemic. Malawi reported a decrease in total health service attendance of up

to 10% from April 2020 to December 2021, with antenatal services specifically falling by 3%.[20] It

is reasonable to assume that, as lockdowns, school closures and other containment measures were

enforced by the Malawian government,[5] women with young children may have had responsibilities

that prevented them from either participating in the study or engage with the health services. These

findings also highlight the importance of understanding the characteristics of the underlying

population when the time periods of study are lengthy.

Our serological findings contribute to understanding immune responses to SARS-CoV-2 in

African pregnant women. A meta-analysis of the seroprevalence of anti-SARS-CoV-2 antibodies

across Africa (excluding Malawi) revealed seroprevalences ranging from 0.0-63.0%,[21]   while

studies in Malawi, estimated prevalences at between 10-85% depending on the sampling strategy

and the population.[6,22,23] In REVAMP, we measured a peak seroprevalence of 39.3%. Although

seropositivity to SARS-CoV-2 does not immediately follow from increases in seroprevalence to

single antigens, the increase of antibody levels to all five SARS-CoV-2 antigens during the

COVID-19 period, coupled with stronger inter-antigenic correlations, suggests true community

transmission occurred in the trial population during this period. This timing agrees with SARS-

CoV-2 seroprevalence data obtained from blood donations surveyed from 4 distinct areas of

Malawi[6] and from pregnant women at first antenatal clinic, in a country-wide study which started

just as REVAMP was recruiting its last participants.[23] The general rise in antibody levels to all the

antigens evaluated suggests a rise in serology due to community transmission of the virus, rather

than through active vaccination campaigns, which only started in March of 2021 and experienced

low uptake until the end of 2021.[5] Notably, the relatively high individual spread of IgG levels

observed against SARS-CoV-2 antigens in samples collected before the emergence of SARS-CoV-

2—a degree of background reactivity not observed in other cohorts—[17] and raises questions about

potential immunological cross-reactivity between SARS-CoV-2 antigens and other antigens in this

population. The clinical significance of this observation remains to be determined.

IgG levels measured against common human coronaviruses as well as Influenza A revealed distinct

patterns to those observed against all SARS-CoV-2 antigens, with little variation across the trial

period. The stability of Tt antibody levels across all periods served as an important internal control,

demonstrating the specificity of the observed changes in antibodies to SARS-CoV-2 and other

control viruses. Tt vaccinations were provided by the Government, not as part of the trial, thus

demonstrating that despite disruptions to the health services during COVID, [20] ANC attendance

and services was still relatively strong in this area of Malawi.

Despite the serological evidence of increasing SARS-CoV-2 exposure in the population, during

the trial period, pregnancies occurring during COVID-19 experienced longer gestational ages and

babies with increased birth weights. As is the case with many respiratory infections,[24] SARS-CoV-

2 infections, and COVID-19 in particular, negatively affect birth outcomes.[25] However,

observations of better outcomes for pregnancy during the pandemic have been made in other

cohorts as well.[26] These results seem to suggest that either 1) exposure levels in our population

may not have reached a significant threshold; or 2) the negative impacts of SARS-CoV-2 infection

were of less magnitude than the improvements due to reductions in other infections during

COVID; or 3) The negative impact of infection is of lesser magnitude than the improvement driven

by a shift in the characteristics of the women recruited to the trial— reduced mobility and limited

physical demands may have conserved maternal energy, and increased presence of partners at

home may have provided added support during pregnancy.

These findings should be interpreted in context. While our study benefits from being embedded

within a clinical trial with standardised follow-up and data collection, this also means that our

findings may not apply to the broader Malawian population. We did not collect specific

information regarding COVID-19 vaccination status of our participants. Our study's temporal

nature makes it challenging to disentangle the effects of SARS-CoV-2 infection from the effects

of pandemic-related social and healthcare changes. While our models accounted for demographics

such as age and primiparity that shifted during our study, these significant demographic shifts

could have introduced unmeasured confounding factors not accounted for in our statistical

adjustments. Additionally, it's important to note that our trial population, receiving regular

antenatal care and nutritional supplementation, may have been relatively protected from pandemic-

related impacts when compared to the general population[20].

In conclusion, we demonstrate that in a population of pregnant women in Malawi, where

symptoms of COVID-19 were absent and there was no recorded evidence of SARS-CoV-2

infection, population-level IgG to SARS-CoV-2 antigens rose in alignment with well-defined

COVID-19 periods. In addition, the increase in inter-antigenic correlation between the antibody

levels to SARS-CoV-2 antigens during the COVID-19 period also sustains the interpretation that

there was true population transmission during the COVID pandemic. This represents additional

evidence to the body of research that proposes pregnancy as a sensitive sentinel period for the

population surveillance of infectious diseases. [11,12,27,28]

References

1   Batista C, Hotez P, Amor Y Ben, *et al.* The silent and dangerous inequity around access to

COVID-19 testing: A call to action Comment. *EClinicalMedicine* 2022; **43**: 101230.

2   Ulaya G, Kabaghe A, Saussier C, *et al.* Monitoring COVID-19 Occurrence in a Resource-

limited Setting – COVID-19 Sentinel Surveillance in Malawi. 2024; published online Dec

26. DOI:10.1101/2024.12.23.24319583.

3   Taylor A, Liwewe T, Todd J, Kankhwali C, Mwale A, Kiwuwa-Muyingo S. Insights into

COVID-19 data collection and management in Malawi: exploring processes, perceptions,

and data discrepancies. *Wellcome Open Res* 2024; **9**.

DOI:10.12688/wellcomeopenres.21131.1.

4   Pasricha S-R, Mwangi MN, Moya E, *et al.* Ferric carboxymaltose versus standard-of-care

oral iron to treat second-trimester anaemia in Malawian pregnant women: a randomised

controlled trial. *Lancet* 2023; **401**: 1595–609.

5   Presidential Task Force on Coronavirus. THE CORONAVIRUS PANDEMIC IN

MALAWI: TRAILING THE WAVES. 2022.

6   Mandolo J, Msefula J, Henrion MYR, *et al.* SARS-CoV-2 exposure in Malawian blood

donors: an analysis of seroprevalence and variant dynamics between January 2020 and July

2021. *BMC Med* 2021; **19**. DOI:10.1186/s12916-021-02187-y.

7   Arnold BF, Scobie HM, Priest JW, Lammie PJ. Integrated serologic surveillance of

population immunity and disease transmission. *Emerg Infect Dis* 2018; **24**: 1188–94.

8   Tedijanto C, Solomon AW, Martin DL, *et al.* Monitoring transmission intensity of trachoma

with serology. *Nat Commun* 2023; **14**. DOI:10.1038/s41467-023-38940-5.

9   Dorigatti I, Lavezzo E, Manuto L, *et al.* SARS-CoV-2 antibody dynamics and transmission

from community-wide serological testing in the Italian municipality of Vo'. *Nat Commun*

2021; **12**. DOI:10.1038/s41467-021-24622-7.

10   van Eijk AM, Hill J, Noor AM, Snow RW, ter Kuile FO. Prevalence of malaria infection in

pregnant women compared with children for tracking malaria transmission in sub-Saharan

Africa: A systematic review and meta-analysis. *Lancet Glob Health* 2015; **3**: e617–28.

11   Pujol A, Brokhattingen N, Matambisso G, *et al.* Detecting temporal and spatial malaria

patterns from first antenatal care visits. *Nat Commun* 2023; **14**. DOI:10.1038/s41467-023-

39662-4.

12   Ataíde R, Mayor A, Rogerson SJ. Malaria, primigravidae, and antibodies: Knowledge

gained and future perspectives. Trends Parasitol. 2014; **30**: 85–94.

13   Mayor A, Menendez C, Walker PGT. Targeting pregnant women for malaria surveillance.

Trends Parasitol 2019; : 1–10.

14   Integrated Management of Pregnancy and Childbirth (IMPAC). Maternal Immunization

against Tetanus. Integrated Management of Pregnancy and Childbirth (IMPAC). .

15   Harding R, Ataide R, Mwangi MN, *et al.* A Randomized controlled trial of the Effect of

intraVenous iron on Anaemia in Malawian Pregnant women (REVAMP): Statistical

analysis plan. *Gates Open Res* 2021; **5**: 174.

16   The Government of Malawi. EPI Comprehensive Multi-Year Plan. 2016.

17   Mazhari R, Ruybal-Pesántez S, Angrisano F, *et al.* SARS-CoV-2 Multi-Antigen Serology

Assay. *Methods Protoc* 2021; **4**. DOI:10.3390/mps4040072.

18   Chambers JM, Cleveland WS, Kleiner B, Tukey PA. Graphical Methods for Data Analysis.

Chapman and Hall/CRC, 2018 DOI:10.1201/9781351072304.

19   Villar J, Ismail LC, Victora CG, *et al.* International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *The Lancet* 2014; **384**: 857–68.

20   She B, Mangal TD, Adjabeng AY, *et al.* The changes in health service utilisation in Malawi during the COVID-19 pandemic. *PLoS One* 2024; **19**. DOI:10.1371/journal.pone.0290823.

21   Chisale MRO, Ramazanu S, Mwale SE, *et al.* Seroprevalence of anti-SARS-CoV-2 antibodies in Africa: A systematic review and meta-analysis. Rev Med Virol. 2022; **32**. DOI:10.1002/rmv.2271.

22   Meinus C, Singer R, Nandi B, *et al.* SARS-CoV-2 prevalence and immunity: a hospital-based study from Malawi. *International Journal of Infectious Diseases* 2022; **116**: 157–65.

23   Tenthani L, Seffren V, Kabaghe AN, *et al.* SARS-CoV-2 Seroprevalence and Vaccine Uptake among Pregnant Women at First Antenatal Care Visits in Malawi. *American Journal of Tropical Medicine and Hygiene* 2024; **110**: 989–93.

24   Di Mascio D, Khalil A, Saccone G, *et al.* Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. Am J Obstet Gynecol MFM. 2020; **2**. DOI:10.1016/j.ajogmf.2020.100107.

25   Male V. SARS-CoV-2 infection and COVID-19 vaccination in pregnancy. *Nat Rev Immunol* 2022; **22**: 277–82.

26   de Knegt VE, Hedley PL, Hedermann G, *et al.* The impact of the COVID-19 lockdown on birthweight among singleton term birth in Denmark. *PLoS One* 2023; **18**. DOI:10.1371/journal.pone.0283909.

27   Gutman J. Pregnant women and children as sentinel populations. 2017.

28   Joint United Nations Programme on HIV/AIDS and World Health Organization. Guidelines

for Conducting HIV Sentinel Serosurveys among Pregnant Women and Other Groups.

2003; : 1–106.

Acknowledgements: The Bill & Melinda Gates Foundation funded the REVAMP trial (INV-010612). S-RP is supported by the Australian National Health and Medical Research Council (NHMRC) Fellowships (GNT1158696 and GNT2009047). We thank the local field workers, participants, and their families for their involvement in the study.

Author contributions: S-RP and KSP conceptualised and designed the trial and secured funding. EM, GMz, GMh and MNM led data collection in the field. RA and EME planned this study. LMR, NK-W, and KLF conducted experimental data collection. NK-W, RM, EME and RA had full access to and analysed the data. RH, ARDM, SB and RA conducted the statistical analysis. LMR, ARDM, EME and RA interpreted the data. RA prepared the initial draft of the manuscript. All authors critically revised the manuscript for intellectual content, edited and approved the final version, and agreed to be accountable for all aspects of the work.

Data sharing: The datasets generated during this study are available from the corresponding author on reasonable request, subject to ethical approval and data sharing agreements with the REVAMP trial investigators.

Competing interests: The authors declare no competing interests.

 Table 1- Baseline characteristics according to the timing of the emergence of COVID-19 in

 Malawi.

	Before emergence of COVID-19 in Malawi[a]	After emergence of COVID-19 in Malawi[b]	Total Cohort[§]	Test p-value[k]
N, %	569 (66.8%)	283 (33.2%)	852 (100.0%)	
Age (years)	21.0 [18.0, 26.0]	19.0 [17.0, 24.0]	20.0 [18.0 25.0]	0.0010
BMI[c] (kg/m^2)	22.8 [21.1, 24.7]	22.5 [21.0, 24.5]	22.6 [21.1, 24.6]	0.20
Primigravid				
No	278 (48.9%)	113 (39.9%)	391 (45.9%)	0.014
Yes	291 (51.1%)	170 (60.1%)	461 (54.1%)	
Religion				
None	0 (0.0%)	2 (0.7%)	2 (0.2%)	0.064
Christian	419 (73.9%)	192 (68.1%)	611 (72.0%)	
Muslim	144 (25.4%)	84 (29.8%)	228 (26.9%)	
Other	4 (0.7%)	4 (1.4%)	8 (0.9%)	
Education				
None	2 (0.4%)	0 (0.0%)	2 (0.2%)	0.0020
Lower Primary	98 (17.9%)	74 (27.0%)	172 (21.0%)	
Upper Primary	215 (39.4%)	122 (44.5%)	337 (41.1%)	
Lower Secondary	85 (15.6%)	28 (10.2%)	113 (13.8%)	
Upper Secondary	132 (24.2%)	45 (16.4%)	177 (21.6%)	
Tertiary	14 (2.6%)	5 (1.8%)	19 (2.3%)	
Marital status				
Single	85 (15.0%)	47 (16.7%)	132 (15.5%)	0.57
Married	474 (83.6%)	228 (80.9%)	702 (82.7%)	
Widowed	3 (0.5%)	1 (0.4%)	4 (0.5%)	
Divorced/Separated	4 (0.7%)	5 (1.8%)	9 (1.1%)	
Others	1 (0.2%)	1 (0.4%)	2 (0.2%)	
Income source				
None	38 (6.7%)	15 (5.3%)	53 (6.2%)	<0.001
Subsistence farming	74 (13.1%)	73 (25.9%)	147 (17.3%)	
Large scale farming	1 (0.2%)	1 (0.4%)	2 (0.2%)	
Employed	120 (21.2%)	32 (11.3%)	152 (17.9%)	
Casual work for wages	162 (28.6%)	99 (35.1%)	261 (30.7%)	
Business	163 (28.7%)	59 (20.9%)	222 (26.1%)	
Other	9 (1.6%)	3 (1.1%)	12 (1.4%)	
HIV positive				
No	462 (81.8%)	239 (85.4%)	701 (83.0%)	0.19
Yes	103 (18.2%)	41 (14.6%)	144 (17.0%)	
Haemoglobin[d] (g/dL)	9.1 [8.1, 9.8]	8.7 [7.9, 9.3]	8.9 [8.0, 9.6]]	<0.001
Anaemia[e]				
No	34 (6.0%)	7 (2.5%)	41 (4.8%)	0.023
Yes	531 (94.0%)	276 (97.5%)	807 (95.2%)	
Ferritin (µg/L)	25.7 [9.4, 64.2]	28.4 [11.6, 86.8]	26.8 [10.0, 72.3]	0.95
CRP (mg/L)[f]	5.3 [2.8, 11.1]	5.0 [2.8, 9.8]	5.2 [2.8, 10.6]	0.85
Inflammation[g]				
No	269 (47.4%)	137 (51.1%)	406 (48.6%)	0.31
Yes	299 (52.6%)	131 (48.9%)	430 (51.4%)	
Iron deficiency[h]				
No	313 (55.1%)	161 (60.1%)	474 (56.7%)	0.18
Yes	255 (44.9%)	107 (39.9%)	362 (43.3%)	

Data are count (%) or median [25% centile, 75% centile]. [a]Women enrolled in the trial before April

2nd 2020. [b]Women enrolled in the trial on or after April 2nd 2020. [c]Body Mass Index.

[d]Haemoglobin levels in venous blood, measured by Symex. [e]Anaemia indicates proportion of

women with haemoglobin <11 g/dL. [f]C-Reactive Protein. [g]Inflammation indicates a CRP >5 mg/L.

[h]Iron deficient indicates a serum ferritin<15 µg/L, or serum ferritin<30 µg/L if CRP >5 mg/L. [k]P-

values relate to testing for differences between before and after emergence of COVID in Malawi.

The analysis included all women with available serological samples at either enrolment or

delivery.

Table 2. Effects of COVID-19 period on Pregnancy and Birth Outcomes

Delivery inside vs outside COVID	Pre-Covid	Covid	OR [95% CI] or mean difference [95% CI]	P-value
Small for gestational age n/N (%)	146/477 (30.6%)	92/266 (34.6%)	1.10 [0.79-1.53]	0.57
Low birth weight (<2,500g) n/N (%)	89/484 (18.4%)	32/265 (12.1%)	0.57 [0.37-0.89]	0.013
Premature birth (<37 weeks) n/N (%)	43/489 (8.8%)	17/270 (6.3%)	0.70 [0.39-1.26]	0.23
Fetal loss n/N (%)	11/522 (2.1%)	5/275 (1.8%)	0.95 [0.32-2.82]	0.93
Gestation duration (weeks) Mean (SD)[#]	39.3 (2.1)	39.8 (1.9)	0.49 [0.20-0.78]	0.0010
Birth weight (grams) Mean (SD)[$]	2889.9 (501.4)	2930 (477.8)	69.36 [-2.39-141.12]	0.06
Entire gestation period inside vs outside COVID				
Small for gestational age n/N (%)	94/335 (28.1%)	40/130 (30.8 %)	0.97 [0.62-1.53]	0.90
Low birth weight (<2,500g) n/N (%)	64/340 (18.8%)	12/127 (9.4%)	0.40 [0.20-0.77]	0.0070
Premature birth (<37 weeks) n/N (%)	35/343 (10.2%)	8/130 (6.2%)	0.59 [0.26-1.33]	0.20
Fetal loss n/N (%)	5/350 (1.4%)	1/132 (0.8%)	0.53 [0.05-5.19]	0.59
Gestation duration (weeks) Mean (SD)[†]	39.2 (2.2)	39.8 (1.6)	0.6 [0.2-0.9]	0.0030
Birth weight (grams) Mean (SD)[‡]	2883.2 (516.3)	3001.5 (487.4)	169.3 [65.9-272.6]	0.0010

Note: Small for gestational age defined as a birthweight below the 10th percentile for gestational age according to INTERGROWTH-21 standards;[19] Low birth weight is defined as a a birth weight < 2500 g; Premature birth is a birth occurring at <37 weeks gestation; Fetal loss represents a composite of pregnancy loss and stillbirth. [#] N=489 Pre-COVID and 270 COVID. [$]N= 484 Pre-COVID and 265 COVID [†]N=343 Pre-COVID and 130 COVID. [‡]N= 340 Pre-COVID and 127 COVID. OR - Odds Ratio; CI - Confidence Interval. All models controlled for treatment group, maternal age, body mass index, primigravid status, and HIV status. Analysis were run on all samples with non-missing data.

Figure 1: Participant timelines and cohort characteristics across pre-pandemic and pandemic periods. (a) Individual participant timeline distribution across the study period. Horizontal lines represent each participant's pregnancy duration, with enrollment dates marked as circles. Blue horizontal lines are pregnancies that occurred exclusively in the pre-COVID-19 period. Magenta lines are pregnancies that occurred partially in the COVID-19 period. Dark red lines are pregnancies entirely occurring within the COVID-19 period. Shaded boxes represent reported COVID-19 waves (wave 1 – green; waves 2 and wave 3 – pink). (b) Anti-tetanus toxoid IgG levels by visit date (enrolment and delivery) (unpaired Welch's t-test). (c) Anti-tetanus toxoid IgG levels at enrollment between first-time pregnancies (G1) and second or higher pregnancies (G2+) (unpaired Welch's t-test). (d) anti-tetanus toxoid IgG levels at delivery between first-time pregnancies (G1) and second or higher pregnancies (G2+) (unpaired t-test). (e) Anti-tetanus toxoid IgG levels between pregnancies occurring entirely within the pre-COVID period (solely pre-COVID) or pregnancies that occurred exclusively during the Malawi COVID-19 period (Solely COVID) (unpaired t-test). All serological values are shown as $\log_e$-transformed median fluorescence intensity (MFI).

Figure 2: Serological patterns of anti-SARS-CoV-2 antibodies during pregnancy. (a) LOWESS smoothing curves illustrating the relationship between time and anti-SARS-CoV-2 antibodies is shown for SARS-CoV-2 S2 and SARS-CoV-2 NP antigens. The analysis was conducted using GraphPad Prism v10. This method fits a non-parametric regression and makes no assumptions about the functional form of the relationship. The graph includes the original data points as scatter markers to allow assessment of the density and distribution of observations across the range of values. (b) Mean percent change in IgG levels per 3 days across the three time periods assessed after fitting a linear regression model. (c) Comparison of mean antibody levels to SARS-CoV-2 antigens between pregnancies occurring outside (entirely pre-COVID) or inside (entirely COVID) the COVID period (unpaired T-test). (d) Correlation coefficient matrix between antibodies measured against SARS-CoV-2 antigens in samples from pregnancies outside (Solely pre-COVID - top) or inside (Solely COVID - bottom) the COVID period. (e) Comparison of mean correlation coefficient between antibodies against SARS-CoV-2 antigens between pregnancies occurring outside (Solely pre-COVID) or inside (Solely COVID) the COVID period (unpaired T-test).

Figure 3: Serological patterns of anti-common Coronavirus and Influenza A (H1N1) antibodies during pregnancy. (a) LOWESS smoothing curves illustrating the relationship between time and antibodies against Influenza A (green) and Coronavirus NL63 (teal) antigens. The analysis was conducted using GraphPad Prism v10. This method fits a non-parametric regression and makes no assumptions about the functional form of the relationship. The graph includes the original data points as scatter markers to allow assessment of the density and distribution of observations across the range of values. (b) Mean percent change in IgG levels per 30 days across the three time periods assessed after fitting a linear regression model. (c) Comparison of mean antibody levels to SARS-CoV-2 antigens between pregnancies occurring outside (entirely pre-COVID) or inside (entirely COVID) the COVID period (unpaired T-test). (d) Correlation coefficient matrix between antibodies measured against SARS-CoV-2 antigens in samples from pregnancies outside (Solely pre-COVID - top) or inside (Solely COVID - bottom) the COVID period. (e) Comparison of mean correlation coefficient between antibodies against SARS-CoV-2 antigens between pregnancies occurring outside (Solely pre-COVID) or inside (Solely COVID) the COVID period (unpaired T-test)