A modular pipeline for natural language processing-screened human abstraction of a pragmatic trial outcome from electronic health records

Robert Y. Lee, MD, MS [1,2]
Kevin S. Li, MS [1,3]
James Sibley, BS [1,4]
Trevor Cohen, MBChB, PhD [1,3]
William B. Lober, MD, MS [1,3,4,5]
Janaki O'Brien, MS [2,6]
Nicole LeDuc, BS [2,7]
Kasey Mallon Andrews, MS [2,8]
Anna Ungar, MPH, MA [1,2]
Jessica Walsh, BA [1,2]
Elizabeth L. Nielsen, MPH [1,2]
Danae G. Dotolo, PhD, MSW, MA [1,2]
Erin K. Kross, MD [1,2]

[1] Division of Pulmonary, Critical Care, and Sleep Medicine, University of Washington, Seattle, USA
[2] Cambia Palliative Care Center of Excellence at UW Medicine, University of Washington, Seattle, USA
[3] Division of Biomedical and Health Informatics, Department of Biomedical Informatics and Medical Education, University of Washington, Seattle, WA
[4] Department of Biobehavioral Nursing and Health Informatics, University of Washington, Seattle, WA
[5] Department of Global Health, University of Washington, Seattle, WA
[6] Norcliffe Foundation Center for Integrative Brain Research, Seattle Children's Research Institute, Seattle Children's Hospital, Seattle, WA
[7] Project ECHO, University of New Mexico Health Sciences Center, Albuquerque, NM
[8] Department of Surgery, University of Washington, Seattle, WA

Address correspondence to:

Robert Y. Lee, MD, MS, Assistant Professor of Medicine
Division of Pulmonary, Critical Care, and Sleep Medicine
Cambia Palliative Care Center of Excellence at UW Medicine
University of Washington School of Medicine, Seattle, WA

Address: Harborview Medical Center, 325 9th Ave., Box 359762, Seattle, WA 98104
Phone: 206-744-6359; fax: 206-744-8584; e-mail: rlee06@uw.edu

Funding/Support: This work was supported by the National Institute on Aging (R01AG062441), the National Heart, Lung, and Blood Institute (K23HL161503, K12HL137940, T32HL125195), the National Palliative Care Research Center, and the Cambia Health Foundation. Infrastructure support was provided by the Institute of Translational Health Science (National Center for Advancing Translational Sciences, UL1TR002319).

Running title: NLP-screened human adjudication of GOC
Trial registries: ClinicalTrials.gov NCT04281784 and NCT04283994

Word count: 3363

ABSTRACT

Background: Natural language processing (NLP) allows efficient extraction of clinical variables and outcomes from electronic health records (EHR). However, measuring pragmatic clinical trial outcomes may demand accuracy that exceeds NLP performance. Combining NLP with human adjudication can address this gap, yet few software solutions support such workflows. We developed a modular, scalable system for NLP-screened human abstraction to measure the primary outcomes of two clinical trials.

Methods: In two clinical trials of hospitalized patients with serious illness, a deep-learning NLP model screened EHR passages for documented goals-of-care discussions. Screen-positive passages were referred for human adjudication using a REDCap-based system to measure the trial outcomes. Dynamic pooling of passages using structured query language (SQL) within the REDCap database reduced unnecessary abstraction while ensuring data completeness.

Results: In the first trial (N=2,512), NLP identified 22,187 screen-positive passages (0.8%) from 2.6 million EHR passages. Human reviewers adjudicated 7,494 passages over 34.3 abstractor-hours to measure the cumulative incidence and time to first documented goals-of-care discussion for all patients with 92.6% patient-level sensitivity. In the second trial (N=617), NLP identified 8,952 screen-positive passages (1.6%) from 559,596 passages at a threshold with near-100% sensitivity. Human reviewers adjudicated 3,509 passages over 27.9 abstractor-hours to measure the same outcome for all patients.

Conclusion: We present the design and source code for a scalable and efficient pipeline for measuring complex EHR-derived outcomes using NLP-screened human abstraction. This implementation is adaptable to diverse research needs, and its modular pipeline represents a practical middle ground between custom software and commercial platforms.

Abstract word count: 250

Keywords: natural language processing, electronic health records, outcome measurement, goals of care, serious illness communication

Background

Advances in natural language processing (NLP) are creating new opportunities for pragmatic clinical trialists to measure important outcomes from free-text electronic health records (EHR). To date, much work in clinical NLP has focused on tasks well-suited to existing methods such as discrete data element extraction (*e.g.*, diagnoses or symptoms), named entity recognition (*e.g.*, persons, organizations, or locations), and text classification (*e.g.*, assigning sentiment or topical categories).[1,2] However, using NLP to measure pragmatic clinical trial outcomes from the EHR presents two key challenges. First, many patient-centered outcomes—such as symptoms, quality of life, adverse events, and functional outcomes—are documented in clinical language that is both linguistically complex and also highly sensitive to context. This makes these outcomes challenging for NLP to measure accurately. Second, the high cost and rigor of clinical trials demands a low tolerance for measurement error and misclassification, which are inherent risks of NLP-based approaches. For these reasons, hybrid measurement strategies that combine NLP with final adjudication of outcomes by human researchers will likely remain essential for clinical trialists. Yet, software solutions are scarce for such hybrid NLP pipelines, which may lead investigators to either forgo these approaches entirely or rely on homegrown software and databases to facilitate their studies.

Our research group has conducted several randomized clinical trials examining the effect of communication-priming interventions on the primary outcome of EHR-documented goals-of-care discussions,[3–5] an important clinician-documented process measure in palliative care research. In many health systems, such discussions are documented in unstructured text and can be difficult and costly to find within voluminous EHRs. Furthermore, the outcome of goals-of-care discussions is of high linguistic and conceptual complexity,[6–8] and it requires extensive training of human abstractors to measure with consistency. To facilitate measurement of this outcome at scale, our research group has developed multiple NLP models (including rule-based and deep learning-based models) to identify documented goals-of-care discussions. Many of these models have shown promising performance but remain insufficiently accurate for stand-alone use in measuring the primary outcome of a clinical trial.[9–12]

Our research group sought to measure EHR-documented goals-of-care discussions in two clinical trials, both evaluating the efficacy of communication-priming interventions designed to promote goals-of-care discussions and their documentation for hospitalized patients with serious illness. To do this, we implemented a two-step outcome measurement process combining an NLP screening instrument with human adjudication. We used a previously-described deep-learning NLP model to screen EHR passages,[11] then fed screen-positive passages into a human adjudication process designed to maximize the efficiency of human adjudication. We implemented the pipeline for NLP-screened human abstraction using commonly available research software and data platforms such as REDCap (Vanderbilt University, project-redcap.org) and Stata (StataCorp, stata.com).[13]

In this report, we describe and provide the source code for a NLP-screened human abstraction pipeline. We also discuss design parameters and pragmatic and statistical implications that may be applicable to other studies. We specifically focus on the design and implementation of human adjudication, which exists separately from the previously-reported NLP model[11] and is portable to different screening models and outcomes.

Design and Implementation

Problem description, and NLP screening of EHR passages

In the United States, healthcare norms often default to providing life-extending treatments to all patients.[14] However, some patients with chronic life-limiting illness may prefer a palliative or comfort-focused approach that emphasizes symptom relief over life extension.[15–17] To better align treatments with patients' values, goals and preferences, experts have highlighted the need for high-quality communication between patients, families, surrogate decision-makers, and clinicians when making decisions about life-sustaining treatments.[18,19] These conversations, commonly called "goals-of-care discussions," play an important role in delivering goal-concordant care.[6,19–22] Given the frequent handoffs and transitions in modern healthcare, experts also agree that documenting goals-of-care discussions in the EHR and communicating them across the continuum of care is essential.[19,23–25] As such, documented goals-of-care discussions are an important process measure and research outcome in the field of serious illness communication.[19,26] However, despite their importance, documented goals-of-care discussions represent a very small proportion of EHR text, and are rarely documented as structured data in the EHR. Additionally, the construct of a goals-of-care discussion is linguistically and medically complex, encompassing multiple dimensions and existing along continua rather than as discrete entities (**Figure 1**).[10,27] These factors make the process of measuring documented goals-of-care discussions through conventional, manual chart abstraction both costly and labor-intensive; and, training human abstractors is further challenged by the inconsistent and infrequent nature of such documentation in the EHR.[9,28]

In this report, we examine the NLP-screened human abstraction pipeline for two trials in the Project to Improve Communication about Serious Illness (PICSI) series: PICSI Hospital Trial 1 (hereafter, "Trial 1"; ClinicalTrials.gov ID NCT04281784), a 2,512-patient pragmatic trial conducted April 2020 – March 2021;[4,5] and, PICSI Hospital Trial 2 (hereafter, "Trial 2"; ClinicalTrials.gov ID NCT04283994), a 617-patient comparative effectiveness trial conducted July 2021 – November 2023.[4] Both trials evaluated the effect of communication-priming interventions on the primary outcome of EHR-documented goals-of-care discussions, as measured by NLP-screened human abstraction. The NLP screening instrument used for both trials was a fine-tuned instance of the freely available BioClinicalBERT[29,30] deep learning model (BERT: Bidirectional Encoder Representations from Transformers)[31] that was trained on a manually labeled corpus of 4,642 clinical notes from the PICSI Pilot Trial (hereafter, "Training dataset"), a 150-patient pilot trial examining the same outcome and conducted November 2018 – February 2020.[3] Notes were tokenized (*i.e.*, split into pre-defined words and sub-words, called tokens) using the BioClinicalBERT dictionary and further divided into non-overlapping passages (hereafter, "BERT segments") of ≤ 512 tokens each (the upper limit for BERT's sequence length) based on common whitespace patterns. Manual labeling of the training dataset was performed by a team of investigators and research staff using a standardized codebook and a qualitative data analysis platform, with regular meetings and co-review of coded passages to reconcile code application differences and ensure consistency. The training dataset was labeled over a total of 287 abstractor-hours during a 4-month period.[11]

Characteristics of the EHR corpora are presented in **Table 1**. The volume of EHR text within which documented goals-of-care discussions could be found was very large in both trials, with Trial 1 containing 44,324 eligible notes, and Trial 2 containing 11,574 eligible notes. Moreover, documented goals-of-care discussions represented only 0.2% of EHR passages (BERT segments) in the training dataset. Because the standalone BERT NLP classifier did not reach

sufficient levels of precision and recall for the study, trial investigators chose to measure the primary outcome using an NLP-screened human abstraction pipeline.[11]

Configuring REDCap for human adjudication of screened passages

Although many software packages exist for annotating unstructured text to support NLP training, there are fewer readily available solutions for rapidly adjudicating NLP-screened text for categorical outcomes. This task requires streamlined, efficient tools that minimize interactional overhead for rapid and scalable adjudication workflows. The most widely cited software for this use, ClinicalRegEx (Lindvall Lab, Dana-Farber Cancer Institute),[32,33] is constrained to regular expression-based screening. Other low-overhead software solutions such as Local Turk (Dan Vanderkam)[34] lack support for multiple abstractors or remote access. Custom-built end-user databases (*e.g.*, Microsoft Access) offer tailored functionality but require substantial effort to configure and rarely meet modern standards for multi-user performance, web accessibility, reliability or security.

Many clinical researchers are already familiar with and have access to REDCap,[13] a robust, web-accessible platform designed for secure, multi-user data collection in research settings. Although not originally intended for unstructured text review, REDCap's flexible form builder supports adaptation to a wide range of research workflows, and its application programming interfaces (APIs) allow for automated dataset management. We developed a REDCap instrument for efficient human adjudication of large volumes of NLP-screened text passages. To import EHR data into REDCap, we encoded screen-positive NLP passages into comma-separated values (CSV) files containing candidate EHR passages, their surrounding context, and pertinent metadata, creating one REDCap record per candidate passage. We also incorporated features to flag passages requiring co-review. Detailed REDCap configurations and other design considerations are provided in **Appendix A**; our configuration is also available from the REDCap Shared Library (access instructions in **Appendix A**).[35,36] Stata source code for preparing segmentized EHR data for contextual display in REDCap is provided in **Appendix B**.

Targeted human adjudication via dynamic record-pooling

The primary outcome of the trial was the occurrence of a documented goals-of-care discussion at any time within 30 days of randomization. The investigators also aimed to measure the number of days between randomization and the *first* documented goals-of-care discussion to inform future analyses. Abstractors did not have access to patient randomization or NLP-predicted probabilities, and we used open-source rule-based NLP software to redact patient, clinician, and family names from passages.[37]

Preliminary data from the training dataset indicated that goals-of-care documentation often appears multiple times in a single patient's record, either due to copy-pasted text or multiple distinct discussions. To minimize unnecessary abstraction, we aimed to refer only those passages essential to the study outcomes for human adjudication—that is, all passages occurring between randomization and the earliest *human-confirmed* documented goals-of-care discussion for each patient, or through the end of the outcome assessment period if no such documentation was found. This created a unique challenge: the set of passages requiring abstraction for the study outcome could not be determined in advance, as it depended on the outcome of the abstraction itself. To address this, we developed a dynamic record-pooling process that ran in parallel with ongoing abstraction and removed passages occurring after a confirmed goals-of-care discussion for the same patient from the abstraction dataset (**Figure 2**).

The efficiency of the record-pooling process also depended on the order in which records were abstracted. To reduce unnecessary abstraction while avoiding a sequential, per-patient approach (*e.g.*, abstracting all records for patient 1, then patient 2, and so on)—which could introduce adverse incentives to code for positive outcomes—we shuffled screen-positive EHR passages across patients while preserving temporal order within each patient. This produced a random-appearing sequence of passages that was still optimized for dynamic record-pooling efficiency (**Figure 3**; Stata source code in **Appendix C**).

A key challenge of dynamic record-pooling is that very few data abstraction platforms have built-in functionality to change the dataset for abstraction in real-time. Owing to the large number of EHR passages in our project, it was impractical to pool records manually. To modify the abstraction dataset at regular intervals, we implemented an automated script that retrieved a deidentified report of human-adjudicated outcomes from the REDCap database, then used this data to determine which unabstracted passages needed to be removed from, or replaced back into, the dataset for abstraction. The script was built to be robust to changes in data (*e.g.*, handling of changing abstraction results following co-review), and leveraged REDCap's sorting behaviors to non-destructively discard or replace records from the abstraction dataset without deleting or re-creating REDCap records. **Appendix D** provides the source code for a Bourne-again shell (`bash`) script implementing dynamic record-pooling using structured query language (SQL) for a REDCap database for time-to-first-event analysis.

Quality control of human adjudications

Abstractors underwent a training curriculum and group abstraction work to achieve consistency in applying the abstraction codebook. All human adjudications were subject to multiple quality-control measures, including regular meetings to discuss a representative sample of human-adjudicated passages, functionality for abstractors to "flag" specific passages for group discussion, periodic audits of individual abstractors' work, and in select instances, referral to clinician-investigators for additional review to arrive at a final adjudication.

Statistical implications

The statistical implications of any two-stage outcome measurement process such as NLP-screened human abstraction must be carefully considered in any study. In our study, we assumed that the specificity of an NLP-screened human-abstracted outcome would be 100% relative to a manually-abstracted gold standard. However, the sensitivity of the outcome measure is dependent on both the performance of the NLP screening instrument and (for non-dichotomous screening instruments) selection of the NLP screening threshold. Outcome misclassification from an imperfect NLP screening instrument may also affect the study power and findings of association for a given study. We have previously reported on the power implications of non-differential outcome misclassification in randomized trials that use NLP-screened human abstraction with imperfect sensitivity.[11]

In considering NLP-screened human abstraction of patient-level outcomes, in which records are screened and adjudicated at a fine level of measurement (*i.e.*, the EHR passage) and patient outcomes are defined by aggregation of passage-level outcomes, it is critically important to note two things: (1) that performance characteristics at passage- and patient-levels are likely to differ substantially, in ways that are dependent on the nature of the aggregation step between passage- and patient-level outcomes; and, (2) that the patient-level sensitivity of the NLP instrument at a given threshold k is <u>not</u> equal to the patient-level sensitivity of NLP-screened human abstraction at the same threshold k. This second observation may seem counterintuitive,

but it is explained by the fact that the posterior testing step in NLP-screened human abstraction only refers screen-positive passages for human adjudication. As a result, some patients who would have true-positive patient-level outcomes when measured by an NLP-only instrument would have false-negative patient-level outcomes when measured by NLP-screened human abstraction using the same threshold, because the NLP instrument may both fail to detect any true-positive passage-level outcomes (thereby concealing positive passage-level outcomes from the human abstractor) while also detecting false-positive passages that are then correctly adjudicated as negative by the human abstractor. These observations lead to the conclusions that (1) estimating patient-level performance from validation data requires either validation data that is collected at the patient level, or advanced statistical methods that may require additional assumptions for validity; and, (2) estimating patient-level sensitivity of NLP-screened human abstraction similarly requires validation data measured at both the passage and patient level.

Implementation results

Over April 2020 to March 2021, Trial 1 enrolled 2,512 patients that were randomized into two arms: 1,255 to a pragmatic clinician-facing-only communication-priming intervention (Jumpstart Guide), and 1,257 to usual care. Over July 2021 to November 2023, Trial 2 enrolled 617 patients that were randomized into three arms: 203 to a bilateral patient- and clinician-facing Jumpstart Guide, 205 to a clinician-facing-only Jumpstart Guide, and 209 to usual care with enrollment surveys. The total number of eligible EHR notes for the primary outcome was 44,324 in Trial 1, and 11,574 in Trial 2 (**Table 1**).

For Trial 1, investigators selected a screening threshold for the NLP screening instrument of 0.5 (an uncalibrated constant).[11] In a 159-patient validation sample drawn from within Trial 1 that underwent conventional whole-chart abstraction, this screening threshold corresponded to NLP passage-level sensitivity of 83.7%, specificity of 99.2%, and precision of 35.9%; note-level sensitivity of 96.2%, specificity of 77.9%, and precision of 37.6%; and, patient-level sensitivity of 99.2%, specificity of 33.2%, and precision of 45.6%. Based on simulations of NLP-screened human abstraction performed within the validation sample, the estimated patient-level sensitivity of NLP-screened human abstraction at the selected screening threshold was 92.6%.[11] Using this NLP screening threshold, our BioClinicalBERT classifier was able to identify 22,187 of 2.6 million (0.8%) EHR passages representing 11,287 of 44,324 (25%) notes and 1,957 of 2,512 patients (80%) that were referred for human adjudication. Of these 22,187 passages, 7,494 (33.8%) were ultimately adjudicated by a team of 3 abstractors over 34.3 abstractor-hours during a 3-week period to achieve complete data for time-to-first-event analysis. For reference, conventional chart abstraction was estimated to require approximately 2,000 abstractor-hours for this same dataset.[11]

For Trial 2, investigators were able to select a lower (more sensitive) screening threshold for the NLP screening instrument due to the smaller number of eligible records for review. To maximize sensitivity, investigators selected a screening threshold corresponding to the 98.5th percentile BERT score (an uncalibrated constant of 0.00279598). The selected threshold was 100% sensitive for documented goals-of-care discussions in the Trial 1 validation sample; as of this writing, validation data from Trial 2 is still undergoing collection. Using this NLP screening threshold, our BioClinicalBERT classifier identified 8,952 of 559,596 (1.6%) EHR passages representing 3,718 of 11,574 (32%) notes and 533 of 617 (86%) patients that were referred for human adjudication. Of these 8,952 passages, 3,509 (39%) were ultimately adjudicated by a team of 8 abstractors over 27.9 abstractor-hours during a 2-month period to achieve complete data for time-to-first-event analysis. For reference, conventional chart abstraction was estimated to require approximately 500 abstractor-hours for this trial dataset.

Discussion, applications and adaptability

By combining commonly available research software and data platforms such as REDCap and Stata with our research center's separately-developed BERT-based NLP infrastructure, we were able to implement a streamlined and efficient workflow for performing large-scale NLP-screened human abstraction of a complex linguistic outcome that leveraged REDCap APIs for automated dynamic record-pooling to achieve complete data with maximal efficiency. Although our implementation is certainly not an "out-of-the-box" solution for all clinical researchers, it provides a starting point for measurement of EHR outcomes that require a two-step measurement strategy with automated screening and subsequent human adjudication. Our approach is portable to different NLP screening methods and models, which are implemented upstream of the human adjudication pipeline; and, it may be further adapted to other types of measures and different essential datasets with nominal programming effort. By implementing this pipeline in a modular fashion using freely available software and platforms, our approach also maximizes flexibility for future deployments in other research projects while maintaining a low-friction user experience and minimizing maintenance development costs. Using common secure platforms such as REDCap ensures a baseline level of data security often overseen at an institutional level while still allowing simultaneous abstraction efforts by multiple individual abstractors over a web-based platform.

In studies where the event of interest is rare, dynamic record-pooling as described herein can significantly reduce the burden of human abstraction. In Trial 1, there were 22,187 screen-positive passages at the selected threshold; human adjudication of all screen-positive passages would have required over 100 abstractor-hours. By dynamically discarding passages following a human-confirmed goals-of-care discussion for the same patient, we were able to achieve complete data for the outcomes of interest (*i.e.*, cumulative incidence and time-to-first event) abstracting only 7,494 passages over 34.3 abstractor-hours—a 66% reduction.[11] Investigators considering such dynamic record-pooling must carefully consider the essential dataset of EHR passages for their type of measure, and how best to prioritize or discard passages for human adjudication to achieve the essential dataset with maximal efficiency and minimal potential bias. In **Table 2**, we discuss process considerations for implementing NLP-screened human abstraction of three common types of measures—time-to-event, cumulative incidence, and incidence rate—alongside general considerations that apply to all types of measures.

Despite the advantages of such a two-stage measurement approach, adopting similar approaches to complex outcome measurement processes also has several important downsides. Although the programming burden of implementing such a pipeline is not high by programmers' standards, substantial technical expertise and access to relevant platforms and servers is nevertheless required for implementation. Furthermore, improper configuration may expose study participants to data security risks, which is of paramount concern when working with human subjects and protected health information. As EHR data becomes an increasingly common source of covariates and outcomes in clinical research, investigators must become familiar with the many ways one may extract data from the EHR and choose their data collection processes wisely. Close collaboration between clinical researchers, informaticists, and front-line research staff is critically important to ensuring fidelity of measurement processes to the clinical variables and outcomes of interest, ensuring an efficient and streamlined user experience for abstractors that maximizes accuracy, and ensuring that all data is stored and transmitted securely.

Conclusion

To facilitate measurement of a linguistically complex EHR-documented outcome in two randomized trials, we combined previous work in NLP with common research software and data platforms to construct a workflow and software pipeline for collecting NLP-screened human-adjudicated outcome measures in large EHR corpora using a streamlined and efficient web-based user interface with dynamic record-pooling to minimize total abstraction time. The methodological details and principles, underlying source code and data instrument design provided herein are easily adaptable to other clinical research projects that aim to measure clinical variables and outcomes using EHR-screened human abstraction or any other two-stage measurement strategy that requires human adjudication. Our implementation of this pipeline represents an appealing middle-ground between custom software solutions and repurposed commercial products for researchers interested in using EHR data.

TABLES

Table 1. Characteristics of electronic health record note corpora.

	Training dataset	Trial 1 dataset	Trial 2 dataset
No. patients	150	2,512	617
No. (%) patients with GOCD	34 (23)	–	–
No. notes per patient, median (IQR)	19 (11-29)	12 (7-22)	14 (9-23)
No. EHR notes	4,642	44,324	11,574
No. (%) notes with GOCD	340 (7)	–	–
Note length, words, median (IQR)	1070 (1578-3202)	1039 (688-1468)	983 (622-1695)
No. BERT segments (≤512 tokens)	201,394	2,644,387	559,596
No. (%) BERT segments with GOCD	446 (0.2)	–	–

Abbreviations: GOCD, [documented] goals-of-care discussion; EHR, electronic health record; IQR, interquartile range; BERT, Bidirectional Encoder Representations from Transformers.

Table 2. Process considerations for NLP-screened human abstraction of rare events.

Type of measure	Essential dataset of EHR passages	Considerations for targeting human adjudication	Additional considerations
Time to first event	All screen-positive passages from observation start to first confirmed event, or to observation end if none found	Randomize patient and passage order to reduce bias. Prioritize passages chronologically, discarding those after a confirmed true-positive.	Selected NLP screening threshold affects both resulting sensitivity and expected size of essential dataset. Pilot abstractions and simulations may estimate burden of human abstraction.

Patient-level sensitivity of an NLP instrument at a given threshold does <u>not</u> equate to the sensitivity of NLP-screened human abstraction at the same threshold (see *Statistical implications*).

Abstraction pipelines must be robust to changes in data (*e.g.*, changes to human abstraction decisions).

Redundancy filtering (*e.g.*, duplicate passage elimination, NLP-based similarity adjudication) reduces abstraction burden in records with high text redundancy. Integrating within-patient active learning into NLP pipelines may further lower abstraction burden with minimal cross-patient bias.

Efficient abstraction approaches may show a decline in apparent precision over time due to preferential discarding of true positives. Measures to mitigate this—at the cost of increased abstraction—may improve abstractor concentration, accuracy and morale. |
| **Cumulative incidence** | At least one confirmed event per patient with an event; and, all screen-positive passages for patients with no confirmed events | Prioritize passages by NLP-predicted probability or other rules to maximize efficiency, discarding passages after a confirmed true-positive. | |
| **Incidence rate** | At least one confirmed event per positive unit of person-time; and, all screen-positive passages for units of person-time with no confirmed events | Aggregate records to discard passages in the same person-time unit as a confirmed true positive. Prioritize passages within time units by NLP-predicted probability or other rules to maximize efficiency. | |

Abbreviations: NLP, natural language processing; EHR, electronic health record.

FIGURES

Figure 1. Dimensions and examples of EHR documentation related to goals of care.

Abbreviations: GOC, goals of care; TLD: tubes, lines, drains; RIJ CVC: right internal jugular central venous catheter; VTE: venous thromboembolism; DNR/DNI: do not resuscitate / intubate; DPOA: durable power of attorney.

Figure 2. Abstraction process for achieving minimum complete dataset for time-to-first-event analysis with dynamic record-pooling.

Data provided is hypothetical, and has been simplified to reflect one electronic health record (EHR) passage per note per hospital day (HD). For this depiction, abstraction has been divided into hypothetical "rounds" of one record per patient, partitioned by dynamic record-pooling events. "NLP+" denotes screen-positive records. Results of human adjudication of screened records are denoted as "TP" (true-positive) or "FP" (false-positive). Records discarded by dynamic record-pooling are shown as hashed. Actual sequence of passages adjudicated in each "round" of abstraction may vary by sort order (see **Figure 3**), and actual dynamic record-pooling may take place at any time (see **Appendix D**), *i.e.*, not necessarily between "rounds."

Figure 3. Temporal-sequence-preserving sort for human adjudication of time-to-event data.

Patient ID	Note date	Passage ID
1	04/06/24	101
1	04/08/24	102
1	04/09/24	103
1	04/11/24	104
2	09/29/24	105
2	09/30/24	106
2	10/03/24	107
3	01/22/24	108
3	01/24/24	109
3	01/27/24	110
3	01/28/24	111

random sort →

Patient ID	Note date	Passage ID
1	04/09/24	103
1	04/11/24	104
3	01/24/24	109
2	10/03/24	107
3	01/28/24	111
3	01/22/24	108
1	04/06/24	101
3	01/27/24	110
2	09/30/24	106
1	04/08/24	102
2	09/29/24	105

restore temporal sequence within each patient →

Patient ID	Note date	Passage ID
1	04/06/24	101
1	04/08/24	102
3	01/22/24	108
2	09/29/24	105
3	01/24/24	109
3	01/27/24	110
1	04/09/24	103
3	01/28/24	111
2	09/30/24	106
1	04/11/24	104
2	10/03/24	107

ACKNOWLEDGEMENTS

Author contributions: Dr. Lee had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Role of the Funder/Support: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Acknowledgement: The authors are grateful for the contributions of the late J. Randall Curtis, MD, MPH, who was a founding principal investigator of this research program.

References

1. Sheikhalishahi S, Miotto R, Dudley JT, Lavelli A, Rinaldi F, Osmani V. Natural Language Processing of Clinical Notes on Chronic Diseases: Systematic Review. *JMIR Med Inform*. 2019;7(2):e12239. doi:10.2196/12239

2. Spasic I, Nenadic G. Clinical Text Data in Machine Learning: Systematic Review. *JMIR Med Inform*. 2020;8(3):e17984. doi:10.2196/17984

3. Lee RY, Kross EK, Downey L, et al. Efficacy of a Communication-Priming Intervention on Documented Goals-of-Care Discussions in Hospitalized Patients With Serious Illness: A Randomized Clinical Trial. *JAMA Netw Open*. 2022;5(4):e225088. doi:10.1001/jamanetworkopen.2022.5088

4. Curtis JR, Lee RY, Brumback LC, et al. Improving communication about goals of care for hospitalized patients with serious illness: Study protocol for two complementary randomized trials. *Contemp Clin Trials*. 2022;120:106879. doi:10.1016/j.cct.2022.106879

5. Curtis JR, Lee RY, Brumback LC, et al. Intervention to Promote Communication About Goals of Care for Hospitalized Patients With Serious Illness: A Randomized Clinical Trial. *JAMA*. 2023;329(23):2028-2037. doi:10.1001/jama.2023.8812

6. Secunda K, Wirpsa MJ, Neely KJ, et al. Use and Meaning of "Goals of Care" in the Healthcare Literature: a Systematic Review and Qualitative Discourse Analysis. *J Gen Intern Med*. 2019;35(5):1559-1566. doi:10.1007/s11606-019-05446-0

7. Edmonds KP, Ajayi TA. Do We Know What We Mean? An Examination of the Use of the Phrase "Goals of Care" in the Literature. *J Palliat Med*. 2019;22(12):1546-1552. doi:10.1089/jpm.2019.0059

8. Myers J, Kim GL, Bytautas J, Webster F. Differing Conceptualizations of the Goals of Care Discussion: A Critical Discourse Analysis. *J Pain Symptom Manage*. 2022;63(4):495-502. doi:10.1016/j.jpainsymman.2021.12.020

9. Lee RY, Brumback LC, Lober WB, et al. Identifying Goals of Care Conversations in the Electronic Health Record Using Natural Language Processing and Machine Learning. *J Pain Symptom Manage*. 2021;61(1):136-142 e2. doi:10.1016/j.jpainsymman.2020.08.024

10. Uyeda AM, Curtis JR, Engelberg RA, et al. Mixed-methods evaluation of three natural language processing modeling approaches for measuring documented goals-of-care discussions in the electronic health record. *J Pain Symptom Manage*. 2022;63(6):e713-e723. doi:10.1016/j.jpainsymman.2022.02.006

11. Lee RY, Kross EK, Torrence J, et al. Assessment of Natural Language Processing of Electronic Health Records to Measure Goals-of-Care Discussions as a Clinical Trial Outcome. *JAMA Netw Open*. 2023;6(3):e231204. doi:10.1001/jamanetworkopen.2023.1204

12. Lee RY, Uyeda AM, Li KS, et al. Performance of Predefined Search Patterns for Identifying Documented Goals-of-Care Discussions in Inpatient Electronic Health Records. *J Palliat Med*. Published online April 4, 2025. doi:10.1089/jpm.2024.0496

13. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inf.* 2009;42(2):377-381. doi:10.1016/j.jbi.2008.08.010

14. Institute of Medicine. *Dying in America: Improving Quality and Honoring Individual Preferences Near the End of Life.* National Academies Press; 2015.

15. Teno JM, Fisher ES, Hamel MB, Coppola K, Dawson NV. Medical care inconsistent with patients' treatment goals: association with 1-year Medicare resource use and survival. *J Am Geriatr Soc.* 2002;50(3):496-500. doi:10.1046/j.1532-5415.2002.50116.x

16. Mack JW, Weeks JC, Wright AA, Block SD, Prigerson HG. End-of-life discussions, goal attainment, and distress at the end of life: predictors and outcomes of receipt of care consistent with preferences. *J Clin Oncol.* 2010;28(7):1203-1208. doi:10.1200/JCO.2009.25.4672

17. Cosgriff JA, Pisani M, Bradley EH, O'Leary JR, Fried TR. The association between treatment preferences and trajectories of care at the end-of-life. *J Gen Intern Med.* 2007;22(11):1566-1571. doi:10.1007/s11606-007-0362-6

18. Turnbull AE, Bosslet GT, Kross EK. Aligning use of intensive care with patient values in the USA: past, present, and future. *Lancet Respir Med.* 2019;7(7):626-638. doi:10.1016/S2213-2600(19)30087-6

19. Bernacki RE, Block SD, American College of Physicians High Value Care Task F. Communication about serious illness care goals: a review and synthesis of best practices. *JAMA Intern Med.* 2014;174(12):1994-2003. doi:10.1001/jamainternmed.2014.5271

20. Comer AR, Hickman SE, Slaven JE, et al. Assessment of Discordance Between Surrogate Care Goals and Medical Treatment Provided to Older Adults With Serious Illness. *JAMA Netw Open.* 2020;3(5):e205179. doi:10.1001/jamanetworkopen.2020.5179

21. Curtis JR, Patrick DL, Shannon SE, Treece PD, Engelberg RA, Rubenfeld GD. The family conference as a focus to improve communication about end-of-life care in the intensive care unit: opportunities for improvement. *Crit Care Med.* 2001;29(2 Suppl):N26-33. doi:10.1097/00003246-200102001-00006

22. Seaman JB, Arnold RM, Scheunemann LP, White DB. An Integrated Framework for Effective and Efficient Communication with Families in the Adult Intensive Care Unit. *Ann Am Thorac Soc.* 2017;14(6):1015-1020. doi:10.1513/AnnalsATS.201612-965OI

23. Dy SM, Apostol C, Martinez KA, Aslakson RA. Continuity, coordination, and transitions of care for patients with serious and advanced illness: a systematic review of interventions. *J Palliat Med.* 2013;16(4):436-445. doi:10.1089/jpm.2012.0317

24. Sinuff T, Dodek P, You JJ, et al. Improving End-of-Life Communication and Decision Making: The Development of a Conceptual Framework and Quality Indicators. *J Pain Symptom Manage.* 2015;49(6):1070-1080. doi:10.1016/j.jpainsymman.2014.12.007

25. You JJ, Fowler RA, Heyland DK, Canadian Researchers at the End of Life N. Just ask: discussing goals of care with patients in hospital with serious illness. *CMAJ*. 2014;186(6):425-432. doi:10.1503/cmaj.121274

26. Jacobsen J, Bernacki R, Paladino J. Shifting to Serious Illness Communication. *JAMA*. 2022;327(4):321. doi:10.1001/jama.2021.23695

27. Chien I, Shi A, Chan A, Lindvall C. Identification of Serious Illness Conversations in Unstructured Clinical Notes Using Deep Neural Networks. In: Fernando Koch, Andrew Koster, Isabelle Bichindaritz, Pau Herrero, et al, eds. *Artificial Intelligence in Health: First International Workshop, AIH 2018, Stockholm, Sweden, July 13-14, 2018, Revised Selected Papers*. Lecture Notes in Artificial Intelligence. Springer Nature Switzerland; 2019:199-212.

28. Wilson CJ, Newman J, Tapper S, et al. Multiple locations of advance care planning documentation in an electronic health record: are they easy to find? *J Palliat Med*. 2013;16(9):1089-1094. doi:10.1089/jpm.2012.0472

29. Alsentzer E, Murphy JR, Boag W, et al. Publicly Available Clinical BERT Embeddings. Published online June 20, 2019. Accessed August 29, 2024. https://arxiv.org/abs/1904.03323

30. Alsentzer E. BioClinicalBERT. 2019. https://huggingface.co/emilyalsentzer/Bio_ClinicalBERT

31. Devlin J, Chang MW, Lee K, Toutanova K. BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding. Published online May 24, 2019. Accessed August 29, 2024. http://arxiv.org/abs/1810.04805

32. Udelsman B, Chien I, Ouchi K, Brizzi K, Tulsky JA, Lindvall C. Needle in a Haystack: Natural Language Processing to Identify Serious Illness. *J Palliat Med*. 2019;22(2):179-182. doi:10.1089/jpm.2018.0294

33. Lindvall C, Deng CY, Moseley E, et al. Natural Language Processing to Identify Advance Care Planning Documentation in a Multisite Pragmatic Clinical Trial. *J Pain Symptom Manage*. 2022;63(1):e29-e36. doi:10.1016/j.jpainsymman.2021.06.025

34. Vanderkam D. Local Turk. 2024. Accessed January 22, 2025. https://github.com/danvk/localturk

35. Lee RY, Li KS, Sibley J, et al. NLP-screened EHR passage adjudication. REDCap Shared Library. January 22, 2025. https://redcap.vumc.org/consortium/library/search.php (requires session authentication via local REDCap project)

36. Obeid JS, McGraw CA, Minor BL, et al. Procurement of shared data instruments for Research Electronic Data Capture (REDCap). *J Biomed Inform*. 2013;46(2):259-265. doi:10.1016/j.jbi.2012.10.006

37. Manning C, Surdeanu M, Bauer J, Finkel J, Bethard S, McClosky D. The Stanford CoreNLP Natural Language Processing Toolkit. In: *Proceedings of 52nd Annual Meeting of*