Original Article

Title: Positive effects of functional electrical stimulation-assisted cycling on perception of effort, cerebral blood flow and cognition in post-stroke patients

Short Title: Functional electrical stimulation-assisted cycling on stroke patients

Author Names: Maël Descollonges (PhD)[1,2,3#], Julie Di Marco (MD)[2,3#], Ehsan Jafari (PhD)[2,4], Pierre-Henri Pouillart (MSc)[3], Julien V. Brugniaux (PhD)[5], Benjamin Pageaux (PhD)[6,7,8] and Gaëlle Deley (PhD)[1]

Affiliations:

[1] INSERM UMR 1093 – Laboratoire CAPS, Université de Bourgogne, UFR des Sciences du Sport, Dijon, France

[2] Kurage, Lyon, France

[3] SMR Val Rosay, Saint-Didier au Mont d'Or, France

[4] Université de Lyon, ENS de Lyon, CNRS, Laboratoire de Physique, F-69342 Lyon, France

[5] HP2 laboratory, Univ. Grenoble Alpes, INSERM, CHU Grenoble Alpes, Grenoble, France

[6] Centre de recherche de l'Institut universitaire de gériatrie de Montréal (CRIUGM), Canada

[7] École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Université de Montréal, Montréal, Canada

[8] Centre Interdisciplinaire de Recherche sur le Cerveau et l'Apprentissage (CIRCA), Montréal, QC, Canada

[#] First co-authors.

Correspondence address: Maël Descollonges, 64b rue de sully, 21000 – Dijon, Inserm CAPS u1093 +33 6 79 64 57 39, descollonges.mael@gmail.com

Submission Summary

[Abstract word count: **172**/200]

[Lay Abstract word count: **129**/150]

[Introduction to conclusion word count: **3583**]

[Manuscript word count: **5474**]

[Number of Tables: **1**]

[Number of Figures: **4**]

[Number of References: **31**]

[Number of Supplementary Materials Submitted: **1** – **1** table, **3** figures]

Abstract

Objective: To test the effect of functional electrical stimulation-assisted cycling on stroke

patients' perception of effort (primary aim), cerebral blood flow, and cognitive performance

(exploratory aims).

Design: Randomized controlled study.

Subjects/Patients: Fifteen post-stroke patients completed functional electrical stimulation-

assisted and traditional cycling sessions separated by 72h.

Methods: Perceived effort, cardio, and cerebrovascular parameters were monitored during

exercise. Cognitive performance was assessed before and after each session. Qualitative data

were reported after both sessions.

Results: Patients reported a lower perceived effort during functional electrical stimulation-

assisted cycling than traditional cycling. Both sessions increased heart rate, end-tidal CO_2,

cardiac output, and cerebral artery blood flow velocity, with higher blood lactate levels after

functional electrical stimulation-assisted cycling. Functional electrical stimulation-assisted

cycling positively impacted cognitive performance.

Conclusion: Traditional and functional electrical stimulation-assisted cycling induced similar

increased cardio and cerebrovascular responses. However, patients perceived functional

electrical stimulation-assisted cycling as less effortful than traditional cycling. As effort is a

barrier to regular exercise engagement and adherence, these results are promising for

implementing functional electrical stimulation-assisted cycling in stroke patients'

rehabilitation.

Key words: Cerebral blood flow, cognition, functional electrical stimulation, rating of

perceived exertion, stroke

Lay Abstract

This study explored whether cycling assisted by functional electrical stimulation (FES) could

make exercise easier for people who have had a stroke, while also providing benefits to the

brain. Fifteen stroke survivors completed two cycling sessions, one with FES and one

without, spaced 72 hours apart. We measured perceived effort, cardiovascular and cerebral

responses during exercise, as well as cognitive performance before and after each session.

Patients reported that FES-assisted cycling felt less difficult and less tiring than traditional

cycling, even though both sessions triggered similar heart and brain blood flow responses.

Cognitive performance also improved more after the FES-assisted session. Since perceived

effort is often a major barrier to rehabilitation after stroke, these findings suggest that FES-

assisted cycling could promote better engagement in physical activity and support recovery.

Introduction

Stroke is a major cause of long-term disability affecting walking, postural control, muscle

strength, cerebrovascular and cardiovascular systems, and cognition. To counteract

cardiorespiratory system deconditioning and experience the positive effect of physical

exercise on cognition (1), stroke patients must engage in regular aerobic exercise during their

rehabilitation. Due to hemiparesis and fatigue leading to increased effort perception (2), and

the proposed key role of effort and its perception in the engagement in physical and cognitive

activities (3), post-stroke individuals experience difficulty engaging in and performing

physical exercise. Therefore, developing and implementing methods of rehabilitation that

minimize the perception of effort experienced during physical exercise is crucial.

To this aim, functional electrical stimulation (FES)-assisted cycling is a promising method.

Clinical evidence shows that FES-assisted cycling activates paralyzed muscles (4) and

improves aerobic capacity and cardiopulmonary function (5). By evoking muscle contraction

superimposed on top of the impaired voluntary contractions, FES is known to assist in

movement execution (4). As the evoked contraction induced by FES pre-activates the

muscles, the individual engaged in FES-assisted cycling must voluntarily produce a lower

muscle force to reach the targeted exercise intensity. As effort perception is proposed to be

generated by signals related to the magnitude of the motor command (6,7) and the lower

muscle force to be voluntarily produced during FES should be associated with a lower motor

command compared to traditional cycling, using FES-assisted cycling is a promising tool to

decrease the level of perceived effort experienced during rehabilitation sessions. To date, it

has been reported that FES application was associated with a reduced perception of effort in

stroke or multiple sclerosis patients during walking tasks (8,9). However, the effectiveness in

reducing perceived effort during cycling in stroke patients remains to be validated. As cycling

is a key aerobic exercise in rehabilitation, there is an urgent need for such a study.

In this context, this study aimed to compare the perceived effort during an FES-assisted

cycling session, compared to a traditional cycling session, in stroke patients. We hypothesized

that stroke patients would perceive a lower effort during FES-assisted cycling compared to

traditional cycling, performed at the same workload. In addition, considering recent research

suggesting a positive effect of electrical muscle stimulation on cerebral blood flow velocity

(CBFv, (10,11)) and potentially cognitive performance, we also explored, in a subsample of

patients, changes in CBFv to the FES- and traditional cycling exercises, and cognitive

performance pre-post exercises.

Methods
106

107   Fifteen post-stroke patients volunteered in this experiment (8♂; 7♀; age: 53 ± 13 years, height:

108   167 ± 0 cm, mass: 69 ± 10 kg, body mass index: 25 ± 3, 148 ± 108 days after stroke).

109   Patients' characteristics stratified by sex are available in Table 1.

110 **Table 1:** Patient's characteristics. I: Ischemic; H: Hemorrhagic.

Females							
Patients	Range of age	Size (cm)	Weight (kg)	BMI	Stroke type	Days after stroke	Hemiparetic side
1	35-40	150	67	29.8	H	189	Left
2	35-40	169	69	24.2	I	40	Right
3	75-80	170	64	22.1	I	90	Right
4	50-55	155	63	26.1	I	90	Right
5	35-40	160	79	30.9	I	80	Right
6	35-40	170	74	25.6	H	237	Right
7	55-60	158	48	19.2	I	385	Left
8	65-70	163	68	25.5	I	213	Left
Mean	51	160	66	25	-	166	-

Males							
9	50-60	175	67	21.9	I	330	Left
10	70-75	165	66	24.2	I	38	Left
11	60-65	175	78	25.5	I	90	Left
12	35-40	162	74	28.1	H	97	Left
13	55-60	167	66	23.7	I	104	Right
14	45-50	186	91	26,4	H	207	Left
15	55-60	180	60	20.1	I	33	Right
Mean	55	173	72	24	-	128	-

Each patient provided written informed consent describing the procedure and potential risks

of the study. The sample size was based on patients' availability at the time of the experiment

and an expected large effect size of FES-assisted cycling on the perception of effort. Inclusion

criteria were diagnosis of an ischemic (n=11) or hemorrhagic (n=4) stroke resulting in

hemiparesis; adequate cognitive abilities to engage in rehabilitation activities; able to sit up to

30 minutes; joint mobility ranges that would not preclude pedaling. The study was supported

by UGECAM Rhône-Alpes and received ethical approval from the Comité de Protection des

Personnes Sud-Est II (French Institutional Review Board). The study was approved under the

clinical trial registration number NCT06230796.

Patients visited the hospital on six occasions. In visit 1, participants performed an incremental

test to determine their peak power output (PPO). The incremental test started at 25 watts, with

the power output increasing by 10 watts every minute until exhaustion. Exhaustion was

defined as voluntary disengagement from the task or an inability to maintain a cadence above

50 revolutions/minute (rpm) for more than 3 seconds, despite verbal encouragement. In visits

2-3-4, patients were extensively familiarized with FES, the cognitive tasks, and the

psychophysical scale used to report their perceived effort. Each familiarization visit was

separated by 48 hours. In visits 5-6, participants performed two testing sessions in a

randomized order: FES-assisted cycling and traditional cycling. Testing sessions were

separated by 72 hours to control for the presence of muscle fatigue and soreness in the second

testing session (visit 6).

During FES-assisted cycling, electrical stimulation was applied to the quadriceps, hamstrings,

and gluteus muscles of the hemiparesis side during cycling using a stimulator (Motimove, 3F-

Fabricando, Belgrade, Serbia) and an electrical stimulation-assisted cycle ergometer (Kurage,

HephaBike, Lyon, France) (Figure 1). The paretic leg was secured in alignment using an

orthotic pedal, ensuring stabilization of the foot and leg during the exercise (Figure 1). A

computerized program linked to an encoder fixed in the pedal axis was employed to stimulate

the muscles in the movement's proper phase. The appropriate phase corresponds to the knee

extension and knee flexion (pedaling movement). Quadriceps and gluteus muscles were

stimulated during extension, and hamstrings during flexion. The stimulation frequency was

fixed at 40 Hz, with a pulse width of 400 μs, and the stimulation intensity (mA) was

constantly adjusted to the individual maximum tolerable level throughout the session. The

maximum tolerable intensity of stimulation was determined by gradually increasing the

current intensity until an effective contraction was achieved and further increased to elicit the

maximal pain intensity tolerated by the participant. Each session began with a 5-minute

warm-up consisting of a progressive increase in stimulation intensity and cycling resistance to

reach 50% PPO. Afterwards, the resistance was fixed at 50% PPO for 20 minutes, and

patients had to maintain a cadence of 50 ± 3 rpm.

Figure 1: Experimental set-up. Electrical stimulation was applied to the paretic leg, on the quadriceps,

hamstrings, and gluteus muscles.

Measurements and data analysis

Perception of effort

The perception of effort is defined as "the conscious sensation of how hard, heavy, and

strenuous the physical task is" (3,12). Participants reported the intensity of their perceived

effort to cycle via the following question: ''How hard is it for you to drive your legs?". The

intensity of the perceived effort was reported every 5 minutes during the cycling exercise,

using the validated CR100 scale (13). The scale was presented to the patient on a laminated

sheet, allowing them to report the intensity of their perceived effort in real time. This scale

ranges from 0 for "nothing at all" to 100 for "maximal". The scale contains verbal anchors

such as light (13), moderate (23), strong (50), or very strong (70). The maximal intensity

(100) was anchored based on previous experience and defined as "the maximal effort

experienced in your life, for most people it corresponds to the most strenuous exercise

previously experienced." Patients were familiarized with the scale during the incremental test

and the familiarization sessions (visits 2-3-4).

Cardio- and cerebrovascular parameters

Heart rate (HR) was recorded continuously using a heart rate monitor (Forerunner 630,

Garmin, USA), mean arterial blood pressure (MAP = Diastolic Pressure + 1/3 (Systolic

Pressure – Diastolic Pressure) and cardiac output (CO) were monitored using finger

photoplethysmography attached to the non-dominant hand (Finometer PRO; Finapres Medical

Systems, Amsterdam, The Netherlands). The blood flow velocity in the right middle cerebral

artery (MCAv) was continuously monitored using a 2 MHz pulsed Doppler ultrasound system

(TCD, Multi-Dop X4, DWL Elektroniche System, Singen, Germany). Nine patients

volunteered for cerebral blood flow velocity monitoring. Each session began with

confirmation of artery insonation, relying on expected signal depth, velocity, visual signal

along, and carotid compression for validation. The Doppler probe was placed on the right

transtemporal window and secured with a DiaMon headband (DWL) throughout all

experimental sessions. The placement of the probe and the depth of the artery were carefully

recorded to be the same for each condition.

Beat-to-beat data were continuously sampled at 1 kHz using an analog-to-digital converter

(PowerLab/16SP ML795; AD Instruments, Oxford, UK) and stored for offline analysis on a

computer (Chart version 7.2.2, AD Instruments, Oxford, UK). Cardio and cerebrovascular

data were averaged for the last 30 seconds of each 5-minute interval period (before, during,

and after the exercise) as done in a previous study (11).

Cognitive tasks

Cognitive tasks were conducted on a table and in a seated position, both before (PRE) and

immediately after (POST) each experimental condition. The selected cognitive tasks included

a Stroop Task and the Trail Making Tests (TMT-A and TMT-B), as performed in previous

studies (11,14). Firstly, participants completed a three-phase Stroop task. These three phases

included three versions of the Stroop tasks, referred to as the congruent, neutral, and

incongruent conditions, respectively (11,14). Each task contained 100 items (words or

crosses). On the first page, the words "blue," "red," "yellow," and "green" were printed in

black ink (congruent task – reading task). On the second page, "XXXX" was printed in blue,

red, yellow, and green colors (neutral task). The last page displayed a color name written in a

different font color from the printed word, and the participant had to name the ink color

(incongruent task). Participants were instructed to read the words from left to right, aloud, as

quickly and accurately as possible for 45 seconds. The number of words and errors was

counted by the experimenter and recorded. Finally, these Stroop tasks were displayed on a 42-

inch screen set to maximum brightness, with a standard distance of 150 cm between

participants and the screen.

Lastly, the Trail Making Test (TMT) was performed. This test is separated into two parts. The

Trail Making Test-A (TMT-A) consists of rapidly connecting a sequence of randomly placed

numbers in ascending order from 1 to 25. The Trail Making Test-B (TMT-B) requires

participants to alternately connect numbers and letters in sequence (e.g., 1 A, 2 B, 3 C, ... 13).

TMT performance was defined as the time (seconds) taken to complete the task. The test was

performed with the patient's unaffected hand due to spasticity in the affected hand.

Blood lactate

Given that blood lactate has been proposed as a potential key factor mediating the cognitive

benefits of physical exercise (15,16), blood lactate concentration was measured. Blood lactate

was collected using an automated analyzer (Lactate Pro, Scout 4), with a 5-μl blood sample

collected from the tip of the non-paretic index finger (due to spasticity in the affected hand),

at the beginning and 2 minutes after the end of exercise to assess peak concentration within an

optimal window (17).

Patients' experience with FES-assisted cycling – Qualitative data

Following both testing sessions, the researcher informed each patient that if they want, they

are free to report to their experience with FES-assisted cycling and traditional cycling. No

predefined questions or interview guide were used. With the patients' explicit consent, their

comments were manually and anonymously transcribed on paper, verbatim.

Statistical analysis

Mean (SD). Statistical significance was determined at $\alpha < 0.05$. The Shapiro–Wilk test

confirmed data normality. Two-way repeated-measures ANOVAs (condition × time) were

used. Post-hoc tests were adjusted for multiple comparisons using the Bonferroni correction.

Effect sizes are reported with the partial eta square (partial $\eta2$), with values of 0.01, 0.06, and

above 0.14 denoting small, medium, and large differences, respectively.

Results
229

The average peak power output was 72 ± 36 watts, and the cycling exercises were performed

at 36 ± 18 watts.

Changes in stimulation intensity during FES-assisted cycling, for each muscle, are available

in the supplemental materials (Figure S1).

The perceived effort to cycle increased over time (time: $P < 0.001$; $\eta_p^2 = 0.497$) and was lower

for FES-assisted cycling compared to traditional cycling (condition: $P < 0.001$; $\eta_p^2 = 0.800$)

(Figure 2).

Figure 2: Perceived effort (A) and associated individual data (B) in response to FES-assisted (black) and traditional (white) cycling. Values are mean (SD). *n*, sample size for the analysis.

MCAv data is available for nine patients, as only these individuals volunteered for cerebral

blood flow velocity monitoring. MCAv increases during both sessions (time: $P = 0.026$; $\eta_p^2 =$

0.407), with no difference between conditions (condition: $P = 0.129$; $\eta_p^2 = 0.263$) (Figure 3).

Figure 3: Middle cerebral artery velocity [MCAv (A)] and MCAv associated individual data (B), in

response to FES-assisted (black) and traditional (white) cycling. (C) MCAv at rest (triangle), during

exercise (rectangle), and associated individual data (white squares). Values are mean (SD). *n*, sample

size for the analysis.

Due to technical issues relative to cable disconnection, PETCO☐ measurements were

available for only 12 patients, and mean arterial pressure (MAP) and cardiac output (CO) for

7 patients. Blood lactate levels were available for 13 patients.

As shown in Figure 4, HR, mean arterial blood pressure (MAP), cardiac output (CO),

$PETCO_2$, and blood lactate increased during exercise (Figure 3, all P < 0.001 and $\eta_p^2 \geq 0.327$).

Blood lactate levels were higher in the FES-assisted cycling session compared to the

traditional cycling session ($P = 0.018$; $\eta_p^2 = 0.384$). CO and MAP are presented in

supplemental materials—Figure S2 and increased in both sessions (all $P \leq 0.001$ and $\eta_p^2 \geq$

0.327).

253

Figure 4: HR (A) and associated heart rate individual data (B), PETCO2 (C) and associated PETCO2

individual data (D), and blood lactate (E) in response to FES-assisted (black) and traditional (white)

cycling. Data are presented as mean (SD). *n*, sample size for the analysis.

Due to fatigue or inability to complete the tasks, cognitive test data were available for 14

participants in the congruent Stroop condition, 13 in the neutral and incongruent conditions,

and 12 for the TMT-B test. Details of the cognitive performance data is available in

supplemental materials. Briefly, the results suggest that FES-assisted cycling induced similar

effects of traditional cycling, with a potential positive effect specific to the Neutral Stroop

task.

Eleven patients freely reported their experience with FES-assisted cycling. The verbatim of

the experience reported is available in the supplemental materials—Table S1. The most

recurring theme among the responses was a perceived benefit of the FES-assisted session

compared to traditional cycling. Most of the participants reported that the FES made pedaling

easier by inducing additional force during cycling, which reduced fatigue, and by improving

coordination and fluidity of movement on the affected limb.

Discussion

The increased perceived effort over the 20 minutes of FES-assisted and traditional cycling is

caused by the progressive development of muscle fatigue, requiring an increased motor

command to recruit additional motor units, leading to higher perceived effort levels (3,6,7).

According to our hypothesis, when cycling at the same workload, the perceived effort was

lower during FES-assisted compared to traditional cycling. To the best of our knowledge, this

study is the first to observe in stroke patients the positive effect of FES-assisted cycling on the

perception of effort. This result replicates previous observations in healthy individuals (18,19)

and conceptually replicates previous observations of a lower perceived effort during a FES-

walking exercise in multiple sclerosis (8) or stroke patients (9). It is well accepted that

applying electrical stimulation to a muscle increases afferent feedback (20). Our results,

combined with the studies observing a lower perception of effort during FES of the lower

limb (i.e., cycling or walking), provide further evidence that the perception of effort scales

with the magnitude of the motor command rather than afferent feedback (for review see (7)).

However, the mechanisms underlying the FES-induced reduction in perceived effort remain

to be fully understood and warrant further investigation. Brain imaging techniques would be

of interest to better elucidate the relationship between the motor command and perceived

effort during physical exercise involving electrical stimulation.

In our study, the lower perceived effort is likely linked to the force evoked by FES, which

helped the patients by requiring only a small additional voluntary engagement to cycle. This

interpretation is further supported by the patients' experience qualitatively reported. Indeed,

most of the patients reported perceiving clear assistance of the stimulation, helping the

movement and facilitating the exercise. A limitation to this observation is that our study was

not a-priori designed for a qualitative analysis. As our qualitative analysis to describe the

patients experience with FES-assisted cycling was exploratory, we encourage future research

to replicate our results by adopting a-priori a mixed-methods design. Doing so will allow a

better interpretation on the relationship between quantitative measurement of effort and its

perception, and the patients' perspectives and experience. Moreover, we were unable to

directly assess the actual contribution of FES to muscle force production and to measure the

force output produced by the stimulator relative to the ergometer resistance. Thus, we cannot

precisely quantify the extent to which FES contributes to force generation at different

intensities. In the future, using force sensors attached directly to the pedals would allow for a

more accurate assessment of this parameter, providing a better understanding of the role of

FES in the perception of effort, in force generation, and its effective contribution to the

cycling motion.

As a high perceived effort is a barrier to engagement in physical exercise and a key

determinant of adherence to rehabilitation programs (3,21), the observed low perceived effort

is of crucial interest for supporting the development and implementation of FES-assisted

cycling in stroke patients' rehabilitation. Future studies should now implement FES-assisted

cycling during a rehabilitation program, and not only a single training session, to test the

possibility that the effects of FES-assisted cycling on stroke patients' rehabilitation could be

superior to those induced by traditional cycling.

As cerebral blood flow has been proposed as one of the mechanisms underlying

neuroplasticity (22), we also explored the CBF responses to FES-assisted cycling and its

potential priming effect on cognitive performance in a subsample of voluntary patients. As

previously observed during an isometric NMES session (11), our results suggest that FES-

assisted cycling can increase CBF in the same proportions as cycling with lower perceived

effort. These increases in blood velocity in the middle cerebral artery are not surprising, since

similar results have been observed on pedaling tasks (23–26) and could be explained by the

increases in cardiovascular parameters observed during exercise. As the literature has reported

that the recruitment of a large muscle volume through electrical stimulation leads to a higher

cardiorespiratory demand (5,27,28), we hypothesized that CBF would be higher in the FES

condition. However, it was not the case in our study, and this could potentially be explained

by the muscle volume stimulated in our FES condition. It is well established that the muscle

mass engaged during physical exercise is a key factor influencing cerebral blood flow (29).

Although a substantial muscle volume of paretic leg was stimulated in this study (quadriceps,

hamstrings, and gluteal muscles), future research could explore the effects of stimulating an

even greater number of muscles, including the entire lower limb (e.g., quadriceps, hamstrings,

gluteus, tibialis anterior, etc.), rather than focusing solely on the paretic limb. Such an

approach could enhance the effects of stimulation on CBF. However, further research is

needed to optimize stimulation parameters and muscle selection, ensuring that potential

adverse effects are avoided. One limitation of this study is that CBF was measured only on the

right MCA, without distinguishing between the affected (ipsilateral) and unaffected

(contralateral) sides. Thus, future research should consider measuring bilateral cerebral blood

flow at the middle cerebral artery level to better understand the differences between the

affected and unaffected sides.

Our results also suggest that cycling, whether assisted by FES or not, has limited and

inconsistent effects on cognitive performance in stroke patients. Although both types of

cycling sessions led to some improvements in cognitive performance, these effects were not

consistently observed across all cognitive tasks (e.g., the incongruent condition of the Stroop

task). These partial effects—improvements seen only in specific tasks—are consistent with

the existing literature reporting mixed findings (10). For instance, recent studies have shown

that an acute NMES protocol (11,14) can enhance cognitive performance, but only under

certain conditions of the Stroop task in healthy individuals. Conversely, other studies have

found no improvement in cognitive performance, particularly in tasks such as the Go/No-Go,

among healthy participants (30,31). To address this discrepancy in the literature, we

encourage future research to focus on the effect of FES-assisted cycling on cognitive

performance as primary outcome of interest and design the study a-priori for this purpose.

Moreover, as shown in the present study, where elevated blood lactate levels were observed

following FES, recent literature has reported that NMES applied to the quadriceps increases

blood lactate levels in both humans and rats (14). Notably, higher blood lactate levels were

associated with improved Stroop Task performance in healthy humans and increased

hippocampal BDNF expression in rats following a single NMES session (14). These findings

support the hypothesis that lactate may act as a key exerkine mediating the cognitive benefits

of electrical stimulation. However, the underlying mechanisms remain poorly understood and

warrant further investigation.

Lastly, we would like to remind the reader that our sample size was determined by our

primary aim, i.e., testing the difference in perceived effort between FES-assisted cycling and

traditional cycling. Due to the likely smallest effect size of FES-assisted cycling on CBF and

cognitive performance, future studies should replicate our results with a specific a priori

approach to CBF and cognitive performance and a higher sample size. Such studies could also

unravel a potentially greater positive effect of FES-assisted cycling compared to traditional

cycling, on CBF and cognitive performance.

To conclude, this study is the first to demonstrate a reduced perceived effort during FES-

assisted cycling, compared to traditional cycling, in stroke patients. This study also suggests a

potential positive effect of FES-assisted cycling on CBF, as observed in traditional cycling.

Given that perceived effort is a significant barrier to engagement in and adherence to

rehabilitation programs, this study highlights the strong potential of implementing FES-

assisted cycling in stroke patients' rehabilitation programs.

Acknowledgements
385

386   We thank Jean-Serein Mbende for co-designing the illustration included in this article.
387

Conflict of interest declaration: The authors declare no conflict of interest.

Funding/ financial support: Inserm CAPS u1093, SMR Val Rosay UGECAM Rhône-Alpes,
Kurage

Contributors:

Conceptualization, M.D, J.DM, J.V.B & G.D; **methodology**, M.D, J.DM, J.V.B, B.P & G.D;

formal analysis, M.D; **investigation**, M.D, J.DM & PH.P; **data curation**, M.D; **writing—**

original draft preparation, M.D, B.P & G.D; **writing—review and editing**, M.D, J.DM,

E.J, J.V.B, B.P & G.D; **supervision**, M.D, J.DM, J.V.B & G.D; **figure designing**, M.D. &

B.P. All authors have read and agreed to the published version of the manuscript.

Declaration of Generative AI and AI-assisted technologies in the writing process: No

generative AI or AI-assisted technologies were used in the writing process, according to the

authors.

Ethical Clearance: The study was supported by UGECAM Rhône-Alpes and received ethical
approval from the Comité de Protection des Personnes Sud-Est II (Institutional Review
Board). The study was approved under the clinical trial registration number NCT06230796.

References

1. Cumming TB, Marshall RS, Lazar RM. Stroke, cognitive deficits, and rehabilitation:
Still an incomplete picture. Int J Stroke. 2013;8(1):38–45.

2. Kuppuswamy A, Clark E V., Turner IF, Rothwell JC, Ward NS. Post-stroke fatigue: a
deficit in corticomotor excitability? Brain. 2015;138(2014):136–48.

3. Pageaux B. Perception of effort in Exercise Science: Definition, measurement and
perspectives. Eur J Sport Sci. 2016;16(8):885–94.

4. Ambrosini E, Ferrante S, Ferrigno G, Molteni F, Pedrocchi A. Cycling induced by
electrical stimulation improves muscle activation and symmetry during pedaling in
hemiparetic patients. IEEE Trans Neural Syst Rehabil Eng. 2012;20(3):320–30.

5. Murillo Frazão, Luis Augusto Werlang, Peltz M. Metabolic, ventilatory and
cardiovascular responses to FES-cycling: A comparison to NMES and passive cycling.
Technol Heal care. 2021;

6. de Morree HM, Klein C, Marcora SM. Perception of effort reflects central motor
command during movement execution. Psychophysiology. 2012;49(9):1242–53.

7. Bergevin M, Steele J, Payen de la Garanderie M, Feral-Basin C, Marcora SM,
Rainville P, et al. Pharmacological Blockade of Muscle Afferents and Perception of
Effort: A Systematic Review with Meta-analysis. Vol. 53, Sports Medicine. 2023. 415–
435 p.

8. Khurana SR, Beranger AG, Felix ER. Perceived Exertion Is Lower When Using a
Functional Electrical Stimulation Neuroprosthesis Compared with an Ankle-Foot
Orthosis in Persons with Multiple Sclerosis: A Preliminary Study. Am J Phys Med
Rehabil. 2017;96(3):133–9.

9. Heller BW, Clarke AJ, Good TR, Healey TJ, Nair S, Pratt EJ, et al. Automated setup of
functional electrical stimulation for drop foot using a novel 64 channel prototype
stimulator and electrode array: Results from a gait-lab based study. Med Eng Phys
[Internet].   2013;35(1):74–81.   Available   from:
http://dx.doi.org/10.1016/j.medengphy.2012.03.012

10. Descollonges M, Chaney R, Garnier P, Pirgent-Tessier A, Brugniaux J V., Deley G.
Electrical stimulation : a potential alternative to positively impact cerebral health ?
Front Physiol. 2024;15:1–11.

11. Descollonges M, Marmier P, Marillier M, Jafari E, Brugniaux J V., Deley G. Effect of
electrical muscle stimulation on cerebrovascular function and cognitive performance.
Am J Physiol - Hear Circ Physiol. 2024;326(4):H923–8.

12. Halperin I, Emanuel A. Rating of Perceived Effort: Methodological Concerns and
Future Directions. Sport Med [Internet]. 2020;50(4):679–87. Available from:

https://doi.org/10.1007/s40279-019-01229-z

13.   Borg E, Kaijser L. A comparison between three rating scales for perceived exertion and
443   two different work tests. Scand J Med Sci Sport. 2006;16(1):57–69.

444   14.   Chaney R, Leger C, Wirtz J, Fontanier E, Méloux A, Quirié A, et al. Cerebral Benefits
445   Induced by Electrical Muscle Stimulation: Evidence from a Human and Rat Study. Int
446   J Mol Sci. 2024;25(3).

447   15.   Hashimoto T, Tsukamoto H, Ando S, Ogoh S. Effect of exercise on brain health: The
448   potential role of lactate as a myokine. Metabolites. 2021;11(12).

449   16.   Huang Z, Zhang Y, Zhou R, Yang L, Pan H. Lactate as Potential Mediators for
450   Exercise-Induced Positive Effects on Neuroplasticity and Cerebrovascular Plasticity.
451   Front Physiol. 2021;12(July).

452   17.   Freund H, Zouloumian P. Lactate After Exercise in Man: Evolution Kinetics in Arterial
453   Blood. Eur J Appl Physiol Occup Physiol. 1981;46(2):121–33.

454   18.   Watanabe K, Yoshimura A, Nojima H, Hirono T, Kunugi S, Takada T, et al.
455   Physiological adaptations following vigorous exercise and moderate exercise with
456   superimposed electrical stimulation. Eur J Appl Physiol [Internet]. 2023;123(1):159–
457   68. Available from: https://doi.org/10.1007/s00421-022-05065-4

458   19.   Pageaux B, Bergevin M, Mangin T, Lepers R, Marcora SM. The central motor
459   command, but not the muscle afferent feedback, is necessary to perceive effort. Int J
460   Psychophysiol   [Internet].   2023;188:29.   Available   from:
461   https://doi.org/10.1016/j.ijpsycho.2023.05.072

462   20.   Bergquist AJ, Clair JM, Lagerquist O, Mang CS, Okuma Y, Collins DF.
463   Neuromuscular electrical stimulation: Implications of the electrically evoked sensory
464   volley. Eur J Appl Physiol. 2011;111(10):2409–26.

465   21.   Cheval B, Boisgontier MP. The Theory of Effort Minimization in Physical Activity.
466   Exerc Sport Sci Rev. 2021;49(3):168–78.

467   22.   Cefis M, Chaney R, Wirtz J, Méloux A, Quirié A, Leger C, et al. Molecular
468   mechanisms underlying physical exercise-induced brain BDNF overproduction.
469   2023;(October):1–18.

470   23.   Willie CK, Colino FL, Bailey DM, Tzeng YC, Binsted G, Jones LW, et al. Utility of
471   transcranial Doppler ultrasound for the integrative assessment of cerebrovascular
472   function. J Neurosci Methods [Internet]. 2011;196(2):221–37. Available from:
473   http://dx.doi.org/10.1016/j.jneumeth.2011.01.011

474   24.   Lucas SJE, Ainslie PN, Murrell CJ, Thomas KN, Franz EA, Cotter JD. Effect of age on
475   exercise-induced alterations in cognitive executive function: Relationship to cerebral
476   perfusion.   Exp   Gerontol   [Internet].   2012;47(8):541–51.   Available   from:
477   http://dx.doi.org/10.1016/j.exger.2011.12.002

478   25.   Brugniaux J V., Marley CJ, Hodson DA, New KJ, Bailey DM. Acute exercise stress
479   reveals cerebrovascular benefits associated with moderate gains in cardiorespiratory
480   fitness. J Cereb Blood Flow Metab [Internet]. 2014;34(12):1873–6. Available from:
481   https://doi.org/10.1038/jcbfm.2014.142

482   26.   Mulser L, Moreau D. Effect of acute cardiovascular exercise on cerebral blood flow: A

483   systematic review. Brain Res. 2023;1809(November).

484   27.   Harrington AT, McRae CGA, Lee SCK. Evaluation of Functional Electrical
485   Stimulation to Assist Cycling in Four Adolescents with Spastic Cerebral Palsy. Int J
486   Pediatr. 2012;2012:1–11.

487   28.   Medrinal C, Combret Y, Prieur G, Robledo Quesada A, Bonnevie T, Gravier FE, et al.
488   Comparison of exercise intensity during four early rehabilitation techniques in sedated
489   and ventilated patients in ICU: A randomised cross-over trial. Crit Care. 2018;22(1):1–
490   8.

491   29.   Montalvo S, Gomez M, Lozano A, Arias S, Rodriguez L, Morales-Acuna F, et al.
492   Differences in Blood Flow Patterns and Endothelial Shear Stress at the Carotid Artery
493   Using Different Exercise Modalities and Intensities. Front Physiol. 2022;13(May):1–
494   10.

495   30.   Sudo M, Kitajima D, Takagi Y, Mochizuki K, Fujibayashi M, Costello JT, et al.
496   Effects of voluntary exercise and electrical muscle stimulation on reaction time in the
497   Go/No-Go task. Eur J Appl Physiol [Internet]. 2024;(0123456789). Available from:
498   https://doi.org/10.1007/s00421-024-05562-8

499   31.   Ando S, Ishioka Y, Kambayashi S, Kano K, Fujibayashi M, Costello JT, et al.
500   Combined effects of electrical muscle stimulation and cycling exercise on cognitive
501   performance. 2024;(May):1–7.

Tables and Figures

Table I. [Patient's characteristics. I: Ischemic; H: Hemorrhagic.]

Fig. 1. [Experimental set-up. Electrical stimulation was applied to the paretic leg, on the
quadriceps, hamstrings, and gluteus muscles.]

Fig. 2. [Perceived effort (A) and associated individual data (B) in response to FES-assisted

(black) and traditional (white) cycling. Values are mean (SD). *n*, sample size for the analysis.]

Fig. 3. [Middle cerebral artery velocity [MCAv (A)] and MCAv associated individual data

(B), in response to FES-assisted (black) and traditional (white) cycling. (C) MCAv at rest

(triangle), during exercise (rectangle), and associated individual data (white squares). Values

are mean (SD). *n*, sample size for the analysis.]

Fig. 4. [HR (A) and associated heart rate individual data (B), PETCO2 (C) and associated

PETCO2 individual data (D), and blood lactate (E) in response to FES-assisted (black) and

traditional (white) cycling. Data are presented as mean (SD). *n*, sample size for the analysis.]

Supplementary Material

SUPPLEMENTRY FIGURE LEGENDS*

Fig. S1. [Stimulation intensities (mA) used over the FES-assisted cycling modality.]

Fig. S2. [MAP (A) and associated individual data (B), CO (C) and associated individual data
(D) in response to FES-assisted (black) and traditional (white) cycling. Data are presented as
mean ± SE. *n*, sample size for the analysis.]

Fig. S3. [Cognitive performances in response to FES-assisted and traditional cycling. A)
Congruent Stroop task; B) Neutral Stroop Task; C) Incongruent Stroop Task; D) TMT-A and
E) TMT-B performances. PRE (triangle) and POST (square). Values are mean ± SE. n,
sample size for the analysis. *: significant difference pre-post within the same condition.]

SUPPLEMENTRY TABLE LEGENDS*

Table SI. [Voluntary patient feedback following completion of both cycling sessions.]