Original Article

Title: Positive effects of functional electrical stimulation-assisted cycling on perception of effort, cerebral blood flow and cognition in post-stroke patients

Short Title: Functional electrical stimulation-assisted cycling on stroke patients

Author Names: Maël Descollonges (PhD)[1,2,3#], Julie Di Marco (MD)[2,3#], Ehsan Jafari (PhD)[2,4], Pierre-Henri Pouillart (MSc)[3], Julien V. Brugniaux (PhD)[5], Benjamin Pageaux (PhD)[6,7,8] and Gaëlle Deley (PhD)[1]

Affiliations:

[1] INSERM UMR 1093 – Laboratoire CAPS, Université de Bourgogne, UFR des Sciences du Sport, Dijon, France

[2] Kurage, Lyon, France

[3] SMR Val Rosay, Saint-Didier au Mont d'Or, France

[4] Université de Lyon, ENS de Lyon, CNRS, Laboratoire de Physique, F-69342 Lyon, France

[5] HP2 laboratory, Univ. Grenoble Alpes, INSERM, CHU Grenoble Alpes, Grenoble, France

[6] Centre de recherche de l'Institut universitaire de gériatrie de Montréal (CRIUGM), Canada

[7] École de kinésiologie et des sciences de l'activité physique (EKSAP), Faculté de médecine, Université de Montréal, Montréal, Canada

[8] Centre Interdisciplinaire de Recherche sur le Cerveau et l'Apprentissage (CIRCA), Montréal, QC, Canada

[#] First co-authors.

Correspondence address: Maël Descollonges, 64b rue de sully, 21000 – Dijon, Inserm CAPS u1093 +33 6 79 64 57 39, descollonges.mael@gmail.com

Submission Summary

[Abstract word count: **237**/350]

[Background to conclusion word count: **3601**]

[Manuscript word count: **5500**]

[Number of Tables: **1**]

[Number of Figures: **4**]

[Number of References: **31**]

[Number of Supplementary Materials Submitted: **1** – **1** table, **3** figures]

____

Abstract

Background: FES-assisted cycling may reduce perceived effort by lowering the required motor

command compared to voluntary cycling. While benefits on effort perception have been shown

during walking in stroke and multiple sclerosis patients, its effectiveness during cycling in

stroke rehabilitation remains unproven. Thus, this work aimed to test the effect of functional

electrical stimulation-assisted cycling on stroke patients' perception of effort (primary aim) in

a randomized controlled study design. In an exploratory way, this work also aimed to examine

the effect of FES-assisted cycling on cerebral blood flow and cognitive performance

(exploratory aims) in a subsample.

Methods: Fifteen post-stroke patients completed functional electrical stimulation-assisted and

traditional cycling sessions separated by 72h. Perceived effort, cardio, and cerebrovascular

parameters were monitored during exercise. Cognitive performance was assessed before and

after each session. Qualitative data were reported after both sessions.

Results: Patients reported a lower perceived effort during functional electrical stimulation-

assisted cycling than traditional cycling. Both sessions increased heart rate, end-tidal CO_2,

cardiac output, and cerebral artery blood flow velocity, with higher blood lactate levels after

functional electrical stimulation-assisted cycling. Functional electrical stimulation-assisted

cycling positively impacted cognitive performance.

Conclusion: Traditional and functional electrical stimulation-assisted cycling induced similar

increased cardio and cerebrovascular responses. However, patients perceived functional

electrical stimulation-assisted cycling as less effortful than traditional cycling. As effort is a

barrier to regular exercise engagement and adherence, these results are promising for

implementing functional electrical stimulation-assisted cycling in stroke patients'

rehabilitation.

Key words: Cerebral blood flow, cognition, functional electrical stimulation, rating of

perceived exertion, stroke

Background

Stroke is a major cause of long-term disability affecting walking, postural control, muscle

strength, cerebrovascular and cardiovascular systems, and cognition. To counteract

cardiorespiratory system deconditioning and experience the positive effect of physical exercise

on cognition (1), stroke patients must engage in regular aerobic exercise during their

rehabilitation. Due to hemiparesis and fatigue leading to increased effort perception (2), and the

proposed key role of effort and its perception in the engagement in physical and cognitive

activities (3), post-stroke individuals experience difficulty engaging in and performing physical

exercise. Therefore, developing and implementing methods of rehabilitation that minimize the

perception of effort experienced during physical exercise is crucial.

To this aim, functional electrical stimulation (FES)-assisted cycling is a promising method.

Clinical evidence shows that FES-assisted cycling activates paralyzed muscles (4) and

improves aerobic capacity and cardiopulmonary function (5). By evoking muscle contraction

superimposed on top of the impaired voluntary contractions, FES is known to assist in

movement execution (4). As the evoked contraction induced by FES pre-activates the muscles,

the individual engaged in FES-assisted cycling must voluntarily produce a lower muscle force

to reach the targeted exercise intensity. As effort perception is proposed to be generated by

signals related to the magnitude of the motor command (6,7) and the lower muscle force to be

voluntarily produced during FES should be associated with a lower motor command compared

to traditional cycling, using FES-assisted cycling is a promising tool to decrease the level of

perceived effort experienced during rehabilitation sessions. To date, it has been reported that

FES application was associated with a reduced perception of effort in stroke or multiple

sclerosis patients during walking tasks (8,9). However, the effectiveness in reducing perceived

effort during cycling in stroke patients remains to be validated. As cycling is a key aerobic

exercise in rehabilitation, there is an urgent need for such a study.

In this context, this study aimed to compare the perceived effort during an FES-assisted cycling

session, compared to a traditional cycling session, in stroke patients. We hypothesized that

stroke patients would perceive a lower effort during FES-assisted cycling compared to

traditional cycling, performed at the same workload. In addition, considering recent research

suggesting a positive effect of electrical muscle stimulation on cerebral blood flow velocity

(CBFv, (10,11)) and potentially cognitive performance, we also explored, in a subsample of

patients, changes in CBFv to the FES- and traditional cycling exercises, and cognitive

performance pre-post exercises.

Methods
98

99   Fifteen post-stroke patients volunteered in this experiment (8♀; 7♂; age: 53 ± 13 years, height:

100 167 ± 0 cm, mass: 69 ± 10 kg, body mass index: 25 ± 3, 148 ± 108 days after stroke). Patients'

101 characteristics stratified by sex are available in Table 1.

103 **Table 1:** Patient's characteristics. I: Ischemic; H: Hemorrhagic.

	Females							
Patients	Age (years)	Size (cm)	Weight (kg)	BMI	Type of stroke	Days after stroke	Hemiparetic side	Assistive device
1	35-40	150	67	29.8	H	189	Left	AFO
2	40-45	169	69	24.2	I	40	Right	-
3	75-80	170	64	22.1	I	90	Right	AFO
4	50-55	155	63	26.1	I	90	Right	Cane
5	35-40	160	79	30.9	I	80	Right	Cane
6	35-40	170	74	25.6	H	237	Right	AFO
7	55-60	158	48	19.2	I	385	Left	AFO
8	65-70	163	68	25.5	I	213	Left	Wheelchair
Mean	51	160	66	25	-	166	-	-
	Males							
9	55-60	175	67	21.9	I	330	Left	AFO
10	65-70	165	66	24.2	I	38	Left	AFO
11	60-65	175	78	25.5	I	90	Left	-
12	35-40	162	74	28.1	H	97	Left	Cane
13	55-60	167	66	23.7	I	104	Right	-
14	45-50	186	91	26,4	H	207	Left	Wheelchair + cane
15	50-55	180	60	20.1	I	33	Right	-
Mean	55	173	72	24	-	128	-	-

Each patient provided written informed consent describing the procedure and potential risks of

the study. The sample size was based on patients' availability at the time of the experiment and

an expected large effect size of FES-assisted cycling on the perception of effort. Inclusion

criteria were diagnosis of an ischemic (n=11) or hemorrhagic (n=4) stroke resulting in

hemiparesis; adequate cognitive abilities to engage in rehabilitation activities; able to sit up to

30 minutes; joint mobility ranges that would not preclude pedaling. The study was supported

by UGECAM Rhône-Alpes and received ethical approval from the Comité de Protection des

Personnes Sud-Est II (French Institutional Review Board). The study was approved under the

clinical trial registration number NCT06230796.

Patients visited the hospital on six occasions. In visit 1, participants performed an incremental

test to determine their peak power output (PPO). The incremental test started at 25 watts, with

the power output increasing by 10 watts every minute until exhaustion. Exhaustion was defined

as voluntary disengagement from the task or an inability to maintain a cadence above 50

revolutions/minute (rpm) for more than 3 seconds, despite verbal encouragement. In visits 2-3-

4, patients were extensively familiarized with FES, the cognitive tasks, and the psychophysical

scale used to report their perceived effort. Each familiarization visit was separated by 48 hours.

In visits 5-6, participants performed two testing sessions in a randomized order: FES-assisted

cycling and traditional cycling. Testing sessions were separated by 72 hours to control for the

presence of muscle fatigue and soreness in the second testing session (visit 6).

During FES-assisted cycling, electrical stimulation was applied to the quadriceps, hamstrings,

and gluteus muscles of the hemiparesis side during cycling using a stimulator (Motimove, 3F-

Fabricando, Belgrade, Serbia) and an electrical stimulation-assisted cycle ergometer (Kurage,

HephaBike, Lyon, France) (Figure 1). The paretic leg was secured in alignment using an

orthotic pedal, ensuring stabilization of the foot and leg during the exercise (Figure 1). A

computerized program linked to an encoder fixed in the pedal axis was employed to stimulate

the muscles in the movement's proper phase. The appropriate phase corresponds to the knee

extension and knee flexion (pedaling movement). Quadriceps and gluteus muscles were

stimulated during extension, and hamstrings during flexion. The stimulation frequency was

fixed at 40 Hz, with a pulse width of 400 μs, and the stimulation intensity (mA) was constantly

adjusted to the individual maximum tolerable level throughout the session. The maximum

tolerable intensity of stimulation was determined by gradually increasing the current intensity

until an effective contraction was achieved and further increased to elicit the maximal pain

intensity tolerated by the participant. Each session began with a 5-minute warm-up consisting

of a progressive increase in stimulation intensity and cycling resistance to reach 50% PPO.

Afterwards, the resistance was fixed at 50% PPO for 20 minutes, and patients had to maintain

a cadence of 50 ± 3 rpm.

Figure 1: Experimental set-up. Electrical stimulation was applied to the paretic leg, on the quadriceps,

hamstrings, and gluteus muscles.

Measurements and data analysis

Perception of effort

The perception of effort is defined as "the conscious sensation of how hard, heavy, and

strenuous the physical task is" (3,12). Participants reported the intensity of their perceived effort

to cycle via the following question: ''How hard is it for you to drive your legs?". The intensity

of the perceived effort was reported every 5 minutes during the cycling exercise, using the

validated CR100 scale (13). The scale was presented to the patient on a laminated sheet,

allowing them to report the intensity of their perceived effort in real time. This scale ranges

from 0 for "nothing at all" to 100 for "maximal". The scale contains verbal anchors such as light

(13), moderate (23), strong (50), or very strong (70). The maximal intensity (100) was anchored

based on previous experience and defined as "the maximal effort experienced in your life, for

most people it corresponds to the most strenuous exercise previously experienced." Patients

were familiarized with the scale during the incremental test and the familiarization sessions

(visits 2-3-4).

Cardio- and cerebrovascular parameters

Heart rate (HR) was recorded continuously using a heart rate monitor (Forerunner 630, Garmin,

USA), mean arterial blood pressure (MAP = Diastolic Pressure + 1/3 (Systolic Pressure

– Diastolic Pressure) and cardiac output (CO) were monitored using finger

photoplethysmography attached to the non-dominant hand (Finometer PRO; Finapres Medical

Systems, Amsterdam, The Netherlands). The blood flow velocity in the right middle cerebral

artery (MCAv) was continuously monitored using a 2 MHz pulsed Doppler ultrasound system

(TCD, Multi-Dop X4, DWL Elektroniche System, Singen, Germany). Nine patients

volunteered for cerebral blood flow velocity monitoring. Each session began with confirmation

of artery insonation, relying on expected signal depth, velocity, visual signal along, and carotid

compression for validation. The Doppler probe was placed on the right transtemporal window

and secured with a DiaMon headband (DWL) throughout all experimental sessions. The

placement of the probe and the depth of the artery were carefully recorded to be the same for

each condition.

Beat-to-beat data were continuously sampled at 1 kHz using an analog-to-digital converter

(PowerLab/16SP ML795; AD Instruments, Oxford, UK) and stored for offline analysis on a

computer (Chart version 7.2.2, AD Instruments, Oxford, UK). Cardio and cerebrovascular data

were averaged for the last 30 seconds of each 5-minute interval period (before, during, and after

the exercise) as done in a previous study (11).

Cognitive tasks

Cognitive tasks were conducted on a table and in a seated position, both before (PRE) and

immediately after (POST) each experimental condition. The selected cognitive tasks included

a Stroop Task and the Trail Making Tests (TMT-A and TMT-B), as performed in previous

studies (11,14). Firstly, participants completed a three-phase Stroop task. These three phases

included three versions of the Stroop tasks, referred to as the congruent, neutral, and

incongruent conditions, respectively (11,14). Each task contained 100 items (words or crosses).

On the first page, the words "blue," "red," "yellow," and "green" were printed in black ink

(congruent task – reading task). On the second page, "XXXX" was printed in blue, red, yellow,

and green colors (neutral task). The last page displayed a color name written in a different font

color from the printed word, and the participant had to name the ink color (incongruent task).

Participants were instructed to read the words from left to right, aloud, as quickly and accurately

as possible for 45 seconds. The number of words and errors was counted by the experimenter

and recorded. Finally, these Stroop tasks were displayed on a 42-inch screen set to maximum

brightness, with a standard distance of 150 cm between participants and the screen.

Lastly, the Trail Making Test (TMT) was performed. This test is separated into two parts. The

Trail Making Test-A (TMT-A) consists of rapidly connecting a sequence of randomly placed

numbers in ascending order from 1 to 25. The Trail Making Test-B (TMT-B) requires

participants to alternately connect numbers and letters in sequence (e.g., 1 A, 2 B, 3 C, ... 13).

TMT performance was defined as the time (seconds) taken to complete the task. The test was

performed with the patient's unaffected hand due to spasticity in the affected hand.

Blood lactate

Given that blood lactate has been proposed as a potential key factor mediating the cognitive

benefits of physical exercise (15,16), blood lactate concentration was measured. Blood lactate

was collected using an automated analyzer (Lactate Pro, Scout 4), with a 5-µl blood sample

collected from the tip of the non-paretic index finger (due to spasticity in the affected hand), at

the beginning and 2 minutes after the end of exercise to assess peak concentration within an

optimal window (17).

Patients' experience with FES-assisted cycling – Qualitative data

Following both testing sessions, the researcher informed each patient that if they want, they are

free to report to their experience with FES-assisted cycling and traditional cycling. No

predefined questions or interview guide were used. With the patients' explicit consent, their

comments were manually and anonymously transcribed on paper, verbatim.

Statistical analysis

Mean (SD). Statistical significance was determined at $\alpha < 0.05$. The Shapiro–Wilk test

confirmed data normality. Two-way repeated-measures ANOVAs (condition × time) were

used. Post-hoc tests were adjusted for multiple comparisons using the Bonferroni correction.

Effect sizes are reported with the partial eta square (partial $\eta2$), with values of 0.01, 0.06, and

above 0.14 denoting small, medium, and large differences, respectively.

Results

The average peak power output was 72 ± 36 watts, and the cycling exercises were performed

at 36 ± 18 watts.

Changes in stimulation intensity during FES-assisted cycling, for each muscle, are available in

the supplemental materials (Figure S1).

The perceived effort to cycle increased over time (time: $P < 0.001$; $\eta_p^2 = 0.497$) and was lower

for FES-assisted cycling compared to traditional cycling (condition: $P < 0.001$; $\eta_p^2 = 0.800$)

(Figure 2).

Figure 2: Perceived effort (A) and associated individual data (B) in response to FES-assisted (black)

and traditional (white) cycling. Values are mean (SD). n, sample size for the analysis.

MCAv data is available for nine patients, as only these individuals volunteered for cerebral

blood flow velocity monitoring. MCAv increases during both sessions (time: $P = 0.026$; $\eta_p^2 =$

0.407), with no difference between conditions (condition: $P = 0.129$; $\eta_p^2 = 0.263$) (Figure 3).

Figure 3: Middle cerebral artery velocity [MCAv (A)] and MCAv associated individual data (B), in

response to FES-assisted (black) and traditional (white) cycling. (C) MCAv at rest (triangle), during

exercise (rectangle), and associated individual data (white squares). Values are mean (SD). *n*, sample

size for the analysis.

Due to technical issues relative to cable disconnection, $PETCO_2$ measurements were available

for only 12 patients, and mean arterial pressure (MAP) and cardiac output (CO) for 7 patients.

Blood lactate levels were available for 13 patients.

As shown in Figure 4, HR, mean arterial blood pressure (MAP), cardiac output (CO), $PETCO_2$,

and blood lactate increased during exercise (Figure 3, all $P < 0.001$ and $\eta_p^2 \geq 0.327$). Blood

lactate levels were higher in the FES-assisted cycling session compared to the traditional

cycling session ($P = 0.018$; $\eta_p^2 = 0.384$). CO and MAP are presented in supplemental

materials—Figure S2 and increased in both sessions (all $P \leq 0.001$ and $\eta_p^2 \geq 0.327$).

Figure 4: HR (A) and associated heart rate individual data (B), PETCO2 (C) and associated PETCO2 individual data (D), and blood lactate (E) in response to FES-assisted (black) and traditional (white) cycling. Data are presented as mean (SD). *n*, sample size for the analysis.

Due to fatigue or inability to complete the tasks, cognitive test data were available for 14 participants in the congruent Stroop condition, 13 in the neutral and incongruent conditions,

and 12 for the TMT-B test. Details of the cognitive performance data is available in

supplemental materials. Briefly, the results suggest that FES-assisted cycling induced similar

effects of traditional cycling, with a potential positive effect specific to the Neutral Stroop task.

Eleven patients freely reported their experience with FES-assisted cycling. The verbatim of the

experience reported is available in the supplemental materials—Table S1. The most recurring

theme among the responses was a perceived benefit of the FES-assisted session compared to

traditional cycling. Most of the participants reported that the FES made pedaling easier by

inducing additional force during cycling, which reduced fatigue, and by improving coordination

and fluidity of movement on the affected limb.

Discussion

The increased perceived effort over the 20 minutes of FES-assisted and traditional cycling is

caused by the progressive development of muscle fatigue, requiring an increased motor

command to recruit additional motor units, leading to higher perceived effort levels (3,6,7).

According to our hypothesis, when cycling at the same workload, the perceived effort was lower

during FES-assisted compared to traditional cycling. To the best of our knowledge, this study

is the first to observe in stroke patients the positive effect of FES-assisted cycling on the

perception of effort. This result replicates previous observations in healthy individuals (18,19)

and conceptually replicates previous observations of a lower perceived effort during a FES-

walking exercise in multiple sclerosis (8) or stroke patients (9). It is well accepted that applying

electrical stimulation to a muscle increases afferent feedback (20). Our results, combined with

the studies observing a lower perception of effort during FES of the lower limb (i.e., cycling or

walking), provide further evidence that the perception of effort scales with the magnitude of the

motor command rather than afferent feedback (for review see (7)). However, the mechanisms

underlying the FES-induced reduction in perceived effort remain to be fully understood and

warrant further investigation. Brain imaging techniques would be of interest to better elucidate

the relationship between the motor command and perceived effort during physical exercise

involving electrical stimulation.

In our study, the lower perceived effort is likely linked to the force evoked by FES, which

helped the patients by requiring only a small additional voluntary engagement to cycle. This

interpretation is further supported by the patients' experience qualitatively reported. Indeed,

most of the patients reported perceiving clear assistance of the stimulation, helping the

movement and facilitating the exercise. A limitation to this observation is that our study was

not a-priori designed for a qualitative analysis. As our qualitative analysis to describe the

patients experience with FES-assisted cycling was exploratory, we encourage future research

to replicate our results by adopting a-priori a mixed-methods design. Doing so will allow a

better interpretation on the relationship between quantitative measurement of effort and its

perception, and the patients' perspectives and experience. Moreover, we were unable to directly

assess the actual contribution of FES to muscle force production and to measure the force output

produced by the stimulator relative to the ergometer resistance. Thus, we cannot precisely

quantify the extent to which FES contributes to force generation at different intensities. In the

future, using force sensors attached directly to the pedals would allow for a more accurate

assessment of this parameter, providing a better understanding of the role of FES in the

perception of effort, in force generation, and its effective contribution to the cycling motion.

As a high perceived effort is a barrier to engagement in physical exercise and a key determinant

of adherence to rehabilitation programs (3,21), the observed low perceived effort is of crucial

interest for supporting the development and implementation of FES-assisted cycling in stroke

patients' rehabilitation. Future studies should now implement FES-assisted cycling during a

rehabilitation program, and not only a single training session, to test the possibility that the

effects of FES-assisted cycling on stroke patients' rehabilitation could be superior to those

induced by traditional cycling.

As cerebral blood flow has been proposed as one of the mechanisms underlying neuroplasticity

(22), we also explored the CBF responses to FES-assisted cycling and its potential priming

effect on cognitive performance in a subsample of voluntary patients. As previously observed

during an isometric NMES session (11), our results suggest that FES-assisted cycling can

increase CBF in the same proportions as cycling with lower perceived effort. These increases

in blood velocity in the middle cerebral artery are not surprising, since similar results have been

observed on pedaling tasks (23–26) and could be explained by the increases in cardiovascular

parameters observed during exercise. As the literature has reported that the recruitment of a

large muscle volume through electrical stimulation leads to a higher cardiorespiratory demand

(5,27,28), we hypothesized that CBF would be higher in the FES condition. However, it was

not the case in our study, and this could potentially be explained by the muscle volume

stimulated in our FES condition. It is well established that the muscle mass engaged during

physical exercise is a key factor influencing cerebral blood flow (29). Although a substantial

muscle volume of paretic leg was stimulated in this study (quadriceps, hamstrings, and gluteal

muscles), future research could explore the effects of stimulating an even greater number of

muscles, including the entire lower limb (e.g., quadriceps, hamstrings, gluteus, tibialis anterior,

etc.), rather than focusing solely on the paretic limb. Such an approach could enhance the effects

of stimulation on CBF. However, further research is needed to optimize stimulation parameters

and muscle selection, ensuring that potential adverse effects are avoided. One limitation of this

study is that CBF was measured only on the right MCA, without distinguishing between the

affected (ipsilateral) and unaffected (contralateral) sides. Thus, future research should consider

measuring bilateral cerebral blood flow at the middle cerebral artery level to better understand

the differences between the affected and unaffected sides.

Our results also suggest that cycling, whether assisted by FES or not, has limited and

inconsistent effects on cognitive performance in stroke patients. Although both types of cycling

sessions led to some improvements in cognitive performance, these effects were not

consistently observed across all cognitive tasks (e.g., the incongruent condition of the Stroop

task). These partial effects—improvements seen only in specific tasks—are consistent with the

existing literature reporting mixed findings (10). For instance, recent studies have shown that

an acute NMES protocol (11,14) can enhance cognitive performance, but only under certain

conditions of the Stroop task in healthy individuals. Conversely, other studies have found no

improvement in cognitive performance, particularly in tasks such as the Go/No-Go, among

healthy participants (30,31). To address this discrepancy in the literature, we encourage future

research to focus on the effect of FES-assisted cycling on cognitive performance as primary

outcome of interest and design the study a-priori for this purpose.

Moreover, as shown in the present study, where elevated blood lactate levels were observed

following FES, recent literature has reported that NMES applied to the quadriceps increases

blood lactate levels in both humans and rats (14). Notably, higher blood lactate levels were

associated with improved Stroop Task performance in healthy humans and increased

hippocampal BDNF expression in rats following a single NMES session (14). These findings

support the hypothesis that lactate may act as a key exerkine mediating the cognitive benefits

of electrical stimulation. However, the underlying mechanisms remain poorly understood and

warrant further investigation.

Lastly, we would like to remind the reader that our sample size was determined by our primary

aim, i.e., testing the difference in perceived effort between FES-assisted cycling and traditional

cycling. Due to the likely smallest effect size of FES-assisted cycling on CBF and cognitive

performance, future studies should replicate our results with a specific a priori approach to CBF

and cognitive performance and a higher sample size. Such studies could also unravel a

potentially greater positive effect of FES-assisted cycling compared to traditional cycling, on

CBF and cognitive performance.

To conclude, this study is the first to demonstrate a reduced perceived effort during FES-

assisted cycling, compared to traditional cycling, in stroke patients. This study also suggests a

potential positive effect of FES-assisted cycling on CBF, as observed in traditional cycling.

Given that perceived effort is a significant barrier to engagement in and adherence to

rehabilitation programs, this study highlights the strong potential of implementing FES-assisted

cycling in stroke patients' rehabilitation programs.

Declaration

Ethics approval and consent to participate: The study was supported by UGECAM Rhône-

Alpes and received ethical approval from the Comité de Protection des Personnes Sud-Est II

(French Institutional Review Board). The study was approved under the clinical trial

registration number NCT06230796 and registered at ClinicalTrials.gov. In addition, written

informed consent was obtained from all subjects. Patients were assured that their information

would only be used for analysis in this study. All methods were carried out under relevant

guidelines and regulations.

Consent for publication: Not applicable as all participants have been de-identified.

Availability of data and materials: The data are available from the corresponding author on

reasonable request.

Competing interests: The authors declare no competing interests.

Funding: Funds for the realization of this work were partly provided by Laboratory Inserm

CAPS u1093, SMR Val Rosay UGECAM Rhône-Alpes, and Kurage.

Authors' contributions: Conceptualization, M.D, J.DM, J.V.B. & G.D; **methodology**, M.D,

J.DM, J.V.B, B.P & G.D; **formal analysis**, M.D; **investigation**, M.D, J.DM & PH.P.; **data**

curation, M.D; **writing—original draft preparation**, M.D, B.P & G.D; **writing—review and**

editing, M.D, J.DM, E.J, J.V.B, B.P & G.D; **supervision**, M.D, J.DM, J.V.B & G.D; **figure**

designing, M.D. & B.P. All authors have read and agreed to the published version of the

manuscript.

Acknowledgements

We thank Jean-Serein Mbende for co-designing the illustration included in this article.

References
397

1. Cumming TB, Marshall RS, Lazar RM. Stroke, cognitive deficits, and rehabilitation:
Still an incomplete picture. Int J Stroke. 2013;8(1):38–45.

2. Kuppuswamy A, Clark E V., Turner IF, Rothwell JC, Ward NS. Post-stroke fatigue: a
deficit in corticomotor excitability? Brain. 2015;138(2014):136–48.

3. Pageaux B. Perception of effort in Exercise Science: Definition, measurement and
perspectives. Eur J Sport Sci. 2016;16(8):885–94.

4. Ambrosini E, Ferrante S, Ferrigno G, Molteni F, Pedrocchi A. Cycling induced by
electrical stimulation improves muscle activation and symmetry during pedaling in
hemiparetic patients. IEEE Trans Neural Syst Rehabil Eng. 2012;20(3):320–30.

5. Murillo Frazão, Luis Augusto Werlang, Peltz M. Metabolic, ventilatory and
cardiovascular responses to FES-cycling: A comparison to NMES and passive cycling.
Technol Heal care. 2021;

6. de Morree HM, Klein C, Marcora SM. Perception of effort reflects central motor
command during movement execution. Psychophysiology. 2012;49(9):1242–53.

7. Bergevin M, Steele J, Payen de la Garanderie M, Feral-Basin C, Marcora SM, Rainville
P, et al. Pharmacological Blockade of Muscle Afferents and Perception of Effort: A
Systematic Review with Meta-analysis. Vol. 53, Sports Medicine. 2023. 415–435 p.

8. Khurana SR, Beranger AG, Felix ER. Perceived Exertion Is Lower When Using a
Functional Electrical Stimulation Neuroprosthesis Compared with an Ankle-Foot
Orthosis in Persons with Multiple Sclerosis: A Preliminary Study. Am J Phys Med
Rehabil. 2017;96(3):133–9.

9. Heller BW, Clarke AJ, Good TR, Healey TJ, Nair S, Pratt EJ, et al. Automated setup of
functional electrical stimulation for drop foot using a novel 64 channel prototype
stimulator and electrode array: Results from a gait-lab based study. Med Eng Phys
[Internet].   2013;35(1):74–81.   Available   from:

423    http://dx.doi.org/10.1016/j.medengphy.2012.03.012

424 10. Descollonges M, Chaney R, Garnier P, Pirgent-Tessier A, Brugniaux J V., Deley G.
425    Electrical stimulation : a potential alternative to positively impact cerebral health ? Front
426    Physiol. 2024;15:1–11.

427 11. Descollonges M, Marmier P, Marillier M, Jafari E, Brugniaux J V., Deley G. Effect of
428    electrical muscle stimulation on cerebrovascular function and cognitive performance.
429    Am J Physiol - Hear Circ Physiol. 2024;326(4):H923–8.

430 12. Halperin I, Emanuel A. Rating of Perceived Effort: Methodological Concerns and Future
431    Directions. Sport Med [Internet]. 2020;50(4):679–87. Available from:
432    https://doi.org/10.1007/s40279-019-01229-z

433 13. Borg E, Kaijser L. A comparison between three rating scales for perceived exertion and
434    two different work tests. Scand J Med Sci Sport. 2006;16(1):57–69.

435 14. Chaney R, Leger C, Wirtz J, Fontanier E, Méloux A, Quirié A, et al. Cerebral Benefits
436    Induced by Electrical Muscle Stimulation: Evidence from a Human and Rat Study. Int J
437    Mol Sci. 2024;25(3).

438 15. Hashimoto T, Tsukamoto H, Ando S, Ogoh S. Effect of exercise on brain health: The
439    potential role of lactate as a myokine. Metabolites. 2021;11(12).

440 16. Huang Z, Zhang Y, Zhou R, Yang L, Pan H. Lactate as Potential Mediators for Exercise-
441    Induced Positive Effects on Neuroplasticity and Cerebrovascular Plasticity. Front
442    Physiol. 2021;12(July).

443 17. Freund H, Zouloumian P. Lactate After Exercise in Man: Evolution Kinetics in Arterial
444    Blood. Eur J Appl Physiol Occup Physiol. 1981;46(2):121–33.

445 18. Watanabe K, Yoshimura A, Nojima H, Hirono T, Kunugi S, Takada T, et al.
446    Physiological adaptations following vigorous exercise and moderate exercise with
447    superimposed electrical stimulation. Eur J Appl Physiol [Internet]. 2023;123(1):159–68.
448    Available from: https://doi.org/10.1007/s00421-022-05065-4

449 19. Pageaux B, Bergevin M, Mangin T, Lepers R, Marcora SM. The central motor command,
450    but not the muscle afferent feedback, is necessary to perceive effort. Int J Psychophysiol
451    [Internet]. 2023;188:29. Available from: https://doi.org/10.1016/j.ijpsycho.2023.05.072

452 20. Bergquist AJ, Clair JM, Lagerquist O, Mang CS, Okuma Y, Collins DF. Neuromuscular
453    electrical stimulation: Implications of the electrically evoked sensory volley. Eur J Appl
454    Physiol. 2011;111(10):2409–26.

455 21. Cheval B, Boisgontier MP. The Theory of Effort Minimization in Physical Activity.
456    Exerc Sport Sci Rev. 2021;49(3):168–78.

457 22. Cefis M, Chaney R, Wirtz J, Méloux A, Quirié A, Leger C, et al. Molecular mechanisms
458    underlying physical exercise-induced brain BDNF overproduction. 2023;(October):1–
459    18.

460 23. Willie CK, Colino FL, Bailey DM, Tzeng YC, Binsted G, Jones LW, et al. Utility of
461    transcranial Doppler ultrasound for the integrative assessment of cerebrovascular
462    function. J Neurosci Methods [Internet]. 2011;196(2):221–37. Available from:
463    http://dx.doi.org/10.1016/j.jneumeth.2011.01.011

464 24. Lucas SJE, Ainslie PN, Murrell CJ, Thomas KN, Franz EA, Cotter JD. Effect of age on

exercise-induced alterations in cognitive executive function: Relationship to cerebral perfusion. Exp Gerontol [Internet]. 2012;47(8):541–51. Available from: http://dx.doi.org/10.1016/j.exger.2011.12.002

25. Brugniaux J V., Marley CJ, Hodson DA, New KJ, Bailey DM. Acute exercise stress reveals cerebrovascular benefits associated with moderate gains in cardiorespiratory fitness. J Cereb Blood Flow Metab [Internet]. 2014;34(12):1873–6. Available from: https://doi.org/10.1038/jcbfm.2014.142

26. Mulser L, Moreau D. Effect of acute cardiovascular exercise on cerebral blood flow: A systematic review. Brain Res. 2023;1809(November).

27. Harrington AT, McRae CGA, Lee SCK. Evaluation of Functional Electrical Stimulation to Assist Cycling in Four Adolescents with Spastic Cerebral Palsy. Int J Pediatr. 2012;2012:1–11.

28. Medrinal C, Combret Y, Prieur G, Robledo Quesada A, Bonnevie T, Gravier FE, et al. Comparison of exercise intensity during four early rehabilitation techniques in sedated and ventilated patients in ICU: A randomised cross-over trial. Crit Care. 2018;22(1):1–8.

29. Montalvo S, Gomez M, Lozano A, Arias S, Rodriguez L, Morales-Acuna F, et al. Differences in Blood Flow Patterns and Endothelial Shear Stress at the Carotid Artery Using Different Exercise Modalities and Intensities. Front Physiol. 2022;13(May):1–10.

30. Sudo M, Kitajima D, Takagi Y, Mochizuki K, Fujibayashi M, Costello JT, et al. Effects of voluntary exercise and electrical muscle stimulation on reaction time in the Go/No-Go task. Eur J Appl Physiol [Internet]. 2024;(0123456789). Available from: https://doi.org/10.1007/s00421-024-05562-8

31. Ando S, Ishioka Y, Kambayashi S, Kano K, Fujibayashi M, Costello JT, et al. Combined effects of electrical muscle stimulation and cycling exercise on cognitive performance. 2024;(May):1–7.

492 **Tables and Figures**
493

Table I. [Patient's characteristics. I: Ischemic; H: Hemorrhagic.]

Fig. 1. [Experimental set-up. Electrical stimulation was applied to the paretic leg, on the
quadriceps, hamstrings, and gluteus muscles.]

Fig. 2. [Perceived effort (A) and associated individual data (B) in response to FES-assisted

(black) and traditional (white) cycling. Values are mean (SD). n, sample size for the analysis.]

Fig. 3. [Middle cerebral artery velocity [MCAv (A)] and MCAv associated individual data (B),

in response to FES-assisted (black) and traditional (white) cycling. (C) MCAv at rest (triangle),

during exercise (rectangle), and associated individual data (white squares). Values are mean

(SD). n, sample size for the analysis.]

Fig. 4. [HR (A) and associated heart rate individual data (B), PETCO2 (C) and associated

PETCO2 individual data (D), and blood lactate (E) in response to FES-assisted (black) and

traditional (white) cycling. Data are presented as mean (SD). n, sample size for the analysis.]

Supplementary Material/Additional file
507

SUPPLEMENTRY FIGURE LEGENDS*

Fig. S1. [Stimulation intensities (mA) used over the FES-assisted cycling modality.]

Fig. S2. [MAP (A) and associated individual data (B), CO (C) and associated individual data
(D) in response to FES-assisted (black) and traditional (white) cycling. Data are presented as
mean ± SE. n, sample size for the analysis.]

Fig. S3. [Cognitive performances in response to FES-assisted and traditional cycling. A)
Congruent Stroop task; B) Neutral Stroop Task; C) Incongruent Stroop Task; D) TMT-A and
E) TMT-B performances. PRE (triangle) and POST (square). Values are mean ± SE. n, sample
size for the analysis. *: significant difference pre-post within the same condition.]

SUPPLEMENTRY TABLE LEGENDS*

Table SI. [Voluntary patient feedback following completion of both cycling sessions.]

A — Heart rate (bpm) vs Time (min): Condition (P = 0.093; ηp² = 0.188); Time (P < .001; ηp² = 0.816); Interaction (P = 0.018; ηp² = 0.205); N = 15

B — Heart rate vs Time (min): FES-assisted cycling; Traditional cycling

C — PETCO₂ (mmHg) vs Time (min): Condition (P = 0.719; ηp² = 0.012); Time (P < .001; ηp² = 0.568); Interaction (P = 0.056; ηp² = 0.165); n = 12

D — PETCO₂ vs Time (min)

E — Blood lactate (mmol/L): PRE; POST; Condition (P = 0.018; ηp² = 0.384); Time (P < .001; ηp² = 0.791); Interaction (P = 0.218; ηp² = 0.124); n = 13

1