Cohort profile update: HUNT4 70+

Skjellegrind, Håvard Kjesbu[1,2]; Thingstad, Pernille[3,4]; Gjøra, Linda[5,6]; Kolberg, Marit[7]; Kjelvik, Grete[6]; Ernstsen, Linda[8,9]; Fagerhaug, Tone Natland[8]; Langhammer, Arnulf[1,2]; Krokstad, Steinar[1,2]; Åsvold, Bjørn Olav[1,2]; Næss, Marit[1,2]; Selbæk, Geir[6,11,12]

HUNT Research Centre, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Levanger, Norway.
Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway.
Department of Neuromedicine and Movement Science, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway.
Department of Health and Welfare, Trondheim Municipality, Trondheim, Norway.
Department of Psychiatry, Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway.
Norwegian National Centre for Ageing and Health, Vestfold Hospital Trust, Tønsberg, Norway.
Center for Oral Health Services and Research Mid-Norway (TkMidt), Trondheim, Norway.
Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway.
Clinic of Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway.
HUNT Center for Molecular and Clinical Epidemiology, Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), Trondheim, Norway.
Department of Geriatric Medicine, Oslo University Hospital, Oslo, Norway.
Institute for Clinical Medicine, University of Oslo, Oslo, Norway.

Word count: 2482

Abstract

The Trøndelag Health Study (HUNT) in Norway has surveyed lifestyle and health in more than 50% of the local adult population every decade since 1984-86. HUNT4 70+ is a sub-cohort of persons aged at least 70 years at the 4[th] survey of the HUNT Study, established to provide data for ageing research. This population-based sample consists of 9956 individuals from the original HUNT catchment area, included between August 2017 - February 2019. Of these, 9589 have been included in one or several of the former surveys. In addition, an urban sample of 1743 persons was included in Trondheim city in the period October 2018 – June 2019. HUNT4 70+ covers comprehensive aspects of ageing health, including clinical examinations, performance-based tests of physical and cognitive function, questionnaires, and biological samples. High participation rates among the old and frail were obtained by examination in private homes and nursing homes when needed (15% of the participants). The data can be linked to all national registers in Norway, such as cause of death, prescription, health care utilization, and diagnosis registries.

Why was the cohort set up

The HUNT4 70+ cohort was set up to survey physical performance, cognitive function, nutritional status, and oral health in a population-based sample of older adults without exclusion criteria, as part of the Trøndelag Health Study (The HUNT Study). The need for high-quality research data from older people in an aging population was an important background. Other population-based studies have reported disease and symptom burden among older adults attending examination sites, but few studies have reported the continuum from the healthy to those in need of home care or nursing homes. The survey was designed to collect age-related endpoints as a follow-up of the main HUNT cohort and as a baseline for the study of trajectories through old age.

Previously, dementia has been found in 83% of nursing home residents in the HUNT catchment area (1). A large-scale survey of cognitive function in community-dwelling older adults was not performed in Norway before. Malnutrition is widespread among older adults (2) and is strongly associated with morbidity and mortality (3, 4). The extent of malnutrition in the older adult population, including both community-dwelling older adults and nursing home residents was unknown. Poor oral health has been reported among nursing home residents (5). Still, there was limited data on oral health at the population-level of older adults including both nursing home residents, home-care recipients, and healthy older adults. Frailty awareness has increased over the last few decades, after seminal publications (6, 7). Still, data on population-wide prevalences were scarce. Thus, the HUNT4 70+ cohort was also designed to allow prevalence assessment of dementia, frailty, malnutrition, and oral health problems, as well as the extent of comorbidities of chronic diseases.

The survey was designed in cooperation with the HUNT ethics committee and approved by the Norwegian Data Protection Authority. Collection and storage of these data are regulated by the "Regulation on population-based health surveys". Participants were included based on informed consent. Individuals unable to consent due to cognitive impairment were allowed participation after informed consent by proxy. All were assumed able to refuse participation. The study protocol was approved by the regional research ethics committee before piloting and commencing the study (REK 2016/1880).

Who is in the cohort?

All adult inhabitants in the HUNT catchment area have been invited to the HUNT surveys. Out of 19 403 invited older adults aged ≥70 years, 12 489 (64.4 %) participated in HUNT4 (8) and 9956 (51.3 %) were included in the HUNT4 70+ cohort described here. To mitigate selection bias, examination at home was offered for those not able to attend a field station. Home visits were available to all, by request. Individuals receiving community-based care were given help by their nurses to book a home visit from a HUNT representative. All nursing home residents were invited to participate in an examination in their nursing home. By performing the examinations in several settings, the survey was able to include more older adults and more persons with age-related health conditions (Table 1).

Furthermore, 5168 persons aged 70+ living in one of four administrative city districts of Trondheim municipality were invited, and 1743 were included (described separately in the supplement, as these were recruited outside the original HUNT catchment area).

Participant profile

To assess representativity of the sample, the point prevalences of common chronic diagnoses were retrieved for both the participant sample and the total invited population, from The Norwegian Registry of Primary Health Care. The relative proportions of diagnosed cases for women and men in 5-year age groups are included in supplementary tables 5 and 6 and illustrated by a heatmap in Figure 1. No systematic difference between the population and the included sample was found. The range of relative proportions spanned from 0.39 to 2.04.

As the sociodemographic composition of the local and included population may differ from the national population, we have established and published a set of inverse probability weights (IPWs) for participation rate grouped on age, sex and education (9). IPWs can improve generalisability and precision of prevalence estimates of conditions that are unevenly distributed among these factors, such as dementia, malnutrition and frailty (10-12).

How often have they been followed up?

The first HUNT survey (HUNT1) was conducted 1984-86, including anthropometric measurements, pulse and blood pressure, as well as questionnaires focusing on hypertension, diabetes, and

influence by disease on quality of life, as well as x-ray screening for tuberculosis (13). HUNT1 included 89.4% of the adult population. HUNT2 (1995-97) included extended questionnaires, blood samples, and clinical measurements as spirometry, bone densitometry, and hearing. The participation rate was 71.1% (14). The HUNT3 survey was performed 2006-2008, including 54.1% of the local adult population (15). HUNT4 was performed in 2017-2019 with participation rate of 54.0 % in the original catchment area. All attending participants aged 70+ were invited to HUNT4 70+ in addition to persons in need of home care or living in nursing homes (8). Out of the 9956 included participants, 9589 were included in one or several of the former surveys (Figure 2).

The HUNT4 70+ cohort was later invited to a 4-year follow-up, as a combined effort between the HUNT Ageing in Trøndelag (AiT) and the HUNT COVID projects. Totally, 5729 individuals from the HUNT4 70+ cohort participated, with questionnaires, clinical assessments and blood samples. The blood samples have been analysed for COVID antibodies. Biomarkers for Alzheimer's disease (AD) (phosphorylated Tau-217) and axonal damage (Neurofilament Light (NfL)) have been analysed in plasma samples from both HUNT3, HUNT4 and HUNT COVID/AiT.

Moreover, the HUNT4 70+ cohort can be followed up by linking data to the comprehensive national registers for cause of death, dispensed drugs, diagnosis-specific registries, and diagnoses from health care services both in primary care and secondary care.

What has been measured?

HUNT4 basic measures

All participants, including those who were examined in their homes or in nursing homes, went through the general HUNT4 examinations as previously described (8). In brief, blood pressure (BP), pulse, and peripheral oxygen saturation (SpO$_2$) were measured three times after a 5-minute rest with 1-minute intervals using an automated BP monitor (GE Carescape V100). At the field stations, body composition, weight, and height were measured using bioelectrical impedance (InBody 770, Cerritos, CA, USA). At examinations in private homes and in nursing homes, weight and height were measured using Seca 813 scale and Seca 217 stadiometer (Seca, Germany). Weight and height were measured with light clothes and without shoes. Blood samples were centrifuged within ninety minutes and transported cooled to HUNT biobank for further inhouse processing within 36 hours (16). Other biomaterials have been collected for parts of the HUNT4 70+ cohort: Urine (44%), saliva (31%), and dried fecal samples (for microbiome) (31%).

Cognitive function

The Montreal Cognitive Assessment (MoCA) assesses executive function, visuospatial abilities, memory, language, attention, abstraction, and orientation, and gives a score ranging 0-30 (17). The Norwegian translation version 7.1 was used. No additional point was given for low educational attainment.

The CERAD Word List Memory Task (WLMT) tests immediate and delayed recall, which are typically impaired in AD (18). The task is also sensitive to mild cognitive impairment (MCI) (19). The WLMT consists of three immediate recall trials of a ten-word list, and then a delayed recall trial. The delayed recall was performed after examination of balance, gait speed, and strength, approximately 5-10 minutes after the immediate recall trials. The WLMT was not applied in cases of MoCA score <22 or zero out of five points in the memory domain of the MoCA.

In nursing homes, for subjects with established moderate to severe cognitive impairment, Severe Impairment Battery-8 (20) was used to assess cognitive function, instead of the MoCA.

The participants were interviewed about subjective cognitive impairments, the development pattern of cognitive impairment, function in activities of daily living, neuropsychiatric symptoms, former examinations on cognitive function, and family history of cognitive impairments.

In nursing homes, a caregiver provided information on symptoms and daily function. Neuropsychiatric symptoms were assessed using the nursing home version of the Neuropsychiatric Inventory (NPI-NH)(21, 22). Activities of daily living (ADL) were assessed using the Physical Self-Maintenance Scale (PSMS)(23).

Proxy interview

Participants were asked for permission to contact a next of kin for supplementary information on their cognitive health if at least one of these criteria were met:
- MoCA score below a threshold (70-80 years: <22/30; 80-90 years: <21; 90+: <20)
- WLMT delayed recall score below threshold (70-80 years: <4/10 words; 80-90 years: <3; 90+: <2)
- Self-reported subjective cognitive impairment: Answering yes to EITHER "Has your memory become significantly worse in the last five years?", OR "Have other cognitive functions (like orientation ability or language) become worse in the last five years?", AND "Do these changes worry you?".

When possible, the proxy was chosen by the participant and was in most cases the spouse or a daughter/son. For participants with consent by proxy, that next of kin was interviewed. All interviews were conducted by telephone and included the following scales:
- Neuropsychiatric Inventory Questionnaire (NPI-Q) (24)
- Instrumental activities in daily living (I-ADL) and physical ADL (P-ADL) (23)
The interview also included questions on symptom debut, symptom development, dementia diagnosis, use of municipal and informal care and family history of dementia.

Clinical dementia rating (CDR)

CDR was scored by the examiner in cooperation with the nursing staff in nursing homes and based on the proxy interviews (25).

Dementia and MCI diagnosis

All available information from questionnaires, interviews, cognitive tests and proxy interviews, were used to establish consensus-based diagnoses of MCI, dementia and dementia subtypes, applying DSM-5 diagnostic criteria (American Psychiatric Association, 12, 26).

Physical function

The Short Physical Performance Battery (SPPB) is a performance-based test of physical function including three timed sub-tests: 4-meter gait speed, five times sit-to-stand at maximum speed and three standing balance positions (27). Each subtest is scored 1-4 points or zero points if the person is unable to perform the subtask. The Norwegian translation of the original test protocol and score sheets were used.

One-leg balance was performed both with open and closed eyes, as a timed test for balancing on the preferred one leg and registered as 0-30 seconds or not able. The time was stopped when moving the standing leg or touching down the elevated leg.

Grip strength was measured by using the Jamar Plus+ Digital Hand Dynamometer® (Performance Health, Warrenwille, USA). Three trials were performed consecutively without resting pauses for each hand with the person in sitting position, the arm hanging free by the body, the shoulder adducted and neutrally rotated, the elbow held in 90° and the wrist in neutral position.

24-hour movement behaviour

Two AX3 accelerometers (Axivity, Newcastle, UK), one placed on the thigh and one on the lower back were worn for seven days (28). Validated machine learning models (ML) have been used to predict the amount of time of different types of physical activity (29) and duration of sleep (30).

Nutritional assessment

In addition to height and weight, the mid-upper arm circumference (MUAC) was measured for nutritional assessment. MUAC was measured twice on the right upper arm at the midpoint between acromion and olecranon. Interview and questionnaire questions included unintentional weight loss,

food intake, loss of appetite, problems with chewing food and oral pain, enabling evaluation of malnutrition (3).

Oral health

The Revised Oral Assessment Guide (ROAG) is a standardized assessment tool for nursing staff to examine and detect oral health problems in older adults (31). ROAG-Jönköping (ROAG-J) is a modified ROAG and includes assessment of voice, swallowing, lips, oral mucosa, tongue, saliva, gums, teeth and dentures. Additionally, information about saliva under the tongue, number of teeth, mobility of teeth, presence of root remnants and dentures were registered. The lower end of a toothbrush was used to test buccal mucosal surface sliding friction. In the main cohort, ROAG-J was performed at nursing homes and home visits only. In HUNT4 Trondheim 70+, ROAG-J was performed for all participants.

The number of included participants with data on different measures are shown in table 2. The distribution of selected key variables are shown in figure 3.

What has it found?

The HUNT4 70+ cohort was used to estimate current and future prevalence of dementia and MCI in Norway (12). Standardized dementia prevalence ranged from 5.6% at 70-74 years, increasing to 48.1% at 90+ years. MCI was found in 33-36% in all ages ≥70. Moreover, we found that only 35.6 % of those with a dementia condition were diagnosed in the health care system (32). The cohort has been used to establish norms for the MoCA (33) and the WLMT (34). We have also described the variation of physical performance with age and education (35). A HUNT Frailty Index (HUNT-FI) has been constructed and, together with the Fried phenotype, used to establish the prevalence of frailty in the Norwegian 70+ population (36). Physical performance was found proportionally impaired with cognitive impairment, also with MCI (37). The prevalence of malnutrition in the population aged 70+ was estimated to 14.3 % (10).

The cohort is well suited to study risk factors for dementia. Using BP from all HUNT surveys, we have found that, on average, BP increases with increasing age in those that maintain cognitive function, while in those developing dementia, average BP declined in advanced age. Still, ten years before dementia diagnosis, average BP was the same. (38). A combination of a low BP trajectory and high physical activity was associated with the lowest dementia risk (39). A study on hearing loss identified in HUNT2 (1995-97) found this associated to dementia risk 22 years later (40).

An overview of published papers is found in the supplement.

What are the main strengths and weaknesses?

The HUNT4 70+ survey comprises a comprehensive dataset covering important aspects of ageing health for a representative sample, including the oldest and frailest part of the population. In total, 51.4 % of the entire population in the original HUNT catchment area was included. This, combined with having both clinical data and biological samples for most participants over several decades is a major strength. In addition, the data may be linked to a wide range of national and local health registries, including cause of death, cancer, primary and secondary care diagnoses, socioeconomic indicators, drug prescriptions, and health care utilization. The very long follow-up time of 33 years from HUNT1 and 22 years from the first blood samples (HUNT2) allows studies with low risk of reverse causation, and the comprehensiveness allows analyses with low levels of residual confounding.

Missing questionnaire data is a potential weakness, especially for individuals with severe cognitive impairments. However, some data, such as education, may be found in questionnaires from previous HUNT surveys. For risk assessment studies, the competing risk of death may be a weakness. Cause of death data may be acquired through the Norwegian Cause of Death register. The participation rate

of approximately 50% may be considered low for some purposes. We find that using weighed analyses compensates this well (11).

Can I get hold of the data? Where can I find out more?

The HUNT4 70+ dataset was collected to provide the public with more extensive epidemiological research on ageing and age-related health. Thus, we invite researchers to apply to HUNT Research Centre to obtain data for projects. Data access requires approval from a Norwegian Research Ethics Committee (REC) following an application submitted by a researcher affiliated with a Norwegian research institution. We advise to contact Håvard Skjellegrind (havard.skjellegrind@ntnu.no) or HUNT Research Centre (kontakt@hunt.ntnu.no) to establish relevant collaborations. A complete catalogue of available variables is found at ntnu.edu/hunt.

Ethics approval

The data collection in the HUNT Study is legally regulated and not subject to external approval requirements from data security, privacy or research ethics authorities. This cohort profile article was approved by the Mid-Norway Regional Committee for Medical and Health Research Ethics (REK 69134).

Acknowledgements

The HUNT4 70+ survey has received funding from the Norwegian National Centre for Ageing and Health and Center for Oral Health Services and Research (TkMidt), Trondheim, in addition to funding of HUNT4 by the Ministry of Health and Care Services, Central Norway Regional Health Authority, Trøndelag county council and NTNU. All municipalities have contributed with staff and field station offices. ROAG was used in HUNT4 70+ with permission from Pia Andersson. The ROAG-J and related guidance manual was used as part of the training of the examiners with permission by Eva Herremo and Charlott Karlsson, Folktandvården, Region Jönköpings län and Senior Alert, Qulturum, Region Jönköpings län. ROAG-J was translated to Norwegian by Center for Oral Health Services and Research (TkMidt) to be used in HUNT4 70+. Data from the Norwegian Registry for Primary Health Care has been used in this publication. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Dep. of Health Registries is intended nor should be inferred.

All data analysis and visualization in this article was performed using R version 4.4.1 "Race for Your Life" in RStudio Version 2024.04.2+764 "Chocolate Cosmos". Tables were made using the table1 and flextable packages, and graphics made using ggplot2.

Conflicts of interest

GS has participated in Advisory Board meetings for Roche, Eli-Lilly and Eisai regarding disease-modifying drugs for Alzheimer's disease. GS has received honoraria for delivering lectures at symposia sponsored by Eisai and Eli-Lilly.

References

1. Bergh S, Holmen J, Gabin J, Stordal E, Fikseaunet A, Selbaek G, et al. Cohort profile: the Health and Memory Study (HMS): a dementia cohort linked to the HUNT study in Norway. Int J Epidemiol. 2014;43(6):1759-68.
2. Leij-Halfwerk S, Verwijs MH, van Houdt S, Borkent JW, Guaitoli PR, Pelgrim T, et al. Prevalence of protein-energy malnutrition risk in European older adults in community, residential and hospital settings, according to 22 malnutrition screening tools validated for use in adults >/=65 years: A systematic review and meta-analysis. Maturitas. 2019;126:80-9.

3. Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. Clin Nutr. 2019;38(1):1-9.
4. Soderstrom L, Rosenblad A, Thors Adolfsson E, Bergkvist L. Malnutrition is associated with increased mortality in older adults regardless of the cause of death. Br J Nutr. 2017;117(4):532-40.
5. Bellander L, Andersson P, Nordvall D, Hägglin C. Oral health among older adults in nursing homes: A survey in a national quality register, the Senior Alert. Nurs Open. 2021;8(3):1262-74.
6. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. J Gerontol A Biol Sci Med Sci. 2001;56(3):M146-56.
7. Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. ScientificWorldJournal. 2001;1:323-36.
8. Åsvold BO, Langhammer A, Rehn TA, Kjelvik G, Grontvedt TV, Sorgjerd EP, et al. Cohort Profile Update: The HUNT Study, Norway. Int J Epidemiol. 2022.
9. Skirbekk V, Bowen CE, Haberg A, Jugessur A, Engdahl B, Bratsberg B, et al. Marital Histories and Associations With Later-Life Dementia and Mild Cognitive Impairment Risk in the HUNT4 70+ Study in Norway. J Aging Health. 2023;35(7-8):543-55.
10. Kolberg M, Paur I, Sun Y-Q, Gjøra L, Skjellegrind HK, Thingstad P, et al. Prevalence of malnutrition among older adults in a population-based study - The HUNT Study. Clinical Nutrition ESPEN. 2023.
11. Gjøra L, Strand BH, Engedal K, Ernstsen L, Myrstad C, Skjellegrind H, et al. Dementia and mild cognitive impairment in older people in Trøndelag. Tidsskr Nor Laegeforen. 2023;143(10).
12. Gjøra L, Strand BH, Bergh S, Borza T, Braekhus A, Engedal K, et al. Current and Future Prevalence Estimates of Mild Cognitive Impairment, Dementia, and Its Subtypes in a Population-Based Sample of People 70 Years and Older in Norway: The HUNT Study. J Alzheimers Dis. 2021;79(3):1213-26.
13. Holmen J, Midthjell K, Forsen L, Skjerve K, Gorseth M, Oseland A. [A health survey in Nord-Trondelag 1984-86. Participation and comparison of attendants and non-attendants]. Tidsskr Nor Laegeforen. 1990;110(15):1973-7.
14. Holmen J, Midthjell K, Krüger Ø, Langhammer A, Holmen TL, Bratberg GH, et al. The Nord-Trøndelag Health Study 1995-97 (HUNT 2). Norsk Epidemiologi. 2003;13(1).
15. Krokstad S, Langhammer A, Hveem K, Holmen TL, Midthjell K, Stene TR, et al. Cohort Profile: the HUNT Study, Norway. Int J Epidemiol. 2013;42(4):968-77.
16. Næss M, Kvaløy K, Sørgjerd EP, Sætermo KS, Norøy L, Røstad AH, et al. Data Resource Profile: The HUNT Biobank. Int J Epidemiol. 2024;53(3).
17. Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. J Am Geriatr Soc. 2005;53(4):695-9.
18. Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillenbaum G, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer's disease. Neurology. 1989;39(9):1159-65.
19. Shankle WR, Romney AK, Hara J, Fortier D, Dick MB, Chen JM, et al. Methods to improve the detection of mild cognitive impairment. Proc Natl Acad Sci U S A. 2005;102(13):4919-24.
20. Schmitt FA, Saxton J, Ferris SH, Mackell J, Sun Y. Evaluation of an 8-item Severe Impairment Battery (SIB-8) vs. the full SIB in moderate to severe Alzheimer's disease patients participating in a donepezil study. Int J Clin Pract. 2013;67(10):1050-6.
21. Cummings JL, Mega M, Gray K, Rosenberg-Thompson S, Carusi DA, Gornbein J. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. Neurology. 1994;44(12):2308-14.
22. Wood S, Cummings JL, Hsu MA, Barclay T, Wheatley MV, Yarema KT, et al. The use of the neuropsychiatric inventory in nursing home residents. Characterization and measurement. Am J Geriatr Psychiatry. 2000;8(1):75-83.
23. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. Gerontologist. 1969;9(3):179-86.
24. Kaufer DI, Cummings JL, Ketchel P, Smith V, MacMillan A, Shelley T, et al. Validation of the NPI-Q, a brief clinical form of the Neuropsychiatric Inventory. J Neuropsychiatry Clin Neurosci. 2000;12(2):233-9.

25. Hughes CP, Berg L, Danziger WL, Coben LA, Martin RL. A new clinical scale for the staging of dementia. Br J Psychiatry. 1982;140:566-72.
26. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-5®): American Psychiatric Publishing; 2013.
27. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. J Gerontol. 1994;49(2):M85-94.
28. Kongsvold A, Flaaten M, Logacjov A, Skarpsno ES, Bach K, Nilsen TIL, et al. Can the bias of self-reported sitting time be corrected? A statistical model validation study based on data from 23 993 adults in the Norwegian HUNT study. Int J Behav Nutr Phys Act. 2023;20(1):139.
29. Logacjov A, Bach K, Kongsvold A, Bårdstu HB, Mork PJ. HARTH: A Human Activity Recognition Dataset for Machine Learning. Sensors. 2021;21(23):7853.
30. Logacjov A, Skarpsno ES, Kongsvold A, Bach K, Mork PJ. A Machine Learning Model for Predicting Sleep and Wakefulness Based on Accelerometry, Skin Temperature and Contextual Information. Nat Sci Sleep. 2024;16:699-710.
31. Andersson P, Hallberg IR, Renvert S. Inter-rater reliability of an oral assessment guide for elderly patients residing in a rehabilitation ward. Spec Care Dentist. 2002;22(5):181-6.
32. Gjøra L, Strand BH, Bergh S, Bosnes I, Johannessen A, Livingston G, et al. Prevalence and Determinants of Diagnosed Dementia: A Registry Linkage Study Linking Diagnosis of Dementia in the Population-Based HUNT Study to Registry Diagnosis of Dementia in Primary Care and Hospitals in Norway. Journal of Alzheimer's Disease. 2024;99:363-75.
33. Engedal K, Gjøra L, Benth J, Wagle J, Rønqvist TK, Selbæk G. The Montreal Cognitive Assessment: Normative Data from a Large, Population-Based Sample of Cognitive Healthy Older Adults in Norway-The HUNT Study. J Alzheimers Dis. 2022.
34. Wagle J, Selbæk G, Benth J, Gjøra L, Rønqvist TK, Bekkhus-Wetterberg P, et al. The CERAD Word List Memory Test: Normative Data Based on a Norwegian Population-Based Sample of Healthy Older Adults 70 Years and Above. The HUNT Study. J Alzheimers Dis. 2022.
35. Melsæter KN, Tangen GG, Skjellegrind HK, Vereijken B, Strand BH, Thingstad P. Physical performance in older age by sex and educational level: the HUNT Study. BMC Geriatrics. 2022;22(1):821.
36. Kyrdalen IL, Strand BH, Selbæk G, Thingstad P, Ormstad H, Hoogendijk EO, et al. Prevalence and future estimates of frailty and pre-frailty in a population-based sample of people 70 years and older in Norway: the HUNT study. Aging Clin Exp Res. 2024;36(1):188.
37. Sverdrup K, Selbæk G, Bergh S, Strand BH, Thingstad P, Skjellegrind HK, et al. Physical performance across the cognitive spectrum and between dementia subtypes in a population-based sample of older adults: The HUNT study. Archives of Gerontology and Geriatrics. 2021;95:104400.
38. Selbaek G, Stuebs J, Engedal K, Hachinski V, Hestad K, Trevino CS, et al. Blood pressure trajectories over 35 years and dementia risk: A retrospective study: The HUNT study. Frontiers in Aging Neuroscience. 2022;14.
39. Lerfald M, Lydersen S, Zotcheva E, Nilsen TIL, Eldholm RS, Martinez-Velilla N, et al. Change in physical activity and systolic blood pressure trajectories throughout mid-life and the development of dementia in older age: the HUNT study. Eur Rev Aging Phys Act. 2023;20(1):18.
40. Myrstad C, Engdahl BL, Costafreda SG, Krokstad S, Lin F, Livingston G, et al. Hearing impairment and risk of dementia in The HUNT Study (HUNT4 70+): a Norwegian cohort study. eClinicalMedicine. 2023;66:102319.

Table 1 Participant characteristics

	Field station (N=8522)	Home visit (N=803)	Nursing home (N=631)	Overall (N=9956)
Sex				
Female	4454 (52.3%)	545 (67.9%)	421 (66.7%)	5420 (54.4%)
Male	4068 (47.7%)	258 (32.1%)	210 (33.3%)	4536 (45.6%)
Age stratum				
70-79	6429 (75.4%)	162 (20.2%)	121 (19.2%)	6712 (67.4%)
80-89	1929 (22.6%)	385 (47.9%)	265 (42.0%)	2579 (25.9%)
90+	164 (1.9%)	256 (31.9%)	245 (38.8%)	665 (6.7%)
Education				
Primary	2355 (27.6%)	468 (58.3%)	350 (55.5%)	3173 (31.9%)
Secondary	3871 (45.4%)	197 (24.5%)	149 (23.6%)	4217 (42.4%)
College/University	2261 (26.5%)	57 (7.1%)	44 (7.0%)	2362 (23.7%)
Smoking Status				
Never	3204 (37.6%)	214 (26.7%)	130 (20.6%)	3548 (35.6%)
Former	4270 (50.1%)	235 (29.3%)	148 (23.5%)	4653 (46.7%)
Current	534 (6.3%)	32 (4.0%)	12 (1.9%)	578 (5.8%)
Occasional	28 (0.3%)	0 (0%)	0 (0%)	28 (0.3%)
Former Occasional	384 (4.5%)	23 (2.9%)	12 (1.9%)	419 (4.2%)
ApoE ε4 carrier				
No	5960 (69.9%)	560 (69.7%)	355 (56.3%)	6875 (69.1%)
Yes	2501 (29.3%)	223 (27.8%)	246 (39.0%)	2970 (29.8%)
BMI (kg/m^2)	27.2 (4.3)	27.1 (5.4)	26.1 (4.8)	27.2 (4.4)
Body fat (%)	33.6 (8.5)			33.6 (8.5)
MUAC (cm)	30.1 (3.5)	28.5 (4.2)	27.9 (4.0)	29.9 (3.6)
Systolic BP (mmHg)	140.2 (19.8)	135.7 (22.7)	130.4 (22.2)	139.4 (20.3)
Grip Strength (kg)	33.7 (10.8)	21.3 (7.8)	18.1 (8.7)	31.9 (11.5)
SPPB	11 [9, 12]	4 [2, 7]	1 [0, 3]	11 [8, 12]
ROAG-J[a]				
Normal	-	167 (20.8%)	81 (12.8%)	-
≥1 Moderate Problem	-	371 (46.2%)	245 (38.8%)	-
≥1 Severe Problem	-	161 (20.0%)	100 (15.8%)	-
MoCA	24 [21, 26]	17 [12, 21]	13 [9, 17]	24 [20, 26]
CERAD 10-word recall	5 [4, 6]	4 [2, 6]		5 [4, 6]
Dementia assessment				
No cognitive impairment	4698 (55.1%)	114 (14.2%)	10 (1.6%)	4822 (48.4%)
aMCI	2658 (31.2%)	211 (26.3%)	51 (8.1%)	2920 (29.3%)
naMCI	438 (5.1%)	37 (4.6%)	22 (3.5%)	497 (5.0%)
Dementia	618 (7.3%)	390 (48.6%)	523 (82.9%)	1531 (15.4%)
Unclassified	110 (1.3%)	51 (6.4%)	25 (4.0%)	186 (1.9%)

Continuous variables are presented as mean (SD), and discrete scores as median [IQR]. Percentages are of total N in column. Missing is not shown. BMI: Body Mass Index, MUAC: Middle Upper Arm Circumference, SPPB: Short Physical Performance Battery, ROAG-J: Revised Oral Assessment Guide – Jönköping, CERAD: Consortium to Establish a Registry for Alzheimer's Disease. aMCI: amnestic mild cognitive impairment (MCI), naMCI: non-amnestic MCI. [a]ROAG-J was only assessed at home visits (n=699) and nursing homes (n=426).

Table 2: Data availability

Data	Field station (N=8522)	Home visit (N=803)	Nursing home (N=631)	Total (N=9956)
Questionnaire				
Education	8487 (99.6 %)	722 (89.9 %)	543 (86.1 %)	9752 (98 %)
Smoking status	8420 (98.8 %)	504 (62.8 %)	302 (47.9 %)	9226 (92.7 %)
Weight loss	7805 (91.6 %)	433 (53.9 %)	209 (33.1 %)	8447 (84.8 %)
Exhaustion	7147 (83.9 %)	455 (56.7 %)	224 (35.5 %)	7826 (78.6 %)
HADS	7002 (82.2 %)	389 (48.4 %)	163 (25.8 %)	7554 (75.9 %)
Clinical examinations				
BMI	8458 (99.2 %)	688 (85.7 %)	312 (49.4 %)	9458 (95 %)
Body Composition	7565 (88.8 %)	-	-	7565 (76 %)
MUAC	7364 (86.4 %)	791 (98.5 %)	527 (83.5 %)	8682 (87.2 %)
Systolic BP	8492 (99.6 %)	775 (96.5 %)	524 (83 %)	9791 (98.3 %)
Gait Speed	8206 (96.3 %)	625 (77.8 %)	311 (49.3 %)	9142 (91.8 %)
Grip Strength	8091 (94.9 %)	765 (95.3 %)	481 (76.2 %)	9337 (93.8 %)
SPPB	7874 (92.4 %)	703 (87.5 %)	490 (77.7 %)	9067 (91.1 %)
1-week Accelerometer recording	4512 (52.9 %)	244 (30.4 %)	107 (17 %)	4863 (48.8 %)
ROAG-J	-	699 (87 %)	426 (67.5 %)	1125 (11.3 %)
MoCA	8116 (95.2 %)	683 (85.1 %)	109 (17.3 %)	8908 (89.5 %)
CERAD 10-word recall	5338 (62.6 %)	101 (12.6 %)	-	5439 (54.6 %)
SIB-8	-	-	356 (56.4 %)	356 (3.6 %)
Dementia assessment	8417 (98.8 %)	752 (93.6 %)	606 (96 %)	9775 (98.2 %)
Lab data				
Cholesterol	8433 (99 %)	637 (79.3 %)	469 (74.3 %)	9539 (95.8 %)
pTau-217	8030 (94.2 %)	531 (66.1 %)	387 (61.3 %)	8948 (89.9 %)
NfL	7888 (92.6 %)	518 (64.5 %)	383 (60.7 %)	8789 (88.3 %)
Genetics	8501 (99.8 %)	785 (97.8 %)	603 (95.6 %)	9889 (99.3 %)
Biosamples				
Serum	8394 (98.5 %)	618 (77 %)	456 (72.3 %)	9468 (95.1 %)
Plasma	8128 (95.4 %)	538 (67 %)	397 (62.9 %)	9063 (91 %)
Urine	4251 (49.9 %)	104 (13 %)	36 (5.7 %)	4391 (44.1 %)
Faeces	3016 (35.4 %)	26 (3.2 %)	9 (1.4 %)	3051 (30.6 %)
Saliva	2939 (34.5 %)	85 (10.6 %)	20 (3.2 %)	3044 (30.6 %)
HUNT3 Serum	7432 (87.2 %)	553 (68.9 %)	400 (63.4 %)	8385 (84.2 %)
HUNT3 Plasma	6593 (77.4 %)	532 (66.3 %)	381 (60.4 %)	7506 (75.4 %)
HUNT2 Serum	7487 (87.9 %)	704 (87.7 %)	529 (83.8 %)	8720 (87.6 %)

HADS: Hospital Anxiety and Depression Scale, BMI: Body Mass Index, MUAC: Middle Upper Arm Circumference, SPPB: Short Physical Performance Battery, ROAG-J: Revised Oral Assessment Guide – Jönköping, MoCA: Montreal Cognitive Assessment, CERAD: Consortium to Establish a Registry for Alzheimer's Disease. SIB-8: Severe Impairment Battery-8. pTau-217: plasma phosphorylated Tau-217, NfL, plasma neurofilament light.

Figure 1. Relative proportions of diagnosis frequency in participants compared to the invited population. Groups with less than 5 cases among participants are missing.

Figure 2. HUNT participation leading up to HUNT4 70+

Figure 3. Distribution of key variables. A Systolic blood pressure (mmHg), B Body mass index (kg/m2), C Montreal Cognitive Assessment, D Gait speed (m/s), E Short Physical Performance Battery, F Grip Strength (kg).