Title: **Metabolic modeling reveals determinants of prebiotic and probiotic treatment efficacy**
across two human intervention trials.

Authors: Nick Quinn-Bohmann[1], Alex V. Carr[1], and Sean M. Gibbons[1-5,*]

Affiliations: [1] Institute for Systems Biology, Seattle, WA 98109, USA; [2] Department of
Bioengineering, University of Washington, Seattle, WA 98195, USA; [3] Department of Genome
Sciences, University of Washington, Seattle, WA 98195, USA; [4] eScience Institute, University of
Washington, Seattle, WA 98195, USA; [5] Department of Environmental and Occupational Health
Sciences, University of Washington, Seattle, WA 98195, USA
[*] correspondence addressed to sgibbons@isbscience.org

Abstract

Prebiotic, probiotic, and combined (synbiotic) interventions often show variable outcomes
across individuals, driven by complex interactions between introduced biotics, the endogenous
microbiota, and the host diet. Predicting individual-specific success or failure of probiotic and
prebiotic therapies remains a major challenge. Here, we leverage microbial community-scale
metabolic models (MCMMs) to forecast probiotic engraftment and microbiota-mediated short-
chain fatty acid (SCFA) production in response to probiotic and prebiotic interventions. Using
data from two clinical trials, testing a five-strain probiotic combined with the prebiotic inulin
designed to improve metabolic health and an eight-strain probiotic designed to treat recurrent
Clostridioides difficile infections, respectively, we show that MCMM-predicted engraftment
largely agrees with measurements, achieving 75-80% accuracy. Engraftment probabilities varied
across taxa, with *Akkermansia muciniphila* and *Bifidobacterium infantis* showing higher
predicted and observed engraftment than the clostridial strains. MCMMs captured treatment-
driven shifts in predicted SCFA production, and higher model-predicted growth rates of *A.
muciniphila* were negatively associated with glucose AUC in the synbiotic trial, providing clues
about the mechanisms underlying treatment efficacy. Finally, extending these models to a
longitudinal cohort undergoing a healthy diet and lifestyle intervention revealed substantial
inter-individual variability in predicted responses to increasing dietary fiber, which were
significantly associated with baseline-to-follow-up changes in cardiometabolic health markers.
Finally, our simulation results suggested that personalized prebiotic selection may further
enhance probiotic efficacy. Together, these findings demonstrate the potential of metabolic
modeling to guide personalized microbiome-mediated interventions.

Introduction

The human colonic microbiota plays a critical role in regulating host physiology and immunity [1–5]. Probiotic, prebiotic, and synbiotic (i.e., combination of prebiotics and probiotics) administration have been shown to influence the metabolic outputs of the endogenous microbiota, inhibit pathogen colonization, and enhance mucosal barrier integrity [6,7]. Emerging evidence suggests that synbiotic interventions may even impact brain function via the gut-brain axis [8–10].

Despite these promising findings, the efficacy of probiotic, prebiotic, and synbiotic interventions can vary widely between individuals, posing a major challenge for their widespread clinical application [11,12]. Engraftment of a probiotic strain—the ability of the administered microbe to grow and persist within the gut ecosystem—depends on multiple factors, including the availability of a suitable metabolic niche, competitive and cooperative interactions with endogenous microbiota, and interactions with the host immune system [13,14]. Beyond the composition of the commensal microbiota, the interplay between factors like total commensal biomass, propagule pressure (i.e., the probiotic dose), available prebiotic substrates, and host diet can impact engraftment [15]. In order to rationally predict probiotic engraftment and prebiotic effects across individuals and dietary backgrounds, we require methods that integrate this complexity.

Genome-scale metabolic models (GEMs) provide a powerful mechanistic framework for estimating microbial growth and metabolism [16,17]. Recent advances in constraint-based modeling have extended this approach to diverse microbial communities, yielding microbial community-scale metabolic models (MCMMs) [18]. By integrating genome-scale metabolic reconstructions from hundreds of gut bacterial taxa with flux balance analysis, MCMMs enable predictions of microbe-microbe interactions, competition for resources, and ecosystem-scale metabolic behaviors [18,19]. These models have demonstrated predictive accuracy in estimating microbial growth rates, short-chain fatty acid production, methane production, and metabolic shifts in response to disease status or dietary interventions [18–22]. Notably, prior work has shown that MCMMs can capture the engraftment potential of opportunistic pathogens, such as *Clostridioides difficile*, suggesting that similar approaches could be used to model probiotic engraftment in individual hosts [19,23].

To validate this approach, we aimed to use MCMMs to predict the enrichment of probiotic species based on data from two human intervention trials. First, we leveraged data from a study conducted by Perradeau et al. (Validation Study A) [24]. In this double-blind, placebo-controlled trial, participants previously diagnosed with type 2 diabetes mellitus were treated for 12 weeks with either a placebo or one of two multi-strain synbiotic cocktail (WBF-010 and WBF-011). WBF-010 contained *Bifidobacterium infantis*, *Clostridium beijerinckii*, and *Clostridium butyricum*, supplemented with a low-dose of inulin (0.3g), a prebiotic fiber known to support probiotic growth. WBF-011 contained the same prebiotic and probiotic contents of

WBF-010, but with the addition of two probiotic strains: *Akkermansia muciniphila* and

Anaerobutyricum hallii. Participants who received the WBF-011 synbiotic exhibited a

significantly greater reduction in the glucose area under the curve (AUC) during a standard

glucose tolerance test from Week 0 to Week 12, compared to the placebo group, indicating a

beneficial population-level effect on glycemic control [24]. For this analysis, only data from

participants in the WBF-011 treatment group were used, given that WBF-010 did not show a

significant clinical effect on the primary outcome variables. In the second study (Validation

Study B) by Dsouza et al., participants were given an 8-strain probiotic cocktail (VE303)

designed to treat recurrent *Clostridioides difficile* infections (rCDI) [12]. VE303 consisted of

Enterocloster bolteae, Anaerotruncus colihominis, Sellimonas intestinalis, Clostridium

symbiosum, Blautia sp001304935*, Dorea longicatena, Longicatena innocuum,* and

Flavonifractor plautii, though *Blautia* sp001304935 and *Longicatena innocuum* were omitted

from the current analysis due to a lack of available GEMs in the AGORA database [25].

Using metagenomic sequencing data, quantitative PCR (qPCR), and immune and

metabolic profiling, we assessed the ability of MCMMs to accurately predict probiotic

engraftment and the subsequent impact on host metabolism and immune function. Specifically,

we evaluated how well MCMM-derived predictions align with observed shifts in microbial

composition and clinical markers. Finally, we expand MCMM simulations to a cross-sectional

cohort of generally-healthy individuals from the Arivale cohort who received a healthy diet and

lifestyle intervention, where they increased their exercise and intake of dietary fiber. MCMM-

predicted increases in butyrate production after switching from a standard European to a high

fiber diet were significantly associated with longitudinal changes in cardiometabolic health

markers in this intervention cohort. Using a subset of this Arivale cohort, we explored

personalized prebiotic, dietary, and probiotic interventions optimized to increase butyrate or

propionate production. Together, these results provide a promising computational platform for

designing personalized prebiotic, probiotic, and synbiotic intervention for human cohorts.

Results

Validation Study A: qPCR shows enrichment of probiotic species in the treatment group

Microbial abundances for focal species were determined using qPCR (see methods).

A. muciniphila and *B. infantis* exhibited significant enrichment from Week 0 to Week 12 across

most participants (Fig. 1A; paired t-test, *A. muciniphila*: p = 3.2×10^{-4}; 1-sample t-test, *B.*

infantis: p = 4.8×10^{-6}). In contrast, *C. beijerinckii, A. hallii,* and *C. butyricum* showed no

comparable enrichment. These trends suggest short-term engraftment of *A. muciniphila* and *B.*

infantis in the majority of treated individuals. Binarized engraftment scores were defined based

on qPCR amplification thresholds (see Methods).

Validation Study A: MCMMs predict species-level enrichment
Models were constructed from baseline taxonomic assignments (N = 21) to assess whether
MCMMs could predict the engraftment patterns observed in the qPCR results (Fig. 1A; see
methods). For each individual, models simulating the addition of 5 probiotic species were used
to predict engraftment patterns observed in Week 12 metagenomic sequencing (see methods)
Across all comparisons (5 predictions for each of 21 samples, total N = 105), model-
predicted growth and binarized enrichment scores showed 84.7% agreement (Fig. 2A, Cohen's κ
= 0.68, indicating substantial agreement). *C. beijerinckii* was correctly predicted as not growing
for all 21 participants, and *A. muciniphila* was correctly predicted as growing in all 21
participants. *B. infantis* had the lowest accuracy, with incorrect prediction in 8 of 21 cases.
Overall, there was significant concordance between predictions and observations (Fig. 2B).
Across all strains, 58 instances of growth were correctly predicted, 31 instances of non-growth
were correctly predicted, 3 cases were incorrectly predicted as growth, and 13 cases were
incorrectly predicted as non-growth (Fisher's exact test, p = 4.4×10^{-13}).
123
Prediction of Microbial Short-Chain Fatty Acid Production
To evaluate the predicted functional impact of probiotic administration, we analyzed the
model-predicted production of butyrate and propionate, two key short-chain fatty acids with
well-documented metabolic health-promoting, anti-inflammatory, and gut barrier-
strengthening properties [26]. Prior work has demonstrated the accuracy of MCMM-based
predictions for SCFA production in the human gut microbiome [19], providing a strong rationale
for employing this approach to assess the mechanistic contributions of both the probiotic and
prebiotic components of the WBF-011.
To systematically dissect the influence of probiotic and prebiotic interventions on SCFA
production, we compared the production of propionate and butyrate between the baseline
condition and the synbiotic condition. Due to the original study using a minimal prebiotic dose
(0.3 grams of inulin), we also simulated the addition of a larger prebiotic dose (30 grams of
inulin). Following simulation, total MCMM-predicted butyrate and propionate production rates
were compared across conditions (Fig. 3A-B).
As expected, the no treatment group exhibited relatively low levels of butyrate
production (Fig. 3A, No Treatment: 14.25 ± 2.27 mmol/gDW/h). The addition of the probiotic
cocktail with a minimal prebiotic did not significantly increase the level of butyrate produced by
the community (Fig. 3A, WBF-011 + Inulin, 0.3g: 14.81 ± 3.72 mmol/gDW/h). In a simulated
synbiotic intervention with a larger dose of inulin, a significant increase was observed (Fig. 3A,
WBF-011 + Inulin, 30g: 23.61 ± 2.99 mmol/gDW/h, Mann-Whitney U test, p = 6.1×10^{-3}).
Similar results were observed for propionate. The no treatment, and probiotic + minimal
prebiotic group showed low levels of propionate production (Fig. 3B, No Treatment: 55.92 ±
5.72 mmol/gDW/h; WBF-011 + Inulin, 0.3g: 45.24 ± 3.30mmol/gDW/h). The higher dose

prebiotic combination significantly increased the production of propionate (Fig. 3B, WBF-011 +

Inulin, 30g: 114.78 ± 5.66 mmol/gDW/h, Mann-Whitney U test, p = 9.3 x 10^{-7}).

Glucose AUC Related to Predicted Akkermansia muciniphila Growth Rate

In Perraudeau et al. [27], glucose AUC was a primary metabolic endpoint. A significant

difference in the change in glucose AUC from Week 0 to Week 12 (ΔglucoseAUC) was observed

between participants receiving WBF-011 from those receiving the placebo (Fig. 3C; Mann–

Whitney U test, p = 5.0×10^{-3}). This effect was not observed for the WBF-010 formulation,

which lacked *A. muciniphila* and *A. hallii*. Combined with the absence of a significant WBF-011–

induced change in MCMM-predicted butyrate or propionate production reported above and

the generally low predicted engraftment of *A. hallii*, we hypothesized that *A. muciniphila*

growth rate may be associated with clinical responsiveness. Indeed, the MCMM-predicted

growth rate of *A. muciniphila* was significantly higher in the WBF-011 group compared to the

placebo group (Fig. 3D; Mann–Whitney U test, p = 9.4×10^{-4}). Furthermore, we found that

ΔglucoseAUC was significantly negatively associated with the predicted growth rate of *A.*

muciniphila within the WBF-011 treatment group (Fig. 3E; linear regression, R^2 = 0.23, p =

0.027).

Validation Study B: Metagenomic Data Indicates Enrichment of Probiotic Species

Microbial relative abundances of focal species were determined using shotgun metagenomics

(see methods). *Entercloster bolteae* and *Flavonifractor plautii* showed an average enrichment

following treatment, from Week 0 to Week 12 (Fig. 1B; paired t-test, *E. bolteae*: p = 5.5 × 10-3;

1-sample t-test, *F. plautii*: p = 4.7 × 10-2). *Sellimonas intestinalis* and *Clostridium symbiosum*

also showed significant enrichment (Fig. 1B; paired t-test, *S. intestinalis*: p = 8.4×10^{-4}; C.

symbiosum: p = 7.4 × 10-3), but stayed below a relative abundance of 0.005 in nearly all cases.

D. longicatena also showed abundance above 0.005 in most cases, but it did not show a

significant increase between timepoints. Binarized engraftment scores were assigned based on

increases in relative abundance between timepoints, and being above a minimal relative

abundance threshold of 0.005 (see Methods).

Validation Study B: MCMMs predict species-level enrichment

Models for Validation Study B were constructed using baseline taxonomic profiles (N = 20). For

each individual, we modeled the addition of 6 of the 8 VE303 species; the remaining two were

omitted because GEMs were unavailable. These models were used to predict the engraftment

patterns observed in the Week 12 metagenomic sequencing data (see methods).

Across all comparisons (6 predictions for each of 20 samples, total N = 120) there was

76.7% agreement between model predicted growth and binarized enrichment scores (Fig. 2C,

Cohen's κ = 0.38, indicating fair agreement). *A. colihominis, S. intestinalis* and *C. symbiosum*

were the best predicted species, with only two incorrect predictions each. *D. longicatena* was

poorly predicted, with 10 incorrect predictions. There was significant agreement between

predictions and observations (Fig. 2D). Across all observations, 16 instances were correctly

predicted as growth, 76 were correctly predicted as non-growth, 13 instances were incorrectly

predicted as growth, and 15 were incorrectly predicted as non-growth (Fisher's exact test, p =

1.3×10^{-4}). The Cohen's κ score was lower than in Validation Study A, largely due to the

imbalance between the two observed engraftment states (31 instances of growth and 89

instances of non-growth), which inflates the expected chance agreement and depresses κ. For

this reason, κ downweights the otherwise strong (76.7%) concordance between MCMMs and

the metagenomic data.

MCMM Predictions Link Dietary Fiber Response to Clinical Health Markers

To evaluate whether MCMM-predicted prebiotic- and probiotic-responsive SCFAs were

associated with changes in host health, we modeled metabolic outputs from microbiome

samples collected from 1,786 individuals enrolled in a former precision-wellness program with

paired 16S amplicon sequencing data, blood-based clinical chemistries, and longitudinal follow-

up data available (Arivale cohort) [27]. Arivale participants were coached to increase their

exercise and to increase their fruit and vegetable intake. Each sample was simulated under two

dietary conditions: a low-fiber standard European diet (i.e., a rough proxy for the baseline, pre-

intervention diet) and a high-fiber diet enriched in resistant starch (i.e., a rough proxy for the

post-intervention diet). Across participants, simulations predicted heterogeneous increases in

butyrate production when transitioning from the low- to the high-fiber intervention (Fig. 4A).

Regression analyses revealed that larger predicted increases in butyrate (Δbutyrate) were

significantly associated with improvements in multiple cardiometabolic markers, including

lower LP-IR, fasting insulin, and HOMA-IR (Fig. 4B). Negative associations were also observed

between Δbutyrate and immune activation markers, such as total white blood cell count,

neutrophils, and lymphocytes.

MCMM-Optimized Prebiotic and Probiotic Interventions Designed to Improve Personalized

Probiotic Engraftment and SCFA Production Profiles

To assess the potential power of our MCMM approach, we applied participant-specific

simulations to a sub-cohort of Arivale participants with available fecal shotgun metagenomic

sequencing data (N = 156), which tend to provide slightly more accurate constraints for

MCMMs than 16S amplicon data [19].

Within the context of this Arivale sub-cohort, interventions that included the WBF-011

strains combined with eight different prebiotic and dietary contexts were simulated: (1) a

standard European diet without prebiotic supplementation, (2) a standard European diet

supplemented with inulin, (3) a standard European diet supplemented with pectin, (4) a

standard European diet supplemented with resistant starch, (5) a standard European diet
supplemented with maltodextrin, (6) a standard European diet supplemented with hemp seed,
(7) a standard European diet supplemented with psyllium husk, and (8) a standard high-fiber
diet. The addition of the probiotic consortia showed significant decreases in predicted butyrate
in every case except the no prebiotic and hemp seed contexts, where a significant increase was
observed (Fig. 5A, Mann-Whitney U test, $p < 0.05$). Similarly, predicted propionate production
decreased significantly when the probiotic cocktail was added, except in the no prebiotic or
hemp seed contexts, where no significant differences were observed (Fig. 5B, Mann-Whitney U
test, $p < 0.05$). Predicted acetate production showed a significant increase when the probiotic
was added in most cases, except in the no prebiotic and hemp seed contexts, where a
significant decrease was observed, and in the starch context, where no significant difference
was observed (Fig. 5C, Mann-Whitney U test, $p < 0.05$). Lactate production showed a significant
increase when the probiotic cocktail was added in concert with pectin, psyllium husk, hemp
seed, or in the no prebiotic context (Fig. 5D, Mann-Whitney U test, $p < 0.05$). To assess whether
changes in predicted SCFA production were driven primarily by increases in overall biomass, we
repeated these analyses using SCFA flux normalized by the predicted community growth rate.
The normalized results (Fig. S3) displayed patterns consistent with the original analysis (Fig. 5A-
D). Probiotic engraftment predictions for the WBF-011 cocktail varied only slightly across
prebiotic and dietary backgrounds, indicating that endogenous microbiota composition and
probiotic identity were stronger drivers of engraftment outcomes than prebiotic and dietary
contexts (Fig. 5E).
For each individual, the optimal prebiotic-probiotic combination on the standard
European diet was identified, in terms of maximizing predicted butyrate or propionate
production rates (Fig. 5F-G). While some patterns emerged, the optimal intervention varied
across individuals (Figs. S4-S5). For butyrate and propionate production, psyllium husk with no
probiotic cocktail was the most common optimal intervention. However, many individuals
showed a synbiotic combination as their optimal intervention (Figs. S4-S5). Overall, a
personalized optimal solution resulted in significantly higher predicted levels of propionate and
butyrate production when compared with the 'standard of care' synbiotic treatment used in
the original WBF-011 trial (i.e., combining inulin with the five-strain probiotic cocktail).

Discussion

The ability of orally administered probiotics to successfully engraft in the gut is influenced by
multiple ecological and host-specific factors, including the pre-existing composition of the gut
microbiome and the background diet. A method for the rational prediction of probiotic
engraftment and health-relevant microbial metabolite production, in the context of a given
microbiota and diet, would open up new avenues for precision prebiotic, probiotic, and
synbiotic design. Prior work by our group has demonstrated that our MCMM platform can be

leveraged to predict personalized SCFA production rates and personalized pathobiont
engraftment risk [19,23]. In order to validate our MCMM methodology in the context of
synbiotic design, we used existing data from Validation Studies A and B to determine whether
or not we can retrospectively predict relevant outcomes. Both studies showed variable
engraftment rates across probiotic organisms, along with variable clinical responses. Given this
heterogeneity, these trial data represented ideal test-cases for our MCMM predictions.

Assessing empirical engraftment—by measuring relative enrichment of a probiotic taxon
post-treatment via qPCR or metagenomics, and comparing these to MCMM predictions—is
challenging. The presence of endogenous strains of the same species can make the detection of
probiotic strains from metagenomic data challenging. Furthermore, species-level enrichment of
probiotic taxa can be difficult to distinguish from increases in the abundances of related strains
already present. In Validation Study A, strain-specific qPCR primers often yielded no baseline
detection, so post-treatment detection above a species-specific threshold was attributed to the
administered strains. In Validation Study B, where baseline metagenomes contained
endogenous strains, engraftment was defined by a two-fold criterion: detection above a global
threshold and an increase relative to the individual baseline. Although these metrics do not
prove long-term engraftment, they provide a rough proxy for short-term enrichment of the
administered probiotic species.

Despite the inherent complexity of predicting probiotic growth, the MCMM-based
framework demonstrated substantial accuracy, showing over 80% agreement with empirical
probiotic engraftment metrics in Validation Study A (Fig. 2A) and over 75% agreement in
Validation Study B (Fig. 2C). These models incorporate key ecological determinants, like
endogenous microbiome composition and the availability of prebiotic and dietary substrates, to
estimate probiotic growth rates. Notably, the high level of accuracy in engraftment predictions
was achieved in spite of the fact that we lacked personalized constraints on dietary intake (i.e.,
a standard European diet was applied to each model). It is also worth noting the instances
where the model failed - for instance, *B. infantis* and *D. longicatena* were poorly predicted in
Validation Study A and Validation Study B, respectively. This could be the result of unannotated
pathways in the GEMs of these poorly-predicted taxa, or a mismatch between the *in silico*
dietary constraints applied in the model and the actual consumption patterns of study
participants.

A fundamental goal of probiotic and prebiotic supplementation is to modulate the
production of key microbial metabolites that support host health. Among these, short-chain
fatty acids (SCFAs)—such as butyrate and propionate—are particularly significant due to their
roles in maintaining gut barrier integrity, regulating proper immune function, and supporting
metabolic homeostasis [19,26]. Synbiotic administration provides a means of enhancing SCFA
production, and MCMMs can be leveraged to predict these metabolic shifts. Notably, our
analysis revealed a synergistic effect between prebiotic administration and the WBF-011

probiotic cocktail. The probiotic cocktail paired with only a small amount of inulin showed a
minimal effect with regard to butyrate and propionate production (Fig. 3A-B). However,
increasing the amount of inulin supplementation resulted in a significant increase in the
production of both butyrate and propionate (Fig. 3A-B).
Given accurate predictions of probiotic engraftment, we sought to identify associations
between probiotic administration and the primary outcomes of the initial study. Perraudeau et
al. reported a significant reduction in postprandial glucose AUC over the 12-week trial in the
WBF-011 treatment group (Fig. 3C), but not in the placebo or WBF-010 groups [27]. The
primary difference between WBF-011 and WBF-010 was the inclusion of two additional strains:
A. muciniphila and *A. hallii.* While *A. hallii* was rarely predicted to grow, *A. muciniphila* was
consistently predicted to grow in most individuals receiving WBF-011 (Fig. 3D). We
hypothesized that *A. muciniphila* growth might be linked to changes in glucose AUC. Consistent
with this hypothesis, MCMM-predicted growth rates of *A. muciniphila* were negatively
associated with ΔglucoseAUC both across the WBF-011 and placebo groups and within the
WBF-011 group (Fig. 3C-E). These findings support the idea that *A. muciniphila* growth and
metabolic activity may contribute substantially to the clinical response observed in WBF-011
recipients. Future intervention studies integrating longitudinal metagenomic and host-response
data could help establish more direct mechanistic links between *A. muciniphila* growth,
microbial metabolism, and host outcomes.
Building on the strong agreement observed between predicted growth and empirically-
estimated engraftment in two probiotic intervention studies, we applied MCMM simulations to
a larger cohort of generally healthy participants from Arivale (N = 1,786) who were coached to
improve their diet and lifestyle. Using MCMMs built from this data set, we simulated a switch
from a low- to a high-fiber diet, and compared how changes in predicted butyrate (Δbutyrate)
were associated with longitudinal changes in cardiometabolic markers (LP-IR, insulin, HOMA-IR)
and immune activation markers (white blood cell, neutrophil, lymphocyte counts). Assuming
that participants were generally guided toward a higher-fiber, butyrogenic diet, these findings
highlight the potential of MCMM-based approaches to predict personalized microbiome-
mediated responses to dietary interventions.
Microbial production of butyrate and propionate increased in response to prebiotic
supplementation and a shift to a standard high-fiber diet. Interestingly, addition of the WBF-
011 probiotic cocktail generally led to decreases in butyrate and propionate production, which
was partially explained by concomitant increases in lactate and acetate. Although the cocktail is
designed to be butyrogenic, the limited growth observed for butyrate producers *A. hallii, C.*
beijerinckii, and *C. butyricum* is consistent with this result. Individual-specific modeling,
however, allows identification of the most effective treatment for each sample, in terms of
SCFA production. Responses varied across individuals, with not all subjects benefiting equally
from the same prebiotic fibers, and some individuals showed higher butyrate production in the

presence of WBF-011, despite the average negative trend across individuals. This highlights the
potential of MCMMs as a tool for personalizing prebiotic and probiotic combinations.
MCMMs have several inherent limitations. Chief among these is that MCMMs rely on
simplifying steady-state assumptions that restrict their ability to fully capture the temporal
dynamics of gut microbial communities. Moreover, the MCMM framework depends on the
availability of accurate GEMs for all of the taxa in a sample, and missing annotations or
reactions in existing GEMs can lead to inaccurate predictions. Existing GEM databases do not
include all extant strains for a given gut bacterial species. The consequence of this is that
MCMM-based analyses constrained by real-world gut microbiome samples are often limited to
species-level resolution, using models that summarize the metabolic capacity of all available
strains within a given species. Our study relied on species-level GEMs from the AGORA2
database, due to their relatively higher levels of curation and quality, compared to *de novo*
constructed models [28,29]. Future work should focus on accounting for strain-level differences
in metabolic capacity. Another limitation was the lack of participant dietary intake data. Better
constraints on personalized dietary intake could further improve the predictive accuracy that
we observed in the current analysis, which assumed that everyone was eating a standard
European diet. Finally, the current methodology for constructing and solving MCMMs does not
account for spatial structure or host physiology within the gxut. For instance, *A. muciniphila*
typically grows within the mucus layer of the colonic epithelium, in a different
microenvironment than many other taxa in the microbiota. For this reason, it is likely to be
more exposed to host-derived compounds. Future work should aim to integrate MCMMs with
metabolic models of the colonic epithelium [30], which may help to further improve
predictions.
Taken together, these findings demonstrate the utility of MCMMs as a predictive
framework for assessing prebiotic, probiotic, and dietary interventions at the individual and
population levels. By integrating metabolic modeling with experimental validation, we highlight
key ecological and metabolic determinants of probiotic engraftment success and SCFA
production, providing a pathway for more targeted microbiome-based therapies. However,
these approaches are still evolving and require further validation in prospective human trials.
Realizing the full potential of these models will require continued refinement, including
improved strain-level resolution, personalized dietary intake constraints, and enhanced host-
microbiome interaction modeling. Future research should aim to bridge these gaps by
incorporating multi-omics approaches, longitudinal sampling, and personalized dietary
profiling. Ultimately, leveraging MCMMs in a clinical setting could enable precision microbiome
therapeutics, optimizing probiotic, prebiotic, and dietary intake to more effectively treat a wide
range of acute and chronic diseases.

Methods

Data collection

Data from Validation Study A were obtained from Perraudeau et al. [24]. The study cohort consisted of 76 participants previously diagnosed with type 2 diabetes. Participants were randomized into two treatment groups (N = 27 and N = 23) and a placebo group (N = 26). 6, 2 and 10 participants were lost to follow-up in each study arm, respectively. Those in the experimental groups received one of two probiotic formulations, WBF-010 or WBF-011. WBF-010 contained strains of *Bifidobacterium longum subsp. infantis*, *Clostridium butyricum*, and *Clostridium beijerinckii*. WBF-011 included the same strains as WBF-010 with the addition of *Akkermansia muciniphila* and *Anaerobutyricum hallii*. Both probiotic formulations were supplemented with a small amount of inulin (0.3 grams), a prebiotic derived from chicory root. Participants in all three study arms followed the intervention for 12 weeks, consuming three capsules twice daily. The primary metabolic endpoint assessed in the original trial was the change in the area under the glucose curve (AUC) during a standard three-hour meal-tolerance test (MTT) conducted at baseline and after 12 weeks. The ΔglucoseAUC values were calculated by subtracting baseline AUC from the 12-week AUC, from the same individual. Analysis revealed that WBF-011, but not WBF-010, led to a significant shift in ΔglucoseAUC compared to the placebo. All data were shared directly by the authors of the original study.

Data from Validation Study B were obtained from Dsouza et al. [12]. This cohort consisted of 32 individuals treated with an 8-strain probiotic cocktail called VE303, containing strains of *Enterocloster bolteae*, *Anaerotruncus colihominis*, *Sellimonas intestinalis*, *Clostridium symbiosum*, *Blautia sp001304935*, *Dorea longicatena*, *Clostridium innocuum,* and *Flavonifractor plautii*. 20 participants who were treated with antibiotics prior to probiotic administration and had followup data at 12 weeks were used to test the modeling approach. Of the strains included in the cocktail, 6 out of 8 (excluding *Clostridium innocuum* and *Blautia sp001304935*) were represented in the AGORA2 database [25], and could be modeled using MCMMs. Shotgun metagenomes from this study were collected from the NCBI Sequence Read Archive under the accession number PRJNA755324.

Additional data used in the analysis were collected from Arivale, Inc. (Seattle, WA). Arivale closed its operations in 2019. For analyzing the shift in butyrate production between a standard European and a high-fiber diet, 16S rRNA amplicon data was used for samples that had paired baseline fecal and blood sampling, as well as a secondary blood sampling between 4 and 8 months later (N = 1786). Participants in this analysis were adults in the United States (1079 female, 707 male, sex assigned at birth), with average age 48.5 ± 0.3 years. 16S data for the Arivale cohort can be found on the NCBI Sequence Read Archive under accession numbers PRJNA826530 and PRJNA826648. For exploratory analysis testing combinatoric prebiotic, probiotic, and dietary interventions, a subset of these participants for whom shotgun metagenomic sequencing was available were used (N = 156). De-identified metagenomic data

410 was collected and used to construct MCMMs as described below. Participants included in this
411 sub-cohort analysis were adults in the United States (92 females and 64 males, sex assigned at
412 birth), with average age 47.4 ± 1.0 years. Arivale metagenomes were deposited at the NCBI
413 Sequence Read Archive under accession number PRJNA1262070. Further details on the Arivale
414 cohort are described in Zubair et al. [31]
415
qPCR Data Processing
qPCR data from Validation Study A were used as ground truth for validating model-based
predictions. These data were provided by the authors of the original study. Custom primer pairs
were designed for each of the five probiotic strains in the WBF-011 treatment [24]. Cycle
threshold (Ct) values were obtained for all five species at baseline (Week 0), during treatment
(Week 4), and at the end of treatment (Week 12). Samples failing quality control were excluded
from analysis. Despite using strain-specific primer pairs, all five probiotic strains were detected
in at least one sample at Week 0, suggesting cross-reactivity with endogenous strains.
Because qPCR standard and melt curves from the original experiments were not
retrievable, the resulting data were semiquantitative and were used to assess within-species
trends, rather than estimate absolute abundances. -Ct was used as a proxy for relative species
abundance. For each species, an engraftment threshold was defined as the mean -Ct at Week 0,
representing the baseline amplification level. Week 12 values were binarized relative to this
threshold to classify each sample as either exhibiting growth or non-growth.
430
Shotgun Metagenome Data Processing
For Validation Study B, metagenomic data from Week 0 and Week 12 was used as ground truth
to validate model predictions. Metagenomic data from Week 0 were used for both Validation
Study A and Validation Study B to construct MCMMs. FASTQ files were collected from the
authors for Validation Study A and from the NCBI Sequence Read Archive for Study B. Raw data
was adapter-trimmed and quality filtered using fastp [32]. Quality-filtered reads were
processed using Kraken2 (v1.01) for taxonomic classification, and Bracken was used to refine
taxonomic abundance estimates [33,34]. The Kraken2 default database (based on Refseq
release 94) was used. Species with fewer than 10 assigned reads were omitted. The analysis
pipeline is available at https://github.com/Gibbons-Lab/pipelines/tree/master/metagenomics.
The same metagenomic processing pipeline was run on the shotgun metagenomic data for
Arivale participants.
For Validation Study B, species relative abundances from Week 12 were used as a
benchmark for comparison with model predictions. An increase in relative abundance of 50%
was used for binarization, subject to an additional relative abundance threshold filter of 0.005
at Week 12. Using this method, each sample/species pair was classified as showing growth or
non-growth.

16S rRNA Amplicon Data Processing

16S rRNA amplicon sequencing data was processed as described by Diener et al. [35]). Briefly, reads were processed using DADA2 to infer amplicon sequence variants, learn run-specific error profiles, merge counts and remove chimeras [36]. Taxonomy was assigned using a naive Bayes classifier and the Silva database (v128) [37]. Approximately 90% of reads were classified to genus level and 32% to species level. Low abundance taxa present in fewer than 50% of samples or with an average of less than 10 reads per sample were excluded.

Model Construction

Read counts were normalized to relative abundance before their incorporation into downstream modeling workflows. MCMMs were constrained using 16S amplicon or metagenomic relative abundance profiles for all baseline samples using MICOM (v0.37.0), a framework for simulating microbial community-scale metabolism [18]. Taxonomic profiles from baseline samples were mapped to genome-scale metabolic models (GEMs) from the AGORA database (v2.01) [25], which were used to construct community-scale metabolic models reflecting the microbial composition of each individual microbiome. A lower relative abundance threshold of 0.001 was applied during model construction to exclude taxa representing less than 0.1% of the community composition.

For samples in Validation Studies A and B, probiotic supplementation was simulated *in silico* by modifying the baseline taxonomic composition. For Validation Study A, the relative abundances of endogenous taxa were proportionally reduced to 80% of their original values, while five probiotic species from WBF-011 were introduced at a total relative abundance of 20% (4% each). For Validation Study B, the relative abundances of endogenous taxa were proportionally reduced to 20% of their original values to reflect vancomycin pre-treatment, while 6 probiotic species from VE303 cocktail were introduced at a total relative abundance of 80% (13.33% each). These models were constructed at the species level.

For the dietary intervention modeling, MCMMs were constructed from 16S rRNA amplicon sequencing data available for 1,786 samples in the Arivale cohort with paired stool and blood samples at baseline and a blood sample from a 4-8 month follow-up sampling. Due to the taxonomic resolution of 16S rRNA amplicon sequencing data, these models were constructed at the genus level.

For the additional analysis of prebiotic, probiotic, and dietary combinations in the Arivale cohort, MCMMs were constructed at the species-level using metagenomic sequencing data available for 156 samples. Probiotic administration was simulated by scaling relative abundances of endogenous taxa to 80% of their original values, and introducing 20% relative abundance of the probiotic cocktail from Study A (4% each). These models were constructed at the species level.

Model Growth

Community growth simulations were performed after selecting MICOM's cooperative tradeoff
parameter using the micom.workflows.grow() workflow, which balances individual taxon
growth with overall community-wide metabolic efficiency. A tradeoff parameter of 0.99 was
applied because it was the closest value to 1.0 that also allowed for >80% of taxa to grow in
Validation Study A. This parameter value was then fixed and applied to all other cohorts in this
study to avoid overfitting. For Validation Studies A and B, models were supplied with a
simulated growth medium reflective of an average European diet collected from the Virtual
Metabolic Human diet database (https://www.vmh.life/#nutrition). Metabolites in the medium
that can be taken up by the host, as defined by uptake reactions in the Recon3D model, were
reduced to 20% of their original flux to account for absorption in the small intestine [38].
Additionally, host-derived compounds such as mucins and bile acids that were missing from the
medium were added. The medium was further supplemented with a minimal set of metabolites
necessary to support growth of all taxa in the model database, using MICOM's
complete_db_medium() function. In the absence of detailed dietary recall data, this can be
used to approximate the average dietary consumption of individuals in North America, where
the cohorts included in this study were located. For Validation Study A, 0.3 g or 30 g of inulin
were included as an additional substrate in the medium to account for the prebiotic component
of the synbiotic intervention. Predicted growth rates were collected from the output of the
micom.workflows.grow() workflow, and prediction metabolic production rates were calculated
using the micom.measures.production_rates() function in MICOM.

For analysis of the Arivale cohort, simulations were conducted using a standard
European diet as well as a high-fiber diet rich in resistant starch. This media was also collected
from the Virtual Metabolic Human database, and augmented in the same fashion as the
standard European medium. The same cooperative tradeoff parameter as in Validation Studies
A and B was used. Metabolic production from the model output was calculated using the
micom.measures.production_rates() function in MICOM. Δbutyrate was calculated as predicted
microbial butyrate production on a standard European diet subtracted from predicted butyrate
production on a high-fiber diet. Changes in clinical chemistry values (Δchemistries) were
calculated as the value at enrollment subtracted from the value at follow-up (4-8 months post
enrollment). These values were used as the independent and dependent variables for
regression analysis, respectively (see Statistical Analysis below).

Inulin, pectin, starch, maltodextrin, cellulose (hemp seed), or arabinoxylan (psyllium
husk) were added to the standard European diet to simulate the addition of various prebiotics.
All prebiotic additions were added in a manner that balanced the overall carbon content to
enable comparison between interventions, equivalent to a prebiotic supplementation of
approximately 30g per day. Predicted growth rates for the five probiotic species MICOM model

outputs, and predicted metabolic production rates were calculated using the

micom.measures.production_rates() function. To facilitate direct comparison with engraftment

scores in the validation studies, predicted growth rates were binarized using a threshold of 10^{-2}

h^{-1} (corresponding to a minimal doubling time of 69.3h). Growth rates below this value were

classified as non-engraftment (0), while those above were classified as successful engraftment

(1).

Statistical Analysis

All statistical analysis conducted in this study was done using Scipy (v1.13.1) [39], scikit-learn

(v1.6.1) [40], or statsmodels (v0.14.4) [41]

To assess significant increases in abundance of probiotic species from qPCR data, paired

t-tests were used to compare Week 0 and Week 12 -Ct values. Statistical analyses were

performed in Python using the scipy.stats.ttest_rel() function. In instances where only a single -

Ct value was available from the Week 0 time point, a one-sample t-test was used to compare

that sample to the 12 week time point using the scipy.stats.ttest_1samp() function.

To assess the concordance between binarized MICOM-predicted engraftment and qPCR

observations, Cohen's Kappa statistic was computed using

sklearn.metrics.cohen_kappa_score(). This measure quantifies the level of agreement beyond

what would be expected by chance, providing a stringent estimate of the predictive accuracy of

MCMMs for probiotic engraftment outcomes. Furthermore, Fisher's exact test was used to

evaluate whether the observed agreement between MICOM predictions and qPCR results in

the confusion matrix was statistically significant, using the scipy.stats.fisher_exact() function.

Significance differences in predicted butyrate and propionate between treatment

groups in both the probiotic trial and the Arivale cohort (Fig. 3A-B, Fig. 5A-D) were assessed

using non-parametric Mann-Whitney U test, using the scipy.stats.mannwhitneyu() function.

Significance differences between predicted butyrate and propionate between optimal and

standard of care interventions in the Arivale cohort (Fig. 5F-G) were also determined using non-

parametric Mann-Whitney U tests.

Significant differences in Δglucose AUC and *A. muciniphila* growth rates between

treatment groups was determined using non-parametric Mann-Whitney U test, using the

scipy.stats.mannwhitneyu() function. The association between *A. muciniphila* growth rate and

change in glucose AUC was assessed using ordinary least squares linear regression, using the

statsmodel.api.ols() function. In the absence of available metadata, no covariates were used.

Regression analysis associating changes in butyrate production between a standard

European and high-fiber diet to changes in a set of 127 blood-based clinical chemistries (Fig. 4)

was conducted using statsmodels, with age, sex and the baseline values of clinical chemistries

as covariates, using the statsmodel.api.ols() function. Correction for multiple comparisons was

performed using the Benjamini-Hochberg method[42], using the
statsmodels.stats.multitest.fdrcorrection() function.
563
Data and Code Availability
qPCR and metagenomic sequencing data from Perraudeau et al. (2020) are available upon
request. Metagenomic sequencing data from Dsouza et al. (2022) are available on the NCBI
Sequence Read Archive under accession number PRJNA755324. Metagenomic sequencing data
from 156 members of the Arivale dataset are available on the NCBI Sequence Read Archive
under accession number PRJNA1262070. The code used for the analyses presented in the
current manuscript can be found at the following GitHub repository:
https://github.com/Gibbons-Lab/2024_probiotic_engraftment.
572

Competing Interests Statement
I have read the journal's policy and the authors of this manuscript have the following competing
interests. This work was funded, in part, by a research grant from Pendulum (to SMG), which is
the company that manufactured the synbiotic tested in Perraudeau et al. (2020). Pendulum had
no part in the design or conduct of this study or the writing of this manuscript. SMG is a paid
member of the scientific advisory board for Thorne. This work is unrelated to Thorne, and
Thorne had no involvement in the design or conduct of this study or in the writing of the
manuscript.
581

582   **Acknowledgements**
583   This work was supported by research funding from Pendulum Therapeutics, Inc., by the
584   National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) of the National
585   Institutes of Health (NIH) under award number R01DK133468, and by a Global Grants for Gut
586   Health Award from Nature Portfolio and Yakult (to SMG).

References

1.   Oliphant K, Allen-Vercoe E. Macronutrient metabolism by the human gut microbiome: major
fermentation by-products and their impact on host health. Microbiome. 2019;7: 91.

2.   Rackerby B, Van De Grift D, Kim JH, Park SH. Effects of Diet on Human Gut Microbiome
and Subsequent Influence on Host Physiology and Metabolism. Gut Microbiome and Its
Impact on Health and Diseases. 2020. pp. 63–84. doi:10.1007/978-3-030-47384-6_3

3.   Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. Cell. 2014;157:
121–141.

4.   Zheng D, Liwinski T, Elinav E. Interaction between microbiota and immunity in health and
disease. Cell Res. 2020;30: 492–506.

5.   Ryan FJ, Clarke M, Lynn MA, Benson SC, McAlister S, Giles LC, et al. Bifidobacteria
support optimal infant vaccine responses. Nature. 2025. doi:10.1038/s41586-025-08796-4

6.   Nagpal R, Wang S, Ahmadi S, Hayes J, Gagliano J, Subashchandrabose S, et al. Human-
origin probiotic cocktail increases short-chain fatty acid production via modulation of mice
and human gut microbiome. Sci Rep. 2018;8: 12649.

7.   Markowiak-Kopeć P, Śliżewska K. The effect of probiotics on the production of short-chain
fatty acids by human intestinal microbiome. Nutrients. 2020;12: 1107.

8.   Snigdha S, Ha K, Tsai P, Dinan TG, Bartos JD, Shahid M. Probiotics: Potential novel
therapeutics for microbiota-gut-brain axis dysfunction across gender and lifespan.
Pharmacology & Therapeutics. 2022;231: 107978.

9.   Mazziotta C, Tognon M, Martini F, Torreggiani E, Rotondo JC. Probiotics mechanism of
action on immune cells and beneficial effects on human health. Cells. 2023;12: 184.

10. Cristofori F, Dargenio VN, Dargenio C, Miniello VL, Barone M, Francavilla R. Anti-
inflammatory and immunomodulatory effects of probiotics in gut inflammation: A door to the
body. Front Immunol. 2021;12: 578386.

11. Maldonado-Gómez MX, Martínez I, Bottacini F, O'Callaghan A, Ventura M, van Sinderen D,
et al. Stable engraftment of Bifidobacterium longum AH1206 in the human gut depends on
individualized features of the resident microbiome. Cell Host Microbe. 2016;20: 515–526.

12. Dsouza M, Menon R, Crossette E, Bhattarai SK, Schneider J, Kim Y-G, et al. Colonization
of the live biotherapeutic product VE303 and modulation of the microbiota and metabolites
in healthy volunteers. Cell Host Microbe. 2022;30: 583–598.e8.

13. Coyte KZ, Rakoff-Nahoum S. Understanding Competition and Cooperation within the
Mammalian Gut Microbiome. Curr Biol. 2019;29: R538–R544.

14. Spencer SP, Fragiadakis GK, Sonnenburg JL. Pursuing Human-Relevant Gut Microbiota-
Immune Interactions. Immunity. 2019;51: 225–239.

15. Fujita H, Ushio M, Suzuki K, Abe MS, Yamamichi M, Iwayama K, et al. Alternative stable
states, nonlinear behavior, and predictability of microbiome dynamics. Microbiome.

624  2023;11: 63.

16. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nat Biotechnol. 2010;28: 245–248.

17. Diener C, Gibbons SM. More is Different: Metabolic Modeling of Diverse Microbial Communities. mSystems. 2023;8: e0127022.

18. Diener C, Gibbons SM, Resendis-Antonio O. MICOM: Metagenome-Scale Modeling To Infer Metabolic Interactions in the Gut Microbiota. mSystems. 2020;5. doi:10.1128/mSystems.00606-19

19. Quinn-Bohmann N, Wilmanski T, Sarmiento KR, Levy L, Lampe JW, Gurry T, et al. Microbial community-scale metabolic modelling predicts personalized short-chain fatty acid production profiles in the human gut. Nature Microbiology. 2024; 1–13.

20. Carr A, Baliga NS, Diener C, Gibbons SM. Personalized Clostridioides difficile engraftment risk prediction and probiotic therapy assessment in the human gut. bioRxiv. 2024. doi:10.1101/2023.04.28.538771

21. Orellana E, Zampieri G, De Bernardini N, Guerrero LD, Erijman L, Campanaro S, et al. Sustainable Food Waste Management in Anaerobic Digesters: Prediction of the Organic Load Impact by Metagenome-Scale Metabolic Modeling. Environ Sci Technol. 2025. doi:10.1021/acs.est.4c11180

22. Marcelino VR, Welsh C, Diener C, Gulliver EL, Rutten EL, Young RB, et al. Disease-specific loss of microbial cross-feeding interactions in the human gut. Nat Commun. 2023;14: 6546.

23. Carr AV, Baliga NS, Diener C, Gibbons SM. Personalized Clostridioides difficile colonization risk prediction and probiotic therapy assessment in the human gut. Cell Syst. 2025;16: 101367.

24. Perraudeau F, McMurdie P, Bullard J, Cheng A, Cutcliffe C, Deo A, et al. Improvements to postprandial glucose control in subjects with type 2 diabetes: a multicenter, double blind, randomized placebo-controlled trial of a novel probiotic formulation. BMJ Open Diabetes Res Care. 2020;8: e001319.

25. Heinken A, Hertel J, Acharya G, Ravcheev DA, Nyga M, Okpala OE, et al. Genome-scale metabolic reconstruction of 7,302 human microorganisms for personalized medicine. Nat Biotechnol. 2023;41: 1320–1331.

26. Tan J, McKenzie C, Potamitis M, Thorburn AN, Mackay CR, Macia L. The role of short-chain fatty acids in health and disease. Adv Immunol. 2014;121: 91–119.

27. Diener C, Qin S, Zhou Y, Patwardhan S, Tang L, Lovejoy JC, et al. Baseline Gut Metagenomic Functional Gene Signature Associated with Variable Weight Loss Responses following a Healthy Lifestyle Intervention in Humans. mSystems. 2021;6: e0096421.

28. Machado D, Andrejev S, Tramontano M, Patil KR. Fast automated reconstruction of genome-scale metabolic models for microbial species and communities. Nucleic Acids Res. 2018;46: 7542–7553.

29. Zimmermann J, Kaleta C, Waschina S. Gapseq: Informed prediction of bacterial metabolic
pathways and reconstruction of accurate metabolic models. Genome Biol. 2021;22: 81.

30. Jiang B, Quinn-Bohmann N, Diener C, Nathan VB, Han-Hallett Y, Reddivari L, et al.
Understanding disease-associated metabolic changes in human colon epithelial cells using
ColonEpithelium metabolic reconstruction. bioRxiv. 2024. doi:10.1101/2024.10.22.619644

31. Zubair N, Conomos MP, Hood L, Omenn GS, Price ND, Spring BJ, et al. Genetic
predisposition impacts clinical changes in a lifestyle coaching program. Sci Rep. 2019;9:
6805.

32. Chen S. fastp 1.0: An ultra-fast all-round tool for FASTQ data quality control and
preprocessing. Imeta. 2025;4: e70078.

33. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome
Biol. 2019;20: 257.

34. Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species abundance in
metagenomics data. PeerJ Comput Sci. 2017;3: e104.

35. Diener C, Dai CL, Wilmanski T, Baloni P, Smith B, Rappaport N, et al. Genome-microbiome
interplay provides insight into the determinants of the human blood metabolome. Nat
Metab. 2022;4: 1560–1572.

36. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13: 581–583.

37. Chuvochina M, Gerken J, Frentrup M, Sandikci Y, Goldmann R, Freese HM, et al. SILVA in
2026: a global core biodata resource for rRNA within the DSMZ digital diversity. Nucleic
Acids Res. 2025; gkaf1247.

38. Brunk E, Sahoo S, Zielinski DC, Altunkaya A, Dräger A, Mih N, et al. Recon3D enables a
three-dimensional view of gene variation in human metabolism. Nat Biotechnol. 2018;36:
272–281.

39. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy
1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17: 261–
272.

40. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. J Mach Learn Res. 2011;12: 2825–2830.

41. Seabold S, Perktold J. Statsmodels: Econometric and statistical modeling with python.
Proceedings of the 9th Python in Science Conference. SciPy; 2010. doi:10.25080/majora-
92bf1922-011

42. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. J R Stat Soc. 1995;57: 289–300.

698
699
700 **Figures and Captions**

702
Figure 1. Probiotic strain abundance quantification in two probiotic intervention trials
(A) Based on qPCR data, there was a significant enrichment in *A. muciniphila* and *B. infantis*
probiotics in Validation Study A at Week 12 compared to Week 0. There was no enrichment of
C. beijerinckii, C. butyricum and *A. hallii* probiotics. (B) Based on shotgun metagenomic data,
there was significant enrichment in *E. bolteae, F. plautii, S. intestinalis,* and *C. symbiosum* in
Validation Study B at Week 12 compared to Week 0. Only *E. bolteae* and *F. plautii* showed
notable enrichment above the relative abundance threshold of 0.005. Significance determined
by Mann-Whitney U test between timepoints, or one-sample t-test when only a single
datapoint was available,* = p < 0.05, ** = p < 0.01, *** = p < 0.001.

Figure 2. MCMM predictions for probiotic growth show significant agreement with observed engraftment across two human intervention trials. (A) Binarized MCMM-predicted growth agreed with qPCR engraftment scores in 89 out of 105 observations in Validation Study A (Cohen's κ = 0.68, indicating moderate agreement). Blue boxes indicate agreement and red boxes indicate disagreement between the model and the qPCR data. (B) Binarized MCMM-predicted growth predictions agree with metagenomic engraftment scores in 92 out of 120 observations in Validation Study B (Cohen's κ = 0.38, indicating fair agreement). Blue boxes indicate agreement and red boxes indicate disagreement between the model and the metagenomic data. (C) Confusion matrices describing results across and within probiotic strains, showing highly significant agreement for Validation Study A (Fisher's Exact Test, aggregate results across all strains, p = 4.4x10^{-13}). (D) Confusion matrices describing results across and within probiotic strains, showing highly significant agreement (Fisher's Exact Test, aggregate results across all strains, p = 1.3x10^{-4}) for Validation Study B.

Figure 3. MCMM–predicted SCFA production rates shift in response to synbiotic treatment
and *Akkermansia muciniphila* growth rates are associated with clinical responses in
Validation Study A. (A) Butyrate production did not increase with the probiotic treatment when
combined with a minimal prebiotic dose. Combining the probiotic treatment with a larger
prebiotic dose results in a significant increase in butyrate production. (B) Propionate production
also does not show significant changes with the probiotic treatment when combined with a
minimal prebiotic dose. A more substantial prebiotic dose results in a significant increase in
propionate production. Each point represents the predicted production level for a single
sample. Significance was determined by Mann-Whitney U-test **: p < 0.01, ***: p < 0.001. (C)
Participants in the WBF-011 treatment group showed a significantly larger decrease in glucose
AUC between Week 0 and Week 12 than those in the placebo group. Significance was
determined by Mann-Whitney U-test **: p < 0.01. (D) *A. muciniphila* showed a significantly
higher predicted growth rate in the WBF-011 treatment group than the placebo group.
Significance was determined by Mann-Whitney U-test ***: p < 0.001 (E) Predicted growth rate
of *A. muciniphila* displays a negative association with change in glucose AUC (Δglucose AUC)
between Week 0 and Week 12 in the WBF-011 treatment group. Each point represents a single
sample, the black line represents a least-squares regression between Δglucose AUC and
predicted growth rate, and the shaded gray region represents the 95% confidence interval.

Figure 4. Switching from a low-fiber to high-fiber diet in the Arivale cohort causes non-uniform shifts in MCMM-predicted butyrate production, which are significantly associated with longitudinal changes in several clinical health markers. (A) Simulating a dietary shift from a low-fiber standard European diet (blue points) to a high-fiber diet (red points) results in a non-uniform increase in butyrate production across the study population (N = 1,786). (B) Change in butyrate production between diets (Δbutyrate) correlates with change in values of clinical chemistries across the duration of a lifestyle coaching program. Associations were determined using multiple regression adjusting for age, sex, and baseline values for each respective chemistry. Correction for multiple comparisons was performed using the Benjamini-Hochberg method. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

Figure 5. MCMM-predicted SCFA & lactate production rates and WBF-011 probiotic species engraftment across simulated probiotic and prebiotic interventions in a cohort of generally-healthy Americans. (A-D) Predicted butyrate, propionate, acetate and lactate production rates were calculated upon treatment with prebiotic substrates on a standard European diet (inulin, pectin, resistant starch, maltodextrin, psyllium or hemp seed), or with a shift from a standard European to a standard high-fiber diet. Addition of a probiotic cocktail further shifted production rates, and tended to decrease butyrate and propionate production rates, relative to the prebiotic-alone treatments. Acetate and lactate production was generally higher in cases where the probiotic cocktail was added. Color encoding demonstrates the effect of adding a probiotic cocktail to the prebiotic/dietary treatments. Significance between the no probiotic and probiotic contexts determined by Mann-Whitney U-test, * = $p<0.05$, ** = $p<0.01$, *** = $p<0.001$. (E) Predicted engraftment of probiotic species, using a growth rate binarization threshold of 10^{-3}, calculated for each prebiotic addition. In all cases, *A. muciniphila* and *B. infantis* showed success in greater than 100 samples, while *A. hallii, C. beijerinckii* and *C. butyricum* did not engraft in more than 15 samples at most. (F) For both predicted butyrate and propionate production, an optimal prebiotic and/or probiotic treatment combination was selected and compared against a standard treatment, which consisted of inulin and the 5-strain probiotic cocktail. In both cases, the individual-specific optimal treatment showed significantly higher levels of production. Significance determined by Mann-Whitney U-test, *** = $p<0.001$.

828 **Supplementary Figure Captions**

829
Figure S1. Sensitivity of Validation Study A to Growth Rate Threshold. Agreement fraction
(red line) and Cohen's κ are shown as a function of the growth rate threshold, around which
model predictions are binarized for growth or non-growth. A black vertical line is shown at
0.01, the value used in this analysis, equivalent to doubling time of ~70 hours.

Figure S2. Sensitivity of Validation Study B to Growth Rate Threshold. Agreement fraction

(red line) and Cohen's κ are shown as a function of the growth rate threshold, around which

model predictions are binarized for growth or non-growth. A black vertical line is shown at

0.01, the value used in this analysis, equivalent to doubling time of ~70 hours.

840
Figure S3. Scaling Metabolic Predictions by Community Growth Rate. To ensure that
observed shifts in metabolite production were not driven by overall changes in biomass,
production rates were normalized by predicted community growth rates which reflect the total
biomass produced by the community. Scaled values for SCFA and lactate production were
largely consistent with the unscaled predictions across prebiotic and probiotic conditions.
Significant differences between ±probiotic treatments within each prebiotic or diet intervention
were assessed using a Mann-Whitney U test (p < 0.05; p < 0.01; p < 0.001).

848
Figure S4. Individual-specific optimal prebiotic, probiotic, and synbiotic treatments for
maximizing butyrate production. MCMM-predicted butyrate production varied substantially
across individuals (N =156) in response to different prebiotic and probiotic combinations. The
combination of psyllium husk with no probiotic cocktail produced the highest butyrate flux in the
greatest number of individuals. However, every prebiotic/probiotic combination was optimal for
at least one individual, with the exceptions of: (1) hemp seed, and (2) the no-probiotic condition.
Heatmap colors indicate predicted butyrate flux (mmol/gDW/h), and black boxes denote the
most effective treatment for each sample.

857
Figure S5. Individual-specific optimal prebiotic, probiotic, and synbiotic treatments for
maximizing propionate production. MCMM-predicted propionate production also varied
substantially across individuals (N = 156) in response to different prebiotic and probiotic
combinations, though slightly less than for butyrate. The combination of psyllium husk without a
probiotic cocktail produced the highest propionate flux in the greatest number of individuals.
However, every other prebiotic/probiotic combination was optimal for at least one individual, with
the exceptions of: (1) hemp seed, and (2) the no-probiotic condition.Heatmap colors indicate
predicted butyrate flux (mmol/gDW/h), and black boxes denote the most effective treatment for
each sample.
867

A.

qPCR Detection Over Time

B.

Metagenomic Abundance Over Time

A.

Change from European to
High-Fiber Diet per Sample

Diet
- European
- High-Fiber

Samples

Butyrate Production
(mmol/gDW/h)

B.

Correlation Heatmap with
FDR-corrected Significance

LP-IR Score *

Insulin **

HOMA-IR *

White Blood Cell Count ***

Absolute Total Neutrophils Count *

Total Protein *

Absolute Lymphocytes Count *

Globulin **

A/G ratio *

Beta coefficient