# Development of an AI-enabled predictive model to identify the 'sick child' at a pediatric telemedicine and medication delivery service in Haiti

## Authors
Ben J. Brintz(PhD)[1†], Molly B. Klarman(MPH)[2†*], Youseline Cajusma(BSN)[2], Lerby Exantus(MD)[3], Jude Ronald Beausejour(MD)[4], Katelyn E. Flaherty (BS)[2], Valery Madsen Beau de Rochars(MD)[5], Chantale Baril(MD)[3], Eric J. Nelson(MD, PhD)[2,6*]

## Affiliations
[1] Division of Epidemiology, University of Utah, Salt Lake City, UT, USA
[2] Departments Pediatrics and Environmental and Global Health, University of Florida, Gainesville, FL, USA
[3] Université d'État d'Haiti- Faculté de Médecine et de Pharmacie, Port-au-Prince, Haiti
[4] Independent clinical practice and affiliation.
[5] Department of Health Services Research, Management and Policy, University of Florida, Gainesville, FL, USA
[6] Department of Environmental and Global Health, University of Florida, Gainesville, FL, USA

[†] Co-first authorship
[*] Corresponding author

## Corresponding authors
Eric J Nelson, MD PhD MS FAAP
2055 Mowry Road, Emerging Pathogens Institute, University of Florida
Gainesville, FL 32611, (352) 294-8536 Email: eric.nelson@ufl.edu

Molly B Klarman, MPH
2055 Mowry Road, Emerging Pathogens Institute, University of Florida
Gainesville, FL 32611, (352) 294-8536 Email: mklarman@ufl.edu

**ABSTRACT**

**Background.** One of the most difficult challenges in pediatric telemedicine is to

accurately discriminate between the 'sick' and 'not sick' child, especially in resource-

limited settings. Models that flag potentially 'sick' cases for additional safety checks

represent an opportunity for telemedicine to reach its potential. However, there are

critical knowledge gaps on how to develop such models and integrate them into

electronic clinical decision support (eCDS) tools.

**Methods.** To address this challenge, we developed a study design that utilized data

from paired virtual and in-person exams at a telemedicine and medication delivery

service (TMDS) in Haiti. Providers were allowed to mark respondent data as potentially

unreliable. Artificial intelligence /machine learning (XGBoost) was applied to analyze

paired data from 683 participants across three consecutive implementation studies.

Model derivation focused on identifying 'sick' patients (not-mild) and those requiring

escalation. An ensemble method, based on gradient boosted decision-trees, was used

given the limited sample size. The area under the receiver operating characteristic

curve (AUC) was the primary outcome measure.

**Results.** A total of 683 paired records were available for this secondary analysis from

2225 participants enrolled. The median age was 15 months and 47% were female. For

prediction of a 'sick' child, we found an AUC of 0.82 (95% CI 0.78-0.86) after 5-fold

cross validation; calibration slope and intercept were 1.31 (95%CI:1.09-1.53) and 0.04

(95%CI:-0.14 – 0.23), respectively. For prediction of escalation, we found an AUC of

0.78 (95%CI:0.74-0.81); calibration slope and intercept were 0.63 (95%CI:0.52-0.74)

and 0.05 (95%CI:0.52-0.74), respectively. Accounting for data marked as potentially

unreliable had mixed effects.

**Interpretation.** These methods and findings offer an innovative and important proof-of-

concept to improve pediatric telemedicine. The models require external validation prior

to eCDS integration and deployment. Once validated, the models are designed to

provide a critical safety check for experienced providers and digitally convey expertise

to new providers.

**Funding.** National Institutes of Health (USA) grants to EJN (R21TW012332;

DP5OD019893), internal funding at UF (Children's Miracle Network), and private

donations.

**RESEARCH IN CONTEXT**

**Evidence before this study**. We conducted two Pubmed searches for reports published in all languages. The first search terms were (telemedicine) AND (artificial intelligence OR machine learning) AND (pediatrics OR paediatrics). The primary search criteria identified 153 publications and reviews were excluded leaving 101 papers all published after 1998. Enumerated results by named subspecialty were neurology (n=2), ophthalmology (n=13), otology (n=3), endocrinology (n=11), cardiology (n=6), pulmonology (n=3), gastroenterology(n=1), dermatology (n=1) and surgery (n=22). Ten of the publications focused on global health or low-middle income countries (LMIC) populations. The second search terms were ((Telemedicine) AND (delivery OR paramedicine) AND (pediatrics OR paediatrics)) AND (global health OR LMIC) which generated 99 publications and 76 papers remained after reviews were excluded, all published after 2015. After manual evaluation of the results from both searches, no publications were identified that fully met the scope of this paper. Examples of telemedicine research for concordance with paired exams does exist [1,2].

**Added value of this study.** To the best of our knowledge, this is the first study that investigated pediatric disease severity prediction using virtual and in-person exams in the context of telemedicine -- for either high or low resourced settings. Therefore, the added value of this study is an innovative and important proof-of-concept to improve telemedicine research and clinical practice beyond the scope of global health.

**Implications of all the available evidence.** Inside the field of global health, there is a need to develop evidence-based approaches to extend care early to pediatric patients who may be isolated by poverty, geography or unrest. This must be done safely and this paper offers an approach to develop and incorporate disease severity prediction models

58    into eCDS tools. In addition, these tools may serve as a welcomed safety check for

59    experienced providers and a method to digitally convey expertise to new providers as

60    these services scale.

## INTRODUCTION

One of the most important and difficult challenges in pediatrics is to correctly

discriminate between the 'sick' and 'not sick' child. This challenge arises in part because

children have a great capacity to maintain normal physiologic function until a tipping

point is reached when children can rapidly decompensate. This ability to mask an

impending crisis is one reason why pediatric training requires mastering how to detect

the subtle early patterns of the 'sick' child. Achieving this level of clinical acumen takes

time and represents one seemingly timeless barrier to providing high-quality pre-

emergency pediatric care at scale.

Telemedicine represents a promising method to improve access to pediatric

care, especially for low-resource settings. However, accurately identifying the 'sick' child

in a telemedicine environment is difficult and there is a lack of pediatric telemedicine

guidelines. The World Health Organization's (WHO) integrated management of

childhood illnesses (IMCI) guidelines were primarily developed for low-resource settings

and are not adapted for telemedicine[3]. The American Academy of Pediatrics (AAP)

developed telemedicine guidelines, however they were designed for high-resource

settings[4]. One additional limitation with the AAP guidelines is that their threshold for

hospital (emergency) referral is low which poses the risk of incurring unnecessary

clinical and financial costs. Essentially, the AAP guidelines operate on a red and green

stoplight workflow that lacks a yellow option which is a key component to cost-

effectiveness[5].

To address these complex challenges, we launched the Improving Access to

Nighttime Care and Treatment (INACT) studies in Haiti[5-9]. We developed a pediatric

telemedicine guideline derived from the WHO IMCI guidelines[3]. The guidelines were

implemented within a telemedicine and medication delivery service (TMDS) called

MotoMeds. The guideline was evaluated for safety by comparing paired virtual and in-

person exams. The TMDS was subsequently configured for scaling by triaging cases

with higher clinical uncertainty to escalated care that might involve hospital referral (red)

or an in-person exam at the household with medication delivery (mostly yellow)[7,8].

Cases with lower clinical uncertainty (mostly green) received medication delivery alone.

Analysis revealed a risk that some patients might not be identified as 'sick' during virtual

exams, despite adherence to the clinical guidelines[6,8,9]. Contributing factors included

parent interpretation of provider questions, rigidity of branching logic within the

guidelines, and non-clinical situational events. These factors are common across

pediatric telemedicine services in both high and low resource settings. They collectively

represent a vulnerability that in select situations parents might falsely be reassured that

a child is 'not sick' based on a virtual exam alone.

We hypothesized that predictive models might address this problem by flagging

cases that need added consideration. When integrated into electronic clinical decision

support (eCDS) tools, these models could serve as a safety check for experienced

providers and digitally convey expertise to new providers. To test this hypothesis, we

used paired virtual and in-person exams from the Haitian INACT study data to derive

and internally validate predictive models to identify cases that require added

consideration. Herein, we demonstrate proofs of concept that data from virtual exams

can be leveraged to identify 'sick' children in a manner that guidelines might miss.

**METHODS**

**Ethics statement.** The INACT studies were reviewed and approved by the University of Florida Institutional Review Board (IRB202002693; IRB201802920, IRB202201220) and the Comité National de Bioéthique (National Bioethics Committee of Haiti; 2021-11; Ref1819-51; Ref1101-41).

**Study design.** This study represents a secondary analysis of pooled data from three prior prospective cohort studies: INACT2-H established the TMDS model and focused on safety, feasibility and evaluation of the clinical guidelines; all non-severe cases had paired virtual and in-person exams[7,10]. INACT3-H positioned the TMDS for scale by reserving in-person exams for cases with higher clinical uncertainty, which mainly included moderate (yellow) cases[11]. Continuing this scalable approach, INACT4-H is an ongoing interrupted time-series study of a digitization of the TMDS clinical guidelines; analysis draws on data from the pre-intervention period.

**Study population and setting.** INACT2-H was conducted in Gressier, Haiti. INACT3-H and INACT 4-H were conducted in both Gressier and Les Cayes, Haiti. The studies were implemented consecutively without interruption. INACT2-H began in September 2019 and concluded in January 2021, followed by INACT3-H, which concluded in September 2022 and the INACT4-H study is ongoing. Gressier has a population of 36,400 and is characterized by rural mountainous regions and more densely populated coastal communities [12]. Les Cayes is approximately 175km west of Gressier. It has a population of 152,000 people [12], and over half live in the urban center. All study periods were marked by political and societal instability with detrimental effects on health and healthcare [13,14].

**Participant recruitment.** The study populations were informed of the TMDS through

radio and print advertisements, plus announcements at schools, clinics and markets.

**Participant inclusion criteria and consent process.** For all studies, children were

eligible to participate if their parent contacted the TMDS regarding their child aged ≤10

years during operational hours (6 PM-5 AM). Written consent was obtained from

parents/guardians, and assent from children ≥7 years, who received an in-person exam.

Parents/guardians whose only contact was a virtual encounter were read a waiver of

documentation of consent over the phone.

**Enrollment.** Enrollment for all three studies was based on call census at the TMDS

which was estimated prior to the INACT2-H and adjusted following each completed

study[10,11].

**Participant incentives and fees.** Across all studies TMDS participants were asked to

pay a fee of 500 Gourdes (5 USD) to contribute to the cost of delivery and medications.

The fee was reduced or waived for those who were unable to pay in full. The funds were

re-invested in the study and the research was not dependent on the fee. The fee was

implemented to avoid incentivizing families to delay daytime care for free nighttime care

and to avoid undercutting daytime providers who charge a fee.

**Implementation.** The TMDS workflow, clinical procedures and TMDS guidelines for

INACT2-H and INACT3-H are described previously[10,11] and the ongoing INACT4-H

protocol is included in appendix 2. The workflow (**Figure 1)** steps are: i) A parent

contacted the call center. ii) The provider gathered information about the child using a

paper case report form (CRF) that included elements of decision support and generated

a treatment plan that consisted of a disease severity classification (mild, moderate, or

severe), disposition (hospital or household), recommended medications and/or fluids,

and advice on follow-up care. iii) Severe cases were immediately referred to the

hospital. iv) For non-severe cases, patients in a delivery zone received a medication

delivery with or without an in-person exam. For INACT2-H, all cases had a paired exam.

For INACT3-H and INACT4-H, only cases with higher clinical uncertainty (mainly

moderate cases) received an in-person exam whereas the remainder received a

medication delivery alone. v) Children residing outside a delivery zone, received a

virtual exam and advice alone. vi) Patients requiring hospital or mandatory clinic follow-

up received a 24-hour follow-up call. All patients received a 10-day call to report the

child's condition and provide feedback.

**Outcome measures**.

*Primary.* The primary outcome measure in this secondary analysis was prediction of

disease severity (mild vs not-mild) based on features collected via virtual exams at the

call center. The primary metric was area under the receiver operating characteristic

curve (AUC). Based on academic standards, a threshold of 0.80 or greater was

considered 'good' performance[15].

*Secondary.* The secondary outcome measure was prediction of the need for escalation

of care (in-person exam at the household or hospital referral).

*Exploratory.* The relative contributions of individual features within the models were

evaluated. Enumerations of cases that would be flagged as non-mild or needing

escalation were estimated from the population based on a given probability generated

by the models.

**Data curation.** We combined data on participants with paired virtual and in-person

exams from the INACT-2H, INACT-3H, and INACT-4H (through August 2023)

studies[10,11]. The date cutoff was chosen because records to this point were complete at

the time model derivation was initiated. Records with missing data for primary outcome

variables (disease severity or disposition) were removed. For all studies, disposition

during in-person exams, and for INACT2-H disposition during virtual exams, was

determined manually based on the clinical guidelines. The severity of cases with missed

'danger signs' during either the virtual or in-person exams were manually reassigned to

severe. 'Danger signs' are clinical indicators (e.g., severe dehydration) outlined in the

clinical guidelines (appendix 3) that signify a case is severe.

Given that data were collected from multiple studies, there were discrepancies

amongst the variables across studies. Moreover, within a study, some questions were

only asked to certain patients based on branching logic. For example, INACT-2H did not

ask whether the parent was concerned about the child's weight and in INACT-3H and

INACT-4H this question was only asked if the child was less than 60 months old. When

appropriate, we used branching logic to fill in some of the missing values. For example,

if the parent did not report diarrhea as a complaint, then the number of diarrheal

episodes in the previous 24 hours was imputed as 0, rather than missing. However,

some instances of missing data could not be resolved through logical imputation, e.g.,

questions that were not asked across all studies. In such cases, we left those values as

missing. The XGBoost's built-in handling procedure which learns the best way to route

missing values during the tree-building process, eliminates the need for explicit

imputation.

Given that virtual exams were conducted over the phone and relied on the parent

correctly hearing and understanding questions, providers were able to document

instances where they felt the information they were receiving might be unreliable by

marking the response as "unsure" on the CRF. In these situations, the question was

omitted when generating the treatment plan. We accounted for 'unsure' responses

using the following methods: For binary variables such as 'blood in stool', we created an

ordinal variable with levels "No", "Yes and unsure" and "Yes". For ordinal variables such

as number of vomiting episodes in the last 24h, we created an "unsure" indicator

variable as a separate feature in the model. For chief complaint, a categorical variable

with more than two categories, we created dummy variables for each level with 0 for all

representing a missing chief complaint.

**Model derivation and validation.** We developed predictive models for two binary

outcomes: 1) Not-mild severity, the primary outcome, was defined as positive for

patients classified as moderate or severe and 2) Escalation of care, defined as positive

if a patient requires an in-person provider visit or referral to emergency services. We

only considered variables that were collected during the virtual exam and those that

were collected during the in-person exam (e.g., oxygen saturation) were excluded. We

tested models that included the "unsure" indicator variables and ordinal variables with

"unsure" as a level, as well as models that excluded these variables where 'yes and

unsure' and 'no and unsure' were treated as 'yes' and 'no', respectively.

We used the XGBoost algorithm, an ensemble approach that uses gradient

boosted decision-trees to derive predictive models for estimating the probability that a

TMDS case would be 'not-mild' or  require escalation. We used five-fold cross-validation

to tune hyperparameters: max depth, $\eta$ (learning rate), subsample, column sample by

tree, lambda, and number of rounds (trees). We considered 100 rounds (or trees) in

each model fit with early stopping if not improved over 10 rounds. Area under the

receiver operating characteristic curve (AUROC or AUC) was used to assess

performance for both cross-validation, model fitting, and importance metrics. Given the

small sample size of our data and the number of features considered (35), we opted to

restrict our analysis to 5-fold cross-validation rather than a separate validation set.

While the model is trained to maximize AUC, we also report the area under the

precision-recall curve (PRAUC) and the calibration of the model which assesses how

well the predicted risk aligns with the observed risk.

We calculated the importance of each predictor using built-in XGBoost output

which calculates the fractional contribution of each predictor to the model performance

(log-loss) based on the total gain in performance when its decision tree splits were

utilized. A higher percentage indicates a more important predictor. Finally, Shapley

Additive Explanations (SHAP) were calculated, a value that expresses the additive

contribution of each feature to the predicted value and is used to explain the direction

and magnitude of a predictor on the outcome dependent on its value.

**RESULTS**

**Participant characteristics.** A total of 2225 cases were enrolled between the three

studies; the median age was 24 months and 47% were female. Given differences in

study designs, there were 683 cases with paired virtual and in-person exams available

for analysis; the median age was 15 months and 47% were female. Additional

participant characteristics are shown (**Table 1**). Features that may have differed

between the virtual (call center) and in-person (household) settings were enumerated.

**Primary outcome.** The models were computed both by accommodating for and not

accommodating for providers' perception of the reliability of responses, or the 'unsure'

designation. When the 'unsure' designation was not considered, the model to predict

the detection of a 'not-mild' status had an AUC of 0.82 (95% CI 0.78-0.86) after 5-fold

cross validation; calibration slope and intercept were 1.09 (95% CI: 0.91-1.26) and 0.16

(95% CI: -0.03 – 0.35), respectively **(Figure 2)**. When the 'unsure' designation was

considered, the model to predict the detection of a 'not-mild' status had an AUC of 0.82

(95% CI 0.78-0.85) after 5-fold cross validation; calibration slope and intercept were

0.91 (95% CI: 0.76 -1.06) and 0.2 (95% CI: 0 – 0.4), respectively (**Figure S1**).

**Secondary outcome.** For prediction of escalation of care that does not consider the

'unsure' designation, the model had an AUC of 0.78 (95% CI 0.74-0.81) after 5-fold

cross validation; the calibration slope and intercept were 0.81 (95% CI: 0.66-0.96) and

0.08 (95%CI: -0.11-0.26), respectively **(Figure 3)**. When considering the 'unsure'

designation, the model had an AUC of 0.78 (95% CI 0.73-0.81) after 5-fold cross

validation; calibration slope and intercept were 0.47 (95% CI:0.39 -0.56) and 0.11

(95%CI: -0.11 – 0.33), respectively (**Figure S2**).

**Exploratory analyses.** SHAP values were used to assess the additive contribution of

each feature to the predicted value of a given model; these values and associated plots

help explain the direction and magnitude of a predictor on the outcome. We show the

top ten most important variables and SHAP values for each model that did not consider

the 'unsure' designation (**Figure S3, Figure S4)**.

Estimated case counts flagged for additional consideration by the DSP models

are shown (**Table 2**). When applying the severity prediction model at a balanced

sensitivity of 75% and specificity of 80%, 11% of cases would be flagged for additional

consideration, of which 38% would be 'true' positives. Similarly, when applying the

escalation prediction model at a balanced sensitivity of 75% and specificity of 71%, 11%

of cases would be flagged for escalation, of which 36% percent would be 'true' based

on in-person.

**DISCUSSION**

Historically, accurately identifying the 'sick' patient early required exceptional bedside

assessment skills combined with an expansive fund of knowledge. How one integrates

technology to improve outcomes and efficiency in a cost-effective manner is today's

challenge. We present a use case specific to telemedicine in resource-limited settings.

Our objective was to develop models that flag potentially 'sick' patients during call

center encounters for additional consideration that might otherwise be missed. To

address this challenge, we developed a study design that utilized data from paired

virtual and in-person exams at a TMDS in Haiti. Using machine learning analytic

techniques, we derived and validated a model that predicted, and therefore

discriminated, patients with moderate or severe disease as well as a model that predicts

if a case needs escalation. Model performance was assessed using the AUC metric,

with results at or above commonly accepted thresholds for inclusion in clinical practice.

The models require external validation prior to integration and evaluation in clinical

trials. Once validated, the models may enable a critical safety check for experienced

providers and a method to digitally convey expertise to new providers.

These findings can be interpreted at three levels: the methodologic proof of

concept, the analytic outcomes, and insights on human-computer interaction. The

formative phase of this research began with an evaluation of pediatric telemedicine

guidelines to understand to what degree the process of identifying the 'sick' child was

evidence-based in a virtual exam environment. There was a paucity of literature in

which guidelines and approaches used gestalt borrowed from in-person exam

experiences to navigate the telemedicine environment. In situations with limited

connectivity and resources, there was a need for a new strategy. Therefore, in the initial

INACT study we developed an approach to conduct paired virtual and in-person exams

on all cases within the delivery zone that did not need immediate hospital referral. We

hypothesized that we could train models using the virtual exam data as predictors and

the in-person exam data as 'truth'. While developed in a low-resource setting, we

challenge teams with similar interests to test this approach in high-resource settings.

The analytic outcomes are difficult to interpret because there is no clear

threshold at which a model is 'proven' accurate and meaningful without incurring

unintended consequences. An AUC threshold of 0.80 or above is generally considered

a reasonable metric to advance to an external validation phase with possible

subsequent deployment. However, initial design work during model derivation exposed

the need for two models. The WHO IMCI guideline is based on a 'stoplight' system of

green (mild), yellow (moderate) and red (severe). In high-risk triage scenarios involving

pediatric patients in a virtual telemedicine environment, an accurate severity

assessment is critical. This was the primary objective and outcome of our study.

However, the first actionable decision following a virtual assessment is deciding whether

the care plan requires escalation through a hospital referral or in-person exam at the

household. Or, whether medication delivery alone with anticipatory guidance is

sufficient. This insight is essential to ensuring clinical safety in telemedicine as it scales.

We explored how probability thresholds for when to take action could be set by

enumerating rates of flagged cases at varying sensitivities and specificities. The next

step is to simulate and model the effects of threshold settings, comparing high

sensitivity, balanced sensitivity and specificity, and high specificity on clinical safety and

monetary costs.

Human-computer interactions within the healthcare space are complex and

evolving. There is an unproductive conversation that AI-enabled eCDS systems will

replace providers. We chose a different approach that recognized the value of both the

provider (biologic intelligence) and the computer (artificial intelligence). Using human

centered design, we developed the TMDS case report form to enable providers to both

record responses (reporter phase) as well as mark those responses that the provider

considers to be unreliable (synthesis phase). This latter step captures a feature of

'biologic intelligence' that is frequently excluded. The clinical guideline follows a

traditional decision tree format, in which variables marked as potentially unreliable were

treated as negative responses. This approach has inherent limitations that the models

can potentially account for. Accordingly, the models were derived with and without

incorporating these data. The results were mixed and did not show a clear advantage,

likely due to limited sample sizes at both the patient and provider levels. Nevertheless,

we consider this an important open question in the field of  human-computer interaction

that warrants further investigation.

These findings should be viewed within the context of the study limitations. The

first limitation relates to a generally small sample size. To address this limitation several

steps were taken that included selecting a subset of predictive variables for model

derivation. This may have created a bias in the models by unintentionally excluding

factors with meaningful effects. Second, the models were derived using datasets pooled

data from three different studies to increase sample size (INACT2-H, INACT3-H,

INACT4-H), however the study population may have changed over the four-year period.

Additionally, differences in study designs between INACT2-H and INACT3/4-H led to

some variation in the types of cases that received paired exams. In INACT2-H, all non-

severe cases had paired exams, whereas in INACT3/4-H, only cases with higher clinical

uncertainty, primarily moderate cases, had paired exams. Although the majority of the

CRF was consistent across studies, variations of certain variables may have affected

model development. Third, the training dataset lacked a balanced distribution of mild,

moderate, and severe cases. This imbalance is partly attributable to the study workflow

which required hospital referral for severe cases identified during virtual exams. As a

result, few severe cases were encountered during in-person exams. Our derivation

methods were selected to account for this limitation. Forth, the decision tree embedded

in the clinical guideline uses a singular triage strategy (green, yellow, red) based on

WHO methods. While simplified, triage is based on both physiologic and logistical

considerations. The models reflect this feature and disaggregating the physiologic and

logistical elements was not possible with the approach taken. This presents a contextual

limitation that may constrain the model's applicability to settings with different

operational or triage workflows.

**CONCLUSION**

This study presents an important proof-of-concept for how to develop evidence-based

pediatric telemedicine guidelines enhanced with predictive models that prioritize safety

for vulnerable populations, especially children in resource limited settings. The models

require external validation prior to eCDS integration. Once validated, the models are

intended to provide a critical safety check for experienced providers and digitally convey

expertise to new providers.

**Abbreviations and acronyms**

TMDS, Telemedicine and medication delivery service; WHO, World Health

Organization; IMCI, Integrated Management for Childhood Illness; CRF, Case report

form; INACT, Improving Nighttime Access to Care and Treatment; IQR, Interquartile

range

**Contributions**

Conceptualization- BJB, MBK, EJN

Data curation- BJB, MBK, YC, KF

Formal analysis- BJB, MBK

Funding acquisition- EJN

Investigation- MBK, YC, LE, JRB, CB

Methodology- BJB, MBK, EJN

Project administration- MBK, EJN, VMBR

Supervision- MBK, YC

Visualization- BJB, MBK, EJN

Writing – original draft- BJB, MBK, EJN

Writing – review & editing- BJB, MBK, EJN, LE, CB, JRB, VMBR, KF, YC

**Declaration of interests.** We declare no competing interests.

**Acknowledgments**. We would like to thank both the study participants and the TMDS

staff for their contributions to the datasets analyzed in this manuscript. We are grateful

to our University of Florida administrator team R Autrey, K Berquist, B Johnson, T Linn,

and N Rushing. We are thankful for the academic support provided by G. Morris at the

Emerging Pathogens Institute, and R. Savani Chair of the Department of Pediatrics, as

well as our colleague T. Becker at the University of Florida for guidance and clinical

guideline reviews. Jason Friesen at Trek Medics International has provided important

guidance and access to logistical software that is used in the operational aspects of

MotoMeds. Finally, we acknowledge that our partnership with the Ministry of Public

Health and Population (Ministère de la Santé Publique et de la Population - MSPP) in

Haiti is invaluable to this work.

**Data sharing**. De-identified individual participant data and the data key that underlie the

results are available in appendix 4.

**Supplementary materials.**

Appendix 2. INACT Study Protocol (INACT4H version)*

Appendix 3. INACT Clinical Guidelines (INACT4H version)*

Appendix 4. De-identified dataset and data key*

Appendix 5. Statistical code*

*AVAILABLE UPON REQUEST AND PROVIDED AT PUBLICATION IN JOURNAL

**FIGURE LEGENDS**

**Figure 1. General workflow of the telemedicine and medication delivery service (TMDS).** The parent of a child who is sick at night contacts the call center. A TMDS provider conducts a virtual exam with questions on clinical context. The provider references clinical guidelines and resources to make and assessment and create a treatment plan. The plan is based on the illness severity assessment which informs the decision to escalate or not escalate the disposition. Escalation includes hospital/ emergency service referral for severe cases, or cases with higher clinical uncertainty (mostly moderate) receive an in-person exam with medication delivery. Cases with low clinical uncertainty (mostly mild) receive medication delivery or advice for home care alone if a delivery is not indicated or feasible. Families outside the delivery zone receive anticipatory advice. All families are contacted by phone at 24-hours and/or 10-days for follow-up. In INACT2-H all non-severe cases residing within the delivery zone received an in-person household exam for research purposes and medication delivery (alone) was not part of the workflow.

**Figure 2. Prediction of disease severity.** Receiver operating characteristic curve. (B) Precision-recall curve. (C) Calibration curve for the disease severity predictive model. In plots A and B, the individual dashed lines represent testing performance of each of the 5-folds and the thick black line represents the combined folds curve. The calibration plot (C) is only based on the combined predictions for all five folds and shows a histogram of prediction density by true outcome (0 – Mild, 1 – Severe) at the bottom. Black circle represents current sensitivity and specificity of provider using the clinical guidelines without the DSP.

**Figure 3. Prediction of need for escalation.** Receiver operating characteristic curve. (B) Precision-recall curve. (C) Calibration curve for the escalation of care predictive model. In plots A and B, the individual dashed lines represent testing performance of each of the 5-folds and the thick black line represents the combined folds curve. The calibration plot (C) is only based on the combined predictions for all five folds and shows a histogram of prediction density by true outcome (0 – No Escalation, 1 – Escalation) at the bottom. Black circle represents current sensitivity and specificity of provider using the clinical guidelines without the DSP.

 **TABLES**

**Table 1.** Characteristics of participants with paired virtual and in-person exams.

| Characteristics of participants from pooled studies (n= 683[a]) | Virtual | In-person |
|---|---|---|
| Participant age; median months (Q1-Q3)[b] | 15 (7-36) | … |
| <2 months (%) | 28 (4) | … |
| 2 months to <2 yrs (%) | 386 (57) | … |
| 2 years to <5 yrs (%) | 174 (25) | … |
| 5 years to 10 yrs (%) | 95 (14) | … |
| Participant sex; female (%) | 318 (47) | … |
| Participant complaints[c] | | |
| Fever | 462 | … |
| Respiratory | 493 | … |
| Gastrointestinal | 364 | … |
| Dermatologic | 158 | … |
| Case severity[d] | | |
| Mild (%) | 429 (63) | 460 (70) |
| Moderate (%) | 236 (35) | 147 (22) |
| Severe (%) | 18 (3) | 53 (8) |
| Disposition[e] | | |
| Virtual exam with delivery (%) | 289 (43) | 413 (60) |
| Virtual exam and escalation to in-person exam with delivery (%) | 362 (52) | 194 (28) |
| Virtual exam and escalation to hospital care (%) | 23 (3) | 54 (8) |
| Virtual exam alone (%) | 9 (1) | 22 (3) |

[a] 14 repeat patients within 30 days were excluded; 1542 out of 2225 (69%) patients did not have
paired exams due to study designs.
[b] (Q1-Q3) = Quartile 1 to quartile 3.
[c] The four most common complaints are represented, multiple complaints per participant were permitted.
Complaints for in-person exams were identified with both parent and nurse input.
[d] 18 and 28 cases with missed danger signs during virtual  and in-person exams respectively were
manually re-assigned as severe.
[e] Disposition for in-person exams for all studies, and for virtual exams for INACT2-H, was determined
manually based on the clinical guidelines.

**Table 2.** Estimated case counts flagged for additional clinical consideration during virtual exams at a TMDS call center

| Assessment /plan decision[b] | Probability | Sensitivity | Specificity | Flagged down[a] n (%) | Flagged down[a] n (%) true | Flagged up[a] n (%) | Flagged up[a] n (%) true |
|---|---|---|---|---|---|---|---|
| Severity level with the DSP | 0.67 | 25 | 99 | 205 (30) | 104 (51) | 7 (1) | 6 (86) |
| | 0.50 | 50 | 92 | 151 (22) | 90 (60) | 36 (5) | 18 (50) |
| | **0.30** | **75** | **80** | **85 (12)** | **62 (73)** | **78 (11)** | **30 (38)** |
| | 0.11 | 90 | 36 | 23 (3) | 16 (70) | 247 (36) | 45 (18) |
| | 0.07 | 95 | 20 | 12 (2) | 8 (67) | 321 (47) | 50 (16) |
| Severity level without the DSP (current) | … | 71 | 77 | … | … | … | … |
| | | | | | | | |
| Disposition decision with the DSP | 0.72 | 25 | 95 | 308 (45) | 178 (58) | 7 (1) | 5 (71) |
| | 0.54 | 50 | 88 | 235 (34) | 154 (66) | 24 (4) | 14 (58) |
| | **0.33** | **75** | **71** | **147 (22)** | **116 (79)** | **74 (11)** | **27 (36)** |
| | 0.11 | 91 | 29 | 42 (6) | 28 (67) | 199 (29) | 50 (25) |
| | 0.07 | 95 | 16 | 23 (3) | 14 (61) | 242 (35) | 55 (23) |
| Disposition decision without the DSP (current) | … | 77 | 55 | … | … | … | … |

[a] Cases identified ('flagged') by the DSP out of 683 total cases as a likely mismatched between the provider assessment at the call center (virtual
exam) and truth as defined by the provider assessment at the household (in-person exam). Percentage values for "n(%)" are calculated based on
the total cases analyzed and those for "n(%) true" are those from the immediate leftward column.
[b] For assessments in which the DSP is used to calculate a probability, the bold text represents values with approximately balanced sensitivity and
specificity. The associated probability can serve as a benchmark to guide decision making using an eCDS. For assessments in without the DSP,
these values represent the current sensitivity and specificity based on use of the clinical guidelines alone.

**REFERENCES**

1. Wagner R, Lima TC, Silva M, et al. Assessment of Pediatric Telemedicine Using Remote Physical Examinations With a Mobile Medical Device: A Nonrandomized Controlled Trial. *JAMA Netw Open* 2023; 6(2): e2252570.

2. Demaerschalk BM, Pines A, Butterfield R, et al. Assessment of Clinician Diagnostic Concordance With Video Telemedicine in the Integrated Multispecialty Practice at Mayo Clinic During the Beginning of COVID-19 Pandemic From March to June 2020. *JAMA Netw Open* 2022; 5(9): e2229958.

3. Handbook:  IMCI Integrated Management of Childhood Illnesses. 4th ed. Geneva, Switzerland: World Health Organization; 2005. ISBN 92 4 154644 1 https://apps.who.int/iris/bitstream/handle/10665/104772/9789241506823_Module-1_eng.pdf?sequence=3 (accessed June 26, 2025).

4. Schmitt BD. Pediatric telephone protocols: office version. 17th edition: Amercian Academy of Pediatrics; 2021.

5. Flaherty KE, Klarman MB, Cajusma Y, et al. A Nighttime Telemedicine and Medication Delivery Service to Avert Pediatric Emergencies in Haiti: An Exploratory Cost-Effectiveness Analysis. *Am J Trop Med Hyg* 2022; 106(4): 1063-71.

6. Garilus FE, Cassion K, Cajusma Y, et al. Pathways to optimize a pediatric telemedicine and medication delivery service: A multi-level qualitative study in Haiti. *medRxiv* 2025.

7. Klarman MB, Chi X, Cajusma Y, et al. Development and evaluation of a clinical guideline for a paediatric telemedicine service in a low-resource setting. *BMJ Paediatr Open* 2024; 8(1).

8.    Klarman MB, Flaherty KE, Chi X, et al. Implementation of a Pediatric

Telemedicine and Medication Delivery Service in a Resource-limited Setting: A Pilot

Study for Clinical Safety and Feasibility. *J Pediatr* 2023; 257: 113304.

9.    Klarman M, Schon J, Cajusma Y, et al. Opportunities to catalyse improved

healthcare access in pluralistic systems: a cross-sectional study in Haiti. *BMJ Open*

2021; 11(11): e047367.

10.    Klarman MB, Flaherty KE, Chi X, et al. Implementation of a pediatric

telemedicine and medication delivery service in a resource-limited setting: A pilot study

for clinical safety and feasibility. *The Journal of Pediatrics* 2022.

11.    Klarman MB, Chi X, Cajusma Y, et al. Evaluation of a Scalable Design for a

Pediatric Telemedicine and Medication Delivery Service: A Prospective Cohort Study in

Haiti. *medRxiv* 2024: 2024.08.16.24312128.

12.    Statistiques Démographiques et Sociales. Populational totale, population de 28

ans et plus ménages et densités estimés en 2015: L'Institut Haitien de Statistique et

d'Informatique, 2015.

13.    Taylor L. Haiti's hospitals plan to close amid fuel shortages. *Bmj* 2022; 379:

o2379.

14.    Severe K, Alcenat N, Rouzier V. Resurgence of Cholera in Haiti amidst

Humanitarian Crises. *N Engl J Med* 2022.

15.    Assessing the Fit of the Model. Applied Logistic Regression; 2000: 143-202.

**Figure 1**

Skin clinical sign

Teledermal assessment

Severity assessment: Severe — Moderate — Mild

Escalation decision:
- Referral to emergency services
- In person visit with medication delivery
- Medication delivery
- Advice for home care

Follow-up call

**FIGURE 2**

**FIGURE 3**