Cost-Effectiveness of Infant and Maternal RSV Immunization Strategies

Javad Taleshi[a,b], Marina Viñeta Paramo[c,d], Allison Watts[c,d], Mark Chilvers[e,f], Jeffrey Man Hay Wong[g], Jolanta Piszczek[h], Lea Separovic[i], Jia Hu[i], Danuta Skowronski[b,i], Pascal M. Lavoie[c,d], Hind Sbihi[a,b]

[a] Data and Analytics Services, BC Centre for Disease Control, Vancouver, BC, Canada.

[b] School of Population and Public Health, Faculty of Medicine, University of British Columbia, Vancouver, BC, Canada.

[c] Department of Pediatrics, University of British Columbia, Vancouver, BC, Canada.

[d] British Columbia Children's Hospital Research Institute, Vancouver, BC, Canada.

[e] Child Health BC, Vancouver, BC, Canada.

[f] Division of Pediatric Respiratory Medicine, BC Children's Hospital, Vancouver, BC, Canada.

[g] Department of Obstetrics and Gynecology, University of British Columbia, Vancouver, BC, Canada.

[h] Faculty of Pharmaceutical Sciences, University of British Columbia, Vancouver, BC, Canada.

[i] Immunization Programs and Vaccine Preventable Diseases Service, BC Centre for Disease Control, Vancouver, BC, Canada.

Abstract

Respiratory syncytial virus (RSV) is a leading cause of lower respiratory tract infections in young children and results in significant healthcare burden and costs. To reduce the impact of RSV in this population, the monoclonal antibody palivizumab has historically been used. Recently, new preventive options have become available, including a longer-acting monoclonal antibody (nirsevimab) and a maternal vaccine (RSVpreF). To compare the cost-effectiveness of various immunization strategies, we developed a discrete-event simulation model using epidemiological and cost data from British Columbia, along with published efficacy estimates. The model simulated a cohort of 100,000 newborns and tracked them up to 24 months. We conducted the analysis from a healthcare system perspective, projecting health outcomes (inpatient and outpatient visits), costs (in Canadian dollars), and quality-adjusted life years (QALYs) under five immunization strategies: (1) palivizumab for high-risk children (reference); (2) nirsevimab for high- and moderate-risk children; (3) in-season maternal RSVpreF vaccination combined with nirsevimab for high-risk children; (4) in-season maternal RSVpreF plus nirsevimab for moderate- and high-risk children; and (5) nirsevimab for all infants. We applied an incremental cost-effectiveness ratio (ICER) threshold of $50,000 per QALY as the willingness-to-pay benchmark for evaluating cost-effectiveness. Compared to strategy 1 (palivizumab), strategy 2 was dominant, improving health outcomes while yielding cost savings. Strategy 3 was also cost saving but was weakly dominated by other options. Strategy 4 was cost-effective, with an ICER less than half the willingness-to-pay threshold. In contrast, strategy 5 was not cost-effective, with an ICER exceeding five times the benchmark value under current assumptions for nirsevimab efficacy and a unit price of $450. These findings support policy recommendations to prioritize a publicly funded seasonal maternal RSVpreF immunization program plus nirsevimab administration for high- and moderate-risk children, as a cost-effective approach for broader reduction in RSV-related healthcare burden in young children.

Keywords: Respiratory Syncytial Virus (RSV), Cost-Effectiveness Analysis (CEA), RSVpreF, nirsevimab, Maternal Immunization, Infant Vaccination Strategy

1. Introduction

RSV is the most common cause of lower respiratory tract illness (LRTI) in young children worldwide [1]. The disease burden is highest in early infancy: nearly all children are infected by RSV by age two, and 1–2% of infants require hospitalization due to RSV infection in the first year of life in high-income countries [2–4]. RSV illness results in significant direct medical costs, with the estimated annual average cost of RSV-related pediatric hospitalizations in Canada reaching $66 million [5]. In the absence of an active pediatric vaccine, efforts to protect young children have relied on passive immunoprophylaxis with the monoclonal antibody palivizumab, given as monthly injections during RSV season to high-risk children (e.g., those born preterm, or with severe chronic lung or cardiac conditions, or immunodeficiency). While palivizumab can reduce severe RSV outcomes in high-risk children, it has notable limitations including high product and administration cost and logistical complexity of its dosing regimen.

In 2024, two new immunization products have become available in Canada for preventing severe RSV in young children: (1) nirsevimab, a long-acting monoclonal antibody given once to protect children during an entire RSV season; and (2) RSVpreF, a prefusion F protein vaccine administered to pregnant individuals, increasing RSV antibody levels in newborns to protect them for six months. The National Advisory Committee on Immunization (NACI) has recommended moving toward a universal, all-infant RSV immunization program in Canada, prioritizing nirsevimab for inclusion in infant programs and considering RSVpreF in pregnant individuals on a case-by-case basis [6]. These recommendations emphasize staged implementation based on supply, cost-effectiveness, and affordability.

Previous evaluations have examined the cost-effectiveness of nirsevimab and maternal RSVpreF immunization strategies within the province of Ontario, Canada [7,8]. However, existing studies have some limitations. The probability of RSV infection in young children is influenced by both age and seasonality [9]. Furthermore, the severity of RSV infection, duration of hospitalization and risk of intensive care unit (ICU) admission also differ in children with co-morbidities [9,10], yet previous studies did not use co-morbidity-adjusted risks and did not account for the joint effect of age and seasonality in modeling RSV risk. Moreover, they relied on 2017 estimates for hospitalization and ICU costs, whereas more recent data are now available [5], providing updated insights into current healthcare expenditures. Additionally, the current negotiated prices for nirsevimab and RSVpreF are nearly half of those assumed in these earlier analyses.

To address these limitations, we developed a discrete-event simulation model that incorporates age- and month-specific RSV infection risks along with co-morbidity-adjusted clinical outcomes. By aligning infection risk with the timing of immunization and accounting for individual vulnerability, the model provides a more granular framework for evaluating the cost-effectiveness of maternal and infant RSV immunization strategies and identifying those that offer the greatest health and economic value. The analysis was also informed by the most current demographic and retrospective healthcare data available from British Columbia, Canada.

2. Methods and data inputs

2.1. Simulation Model

We developed a discrete-event simulation model to track RSV infections and associated outcomes in a birth cohort of 100,000 children. The simulation began on May 1st, immediately after the typical end of the RSV season in British Columbia, and ran for two full RSV seasons, concluding on April 30th. To account for seasonal variation in birth timing, the cohort was divided into 12 monthly sub-cohorts, each representing approximately one-twelfth of all births. Each child was followed from birth until the end of their second RSV season, enabling the model to capture exposure to RSV over two consecutive seasons. This extended time horizon was selected to reflect the potential long-term impact of immunization strategies, especially those that provide protection beyond the first year of life.

The model advanced in monthly time steps, during which individuals transitioned between health states, as illustrated in Figure 1A. All children entered the simulation in a healthy state, which denotes not infected with RSV. In any given month, a child could remain healthy, acquire an RSV

infection, or—rarely—die from non-RSV causes. If infected with RSV, the child could either recover and return to the healthy state or die.

Figure 1 (A) Overview of health states. (B) Structure of treatment pathways for RSV infections.

In the model, an RSV infection can lead to different levels of healthcare utilization, as illustrated in Figure 1B. Children with RSV illness may receive medical care or not, and those receiving such care are further categorized into outpatient vs. inpatient management. Outpatients can be seen either in an emergency department or by a primary care provider, while inpatients are treated in hospital pediatric wards or in the ICU. Preventive interventions (monoclonal antibodies or vaccines) are assumed to reduce the probability of RSV infection and subsequent healthcare utilization, including the likelihood of severe and critical disease requiring pediatric ward and ICU admission.

2.2. Population data input and Hospitalization Modeling

The primary data for this analysis were drawn from a population-based cohort of children under two years of age, born between 2013 and 2023 and residing in British Columbia (BC). This cohort is maintained within the Provincial Health Services Authority's (PHSA) Platform for Analytics and Data (PANDA) [11] and integrates information from provincial registries and administrative datasets,

including hospitalization records and Vital Statistics [12–14]. We stratified the study cohort into three risk groups based on perinatal factors and chronic conditions, consistent with previous analyses of RSV severity during the first two seasons [9]:

- **High-risk:** Young children who would have been eligible for palivizumab under previous BC criteria (e.g., very preterm births or severe chronic medical conditions) [15]. This group represents just over 1% of children and has the highest RSV hospitalization risk.
- **Moderate-risk:** Young children born moderately preterm (28–36 weeks gestation) or those with chronic medical conditions, as defined by the Pediatric Complex Chronic Conditions classification [16], who are not classified as high-risk under the previous BC palivizumab program. Approximately 11% of BC children fall into this group.
- **Low-risk:** The remaining young children (~88%) who are full-term without chronic medical conditions.

Summary statistics for demographic variables used in the simulation are presented in Table 1.

We estimated the monthly RSV-related hospitalization rate among children under two years of age in the study cohort to define the RSV season for the microsimulation. Rates reflect the monthly averages over the 2013–2023 period, excluding data from May 1, 2020 to April 30, 2021, during which RSV activity collapsed due to public health measures implemented during the COVID-19 pandemic. As shown in Figure 2, the RSV season in BC typically spans from early November to the end of April.

Figure 2 Monthly RSV hospitalization rate (per child) among children under two years of age, by calendar month of infection. Rates represent the monthly averages from 2013 to 2023.

We modeled the monthly probability of RSV hospitalization using logistic regression, incorporating age, a key determinant of RSV severity [9], palivizumab prophylaxis status, risk group, and month of

infection. Model specifications and coefficients are detailed in the supplementary material (see Figure S1, Equation S1, and Table S1). The predicted probabilities showed strong agreement with observed age-specific seasonal patterns, as illustrated in Figure 3.

Figure 3 Predicted monthly probability of RSV hospitalization by age for low-risk children, without receiving palivizumab.

For high-risk children who received palivizumab prophylaxis, we adjusted the hospitalization probability based on palivizumab efficacy, reflecting the reduced observed hospitalization rates in this group.

For each risk group, we modeled the probability of ICU admission conditional on hospitalization (Table 1). We also estimated the length of stay for both pediatric ward and ICU admissions. A small monthly background mortality rate was applied, and we assumed that the mortality rate among hospitalized children was equivalent to this background rate, reflecting the very low RSV-specific mortality observed in BC [14]. Each death was associated with a discounted loss of 45.3 QALYs [7].

*Table 1 Clinical risk parameters (BC data). * Upper range value is from literature.*

Risk parameters	Value	Range
Probability of high risk	1.36 %	(1.32 – 1.41)
Probability of low risk	88.07 %	(87.97 – 88.17)
Probability of ICU admission if hospitalized – low risk	10.03 %	(8.93 – 11.20)
Probability of ICU admission if hospitalized – moderate risk	18.99 %	(16.02 – 22.03)
Probability of ICU admission if hospitalized – high risk	31.67 %	(20.37 – 43.55)
Relative risk of ICU admission in second year of life (vs. first year)	0.69	(0.55 – 0.84)
Average length of hospital stay – pediatric ward, low risk	3.44 days	(3.36 – 3.53)
Average length of hospital stay – pediatric ward, moderate risk	4.69 days	(4.41 – 4.96)
Average length of hospital stay – pediatric ward, high risk	8.16 days	(6.68 – 9.97)

Average length of hospital stay – ICU, low risk	4.09 days	(3.74 – 4.47)
Average length of hospital stay – ICU, moderate risk	5.33 days	(4.72 – 6.00)
Average length of hospital stay – ICU, high risk	8.50 days	(4.95 – 13.55)
Background mortality rate (per month)	0.000132	(0.000123-0.00014)
Mortality rate for hospitalized children (per month)	0.000132	(0.000123-0.00192*)

We also used published literature to obtain age-specific rates of medically attended RSV infections, which informed the distribution of RSV cases managed in outpatient settings [17]. Using these data, we estimated the proportion of RSV illnesses resulting in hospital admissions versus outpatient care by child age. These proportions enabled us to calibrate the overall probability of any medically attended RSV illness and the probability of an emergency department visit among outpatient cases.

2.3. Costs and Utilities

The cost-effectiveness analysis was conducted from the healthcare perspective, specifically a universal public payer system, considering only direct medical costs and utility losses experienced by children up to two years with RSV. Costs were assigned to RSV-related healthcare encounters, and immunization programs. Table 2 summarizes the cost inputs and health state utilities (QALY loss) applied in the model. All costs are in Canadian dollars (CAD).

*Table 2 Direct medical costs and QALY losses associated with RSV-related healthcare encounters and immunization. All costs are reported in Canadian dollars (CAD). * Range defined as ±25% of the base value.*

Cost and disutility parameter	Value	Range	Reference
Direct healthcare costs			
ICU (per day)	$4,615	(3461-5769) *	[5]
Pediatric ward (per day)	$2,283	(1712- 2854) *	[5]
Emergency department (per visit)	$342	(257- 428) *	[7,18]
Primary care (per visit)	$ 229	(172- 286) *	[7]
Immunization costs			
RSVpreF and nirsevimab administration cost	$15	(0-30)	[19,20]
RSVpreF maternal vaccine cost per dose	$125	(100- 250)	Estimated
Nirsevimab cost per dose	$450	(400- 950)	Estimated
Palivizumab administration cost	$150	(0- 188*)	Estimated
Palivizumab cost per RSV season	$3,600	(2700- 4500) *	Estimated
QALY loss			
QALY loss per ICU admission	0.0245	(0.0145–0.105)	[8]
QALY loss per pediatric pard admission	0.0169	(0.01–0.0726)	[21,22]
QALY loss per emergency department visit	0.0135	(0.008– 0.0454)	[21,22]
QALY loss per primary care visit	0.00845	(0.005–0.0454)	[21,22]

Costs for hospital care are based on recent Canadian estimates of pediatric hospitalization costs [5]. Outpatient costs represent the estimated 30-day healthcare expenditures associated with an RSV infection not requiring hospitalization. These costs include physician visits, diagnostics, and any short-term treatments within a month of illness.

Unit prices for nirsevimab and RSVpreF are not publicly available and subject to change but we approximated their anticipated current cost as specified in Table 2 plus a $15 administration cost per dose.

Health outcomes were measured in QALYs. We assigned a small decrement in QALYs for each RSV illness, proportional to its severity, based on disutility values reported in prior cost-effectiveness analyses [8].

2.4. Immunization Strategies and Assumptions

The efficacy of nirsevimab and RSVpreF was drawn from phase 3 clinical trial estimates, using efficacy against severe medically attended RSV illness as a proxy for protection against ICU-level outcomes [23,24]. We modeled efficacy of both RSVpreF and nirsevimab as constant for the first five months post-immunization, followed by a linear decline to zero by ten months (see Figure S2 in the supplementary material). This waning pattern is consistent with assumptions made in prior studies [7]. Detailed information on vaccine efficacy and coverage is provided in Table S2 of the supplementary material.

Currently, all high-risk children are eligible for palivizumab prophylaxis in their first RSV season, with some also receiving it in their second season [9]. Under the evaluated strategies, nirsevimab would fully replace palivizumab prophylaxis for high-risk children, covering both their first and, as indicated by previous palivizumab eligibility, second RSV seasons. Moderate- and low-risk children, who currently do not receive palivizumab prophylaxis would become eligible for nirsevimab in select strategies, but only during their first RSV season.

We modeled the timing of prophylaxis to maximize protection during the RSV season. If a child was born during the RSV season (Nov–Apr), nirsevimab (or palivizumab) was administered in the birth month. If a child was born outside the season (May–Oct), prophylaxis was given at the season's start in November. Similarly, RSVpreF maternal vaccination was assumed to be given late in pregnancy (around 32–36 weeks gestation) if the child's birth would occur during the season. This seasonal approach aligns with NACI guidance and ensures stronger protection during periods of highest RSV risk.

We compared five immunization strategies in the analysis, as described below (the first being the status quo). Each strategy's target population and timing are defined based on the above criteria:

- **Palivizumab (Current Program):** Only high-risk children born during the RSV season are eligible for palivizumab. This represents the standard of care in BC up to 2024 and serves as the main referent strategy against which alternative options listed below are compared.
- **Nirsevimab (High & Moderate):** All high-risk and moderate-risk children are eligible for nirsevimab. Children born before the RSV season receive nirsevimab at the start of the season, while those born during the season are immunized at birth.
- **RSVpreF + Nirsevimab (High):** Maternal RSVpreF vaccination is offered to pregnant individuals with expected delivery during the RSV season. In addition, high-risk children receive nirsevimab irrespective of maternal vaccination status.
- **RSVpreF + Nirsevimab (High & Moderate):** Maternal RSVpreF vaccination is offered to pregnant individuals with expected in-season births. Additionally, all high-risk children are

eligible for nirsevimab regardless of maternal vaccination status, while moderate-risk children are eligible only if they are not protected through maternal vaccination.

- **Nirsevimab (All-infant):** All children, regardless of risk status, are eligible for nirsevimab. Young children born before the RSV season are immunized at the start of the season, while those born during the season receive nirsevimab at birth.

We assumed coverage rates for these immunization programs based on emerging data (see Table S2 in the supplementary material). For maternal RSV vaccination, we assumed about 65% coverage of eligible pregnancies, reflecting uptake in a newly recommended maternal vaccine program [25]. For nirsevimab, we assumed 70% coverage among eligible low-risk and moderate-risk children [8], and 100% uptake among young children identified as high-risk based on observed experience with palivizumab acceptance rates.

For each strategy, the simulation was run to tally outcomes over the cohort's first two RSV seasons. Key outcomes included the number of RSV-related medical visits (primary care visits, emergency department visits), hospitalizations (ward and ICU), total costs (healthcare costs plus immunization-related costs), total QALYs, and ICERs relative to the status quo.

2.5. Input parameters for sensitivity analyses

To assess the robustness of our base-case findings, we conducted a series of sensitivity analyses using net monetary benefit (NMB) as the primary outcome. NMB is defined as:

$$NMB \ = \ (\text{Incremental QALYs} \times \text{WTP}) - \text{Incremental Cost}$$

where WTP represents the willingness-to-pay threshold, set at $50,000 per QALY in this analysis. An intervention demonstrating a positive NMB is thus considered cost-effective at the specified WTP threshold.

We performed one-way sensitivity analyses by varying individual input parameters across their plausible ranges, focusing on the strategy identified as most favorable in the base-case analysis. For demographic and clinical parameters, the ranges were generally derived from 95% confidence intervals of estimates informed by provincial administrative data. An exception was the mortality rate for hospitalized children, for which the upper bound (0.00192) was based on published literature estimates [8]. For cost parameters, we varied inputs by ±25% around their base-case values unless more reliable estimates were available. Utility inputs were sourced from the literature, with ranges reflecting reported uncertainty. A complete listing of parameter ranges and sources is provided in Tables 1 and 2.

In addition to the one-way analysis, we conducted a two-way sensitivity analysis to evaluate the combined effect of varying the unit prices of nirsevimab and RSVpreF across all strategies. A broader range of prices was selected to identify the price combinations under which each strategy yields the highest net monetary benefit (NMB) relative to the alternatives.

3. Results

3.1. Health Outcomes and cost

We simulated all five immunization strategies and estimated RSV-related health outcomes for the birth cohort under each scenario. Table 3 summarizes the projected number of RSV cases requiring different levels of healthcare utilization, along with the associated costs and QALYs for each strategy.

Table 3 RSV-related health outcomes and costs for a simulated cohort of 100,000 infants through the end of the second RSV season, by immunization strategy. The health outcomes include ICU admissions, Pediatric Ward admissions, Emergency Department visits, Primary Care visits, and total QALYs.

Strategies	Total Cost	Vaccine Cost	ICU	PW[1]	ED[2]	PC[3]	Total QALY
Palivizumab (Current)	$22,401,120	$2,942,531	137	1044	7510	19047	143733
Nirsevimab (High & Mod)	$21,242,896	$4,119,551	111	956	6948	17504	143756
RSVpreF + Nirsevimab (High)	$21,869,408	$5,166,464	107	894	6867	17411	143759
RSVpreF + Nirsevimab (High & Mod[4])	$23,071,289	$7,481,164	95	846	6570	16565	143772
Nirsevimab (All-infant)	$45,327,835	$32,799,077	74	647	5437	13347	143818

The current strategy using palivizumab for high-risk children results in the highest RSV-related healthcare burden among the evaluated options. Expanding coverage with nirsevimab to include moderate-risk children reduces hospitalizations and overall costs, yielding health gains at a lower total cost.

Combining maternal RSVpreF vaccination for in-season births with nirsevimab for high-risk children offers further improvements in health outcomes, with only a modest increase in total cost. Adding moderate-risk children to this combined strategy reduces the disease burden even more but raises overall expenditures above the current program.

An all-infant nirsevimab strategy results in the largest health benefits, substantially lowering RSV-related disease burden. However, it also incurs significantly higher vaccination costs, making it the most expensive option evaluated.

---

[1] Pediatric Ward
[2] Emergency Department
[3] Primary Care
[4] Moderate

3.2. Cost-Effectiveness of Immunization Program Options

Table 4 presents the incremental costs, QALYs, and ICERs (cost per QALY gained) for each strategy compared to baseline (palivizumab strategy).

Table 4 Incremental cost-effectiveness of RSV immunization strategies (vs. current program).

Strategies	Incremental Cost	Incremental QALY	ICER
Nirsevimab (High & Mod)	-$1,158,224	23	Dominant
RSVpreF + Nirsevimab (High)	-$531,712	26	Dominant
RSVpreF + Nirsevimab (High & Mod)	$670,168	38	$ 17,611
Nirsevimab (All-infant)	$22,926,715	84	$ 271,651

Two of the evaluated strategies emerge as dominant, meaning they both improve health outcomes and reduce healthcare costs relative to the current program. Specifically, replacing palivizumab with nirsevimab for high-risk and moderate-risk children generates savings of approximately $1.2 million and yields an additional 23 QALYs. Implementing maternal RSVpreF vaccination for in-season births combined with nirsevimab for high-risk children is also cost-saving. Although the net savings are smaller (around $532,000), this strategy yields a slightly larger health benefit (gaining 26 QALYs).

The combined strategy of RSVpreF vaccination for in-season births plus nirsevimab for both high- and moderate-risk children results in an incremental cost of approximately $670,000 and generates an additional 38 QALYs compared to the current program. With an ICER of about $18,000 per QALY gained, this strategy can be considered cost-effective, requiring additional investment but delivering health gains within conventional willingness-to-pay standards. In contrast, the all-infant nirsevimab strategy is not cost-effective under current model assumptions, albeit providing the greatest health benefits. Its ICER is approximately $272,000 per QALY, exceeding conventional thresholds for willingness to pay by more than five-fold.

Figure 4 presents the sequential cost-effectiveness analysis comparing all strategies to each other and to the current standard of care, from the health system perspective. Strategies not on the efficient frontier are considered dominated. The nirsevimab strategy for high- and moderate-risk children is cost-saving relative to the current program and lies on the efficient frontier. In contrast, the maternal RSVpreF plus high-risk nirsevimab strategy is weakly dominated. The combined maternal and child strategy (targeting both high- and moderate-risk groups) is also on the efficient frontier, with a positive ICER within the acceptable willingness-to-pay threshold.

Figure 4 Results of the sequential cost-effectiveness analysis comparing all RSV prevention strategies to each other and to the standard of care, from the health system perspective.

3.3. Sensitivity analysis

We conducted additional analyses to assess the robustness of our base-case results. The one-way sensitivity analysis assessed the impact of uncertainty in key parameters on the cost-effectiveness of the combined strategy—maternal RSVpreF vaccination for in-season births plus nirsevimab for high- and moderate-risk children—compared to the current baseline strategy (palivizumab for high-risk children only). Figure 5 illustrates how NMB varies when each parameter is adjusted across its range while holding all others constant. The most influential parameters were RSVpreF efficacy and unit costs of RSVpreF and nirsevimab (see Figure S3 in the supplementary material for a visual summary of parameter influence). Other key parameters included the efficacy of palivizumab and nirsevimab, as well as the per-day pediatric ward cost. The strategy was found to be no longer cost-effective if the price of nirsevimab exceeds $600, the price of RSVpreF exceeds $160, or the efficacy of RSVpreF drops below 30%.

Figure 5 One-way sensitivity analysis for the combined maternal RSVpreF and infant nirsevimab (High & Moderate) strategy. Each parameter was varied independently across its plausible range, while all other parameters remained constant. The outcome is Net Monetary Benefit (NMB), expressed in thousands of Canadian dollars.

Additionally, we performed a two-way sensitivity analysis to explore the simultaneous variation of RSVpreF and nirsevimab unit costs. Figure 6 indicates the immunization strategy yielding the highest NMB across a range of plausible price combinations (nirsevimab: $0–$1000 per dose; RSVpreF: $0–$300 per dose). For most price scenarios, the highest NMB was achieved by either (a) maternal RSV vaccination for in-season births combined with nirsevimab for high-risk children, or (b) nirsevimab for both high- and moderate-risk children. The fully combined strategy—maternal vaccination plus nirsevimab for both high- and moderate-risk children— generated the highest NMB within only a narrow price range. Notably, for all-infant nirsevimab to yield the highest NMB, its price would need to drop below approximately $110 per dose, well below current estimates.

Figure 6 Two-way sensitivity analysis varying the unit prices of nirsevimab and RSVpreF. Each region on the plot indicates the strategy yielding the highest NMB given a specific combination of nirsevimab and RSVpreF prices, including the current standard of care for reference.

4. Discussion

Our evaluation provides compelling evidence that targeted RSV immunization strategies offer significantly greater economic value than both the historical palivizumab program and an all-infant prophylaxis approach. Specifically, replacing the current palivizumab regimen, which is limited to high-risk children, with nirsevimab for both high- and moderate-risk children led to improved health outcomes while reducing overall healthcare expenditures. The combined maternal plus child strategy for these risk groups yielded substantial health gains, with an ICER well below the conventional $50,000/QALY willingness-to-pay threshold, in fact, less than half that benchmark. In contrast, all-infant nirsevimab immunization was not cost-effective under base-case assumptions, yielding an ICER of approximately $270,000/QALY.

These findings partially align with the current direction of national immunization policy in Canada. NACI has recently recommended a staged expansion of RSV immunization programs, beginning with high-risk infants and working toward an all-infant nirsevimab program for all infants, while suggesting maternal vaccination be considered on a case-by-case basis [6]. This approach reflects constraints related to supply, cost-effectiveness, and fiscal sustainability. However, our analysis diverges from NACI's recommendations in two key areas: maternal RSVpreF immunization and universal nirsevimab. First, we found that combining maternal vaccination with nirsevimab for high-

and moderate-risk infants is cost-effective. Second, our analysis indicates that universal infant nirsevimab immunization, as envisioned by NACI, would only be economically justified if the price dropped substantially. Thus, while our results support a targeted and phased rollout, they do not justify a shift to universal prophylaxis at current prices.

Our study builds on prior economic evaluations of RSV immunization in Canada and internationally. Shoukat et al. employed a similar simulation-based approach based on data from Ontario, and concluded that all-infant nirsevimab prophylaxis would only be cost-effective if the unit price fell to roughly $290 per dose or lower [7]. They found that a strategy combining maternal RSV vaccination with targeted nirsevimab for high-risk infants achieved substantial reductions in infant mortality (~76–85%) and was more economically efficient than all-infant immunization. These findings parallel our own, with both studies highlighting that targeted strategies can provide nearly equivalent health benefits at a lower cost. Similarly, Shin et al. used a static decision tree model to assess an all-infant nirsevimab program across Canadian infants, reporting that such a strategy would be cost-effective if the price per dose remained below approximately $550 under a $50,000/QALY willingness-to-pay threshold [26]. Their model projected large reductions in RSV-related healthcare events and direct costs; however, the inputs for RSV mortality and intensive care utilization were higher than those observed at the population level in BC. Another national-level analysis by Gebretekle et al. examined multiple immunization strategies across the first RSV season and found that neither an all-infant nirsevimab approach nor a year-round maternal RSVpreF program were cost-effective at list prices [8]. Their model identified seasonal nirsevimab administration with "catch-up" dosing for infants born just before the RSV season as the most economically favorable strategy, especially when focused on moderate- and high-risk infants. Notably, the list prices assumed for nirsevimab and RSVpreF in their analysis were nearly double the prices used in our study, which likely contributed to their more conservative cost-effectiveness conclusions for broader strategies. Their findings are particularly notable in remote and high-cost settings, where ICERs were as low as $5,700 per QALY. These Canadian studies, together with ours, support the conclusion that risk-based targeting of RSV immunoprophylaxis maximizes value for money.

Although these studies assumed higher mortality rates among hospitalized infants, in-hospital case fatality attributable to RSV remains very low in high-income countries [1,27], consistent with BC's recent experience of approximately one RSV-associated infant death every 2–3 years [10]. This low mortality likely reflects the impact of longstanding preventive programs (e.g., palivizumab for high-risk infants), advanced pediatric care, and potentially lower viral transmission due to demographic or geographic factors [9]. As such, models assuming uniformly high mortality among hospitalized infants may overestimate the number of deaths averted by immunization. To avoid overstating the potential life-saving impact and cost-effectiveness of RSV interventions, we applied a conservative baseline mortality assumption.

International analyses also reinforce the prioritization of targeted RSV immunization and highlight the influence of health system context. In the United States, where both nirsevimab and RSVpreF have recently been authorized, Averin et al. evaluated cost-effectiveness under multiple program designs. Their analysis showed that a year-round maternal vaccination strategy would prevent about 12% of medically attended RSV cases only yielding an ICER of $90,000 USD per QALY [28]. By contrast, a seasonal maternal program—targeting only pregnancies due during the RSV season—

was found to be cost-saving from a societal perspective despite preventing fewer total cases. These results support a seasonal maternal immunization strategy, emphasizing the importance of targeting protection to periods of peak vulnerability. In Europe, Getaneh et al. assessed RSV prevention strategies in six countries and found that seasonal all-infant immunization with a catch-up dose was generally the most cost-effective strategy when the prophylaxis price was low (€50 per dose) [29]. While maternal vaccination had more limited impact on infant outcomes, it became the preferred strategy in specific contexts (e.g., Veneto, Italy) or under more generous willingness-to-pay thresholds.

From a methodological standpoint, our study introduces several important innovations relative to earlier work. First, we addressed a key limitation in prior Canadian models by incorporating both age and seasonality into RSV risk estimation. Risk of infection and severity is highest during the first months of life and peaks in winter, but earlier models did not fully capture this interaction. Previous studies also did not consider that the severity of RSV illnesses is not uniform across children's groups, and that children with co-morbidities require significantly longer hospitalizations stays and a higher proportion of them required ICU admissions [9]. By implementing a discrete-event simulation with monthly time steps and stratifying children by birth month, we were able to apply joint age–season risk profiles when estimating probabilities of hospitalization and ICU admission. This modeling structure allowed for a more precise evaluation of interventions such as in-season maternal vaccination, which is most effective for children born just before or during the RSV season. It also adjusts for the potential additional health gains of immunization in children's groups who are most vulnerable. Second, by following children over two full RSV seasons, our model captures benefits that extend into the second year of life. Third, we used actual RSV outcome data representing the entire BC population, which provide population-level RSV hospitalization and ICU risks. This contrasts with previous studies that relied on tertiary care hospital data drawn from diverse sources and regions, which may not reflect consistent patterns [7,8]. Additionally, the use of current healthcare cost data, including estimates of pediatric ward and ICU costs provide more contemporary estimates of costs, which have increased substantially over the past decade. By contrast, many prior analyses relied on cost estimates from 2017, potentially underestimating the economic burden of severe RSV illness and thus the potential savings from prevention [7,8].

Our sensitivity analyses further bolster the robustness of our findings. One-way analyses of key parameters—including vaccine efficacy, hospitalization costs, and unit prices—showed that the qualitative result remained unchanged: targeted strategies consistently outperformed both the status quo and all-infant immunization. Even under conservative assumptions, the maternal plus moderate/high-risk nirsevimab strategy yielded a higher NMB compared to the status quo. Our two-way sensitivity analysis on intervention prices was particularly informative, identifying the price combinations at which the preferred strategy would shift. The cost threshold for nirsevimab for all-infant programs to be considered cost-effective is consistent with findings from Gebretekle and Shoukat. This alignment across studies enhances the credibility of our conclusions and affirms the central role of price and risk stratification in determining the value of RSV immunization strategies.

Nevertheless, our analysis has some limitations. First, we assumed a simplified waning pattern for immunity: constant protection for five months post-immunization, followed by linear decline to zero by ten months. While this assumption aligns with clinical trial follow-up durations, real-world immunogenicity may differ. If protection extends beyond ten months, the cost-effectiveness of

interventions would be underestimated in our model. Conversely, if immunity wanes more rapidly, particularly in specific subgroups, our results may overstate benefits. More empirical data on real-world durability of RSVpreF and nirsevimab will be essential to refine future models. Second, we hypothesized a higher uptake rate for infant nirsevimab prophylaxis than for maternal RSVpreF vaccination, whereas emerging Canadian data indicate that families may prefer maternal immunization [30]. This preference could affect the relative cost-effectiveness of combined maternal-plus-infant immunization strategies.

In addition, our model used a static cohort design and did not account for indirect effects on RSV transmission. While static models assume each child's infection risk is independent of others, dynamic models allow for herd effects, age shifts in incidence, and broader population-level impacts. Previous work has shown that dynamic transmission models often predict different patterns of benefit. For instance, Li et al. observed that infant-focused immunization may delay, rather than prevent, some RSV infections, shifting disease burden to older children [31]. Similarly, Hodgson et al. used a dynamic transmission model in the UK and found that while large-scale RSV immunization programs can reduce overall virus circulation, they may also lead to indirect effects— such as changes in the age distribution of infections or transmission patterns—that must be carefully considered in policy decisions [32]. Our static model likely overestimates the total number of infections averted, but severe outcomes in children (hospitalizations and deaths) are more robust to model choice. Since our analysis focuses on cost and QALYs from the healthcare payer perspective over a two-year child horizon, we expect the core conclusions about cost-effectiveness of targeted strategies to remain valid.

Finally, we adopted a health system perspective, excluding societal costs such as caregiver productivity losses, travel expenses, and the psychosocial impact on families. This is a conservative approach that likely underestimates the full value of immunization. RSV-related hospitalizations impose substantial non-medical costs, particularly in rural or remote regions where medical evacuations are frequent and costly. Including these broader costs—as done in Gebretekle et al. [8] and Getaneh et al. [29]—would likely make prophylactic interventions appear even more cost-effective. Our QALY estimates also captured only the health-related utility loss in children, not the quality-of-life burden on caregivers or potential long-term morbidity. Including these outcomes in future models would provide a more comprehensive assessment of value.

5. Conclusion

This study demonstrates that replacing the current palivizumab program with nirsevimab for high- and moderate-risk children is both cost-saving and more effective in preventing RSV-related illness. Adding seasonal maternal RSVpreF vaccination to this strategy yields additional health benefits at an acceptable incremental cost, making it a cost-effective next step if resources allow. In contrast, all-infant immunization with nirsevimab is not economically justified without a substantial price reduction. These findings support a phased approach to RSV immunization, starting with targeted nirsevimab use and expanding to maternal vaccination, aligning with national recommendations and offering a pathway to optimize both health outcomes and healthcare resource use.

CRediT authorship contribution statement

Javad Taleshi: Conceptualization, Methodology, Software, Formal analysis, Data Curation, Writing - Original Draft, Writing - Review & Editing, Supervision. **Marina Viñeta Paramo:** Software, Formal analysis, Data Curation, Writing - Review & Editing. **Allison Watts:** Data Curation, Writing - Review & Editing. **Mark Chilvers** Formal analysis, Writing - Review & Editing. **Jeffrey ManHay Wong** Formal analysis, Writing - Review & Editing. **Jolanta Piszczek** Formal analysis, Writing - Review & Editing. **Lea Separovic** Formal analysis, Writing - Review & Editing. **Jia Hu** Formal analysis, Writing - Review & Editing. **Danuta Skowronski** Formal analysis, Writing - Review & Editing. **Pascal M. Lavoie:** Conceptualization, Formal analysis, Writing - Review & Editing, Supervision. **Hind Sbihi:** Conceptualization, Methodology, Formal analysis, Data Curation, Writing - Review & Editing, Supervision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Access to data provided by the Data Stewards is subject to approval but can be requested for research projects through the Data Stewards or their designated service providers. The following data sets were used in this study: Client Roster (CR), Discharge Abstracts Database (DAD), BC Vital Statistics (VS) [12–14]. All inferences, opinions, and conclusions drawn in this publication are those of the author(s), and do not reflect the opinions or policies of the Data Steward(s).

Reference

1. Li Y, Wang X, Blau DM, Caballero MT, Feikin DR, Gill CJ, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in children younger than 5 years in 2019: a systematic analysis. The Lancet. 2022 May 28;399(10340):2047–64.

2. Borchers AT, Chang C, Gershwin ME, Gershwin LJ. Respiratory syncytial virus—a comprehensive review. Clin Rev Allergy Immunol. 2013;45:331–79.

3. Johannesen CK, van Wijhe M, Tong S, Fernández LV, Heikkinen T, van Boven M, et al. Age-specific estimates of respiratory syncytial virus-associated hospitalizations in 6 European countries: a time series analysis. J Infect Dis. 2022;226(Supplement_1):S29–37.

4. Curns AT, Rha B, Lively JY, Sahni LC, Englund JA, Weinberg GA, et al. Respiratory syncytial virus-associated hospitalizations among children< 5 years old: 2016 to 2020. Pediatrics. 2024;153(3).

5. Vadlamudi NK, Gomes K, Chow E, Bourdeau M, Embree J, Halperin SA, et al. Healthcare Costs and Resource Utilization for Acute Respiratory Syncytial Virus Pediatric Hospitalizations in

Canada [Internet]. Rochester, NY: Social Science Research Network; 2025 [cited 2025 Jun 2]. Available from: https://papers.ssrn.com/abstract=5199070

6. National Advisory Committee on Immunization (NACI). Statement on the prevention of respiratory syncytial virus disease in infants. Public Health Canada; 2024.

7. Shoukat A, Abdollahi E, Galvani AP, Halperin SA, Langley JM, Moghadas SM. Cost-effectiveness analysis of nirsevimab and maternal RSVpreF vaccine strategies for prevention of Respiratory Syncytial Virus disease among infants in Canada: a simulation study. Lancet Reg Heal. 2023;28.

8. Gebretekle GB, Yeung MW, Ximenes R, Cernat A, Simmons AE, Killikelly A, et al. Cost-effectiveness of RSVpreF vaccine and nirsevimab for the prevention of respiratory syncytial virus disease in Canadian infants. Vaccine. 2024;42(21):126164.

9. Viñeta Paramo M, Watts AW, Bone JN, Sadarangani M, Bettinger JA, Seaton C, et al. RSV Hospital Admissions During the First 2 Seasons Among Children With Chronic Medical Conditions. JAMA Netw Open. 2025 Jul 8;8(7):e2519410.

10. Viñeta Paramo M, Ngo LPL, Abu-Raya B, Reicherz F, Xu RY, Bone JN, et al. Respiratory syncytial virus epidemiology and clinical severity before and during the COVID-19 pandemic in British Columbia, Canada: a retrospective observational study. Lancet Reg Health Am. 2023 Sep;25:100582.

11. Wilton J, Abdulmenan J, Chong M, Becerra A, Najmul Hussain M, Harrigan SP, et al. Cohort profile: the British Columbia COVID-19 Cohort (BCC19C)—a dynamic, linked population-based cohort. Front Public Health. 2024;12:1248905.

12. British Columbia Ministry of Health [creator]. Client Roster (Client Registry System/Enterprise Master Patient Index) [Internet]. British Columbia Ministry of Health [publisher]. Data Extract. Ministry of Health (2021); 2020. Available from: https://www2.gov.bc.ca/gov/content/health/health-forms/online-services

13. Canadian Institute of Health Information [creator]. Discharge Abstract Database (Hospital Separations) [Internet]. British Columbia Ministry of Health [publisher]. Data Extract. MOH (2020); 2020. Available from: http://www2.gov.bc.ca/gov/content/health/conducting-health-researchevaluation/data-access-health-data-central

14. BC Ministry of Health [creator]. Vital Statistics Deaths [Internet]. BC Vital Statistics Agency [publisher]. Data Extract. BC Vital Statistics Agency (2021); 2020. Available from: https://www2.gov.bc.ca/gov/content/health/health-forms/online-services

15. Lavoie PM, Solimano A, Taylor R, Kwan E, Claydon J, Turvey SE, et al. Outcomes of Respiratory Syncytial Virus Immunoprophylaxis in Infants Using an Abbreviated Dosing Regimen of Palivizumab. JAMA Pediatr. 2016 Feb 1;170(2):174–6.

16. Feudtner C, Feinstein JA, Zhong W, Hall M, Dai D. Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. BMC Pediatr. 2014;14:1–7.

17. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. N Engl J Med. 2009;360(6):588–98.

18. Canadian Institute for Health Information. Hospital spending: Focus on the emergency department [Internet]. CIHI Ottawa, ON; 2020. Available from: chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.cihi.ca/sites/default/files/docu ment/hospital-spending-highlights-2020-en.pdf

19. Papenburg J, Saleem M, Teselink J, Li A, Caouette G, Massé É, et al. Cost-analysis of Withdrawing Immunoprophylaxis for Respiratory Syncytial Virus in Infants Born at 33–35 Weeks Gestational Age in Quebec: A Multicenter Retrospective Study. Pediatr Infect Dis J. 2020 Aug;39(8):694.

20. Abu-Raya B, Coyle D, Bettinger JA, Vaudry W, Halperin SA, Sadarangani M. Pertussis vaccination in pregnancy in Canada: a cost-utility analysis. Can Med Assoc Open Access J. 2020 Oct 1;8(4):E651–8.

21. Glaser EL, Hariharan D, Bowser DM, Gervasio RM, Rowlands KR, Buckley L, et al. Impact of Respiratory Syncytial Virus on Child, Caregiver, and Family Quality of Life in the United States: Systematic Literature Review and Analysis. J Infect Dis. 2022 Aug 15;226(Supplement_2):S236–45.

22. Régnier SA. Respiratory syncytial virus immunization program for the United States: Impact of performance determinants of a theoretical vaccine. Vaccine. 2013 Sep 13;31(40):4347–54.

23. Simões EA, Madhi SA, Muller WJ, Atanasova V, Bosheva M, Cabañas F, et al. Efficacy of nirsevimab against respiratory syncytial virus lower respiratory tract infections in preterm and term infants, and pharmacokinetic extrapolation to infants with congenital heart disease and chronic lung disease: a pooled analysis of randomised controlled trials. Lancet Child Adolesc Health. 2023;7(3):180–9.

24. Kampmann B, Madhi SA, Munjal I, Simões EA, Pahud BA, Llapur C, et al. Bivalent prefusion F vaccine in pregnancy to prevent RSV illness in infants. N Engl J Med. 2023;388(16):1451–64.

25. Public Health Agency of Canada. Results of the survey on vaccination during pregnancy 2021. Ottawa: Government of Canada; 2022.

26. Shin T, Lee JK, Kieffer A, Greenberg M, Wu J. Health economic evaluation of implementing a universal immunization program with nirsevimab compared to standard of care for the prevention of respiratory syncytial virus disease in Canadian infants. Hum Vaccines Immunother. 2025 Dec;21(1):2480875.

27. Hansen CL, Chaves SS, Demont C, Viboud C. Mortality Associated With Influenza and Respiratory Syncytial Virus in the US, 1999-2018. JAMA Netw Open. 2022 Feb 28;5(2):e220527.

28. Averin A, Quinn E, Atwood M, Weycker D, Shea KM, Law AW. Cost-effectiveness of bivalent respiratory syncytial virus Prefusion F (RSVpreF) maternal vaccine among infants in the United States. Vaccine. 2025 Jun 11;58:127191.

29.   Getaneh AM, Li X, Mao Z, Johannesen CK, Barbieri E, van Summeren J, et al. Cost-effectiveness of monoclonal antibody and maternal immunization against respiratory syncytial virus (RSV) in infants: Evaluation for six European countries. Vaccine. 2023 Feb 24;41(9):1623–31.

30.   McClymont E, Wong JMH, Forward L, Blitz S, Barrett J, Bogler T, et al. Acceptance and preference between respiratory syncytial virus vaccination during pregnancy and infant monoclonal antibody among pregnant and postpartum persons in Canada. Vaccine. 2025 Mar 19;50:126818.

31.   Li X, Hodgson D, Flaig J, Kieffer A, Herring WL, Beyhaghi H, et al. Cost-Effectiveness of Respiratory Syncytial Virus Preventive Interventions in Children: A Model Comparison Study. Value Health. 2023 Apr 1;26(4):508–18.

32.   Hodgson D, Wilkins N, Leeuwen E van, Watson CH, Crofts J, Flasche S, et al. Protecting infants against RSV disease: an impact and cost-effectiveness comparison of long-acting monoclonal antibodies and maternal vaccination. Lancet Reg Health – Eur [Internet]. 2024 Mar 1 [cited 2025 Jun 18];38. Available from: https://www.thelancet.com/journals/lanepe/article/PIIS2666-7762(23)00248-X/fulltext