TITLE

Genetic insights into foveal morphology and its associations with

pigmentation and age-related macular degeneration

AUTHOR LIST & AFFILIATIONS

David J. Green[1], David Romero-Bascones[2], Thomas H. Julian[1,3-5], Sofia

Torchia[5], Heer N.V. Joisher[6-8], UK Biobank Eye and Vision Consortium, Unai

Ayala[2], Maitane Barrenechea[2], Jay E. Self[9,10], Graeme C. Black[1,11], Tomas

Fitzgerald[5], Ewan Birney[5], Constance L. Cepko[6-8], Joseph Carroll[12,13], Panagiotis

I. Sergouniotis[1,3,5,11]

[1] Division of Evolution, Infection and Genomics, School of Biological Sciences,

Faculty of Biology, Medicine and Health, University of Manchester, Manchester,

UK.

[2] Biomedical Engineering Department, Faculty of Engineering (MU-ENG),

Mondragon Unibertsitatea, Mondragón, Spain.

[3] Manchester Royal Eye Hospital, Manchester University NHS Foundation Trust,

Manchester, UK.

[4] Christabel Pankhurst Institute, The University of Manchester, Manchester, UK

[5] European Molecular Biology Laboratory, European Bioinformatics Institute

(EMBL-EBI), Wellcome Genome Campus, Cambridge, UK

[6] Department of Genetics, Blavatnik Institute, Boston, MA, USA

[7] Department of Ophthalmology, Harvard Medical School, Boston, MA, USA

[8] Howard Hughes Medical Institute, Chevy Chase, MD, USA

[9] Clinical and Experimental Sciences, Faculty of Medicine, University of

Southampton, Southampton, UK

[10] Southampton Eye Unit, University Hospital Southampton NHS Foundation

Trust, Southampton, UK

[11] Manchester Centre for Genomic Medicine, Saint Mary's Hospital, Manchester

University NHS Foundation Trust, Manchester, UK

[12] Cell Biology, Neurobiology and Anatomy, Medical College of Wisconsin,

Milwaukee, WI, USA

[13] Ophthalmology and Visual Sciences, Medical College of Wisconsin,

Milwaukee, WI, USA

Correspondence to Panagiotis I. Sergouniotis

(panagiotis.sergouniotis@manchester.ac.uk).

ABSTRACT

The fovea is the small depression at the neurosensory retina that underlies high-

resolution central vision. It is vulnerable to disease and disruption of its

architecture causes visual disability. Foveal morphology varies significantly

across individuals. The molecular causes and functional consequences of this

anatomical diversity are incompletely understood. Here, we extracted six foveal

morphological parameters from Optical Coherence Tomography (OCT) images of

55,604 UK Biobank participants. We found notable variability in foveal

morphology, and detected significant links with sex and genetic ancestry.

Genome-wide association studies identified 197 lead loci across the six foveal

morphological parameters, implicating genes involved in pigmentation (e.g., *TYR*,

TSPAN10, GPR143) and patterning (e.g., *FGFR2*, *PTPRD, CYP1A1*). Heritability

estimates ranged from 29-43%. Foveal pit volume was associated with future risk

of age-related macular degeneration (HR>3.5, p=0.001), a finding supported by

Mendelian randomization. These results establish foveal morphology as a highly-

heritable trait with substantial influence over retinal disease risk.

INTRODUCTION

The fovea is a specialized structure at the center of the retina. It has a key role in high-acuity vision, supports most daily life activities, and enables tasks such as reading and face recognition.[1,2] The fovea is affected in a range of retinal disorders, including common adult-onset diseases such as age-related macular degeneration (AMD) (which is associated with loss of foveal cells) and pediatric conditions like albinism (which is characterized by foveal hypoplasia).[2]

The fovea has been the subject of study for more than a century. Despite this, there are significant gaps in our knowledge, partly because animal models that are routinely used in research (e.g. most murine and teleost models) lack this morphological specialization.[2] Recent advances in genomics, data science and *in vivo* imaging technologies have transformed our ability to study complex biological structures like the fovea.[3] Another major enabler has been the increasing availability of large-scale cohorts with a rich phenotypic scope. One example is the UK Biobank, a biomedical resource containing in-depth genetic and health information from >500,000 individuals.[4] Many UK Biobank participants underwent enhanced phenotyping including visual function assessment (>130,000 volunteers) and imaging of the central retina (>84,000 volunteers);[5] the latter was obtained using optical coherence tomography (OCT), a non-invasive imaging modality that rapidly generates cross-sectional retinal scans at micrometre-resolution.[6]

The most prominent morphological feature of the human fovea is the pit,

corresponding to a central excavation of the inner retinal layers. Previous

imaging studies have highlighted substantial morphological variability in respect

to the depth, width and volume of the pit.[1,7–11] This anatomical diversity is thought

to arise from variations in foveal development. Indeed, a blunted or absent foveal

pit is often seen in individuals born prematurely.[12] Differences in foveal

morphological parameters (hereafter, foveal traits) have also been reported

between male and female and among individuals with different ancestral

backgrounds.[7,13,14]

Previous studies have sought to investigate the genetic architecture of OCT-

derived retinal traits, including central retinal thickness, and inner and outer

retinal layer thickness.[15–20] However, to date, a comprehensive analysis of the

genetic determinants of foveal morphology has not been undertaken. Here, we

sought to address this knowledge gap and to explore the association between

foveal traits and common retinal conditions (Fig.1).

RESULTS

Fig.1. Outline of the study design. First, we used the RETIMAT tool to extract six key foveal traits from UK Biobank OCT scans (pit volume, rim radius, rim height, mean slope, pit depth and CFT). Subsequently, we performed phenotypic and genetic studies to gain insights into these foveal traits. Phenotypic analyses included the

investigation of the relationships between the studied foveal traits and age, spherical equivalent refractive error, sex, genetic ancestry, and visual acuity. Time-to-event analyses involved assessing the relationships between the six studied foveal traits and two major causes of visual impairment, age-related macular degeneration and glaucoma. Genetic investigations included common-variant genome-wide association studies (primary and replication), heritability estimation, and genetic correlation analyses.

OCT, optical coherence tomography; CFT, central foveal thickness.

113   **Phenotypic variability in foveal traits**

After applying standard OCT quality control filters,[20] we defined a subset of the

UK Biobank population with high-quality OCT scans. This cohort included 55,604

individuals and had similar sex and age profile to the overall UK Biobank

population.[15] Most study subjects were female (52%) and the mean age at OCT

imaging was 57.7 years (standard deviation: 8.1 years).

For each study subject, an OCT 'volume scan' (containing 128 cross-sectional

images) was available from the central retina of each eye. We used left eye

images as input to RETIMAT, an automated pipeline designed to extract foveal

traits from OCTs.[14] We focused on the following six parameters: foveal pit

volume, rim radius, rim height, mean slope, pit depth and central foveal thickness

(CFT) (Fig.1). The greatest fold range across UK Biobank participants was

observed for pit volume (15.1-fold), followed by pit depth (6.5-fold) and mean

slope (6.3-fold), indicating large differences between the most extreme

phenotypes in the population. In contrast, parameters such as rim height (1.4-

fold) and rim radius (2.1-fold) were more constrained (Fig.2).

Fig.2. Histograms showing the distribution of six foveal traits in UK Biobank participants (after removal of 1% from the upper and lower tails of the data). Also shown are six examples of OCT scans from the extreme ends of each distribution. The following mean values were obtained: pit volume 0.080 mm^3 (range: 0.024–0.356 mm^3), rim radius 0.98 mm (range: 0.71–1.49), rim height 353 µm (range: 291–394 µm), mean slope 6.61° (range: 1.88–11.86°), pit depth 113 µm (range: 32–213 µm), central foveal thickness (CFT) 240 µm (range: 102–312 µm).

Fig.3. Violin plots showing the distribution of six foveal traits between male and female UK Biobank participants. Females tend to have larger pit volumes (0.085 mm^3 vs. 0.073 mm^3), deeper pits (114 µm vs. 111 µm) and slightly larger rim radii (1.00 vs. 0.954) than male. In contrast, male had higher central foveal thickness (CFT) (244 µm vs. 236 µm), steeper mean pit slopes (6.69° vs. 6.54°) and greater rim height (355 µm vs. 350 µm) than females. The Wilcoxon signed rank test was used to make comparisons and all p-values were found to be < 2.3 x 10^{-10}. It is noted that the units for the foveal traits differ and, as such, the scales should not be directly compared. Further information, including numerical data, can be found in Supplementary Table S3.

First, we explored associations between the six studied foveal traits and age at

image acquisition and found no notable correlations (Supplementary Table S1).

We then examined associations between the same six foveal traits and spherical

equivalent refractive error (a proxy for eye size) and found these to be variably

correlated (Supplementary Table S2). Next, we inspected sex-related differences

in foveal morphology, identifying statistically significant variation for all six studied

traits. On average, females were found to have deeper, broader and larger foveal

pits than males (Fig.3; Supplementary Table S3). We also compared UK Biobank

participants of different genetic ancestries, focusing primarily on individuals of

European- and African-like ancestries (as these are the two largest groups in the

dataset and the only two to contain >1,000 individuals; Fig.4; Supplementary

Fig.S1; Supplementary Table S4). Again, we found statistically significant

differences for all six studied foveal traits. On average, individuals of African-like

genetic ancestries had deeper, broader and larger foveal pits that their European

counterparts. These findings reinforce the role of sex and ancestry in shaping

foveal morphology, and highlight the importance of including these parameters

as covariates in subsequent genetic and phenotypic analyses of foveal traits.

Fig.4. Violin plots showing the distributions of six foveal traits in individuals with European- and African-like genetic ancestries (EUR and AFR respectively). The AFR group had notably larger pit volumes (0.11 mm^3 vs. 0.08 mm^3), deeper foveal pits (121.37 µm vs. 112.19 µm) and greater rim radii (1.09 mm vs. 0.97 mm) than the EUR group. In contrast, the EUR group had higher central foveal thickness (240.65 µm vs 221.26 µm), steeper mean pit slope (6.62 ° vs. 6.41 °) and greater rim height (353.17 µm

vs. 342.80 µm) than the AFR group. The Wilcoxon signed rank test was used to make comparisons and all p-values were found to be $< 1 \times 10^{-10}$. It is noted that he units for the foveal traits differ and, as such, the scales should not be directly compared. Further information, including numerical data, can be found in Supplementary Table S4.

Aiming to gain insights into the factors underlying the detected differences

between ancestral groups, we investigated the role of fundus (principally

choroidal) pigmentation. To evaluate this, we used a fundus imaging-derived

proxy of pigmentation, the 'Retinal Pigment Score' (RPS).[21] We then calculated

the unique variance explained (adjusted R^2) values by RPS and ancestry across

the six studied foveal traits, adjusting for age, sex and spherical equivalent

refractive error. Overall, genetic ancestry explained a greater proportion of trait

variance than fundus pigmentation for most traits, particularly pit volume and rim

radius. It is noted that pigmentation explained negligible unique variance for rim

radius, suggesting little independent contribution beyond genetic ancestry (Fig.5;

Supplementary Table S5).

Fig.5. Unique contributions of fundus pigmentation and genetic ancestry to foveal morphology. Bar plots show the unique adjusted R² values for the six studied foveal traits, representing the proportion of variance explained by Retinal Pigment Score (RPS) and genetic ancestry, after adjusting for age, sex and spherical equivalent refractive error. Unique R² values were calculated from linear models by subtracting the R² of ancestry-only or RPS-only models from the combined model. Genetic ancestry accounted for a larger share of variance across most traits, particularly rim radius and pit volume (2.7% and 2.4% of variance, respectively). Pigmentation (RPS) explained modest additional variance, most notably for rim height and pit depth (0.9% and 0.6% of variance, respectively). For some traits, such as rim radius, the unique contribution of RPS was slightly negative, likely reflecting noise or model penalization rather than a true inverse effect. Further information, including numerical data, can be found in Supplementary Table S5.

To assess the extent to which foveal morphology predicts visual acuity, we

performed linear regression with LogMAR vision as the outcome, and age, sex,

spherical equivalent refractive error, and genetic ancestry as covariates. Several

traits were significantly associated with acuity, but all effect sizes were small (R²

< 1.3%). Mean slope was the strongest predictor, with steeper pits linked to

better vision. In contrast, greater pit volume was associated with worse acuity,

and central foveal thickness showed no effect (p = 0.30) (Supplementary Fig.S2).

Association between foveal traits and risk of retinal disease

To investigate whether foveal morphology constitutes a risk factor for future

retinal conditions, we performed Cox proportional hazards modeling using the six

studied foveal traits, along with time-to-event data for two leading causes of

visual impairment, AMD and glaucoma. We adjusted for age, sex, spherical

equivalent refractive error, and genetic ancestry in these analyses.

For AMD (n=1,345 cases), there was a notable association with larger pit volume

(hazard ratio 17.4; 95% confidence intervals [CI] 3.5-85.7) and wider rim radius

(hazard ratio 2.0; 95% CI 1.2-3.5). For glaucoma (n=1,166 cases), there were

statistically significant associations with mean slope, rim height and pit depth.

However, the corresponding hazard ratios indicated only small effect sizes

(ranging from 0.255 to 0.999) (Fig.6, Supplementary Table S6).

Fig.6. Cox regression analysis of associations between foveal traits and post-OCT-scan assigned codes for age-related macular degeneration (AMD) (H35) and glaucoma (H40). Statistically significant associations (after correcting for multiple comparisons using a 5% false-discovery rate [FDR] threshold) are highlighted with red. Further information, including numerical data, can be found in Supplementary Table S6.

Genetic associations of foveal traits

To investigate the genetic factors influencing the six studied foveal traits, we

performed common-variant genome-wide association studies (GWAS) using

REGENIE.[22] For these analyses, we defined a subset of the UK Biobank

population that can be considered genetically well-mixed (*i.e.* includes

participants that were assigned by genotype principal component analysis to a

cluster with subjects of mostly European-like ancestries). After applying relevant

imaging, genetic, and foveal trait quality controls, this cohort included 36,205

individuals; 29,710 of these were recruited between 2006 and 2010 (Instance 0,

'Initial assessment') and were used in the primary analysis while 6,495 were

independently recruited between 2012 and 2013 (Instance 1, 'First repeat

assessment visit') and were included in the replication study. The following

covariates were incorporated into the models: age, sex, height, weight, spherical

equivalent refractive error, and genetic principal components 1 to 20. We utilized

a conventional threshold for multiple testing correction in GWAS studies, $p < 5 \times$

10^{-8} ('genome-wide significance').

Genome-wide significant variants in the primary GWAS included: 5,933 for pit

volume; 6,251 for rim radius; 7,694 for rim height; 4,117 for mean slope; 4,710

for pit depth; and 2,723 for CFT. Following analysis with GCTA-COJO

(conditional and joint multiple-variant analysis),[23] these merged into the following

lead loci: 64 for pit volume, 65 for rim radius, 45 for rim height, 29 for mean

slope, 40 for pit depth, and 28 for CFT (Fig.7, Supplementary Fig.S3-8;

Supplementary Dataset S1). We subsequently performed a replication GWAS

using UK Biobank subjects that were not included in the primary analysis,

establishing a high level of concordance between the primary and replication

studies with all traits having a correlation coefficient greater than 0.9

(Supplementary Fig.S9–14). Although the replication sample was considerably

smaller, several lead loci still reached genome-wide significance in both studies:

6 for pit volume, 4 for rim radius, 1 for rim height, 1 for mean slope, 2 for pit

depth, and 2 for CFT (Supplementary Dataset S1).

We identified a combination of trait-specific and shared loci influencing foveal

morphology. Many of these loci encompass variants previously linked to retinal

layer thickness parameters (including *TSPAN10* and *LINC00461*) while a subset

of them has been linked to monogenic disorders, most notably, albinism

(including *TYR*, *OCA2* and *GPR143*) (Table 1 and Supplementary Dataset S1).

Our primary analysis highlighted several loci that have not been previously

associated with retinal phenotypes, including *CYP1A1, PPP1R9A, KMT2A,*

CASZ1 and *CCDC122*. We also identified variants in *FGFR2*, *PTPRD*, *SPRY2,*

FLT1 and LATS1, for which associations have only been detected for retinal

thickness measurements but not for foveal/macular traits.

MAGMA gene set enrichment analysis did not yield any statistically significant

hits after correction for multiple testing.

It is noted that the genomic inflation factors (λGC) ranged from 1.12 to 1.17, with

corresponding linkage disequilibrium (LD) score intercepts close to 1 (range:

1.005–1.027). These results suggest minimal confounding from population

structure or cryptic relatedness. The attenuation ratios were uniformly low (all <

0.11), consistent with polygenic architecture and well-controlled inflation across

all six studied traits (Supplementary Table S7).

Fig.7. A Miami plot, illustrating the results of the primary genome-wide association study for foveal pit volume (top) and rim radius (bottom). The red line indicates the threshold for genome-wide significance, and 5 key loci exceeding this value have been annotated.

Heritability estimates for foveal traits

We used linkage disequilibrium (LD) score regression analysis[24] to obtain

heritability estimates for the six studied foveal traits. We found moderate to high

heritability, ranging from $h^2 = 0.29$ (standard error [SE] = 0.03) for CFT to h^2 =

0.43 (SE = 0.04) for pit volume. Rim height ($h^2 = 0.41$), rim radius ($h^2 = 0.39$), pit

depth ($h^2 = 0.34$), and mean slope ($h^2 = 0.31$) were also significantly heritable

(Supplementary Table S7).

Global genetic correlation between foveal traits

LD score regression was also used to assess the global genetic correlation

between the studied traits. We found significant interdependence between many

foveal traits. The strongest correlations (rg > 0.8) were observed between pit

depth and mean slope (rg = 0.88) and between pit volume and rim radius (rg =

0.85), suggesting that these pairs reflect closely related biological processes.

Moderate correlations (rg = 0.6-0.8) included pit depth and pit volume (rg = 0.76),

mean slope and rim height (rg = 0.65), and pit depth and rim height (rg = 0.57),

indicating shared underlying variation across several morphological aspects of

the fovea.

In contrast, correlations between CFT and other parameters were more variable.

We observed strong negative correlations between CFT and pit volume (rg = –

0.79), pit depth (rg = –0.69), and rim radius (rg = –0.51), consistent with an

inverse relationship between retinal thickness and pronounced foveal pit

excavation (Fig.8).

Correlations between foveal traits and fundus pigmentation (RPS) were generally

weak (rg = -0.23 with pit depth, -0.22 with mean slope), with the strongest

positive correlation seen with CFT (rg = 0.23) (Supplementary Table S8).

Fig.8. Genetic correlations between foveal traits. Asterisks indicate levels of statistical significance (* p < 0.05, ** p < 0.01, *** p < 0.001). Further information, including numerical data, can be found in Supplementary Table S8.

CFT, central foveal thickness.

Table 1. Summary of the 10 top-ranking loci associated with foveal traits.

Top-ranking common variant in locus	Chr: position (GRCh37)	Key gene(s)	Allele freq	Foveal trait with significant result (minimum p-value)	Selected previous associations in the GWAS Catalog; [Panelapp association(s)]
rs6420484	17:79612397	*TSPAN10 / NPLOC4 / PDE6G*	0.65	Pit depth (2.01×10^{-107}); Pit volume (7.41×10^{-105}); Rim radius (3.54×10^{-59}); Mean slope (3.16×10^{-55}); Rim height (4.88×10^{-48})	Retinal thickness measurements, refractive error, eye color, hair color
rs3138142	12:56115585	*RDH5 / CD63*	0.24	Rim height (1.98×10^{-77}); Pit depth (2.86×10^{-14}); Mean slope (6.21×10^{-13})	Retinal thickness measurements, refractive error, retinal vascular fractal density; [retinal dystrophy]
rs11708067	3:123065778	*ADCY5*	0.24	Rim radius (6.34×10^{-69}); Pit volume (7.95×10^{-56})	Retinal thickness measurements, height, birth weight, type 2 diabetes, glucose levels, HbA1c levels, cholesterol levels; [dyskinesia]
rs1042602	11:88911696	*TYR*	0.37	Pit depth (1.67×10^{-62}); Pit volume (8.26×10^{-58}); Rim radius (1.16×10^{-22}; Mean slope (3.48×10^{-39}); CFT (9.32×10^{-40});	Retinal thickness measurements, brain measurements, skin color, hair color; [albinism]
rs9298817	9:21576591	*MIR31HG*	0.66	Pit volume (1.43×10^{-61}); Rim radius (1.44×10^{-21}); Rim height (2.30×10^{-20})	Retinal thickness measurements, thyroid medication use
rs17421627	5:87847586	*LINC00461*	0.07	Pit volume (3.12×10^{-46}); CFT (1.34×10^{-35}); Rim radius (3.87×10^{-32}); Rim height (1.60×10^{-27})	Retinal thickness measurements, retinal vascular fractal density
rs11051145	12:31050201	*TSPAN11*	0.16	Pit volume (5.02×10^{-46}); CFT (5.22×10^{-34})	Retinal thickness measurements, optic disc size

rs7020703	9:8632149	*PTPRD*	0.22	Rim radius (1.25 x 10^{-37})	Retinal thickness measurements, bone mineral density, height, restless leg syndrome
rs67138036	4:187683202	*FAT1*	0.26	Rim radius (4.58 x 10^{-37})	Retinal thickness measurements; [microphthalmia/ coloboma, nephropathy]
rs1493523	15:44345614	*CCDC122*	0.68	Pit volume (1.86 x10^{-36})	-

CFT, central foveal thickness

Mendelian randomization

We performed Mendelian randomization (MR) analysis on the two foveal traits

that we found to be associated with significantly increased risk of AMD and

glaucoma. This analysis revealed causal relationships between foveal

morphology and retinal disease risk. There was evidence of causal association

between increased pit volume and AMD (OR = 1.12, 95% CI: 1.04–1.21, inverse

variance weighted [IVW] p = 0.004). This was robust to a leave-one-out test

using the IVW method. In contrast, rim radius showed no evidence of a causal

effect on AMD or glaucoma (p > 0.2 across all methods). Pit volume was also

nominally associated with decreased risk of glaucoma (IVW p = 0.02; OR = 0.94,

95% CI: 0.89–0.99). This result remained significant in the MR Egger (p=0.02)

and weighted mode (p=0.03) but did not quite exceed significance in the

weighted median (p=0.056) nor persist throughout leave-one-out testing using

the IVW method (Supplementary Table S9). Reverse Mendelian randomization

analyses yielded no evidence of AMD or glaucoma influencing foveal

morphology. Overall, the findings provide evidence that pit morphology causally

influences the risk of AMD and primary open-angle glaucoma, but these results

should be interpreted in the context of their reliance on several key assumptions

of Mendelian randomization being met (see Methods for further details on this).

DISCUSSION

In the present study, we performed large-scale phenotypic and genetic analyses

to better understand the variability and determinants of foveal morphology. By

leveraging retinal imaging data from UK Biobank participants and a

comprehensive foveal trait calculation tool (RETIMAT), we were able to detect

differences in foveal morphology influenced by sex, genetic ancestry, and genetic

factors. We also found a notable association between foveal pit volume and

AMD.

We first analyzed how foveal architecture varies at the population level. As the

number of subjects in this study is an order of magnitude greater than in most

prior reports, we were able to identify previously unrecognized variability in foveal

pit morphology (e.g. 15-fold for pit volume). Sex and ancestry were found to

significantly influence foveal morphology, an observation that is in keeping with

the findings of previous smaller-scale studies.[7,9,13,14] In line with previous

research, males exhibited greater rim height, shorter rim radii, and a steeper

mean slope.[14] Similarly, we found that individuals with African ancestries had

wider, deeper, and more voluminous foveal pits than their European

counterparts, in line with findings by Zouache *et al.*[7] The smaller East- and

South-Asian ancestral groups fell between these two, consistent with the limited

normative datasets available for Asian populations.[25,26] It is highlighted that, in

contrast to most previous studies, we used genetic ancestry, a metric that is

more accurate and objective than ethnicity or self-reported ancestry.[27] We also

performed additional analyses to gain insights into the extent to which the

observed difference between continental groups is driven by pigmentation. Our

observations suggest that genetic ancestry tends to be an independent

contributor irrespective of fundus pigmentation, especially for foveal pit volume

and rim radius.

We found that a larger pit volume is a key risk factor for AMD, and that there is a

33% increase in risk per $0.1\,mm^3$ increase in pit volume. This observation was

supported by two orthogonal approaches, Cox regression analysis and

Mendelian randomization. The results of the latter substantially reduce the risk

that the observed association is due to reverse causation or confounding, and

support the assertion that pit volume is a risk factor for AMD rather than a

correlate. While the hazard ratio was 17.4 (95% CI 3.5-85.7), the corresponding

odds ratio (OR) was 26.1 (95% CI 6.1-112.0, p = 1.1×10^{-5}) (Fig.6,

Supplementary Tables S6 and S10). It is noted that several environmental and

genetic risk factors have been previously associated with AMD, including: age

(OR 1.1), gender (OR 1.6), smoking (OR 1.9)[28], *CFH* gene variants (e.g.

homozygosity for rs1061170 corresponds to OR of 4.7)[29], *ARMS2* locus variants

(e.g. homozygosity for rs10490924 corresponds to OR of 8.6)[30] and family history

(OR 10.8).[31] Although AMD selectively affects the foveal/macular region, to date,

anatomical risk factors have been understudied[9] and, to our knowledge, this is

the first time that specific foveal morphological parameters are found to set the

stage for susceptibility to AMD. It can be speculated that the link between a

larger pit and AMD risk is driven by differences in the distribution of macular

xanthophyll pigments and/or by alterations in photoreceptor cell densities in the

vulnerable parafoveal area of low rod-to-cone ratio.[32] Future advanced

ophthalmic imaging and mathematical modelling studies will provide further

insights into how foveal anatomy predisposes to this leading cause of visual

impairment.

Following phenotypic analyses, including of foveal traits that have received little

attention so far (such as foveal volume and radius), we performed genetic

association studies. We found that the observed morphological diversity has a

strong genetic underpinning, with a high level of heritability across most traits

(Supplementary Table 7). Pigmentation-related genes such as *TYR* and

TSPAN10 surfaced across multiple foveal traits, underscoring the role of melanin

synthesis in the shared developmental pathways that shape foveal morphology

(Table 1). This is in keeping with previous observations in albinism[33,34] and in a

smaller cohort where a link between the common functional *TYR:*c.1205G>A

(p.Arg402Gln) [rs1126809] variant and decreased rim radius was reported.[35]

While many of the lead loci detected here have previously been linked to retinal

traits, at least 124 of them have no such prior associations (Supplementary

Dataset S1). Two examples are *CYP1A1* and *PPP1R9A. CYP1A1* (lead marker:

rs200391170; association with mean slope; $p = 2.2 \times 10^{-21}$) encodes a major

cytochrome P450 enzyme and is involved in the metabolism of both endogenous

substances and environmental chemicals. It has been implicated in the

conversion of all-trans retinol to all-trans retinoic acid in human fetal tissues.[36]

Notably, retinoic acid gradients are essential for dorsoventral patterning, and

retinoic acid signaling has been directly linked to the positioning of the fovea-like

area in the chick retina.[37,38] Another gene that has not been previously linked to

retinal phenotypes is *PPP1R9A* (lead marker: rs9649194; association with pit

depth and mean slope; minimum $p = 6.5 \times 10^{-14}$). This gene encodes a protein

phosphatase that modulates actin cytoskeleton organization,[39] and one

possibility is that it plays a role in the formation of the foveal pit (rather than the

patterning/positioning of this high-acuity structure to the correct retinal region, a

process that occurs earlier in development).

A subset of the detected lead loci (n=36) has been previously linked to retinal

thickness measurements (either clinically-used[15–17] or deep learning based[5,19,40])

but not with foveal/macular traits (Supplementary Dataset S1). These novel

associations that we report here enhance the interpretability of the signals and

allow more robust links to be drawn with central retinal development/patterning.

Two notable examples are *FGFR2* and *PTPRD. FGFR2* (lead marker:

rs146727842; association with mean slope; $p = 1.4 \times 10^{-20}$) encodes a receptor

tyrosine kinase that is essential for embryonic development.[41] *FGFR2* has been

shown to interact with *FGF8*, whose expression is required to pattern the chick

fovea-like area.[37,38,42] *PTPRD* (lead marker: rs7020703; association with rim

radius; $p = 1.3 \times 10^{-37}$) encodes a tyrosine phosphatase that regulates neural

development by modulating receptor tyrosine kinase activity.[16,43] A related

protein (cRPTPlambda) is found to be absent from the chick fovea-like area, and

it can be speculated that *PTPRD* is involved in regulating FGF signaling early to

set up the foveal region.[44]

This study has several limitations. First, obtaining accurate quantitative

measurements of OCT-derived retinal features is inherently challenging. One

issue is image scaling, as precise measurements require knowledge of the exact

lateral scale of each scan. To address lateral magnification, we included

spherical equivalent refractive error as a covariate in our analyses. While this

method is less accurate than using ocular axial length[31,48], such measurements

are not available in the UK Biobank dataset. It is noted though that our findings

showed good agreement with those from smaller, more targeted studies.[47,49] This

approach is also consistent with most previous genetic association studies

involving retinal imaging-derived phenotypes.[15,16,18–20] Second, we relied on UK

Biobank healthcare record data to identify cases of AMD and glaucoma. These

records are known to be incomplete, and several alternative methods using

retinal imaging data have been proposed.[50,51] Although these have shown

positive diagnostic performance, they can overestimate prevalence,[51] and we

opted to use a conventional approach focused on record-based diagnoses.

Finally, while the UK Biobank was designed to be broadly representative of the

ageing UK population, it is subject to selection biases. All reported associations

should therefore be interpreted considering this limitation.[52,53]

In conclusion, we show that (i) foveal architecture is patterned by sex, genetic

ancestry and genes related to pigmentation, and (ii) one aspect of that

architecture, pit volume, appears to lie on the causal pathway to AMD. These

findings position foveal pit metrics as potential biomarkers of macular

vulnerability and invite mechanistic work to dissect how developmental pathways

intersect with ageing to drive retinal degeneration.

MATERIALS AND METHODS

Ethical approval

The UK Biobank has received approval from the UK National Information Governance Board for Health and Social Care and the National Health Service North West Centre for Research Ethics Committee (Ref: 11/NW/0382). This research was conducted using the UK Biobank Resource under projects 53144 and 49978. All investigations were conducted in accordance with the tenets of the Declaration of Helsinki.

UK Biobank: cohort characteristics and ophthalmic phenotyping

We used data from the UK Biobank, a resource containing genomic and health information from 502,355 individuals from across the United Kingdom, aged between 40 and 69 years at the time of recruitment.[3] Subsets of UK Biobank volunteers underwent enhanced phenotyping including visual acuity testing (>130,000 individuals), non-cycloplegic autorefraction (>125,000 individuals) and imaging of the central retina using color fundus photography and OCT (>84,000 individuals). Baseline ophthalmic assessment was conducted between 2009 and 2010 for most participants and between 2012 and 2013 for a smaller subset.[4]

Best-corrected visual acuity was assessed using a computer-displayed LogMAR chart (Precision Vision, LaSalle, IL, US) (UK Biobank data-fields: 5208 and 5201). Participants wore their habitual distance correction and were tested at 4 m; if unable to identify any letters, the chart was moved to 1 m. Starting from the top line, testing stopped once ≥2 letters on a line were misread.[4]

Non-cycloplegic autorefraction was carried out using a Tomey RC 5000 device (Tomey Corp., Nagoya, Japan). The spherical equivalent refractive error (spherical error + 0.5 ×

cylindrical error) was subsequently calculated for each participant (data-fields 5084-

5085; 5086-5087).[4]

Color fundus photographs and spectral-domain OCT images were acquired using the

Topcon 3D-OCT 1000 Mark II device (Topcon Corp., Tokyo, Japan), without

pharmacological dilation (data-fields 21015, 21016, 21011, 21013). OCTs were obtained

using a '3D macular volume' protocol (6 × 6 mm; 128 horizontal B-scans, each

comprising 512 A-scans arranged in a raster pattern).[5,40,54] The right eye was imaged

first. Our analysis focused on left eye images as we assumed that familiarity with the test

would have led to scans that, on average, had higher overall quality.[5,18]

Fundus pigmentation quantification

We used a recently described method[21] to evaluate fundus pigmentation in UK Biobank

participants. This involved calculating the Retinal Pigment Score (RPS), a continuous

quantitative metric that is thought to primarily reflect choroidal pigmentation.[55]

RPS was derived using the approach described by Rajesh *et al.* which entails selecting

fundus photographs of sufficient quality (*i.e.* those labeled as 'Good' by Automorph, an

established deep learning tool)[56] and processing them through the RPS pipeline to

quantify background pigmentation.[17]

OCT quality control and foveal morphological parameter extraction

OCT scans were filtered for overall quality in line with previous studies.[18,20] Briefly, we

discarded OCTs that (j) had a quality score below 40 and (ii) fell within the lowest 10%

of the inner limiting membrane (ILM) edge-strength indicator (a metric that captures the

sharpness and continuity of the ILM boundary and flags blinks, signal dropouts, and

major segmentation errors).[20]

OCT images meeting the above quality control criteria were processed using RETIMAT,

an open-source MATLAB toolbox for OCT image analysis. The processing pipeline

included the following steps:

• parsing Topcon (.fda) files to extract the ILM and Bruch's membrane (BM) surfaces

delineated by the device software;

• computing total retinal thickness maps;

• automatically aligning thickness maps to the foveal center;

• transforming raster thickness maps into a radial pattern; and

• computing morphometric values.

RETIMAT has been used successfully in previous studies and its fully automated

methods for fovea location and analysis have been independently assessed, making it

well-suited for high-throughput analysis.[14,57] We focused on six RETIMAT-derived foveal

traits: foveal pit volume, rim radius, rim height, mean slope, pit depth and central foveal

thickness (CFT) (Fig.1, Table 2).

Aiming to study the phenotypic variability of these parameters, we assembled a broad

'phenotypic analysis' cohort. This included UK Biobank participants whose OCT scans

passed the imaging quality control criteria for this study; whose RETIMAT-derived foveal

trait values did not fall in the 1% upper or lower extreme; and who were assigned to one

of the four main genetically-inferred ancestral groups (EUR, AFR, SEA, and SAS; see

'Genetic ancestry assignment' section below). Aiming to study the genetic architecture of

foveal traits, a second, more focused 'genetic analysis' cohort was defined. This was

limited to unrelated participants of European-like ancestries with OCT scans; no sex

chromosome aneuploidy; and not flagged as outliers for heterozygosity nor missingness.

Individuals with RETIMAT-derived foveal trait values exceeding 2.5 standard deviations

were also excluded (Supplementary Fig.S15); this cut-off was selected in accordance

with the approach used in previous genetic association studies of retinal thickness

measurements. [15,16,18–20] The final GWAS sample comprised 36,205 unrelated

individuals with European-like ancestries (29,710 primary plus 6,495 replication); the

broader phenotypic cohort totaled 55,604 individuals across ancestries.

Table 2. Definitions of the six RETIMAT-derived foveal traits analysed in this study

Foveal trait	Definition
Pit volume (mm^3)	Integrated volume between rim height and the full thickness profile
Rim radius (mm)	Radial distance from center to point of rim height
Rim height (µm)	Maximum thickness at the foveal rim
Mean slope (°)	Average inclination from center to rim
Pit depth (µm)	Rim height minus central foveal thickness
Central foveal thickness (µm)	Thickness at the foveal center
A pictorial representation of rim radius, rim height and mean slope can be found in Fig.1	

Genetic ancestry assignment

To facilitate inter-ancestry analyses of foveal traits, we sought to characterize the

genetic ancestry of UK Biobank participants. A subset of 409,373 participants with

European-like ancestries is readily-available (data-field 22006) and we aimed to obtain

genetic ancestry estimates for the remaining 78,296 individuals. Using a previously-

described approach[58] we projected these UK Biobank participants onto reference

genetic data from the 1000 Genomes Project.[59] In brief, we used directly genotyped

single-nucleotide variants from the UK Biobank and genetic data from the 1000

Genomes Project (Phase 3, v5a.20130502). After matching variants between these two

resources, we merged the two datasets using PLINK v2.0,[60] retaining 718,487 high-

quality variants. We then performed linkage disequilibrium (LD) pruning and removed

regions with long-range LD, resulting in 30,320 independent variants. Ancestry

proportions were estimated using ADMIXTURE (v1.3.0),[61] with four reference groups

from the 1000 Genomes Project dataset: GBR (European), YRI (African), ITU (South

Asian), and CHS (East Asian). Only individuals with at least 80% ancestry probability

assigned to one of these four groups were retained for further phenotypic analyses.

Statistical analyses of foveal phenotypes

The relationships between foveal traits, age of OCT image acquisition and spherical

equivalent refractive error were studied using Spearman's correlation analyses. Further,

comparisons were made between sex and genetic ancestry groups for each of the

studied foveal traits using the Wilcoxon rank-sum test (when comparing two groups) and

the Kruskal–Wallis test (when comparing more than two groups).

To quantify the contributions of fundus pigmentation (RPS) and genetic ancestry to

variability in foveal morphology, we fitted linear regression models for each trait, as

follows: The two predictors were analyzed together given their presumed biological

relationship (as pigmentation traits vary by ancestry and may capture overlapping

sources of variation).

Model 1: included RPS, spherical equivalent refractive error, and sex.

Model 2 included genetic ancestry (as a categorical variable), spherical equivalent

refractive error, and sex.

Model 3: included both RPS and genetic ancestry terms, along with spherical equivalent

refractive error and sex.

From each model, we extracted adjusted R^2 values to assess explained variance.

Unique contributions of RPS and ancestry were derived by comparing the full model

against models excluding one predictor set at a time.

Finally, the association between foveal morphology and visual acuity was studied using

a linear regression approach with age, sex, spherical equivalent refractive error and

genetic ancestry used as covariates.

Genome-wide association studies (GWAS)

We performed a common-variant GWAS using an additive linear model in REGENIE

v3.1.1.[22] To meet the normality assumptions of GWAS, we applied rank-based inverse

normal transformation to any foveal traits that deviated from a normal distribution. For

this analysis, we focused only on individuals of European-like genetic ancestries (data-

field 22006). We performed a primary and a replication GWAS. The focus of the former

was on the 67,250 individuals that were imaged at the time of their baseline visit

(Instance 0, 'Initial assessment visit (2006–2010)'). The focus of the replication study

was on the 15,568 different participants that were imaged for the first time during their

first repeat assessment (Instance 1, 'First repeat assessment visit (2012–2013)').

To process the imputed genotype data, we applied conventional quality control filters

using PLINK2.[60] Variants were retained if they had: an imputation quality score above

0.8; a minor allele frequency (MAF) of 5% or more; and a minor allele count (MAC) of 20

or more. Variants with a Hardy-Weinberg equilibrium p-value below 1×10^{-15} were

excluded, along with those with missingness greater than 10%. Individuals with more

than 10% missing genotypes were also excluded.[22]

The GWAS model was adjusted for age at recruitment (data-field 21022), sex (data-field

31), height (data-field 50), weight (data-field 21002), spherical equivalent refractive error

(calculated from data-fields 5085 and 5086 and defined as spherical error + 0.5 ×

cylindrical error), and the first 20 genetic principal components (data-field 22009).

Genome-wide statistical significance was set at $p < 5 \times 10^{-8}$.

To refine the obtained association signals, further analyses were performed using the

GCTA-COJO tool (conditional and joint multiple-variant analysis).[23] Variants located on

different chromosomes, or more than 10 Mb apart, were treated as unlinked. Lead

genetic variants were then annotated using Ensembl,[62] Open Targets,[63] and GWAS

Catalog[64] data (accessed on June 24, 2025). To accurately summarize the strongest

signals, the LD metrics of the changes that were highlighted as lead variants by GCTA-

COJO analysis and were within 1 Mb of one another were manually inspected using the

LDlink tool.[65]

LD score regression: heritability estimation and global genetic correlation

analysis

The global genetic correlations between the six studied foveal traits (pit volume, rim

radius, rim height, mean slope, pit depth, and CFT) and RPS were calculated using LD

score regression.[66] The GWAS summary statistics for RPS were obtained from a study

by Julien *at al.*[55] The analysis adhered closely to the methodology described by Bulik-

Sullivan *et al.*,[66] with genetic variants filtered according to the following criteria: inclusion

in HapMap3; minor allele frequency > 0.01; imputation score > 0.9, exclusion of strand-

ambiguous variants; and removal of duplicated variants. LD scores were derived from

the European subset of the 1000 Genomes Project dataset.[59]

LD score regression was also used to calculate heritability, λ_G (genomic inflation factor),

intercept, and ratio (denoting the proportion of inflation resulting from confounding or

bias) for each of the six studied foveal traits.

Cox regression analysis

We performed Cox proportional hazards regression to investigate the association

between foveal traits and incident age-related macular degeneration (AMD) and

glaucoma. Participants with available OCT-derived foveal traits and outcome data were

included in the analysis (n = 55,604). Disease status was defined using hospital inpatient

ICD-10 diagnosis fields (data-field: 41202; ICD-10 H35 and H40). The date of baseline

imaging (Year of Scan, YOS) was extracted from UK Biobank data-field 21836 and used

to define time-zero for each participant.

Age at scan was computed from YOS and the month and year of birth for each

participant. We defined disease onset as the earliest recorded date for the relevant ICD-

10 code. Participants were censored at the last available follow-up date (May 31, 2022),

if their imaging date was missing, or if disease onset occurred before the scan.

Cox regression models were fitted using the 'lifelines' Python package,[67] with time-to-

event (in days) as the outcome, and foveal traits as the predictors. All models were

adjusted for age at image acquisition, sex, spherical equivalent refractive error, and

genetic ancestry. Ancestry was modelled using one-hot encoded genetic ancestry

categories with individuals of European-like ancestries as the reference group. Models

were fit for each trait–disease pair. Hazard ratios (HR), 95% confidence intervals, and p-

values were extracted. The p-values were corrected for multiple testing using the

Benjamini–Hochberg false discovery rate (FDR) method.

To enhance the interpretability of the detected associations, we performed logistic

regression using a binary outcome defined as the presence or absence of disease onset

after the scan date and before the censoring date (May 31, 2022). Individuals who had

not developed the disease by the censoring date were treated as controls. Covariates

included age at image acquisition, sex, spherical equivalent refractive error, and genetic

ancestry (encoded as dummy variables). Analyses were run separately for each foveal

trait. Logistic regression models were fitted using the 'statsmodels' Logit function in

Python[68], and odds ratios (with 95% confidence intervals) and p-values were extracted.

Causal inference using two-sample Mendelian randomization

Causal associations between AMD, glaucoma and the six studied foveal traits were

assessed using univariable two-sample Mendelian randomization (MR).[69,70] Population

overlap biases univariable two-sample MR in the direction of causation. Accordingly, the

populations contributing toward traits determined to be causal were inspected to ensure

that there was no overlap. Genetic instruments were selected based on $p < 5 \times 10^{-8}$ in

relation to each exposure trait. Where an exposure instrument was not present in the

outcome dataset, we sought to identify a suitable proxy. Proxy variants were single-

nucleotide variants that were in high LD with exposure-associated variants which were

not present in the outcome dataset. Proxies were identified using the Ensembl server,[62]

selecting variants with an LD R^2 value of 0.9 or above. Genetic variants were clumped

using an LD R^2 value of 0.001 and a genetic distance cut-off of 10,000 Kb. This was

achieved using the 1000 Genomes Project European superpopulation reference panel

and the TwoSampleMR PLINK clumping function.[59,60] The effects of instruments on

outcomes and exposures were harmonized to ensure that the β values (*i.e.* the

regression analysis estimates of effect size) were signed with respect to the same

alleles. For palindromic alleles (*i.e.* alleles that are the same on the forward as on the

reverse strand), those with minor allele frequency (MAF) > 0.42 were omitted from the

analysis to reduce the risk of errors due to strand issues. In addition to using a range of

Mendelian randomization methods and quality control measures (as detailed below), we

endeavored to remove pleiotropic instruments and outliers from the analysis. To achieve

this, instruments that were more statistically significant (in terms of p-value) for the

outcome than the exposure were removed. Radial Mendelian randomization,[71] an

approach that detects outlying instruments, was also used for outlier identification, and

outliers were subsequently removed.[71] The Cochran's Q test was performed for each

analysis. The MR Egger intercept[72] was used to detect horizontal pleiotropy. The I^2

statistic was calculated as a measure of heterogeneity between variant specific causal

estimates. An $I^2 < 0.9$ indicates that MR Egger is more likely to be biased towards the

null through violation of the "NO Measurement Error" (NOME) assumption. Our primary

analysis was a multiplicative random effects inverse variance weighted (IVW) test, which

is the most efficient analysis method with valid instrumental variables.[73] It is noted that

Mendelian randomization relies upon several assumptions holding true to provide an

accurate assessment of causation. The core assumptions are that (i) the instrumental

variables (single-nucleotide polymorphisms) utilized are associated with the exposure;

(ii) the instrumental variables are not associated with the outcome via a confounder; and

(iii) the instrumental variables do not impact the outcome other than through the

exposure being considered (*i.e.* no horizontal pleiotropy).[73] Robust analyses are those

which can provide valid causal inference under weaker assumptions than an IVW test,

and are important for a complete Mendelian randomization study. Robust tests included

the weighted median (robust to up to 50% of invalid instruments), weighted mode (robust

to >50% of invalid instruments but assumes more variants estimate the true causal effect

than estimate any other quantity), MR Egger (robust to pleiotropic effects of variants,

provided the effects are independent of the variant-exposure associations) and a leave-

one-out analysis using the MRE IVW method (to test for highly influential variants).[73]

When appraising the results of a Mendelian randomization study, the confidence with

which one can assert causality is influenced by both the strength of association in the

primary analysis (IVW) and the range of robust tests in which the results are consistent.

Where the IVW is significant but robust tests are not, this may be in keeping with

causality, but under stricter assumptions of validity across the instrumental variables

utilized (and therefore may warrant more cautious interpretation).

DATA AVAILABILITY

UK Biobank data are available under restricted access through a procedure

described at http://www.ukbiobank.ac.uk/using-the-resource/. All other data

supporting the findings of this study are available within the article (including its

Supplementary Information files).

CODE AVAILABILITY

The scripts used to analyse the datasets included in this study are available

at https://github.com/davidjohngreen/fovea and https://github.com/mu-

biomed/retimat.

REFERENCES

1. Bringmann, A. *et al.* The primate fovea: Structure, function and development. *Prog Retin Eye Res* **66**, 49–84 (2018).

2. Provis, J. M., Dubis, A. M., Maddess, T. & Carroll, J. Adaptation of the central retina for high acuity vision: Cones, the fovea and the avascular zone. *Prog Retin Eye Res* **35**, 63–81 (2013).

3. Meyer, H. V. *et al.* Genetic and functional insights into the fractal structure of the heart. *Nature* **584**, 589–594 (2020).

4. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).

5. Chua, S. Y. L. *et al.* Cohort profile: design and methods in the eye and vision consortium of UK Biobank. *BMJ Open* **9**, e025077 (2019).

6. Bouma, B. E. *et al.* Optical coherence tomography. *Nature Reviews Methods Primers* **2**, (2022).

7. Zouache, M. A. *et al.* Comparison of the morphology of the foveal pit between african and caucasian populations. *Transl Vis Sci Technol* **9**, 1–11 (2020).

8. Tick, S. *et al.* Foveal shape and structure in a normal population. *Invest Ophthalmol Vis Sci* **52**, 5105–5110 (2011).

9. Wagner-Schuman, M. *et al.* Race- and sex-related differences in retinal thickness and foveal pit morphology. *Invest Ophthalmol Vis Sci* **52**, 625–634 (2011).

10. Dubis, A. M., McAllister, J. T. & Carroll, J. Reconstructing foveal pit morphology from optical coherence tomography imaging. *British Journal of Ophthalmology* **93**, 1223–1227 (2009).

11. De Moraes, C. G. *et al.* Interindividual variations in foveal anatomy and
artifacts seen on inner retinal probability maps from spectral domain OCT
scans of the macula. *Transl Vis Sci Technol* **7**, (2018).

12. O'Sullivan, M. L. *et al.* Foveal differentiation and inner retinal displacement
are arrested in extremely premature infants. *Invest Ophthalmol Vis Sci* **62**,
(2021).

13. Olvera-Barrios, A. *et al.* Foveal Curvature and Its Associations in UK
Biobank Participants. *Invest Ophthalmol Vis Sci* **63**, (2022).

14. Romero-Bascones, D. *et al.* Spatial characterization of the effect of age
and sex on macular layer thicknesses and foveal pit morphology. *PLoS*
One **17**, (2022).

15. Currant, H. *et al.* Sub-cellular level resolution of common genetic variation
in the photoreceptor layer identifies continuum between rare disease and
common variation. *PLoS Genet* **19**, (2023).

16. Currant, H. *et al.* Genetic variation affects morphological retinal phenotypes
extracted from UK Biobank optical coherence tomography images. *PLoS*
Genet **17**, (2021).

17. Gao, X. R., Huang, H. & Kim, H. Genome-wide association analyses
identify 139 loci associated with macular thickness in the UK Biobank
cohort. *Hum Mol Genet* **28**, 1162–1172 (2019).

18. Sergouniotis, P. I. *et al.* Autoencoder-based phenotyping of ophthalmic
images highlights genetic loci influencing retinal morphology and provides
informative biomarkers. *Bioinformatics* **41**, (2024).

19. Jackson, V. E. *et al.* Multi-omic spatial effects on high-resolution AI-derived
retinal thickness. *Nat Commun* **16**, 1317 (2025).

20. Maryam Zekavat, S. *et al.* Phenome-and genome-wide analyses of retinal
optical coherence tomography images identify links between ocular and
systemic health. *Sci. Transl. Med* **16**, 4517 (2024).

21. Rajesh, A. E. *et al.* Machine learning derived retinal pigment score from
ophthalmic imaging shows ethnicity is not biology. *Nature Communications*
16, (2025).

22. Mbatchou, J. *et al.* Computationally efficient whole-genome regression for
quantitative and binary traits. *Nat Genet* **53**, 1097–1103 (2021).

23. Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS
summary statistics identifies additional variants influencing complex traits.
Nat Genet **44**, 369–375 (2012).

24. Bulik-Sullivan, B. *et al.* LD score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295
(2015).

25. Appukuttan, B., Giridhar, A., Gopalakrishnan, M. & Sivaprasad, S.
Normative spectral domain optical coherence tomography data on macular
and retinal nerve fiber layer thickness in Indians. *Indian J Ophthalmol* **62**,
316–321 (2014).

26. Wu, J. *et al.* Clinical science Macular thickness and its associated factors
in a Chinese rural adult population: the Handan Eye Study. *Br J*
Ophthalmol **107**, 1864–1872 (2023).

27. *Using Population Descriptors in Genetics and Genomics Research*.
(National Academies Press, Washington, D.C., 2023). doi:10.17226/26902.

28. Babaker, R. *et al.* Risk factors for age-related macular degeneration:
Updated systematic review and meta-analysis. *Medicine (United States)*
104, e41599 (2025).

29.   Maugeri, A., Barchitta, M. & Agodi, A. The association between
complement factor H rs1061170 polymorphism and age-related macular
degeneration: a comprehensive meta-analysis stratified by stage of
disease and ethnicity. *Acta Ophthalmol* **97**, e8–e21 (2019).

30.   Ross, R. J. *et al.* The LOC387715 Polymorphism and Age-Related Macular
Degeneration: Replication in Three Case-Control Samples. *Invest*
Ophthalmol Vis Sci **48**, 1128–1132 (2007).

31.   Shahid, H. *et al.* Age-related macular degeneration: the importance of
family history as a risk factor. *British Journal of Ophthalmology* **96**, 427–
431 (2012).

32.   Curcio, C. A., Kar, D., Owsley, C., Sloan, K. R. & Ach, T. Age-Related
Macular Degeneration, a Mathematically Tractable Disease. *Invest*
Ophthalmol Vis Sci **65**, (2024).

33.   Lejoyeux, R. *et al.* Genetic analysis of participants with foveal hypoplasia.
Ophthalmic Genet 1–4 (2025) doi:10.1080/13816810.2025.2520411.

34.   Bakker, R. *et al.* The retinal pigmentation pathway in human albinism: Not
so black and white. *Prog Retin Eye Res* **91**, (2022).

35.   Ayala, G. D. *et al.* Assessing foveal structure in individuals with TYR
R402Q and S192Y hypomorphic alleles. *Ophthalmology Science* **1**,
100077 (2021).

36.   Jones, W. D. *et al.* De novo mutations in MLL cause Wiedemann-Steiner
syndrome. *Am J Hum Genet* **91**, 358–364 (2012).

37.   Choi, J., Joisher, H. N. V., Gill, H. K., Lin, L. & Cepko, C. Characterization
of the development of the high-acuity area of the chick retina. *Dev Biol* **511**,
39–52 (2024).

38.   da Silva, S. & Cepko, C. L. Fgf8 expression and degradation of retinoic
acid are required for patterning a high-acuity area in the retina. *Dev Cell*
42, 68-81.e6 (2017).

39.   Taneera, J. *et al.* Unraveling the significance of PPP1R1A gene in
pancreatic β-cell function: A study in INS-1 cells and human pancreatic
islets. *Life Sci* **345**, (2024).

40.   Keane, P. A. *et al.* Optical coherence tomography in the UK Biobank study
- Rapid automated analysis of retinal thickness for large population-based
studies. *PLoS One* **11**, (2016).

41.   Xie, Y. *et al.* FGF/FGFR signaling in health and disease. *Signal Transduct*
Target Ther **5**, (2020).

42.   Xu, X. *et al.* Fibroblast growth factor receptor 2 (FGFR2)-mediated
reciprocal regulation loop between FGF8 and FGF10 is essential for limb
induction. *Development* **125**, 753–765 (1998).

43.   Tomita, H. *et al.* The Protein Tyrosine Phosphatase Receptor Delta
Regulates Developmental Neurogenesis. *Cell Rep* **30**, 215-228.e5 (2020).

44.   Badde, A., Bumsted-O'Brien, K. M. & Schulte, D. Chick receptor protein
tyrosine phosphatase λ/ψ (cRPTPλ/cRPTPψ) is dynamically expressed at
the midbrain-hindbrain boundary and in the embryonic neural retina. *Gene*
Expression Patterns **5**, 786–791 (2005).

45.   Woertz, E. N. *et al.* Quantitative Foveal Structural Metrics as Predictors of
Visual Acuity in Human Albinism. *Invest Ophthalmol Vis Sci* **65**, (2024).

46.   Wilk, M. A. *et al.* Relationship between foveal cone specialization and pit
morphology in Albinism. *Invest Ophthalmol Vis Sci* **55**, 4186–4198 (2014).

47.   Kreis, J. & Carroll, J. Applications of Adaptive Optics Imaging for Studying
Conditions Affecting the Fovea. *Annu Rev Vis Sci* **35**, 239–262 (2024).

48.   Sampson, D. M., Dubis, A. M., Chen, F. K., Zawadzki, R. J. & Sampson, D.
815   D. Towards standardizing retinal optical coherence tomography
816   angiography: a review. *Light Sci Appl* **11**, 63 (2022).

817   49.   Dubis, A. M. *et al.* Relationship between the foveal avascular zone and
818   foveal pit morphology. *Invest Ophthalmol Vis Sci* **53**, 1628–1636 (2012).

819   50.   Yan, Q. *et al.* Deep-learning-based prediction of late age-related macular
820   degeneration progression. *Nat Mach Intell* **2**, 141–150 (2020).

821   51.   Warwick, A. N. *et al.* UK Biobank retinal imaging grading: methodology,
822   baseline characteristics and findings for common ocular diseases. *Eye*
823   *(Basingstoke)* **37**, 2109–2116 (2023).

824   52.   Munafò, M. R., Tilling, K., Taylor, A. E., Evans, D. M. & Smith, G. D.
825   Collider scope: When selection bias can substantially influence observed
826   associations. *Int J Epidemiol* **47**, 226–235 (2018).

827   53.   Schoeler, T. *et al.* Participation bias in the UK Biobank distorts genetic
828   associations and downstream analyses. *Nat Hum Behav* **7**, 1216–1227
829   (2023).

830   54.   Patel, P. J. *et al.* Spectral-domain optical coherence tomography imaging
831   in 67 321 adults: Associations with macular thickness in the UK biobank
832   study. *Ophthalmology* **123**, 829–840 (2016).

833   55.   Julian, T., Fitzgerald, T., Birney, E. & Sergouniotis, P. I. Pigmentation and
834   retinal pigment epithelium thickness: a study of the phenotypic and
835   genotypic relationships between ocular and extraocular pigmented tissues.
836   Preprint at https://doi.org/10.1101/2024.12.04.626809 (2024).

837   56.   Zhou, Y. *et al.* AutoMorph: automated retinal vascular morphology
838   quantification via a deep learning pipeline. *Transl Vis Sci Technol* **11**,
839   (2022).

840   57.   Romero-Bascones, D. *et al.* Foveal pit morphology characterization: A
841   quantitative analysis of the key methodological steps. *Entropy* **23**, (2021).

842   58.   Constantinescu, A. E. *et al.* A framework for research into continental
843   ancestry groups of the UK Biobank. *Hum Genomics* **16**, (2022).

844   59.   Auton, A. *et al.* A global reference for human genetic variation. *Nature* vol.
845   526 68–74 Preprint at https://doi.org/10.1038/nature15393 (2015).

846   60.   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and
847   population-based linkage analyses. *Am J Hum Genet* **81**, 559–575 (2007).

848   61.   Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation
849   of ancestry in unrelated individuals. *Genome Res* **19**, 1655–1664 (2009).

850   62.   Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Res* **50**, D988–D995
851   (2022).

852   63.   Ochoa, D. *et al.* Open Targets Platform: Supporting systematic drug-target
853   identification and prioritisation. *Nucleic Acids Res* **49**, D1302–D1310
854   (2021).

855   64.   Sollis, E. *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and
856   deposition resource. *Nucleic Acids Res* **51**, D977–D985 (2023).

857   65.   Myers, T. A., Chanock, S. J. & Machiela, M. J. LDlinkR: An R Package for
858   Rapidly Calculating Linkage Disequilibrium Statistics in Diverse
859   Populations. *Front Genet* **11**, (2020).

860   66.   Bulik-Sullivan, B. *et al.* LD score regression distinguishes confounding from
861   polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295
862   (2015).

863   67.   Davidson-Pilon, C. lifelines: survival analysis in Python. *J Open Source*
864   *Softw* **4**, 1317 (2019).

865   68.   Seabold, S. & Perktold, J. Statsmodels: Econometric and Statistical
866   Modeling with Python. *Proceeding of the 9th Python in Science Conference*
867   (2010).

868   69.   Burgess, S., Butterworth, A. & Thompson, S. G. Mendelian randomization
869   analysis with multiple genetic variants using summarized data. *Genet*
870   *Epidemiol* **37**, 658–665 (2013).

871   70.   Hemani, G. *et al.* The MR-Base platform supports systematic causal
872   inference across the human phenome. *Elife* **7**, (2018).

873   71.   Bowden, J. *et al.* Improving the visualization, interpretation and analysis of
874   two-sample summary data Mendelian randomization via the Radial plot
875   and Radial regression. *Int J Epidemiol* **47**, 1264–1278 (2018).

876   72.   Burgess, S. & Thompson, S. G. Interpreting findings from Mendelian
877   randomization using the MR-Egger method. *Eur J Epidemiol* **32**, 377–389
878   (2017).

879   73.   Burgess, S. *et al.* Guidelines for performing Mendelian randomization
880   investigations. *Wellcome Open Res* **4**, 186 (2019).

881
882

ACKNOWLEDGEMENTS

We acknowledge the following sources of funding: the Wellcome Trust

(224643/Z/21/Z, Clinical Research Career Development Fellowship to P.I.S.); the

UK National Institute for Health Research (NIHR) Clinical Lecturer Programme

(CL-2017-06-001 to P.I.S.); the NIHR Manchester Biomedical Research Centre

(NIHR 203308 to P.I.S. and G.C.B.); the EMBL European Bioinformatics Institute

(EMBL-EBI) (S.T., T.F. and E.B.); and the Howard Hughes Medical Institute

(C.L.C). Research reported in this publication was supported in part by the

National Eye Institute of the National Institutes of Health (NIH) under award

numbers R01EY017607 and R01EY033580. The content is solely the

responsibility of the authors and does not necessarily represent the official views

of the NIH.

The UK Biobank Eye and Vision Consortium is supported by funding from the

NIHR Biomedical Research Centre at Moorfields Eye Hospital and UCL Institute

of Ophthalmology, the Alcon Foundation and the Desmond Foundation. The

complete list of members of this Consortium can be found in the Supplementary

Information.

AUTHOR CONTRIBUTIONS STATEMENT

P.I.S. conceived and designed the experiments. The UK Biobank Eye and Vision

Consortium, D.R., H.N.V.J., U.A., M.B., T.F., E.B., C.L.C., J.C. and P.I.S

provided datasets and analytical tools. D.J.G., D.R., T.H.J., S.T., H.N.V.J., T.F.,

C.L.C., and P.I.S. analyzed the data. D.J.G. wrote the manuscript with support

from P.I.S. All authors critically revised and approved the manuscript.

COMPETING INTERESTS STATEMENT

E.B. is a paid consultant and equity holder of Oxford Nanopore, a paid consultant

to Dovetail, and a non-executive director of Genomics England, a limited

company wholly owned by the UK Department of Health and Social Care. All

other authors declare no competing interests. D.R. is an employee of RetinAI, but

performed this work independently of RetinAI.

UK Biobank Eye and Vision Consortium

Jay E. Self,[9,10] Graeme C. Black[1,11], Panagiotis I. Sergouniotis[1,3,5,11]

[1] Division of Evolution, Infection and Genomics, School of Biological Sciences,

Faculty of Biology, Medicine and Health, University of Manchester, Manchester,

UK.

[3] Manchester Royal Eye Hospital, Manchester University NHS Foundation Trust,

Manchester, UK.

[9] Clinical and Experimental Sciences, Faculty of Medicine, University of

Southampton, Southampton, UK

[10] Southampton Eye Unit, University Hospital Southampton NHS Foundation

Trust, Southampton, UK

[11] Manchester Centre for Genomic Medicine, Saint Mary's Hospital, Manchester

University NHS Foundation Trust, Manchester, UK

933
934 A full list of members appears in the Supplementary Information.

1

En Face

Central Retina ⌀ 5.5mm
Fovea ⌀ 0.8-1.5mm

Section

Fovea

OCT

2

biobank uk

Quality control

3

RETIMAT
Retinal image analysis in MATLAB

Radius

Slope

Height

Foveal morphological parameter extraction:
pit volume, rim radius, rim height, mean slope, pit depth, CFT

4

Low Risk Score
High Risk Score

Disease Free Survival

Time

Foveal phenotype studies:
- Phenotypic associations
- Time-to-event analysis

Genetic studies:
- Genetic associations
- Heritability estimates
- Genetic correlation