Quantitative Microbial Risk Assessment of Extended-Spectrum β-Lactamase-Producing
Escherichia coli **Transfer from Broiler Litter to Fresh Lettuce Consumption**

Nunzio Sarnino*[1], Subhasish Basak[2], Lucie Collineau[2], Roswitha Merle[1]

Nunzio.Sarnino@fu-berlin.de

[1]*Freie Universität Berlin, Institute of Veterinary Epidemiology and Biostatistics, Veterinary*
Centre for Resistance Research Berlin, Germany

[2]*University of Lyon - French Agency for Food, Environmental and Occupational Health and*
Safety (ANSES), Epidemiology and Surveillance Support Unit, Lyon, France

Abstract

Background. Extended-spectrum β-lactamase-producing *Escherichia coli* (ESBL *E. coli*) from
broiler chicken production pose potential public health risks via multiple environmental and
foodborne pathways. We developed a modular quantitative microbial risk assessment (QMRA)
model linking four components, namely farm, soil, river, and lettuce consumption, to predict
human environmental exposure to ESBL *E. coli* originating from broiler flocks.

Methods. A stochastic farm module simulated broiler colonization over a 36-day cycle and
generated end-cycle litter loads. Field modules represented first-order decay,
attachment/partitioning, and runoff to rivers; irrigation transfer yielded lettuce contamination for
a 100 g serving. Disability-Adjusted Life Year (DALY) estimation characterized health risk and
global sensitivity analyses identified main drivers.

Results. The farm model produced mean end-cycle litter of 1.6×10^4 (SD 16.1, UI 1.60–1.61 ×
10^4) CFU/g and near-complete flock colonization within one week. Soil surface loads declined
from 3.2×10^7 (SD 3.2×10^4, UI 3.2–3.2 × 10^7) CFU/m² to 8.6×10^5 (SD 8.8×10^2, UI 8.6–8.6 ×
10^5) CFU/m² by day 100. Runoff yielded river concentrations of 6.0×10^{-2} (SD 3.5×10^{-3}, UI
5.5–6.6 × 10^{-2}) CFU/mL after ten days. Lettuce contamination was highest at short intervals after
land application 1.7 (SD 8.5×10^{-2}, UI 1.5–1.8) CFU/100 g at one day interval, dropping to 0.85
(SD 5.0×10^{-2}, UI 0.76–0.95) CFU/100 g at day 20 interval; simple household washing cut
exposure by ~90 %. Global sensitivity analysis identified soil-water partitioning and decay rates
as most important parameters of exposure variability. Estimated Disability-Adjusted Life Year
(DALYs) lost ranged from 10^{-8} to 10^{-10} per serving.

Conclusion. In our scenarios, predicted health burdens varied with ESBL *E. coli* concentration in
irrigation water and intervals between litter application and lettuce planting. While environmental
decay and simple measures such as household washing substantially reduce exposure, residual
contamination persists. Future studies should evaluate the effectiveness of manure treatments and
irrigation-water quality interventions on reducing environmental loads and human risk.

Introduction

Antimicrobial resistance (AMR) is steadily reducing the effectiveness of antibiotics in human and veterinary medicine. In broiler farming, antimicrobial use may create strong selective pressure and accelerate the emergence and spread of resistant bacteria (Chantziaras et al. 2014). A prime example is extended-spectrum β-lactamase (ESBL)-producing *Escherichia coli (E. coli)*, which can break down β-lactam antibiotics. These organisms pose a serious public health threat and could increase healthcare costs (Leistner et al. 2014; MacKinnon et al. 2020; Apostolakos et al. 2019)

Resistant strains may enter broiler houses in multiple ways. Day-old chicks may arrive already colonized (Moreno et al. 2019), or resistance may pass vertically from breeder flocks (Dierikx et al. 2013; Agersø et al. 2014; Nilsson et al. 2014). Environmental contamination and biosecurity lapses offer additional routes (Mallioris et al. 2023; Robé et al. 2024). Although the prevalence of resistance at chick placement is generally low (Oikarainen et al. 2019), within-flock spread frequently leads to near complete colonization within a few days (Robé et al. 2019).

Once broiler houses are emptied, litter used as fertilizer potentially serves as a reservoir of ESBL *E. coli.* These bacteria can survive in soil for extended periods (van Overbeek et al. 2021), enter surface and groundwater through runoff (Sowah et al. 2020), and contaminate produce (Jensen et al. 2013; Mootian et al. 2009). Humans may then be exposed via contaminated food, recreational water contact, or other environmental pathways (O'Flaherty et al. 2018; O'Flaherty et al. 2019a; O'Flaherty et al. 2019b; Chapman et al. 2018). Environmental factors, notably rainfall events, can significantly influence bacterial concentrations in rivers, often leading to transient peaks (Alegbeleye and Sant'Ana 2020; Jacobs et al. 2019). Furthermore, the proximity to animal production sites seems to be an additional risk factor (Hubbard et al. 2020; Chapman et al. 2018).

Litter management may reduce ESBL *E. coli* loads. Thermophilic composting inactivates pathogens and lowers bacterial counts (Thomas et al. 2024; Okada et al. 2024). Respecting pre-harvest intervals limits the environmental spread (Subirats et al. 2021). Advanced composting and short-term litter storage further reduce resistant bacterial populations (Atanasova et al. 2025; Siller et al. 2020; Howard et al. 2016; Suzuki et al. 2024). However, bacterial survival depends on environmental conditions, underscoring the need for site-specific risk assessments and tailored interventions (Çekiç et al. 2017; van Overbeek et al. 2021; Sarnino et al. 2025).

Quantitative microbial risk assessment (QMRA) frameworks offer a systematic approach to characterize the propagation of resistant pathogens and to evaluate the potential effect of control measures (Collineau et al. 2020; Rawson et al. 2019; Fastl et al. 2023).

Within the ENVIRE project (https://www.envire-project.de/), we developed an integrated QMRA model to simulate how ESBL *E. coli* move from broiler farms under European conditions into the environment and reach humans. Our objectives were to:

1. Predict ESBL *E. coli* loads in raw broiler litter at the end of a conventional production
cycle,

2. Characterize transmission pathways from litter to soil, water, and crops,

3. Identify the dominant drivers of exposure at consumption and how the time between litter
application and planting shapes contamination at harvest,

4. Estimate human exposure and Disability-Adjusted Life Year (DALYs) lost via lettuce
consumption.

Other potential routes (for example direct crop contamination from soil) were not included
because our previous work (Sarnino et al. 2025) found that crop contamination primarily occurs
via irrigation water.
This study is, to our knowledge, the first modular Quantitative Microbial Risk Assessment
(QMRA) to connect broiler farm, amended soil, receiving water, lettuce consumption for ESBL
E. coli, using one internally consistent set of assumptions across modules. We (i) link flock
colonization dynamics to end-cycle litter loads, (ii) propagate soil partitioning and hydrological
wash-off to river concentrations, (iii) map irrigation transfer to edible portions, and (iv) quantify
health burden (DALY). This integrated framing extends prior single-pathway models and generic
E. coli assessments by targeting ESBL strains and explicitly testing time-since-application
scenarios relevant to produce safety.

Materials and Methods

Model Description

94 We built a modular QMRA model to estimate human exposure to ESBL *E. coli* from broiler
95 production. The model links four modules - farm, soil, river and lettuce consumption - to trace
96 contamination from its source all the way to human exposure (Figure 1).

97 **Figure 1.** Diagram of the QMRA model linking the farm, soil, river, and lettuce modules.

In the farm module, we simulated a 36-day broiler cycle with a stochastic susceptible-infectious compartmental model. Broilers enter as susceptible (S) or colonized (C) at a set prevalence (p_{init}) (Table 1). Gut bacterial growth (r) follows a logistic curve; infection may spread via ingestion (ρ_{ingest}) of contaminated feces (F_{barn}^{cont}). Each day, the model updates broiler age, infection duration, fecal output, and bacterial shedding. ESBL *E. coli* levels accumulate in the barn litter and then decline by natural decay.

Figure 2. Diagram of the farm module simulating the transmission dynamics of ESBL-producing *E. coli* within a broiler flock.

Throughout this article, we use the term "litter" to refer to the mixture of broiler manure, bedding, feathers, and other materials commonly removed at the end of a production cycle, while "clean litter" is the fresh bedding material added at the start of a production cycle.

The soil module applies broiler litter to an agricultural field - assuming zero delay between chicken harvest and litter spread - and models ESBL *E. coli* decay using an exponential rate fitted to experimental data from Sharma Manan et al. (2019). While poultry litter is often treated, this is not a universal or legally mandated practice across the EU. There are no specific regulations requiring all poultry litter to be treated or specifying an interval between removal from the barn and field application. Therefore, in our model, we consider a worst-case scenario in which broiler litter is applied to fields immediately after removal from poultry houses.

The river module estimates daily bacterial runoff from the field to the river using a simplified SWAT (soil and water assessment tool)-inspired approach (Douglas-Mankin et al. 2010). We considered water temperature, depth, solar radiation, and salinity, then applied wash-off fractions and in-water decay to predict river concentrations.

The lettuce module simulates multiple planting events over a growing season. On irrigation days, bacteria adhere to leaves in proportion to river concentration and retention volume. From irrigation to harvest, loads decrease according to a biphasic decay model, and post-harvest washing further reduces bacterial counts. Exposure is reported as colony-forming units per 100 g

128 serving. In this module, we assume that the lettuce is consumed on harvest day.
129

Across all modules, we employed a stochastic approach; each module ran 1000 simulations to
reflect natural variability in environmental conditions and individual behaviour. For each result,
we defined the standard deviation (SD) as descriptive measure of dispersion around the mean and
the 95% uncertainty interval (UI) as the 2.5th and 97.5th percentiles of the resulting empirical
distribution.

Model credibility. We harmonized units and variable definitions across module, constrained
inputs to ranges reported in peer-reviewed studies or agronomic practice, and ran all analyses
from scripted, version-controlled code. To support external validity, we planned piecewise checks
against independent evidence, comparing our results with experimental studies and other QMRA.

Farm Module

The farm module simulates the transmission dynamics of ESBL *E. coli* between two primary
reservoirs: the broilers' gastrointestinal tract and the litter within the barn environment. A single
simulation of this stochastic module corresponds to a complete broiler production cycle of $D_{\max}$
days of a single flock of broilers. The module operates within a discrete-time framework with a
daily time step. For each day $d = 1, 2, ..., D_{\max}$ of the production cycle, a series of barn-level
events, each representing distinct mechanisms in the transmission dynamics of ESBL *E. coli*
within the barn environment, is executed once per day. These events form the core components of
the farm module and are detailed following the model initialization.

Table 1. Key input variables for the farm module

Variable	Description	Source	Unit	Value
α	Proportion of ingested water lost by metabolism	Becker et al. (2022)	proportion	0.7
ϵ	Excretion rate of ESBL *E. coli*	Becker et al. (2022)	%	0.3
$D_{\max}$	Total duration of the fattening period	Assumed	day	36
$r_{\min}$	Minimum growth rate for ESBL *E. coli* in the broiler's intestine	Becker et al. (2022)	log10 CFU	0
$r_{\max}$	Maximum growth rate for ESBL *E. coli* in the broiler's intestine	Becker et al. (2022)	log10 CFU	5
K	Maximum amount of bacteria in the substrate (the intestinal content in this case)	Robé et al. (2019)	CFU/g	10^6
ϕ	Environmental decay rate of ESBL *E. coli* in broiler litter	Becker et al. (2022)	%	0.5

ρ_{ingest}	Proportion of ingested contaminated feces along with feed	Becker et al. (2022)	%	0.014
β	Transmission rate	Dame-Korevaar et al. (2020)	CFU/day	0.31
$w_{\text{min}}(d)$	Minimum water consumption on day d	Avian advice 11(2), University of Arkansas. (2009)	g	21.16–10.35
$w_{\text{max}}(d)$	Maximum water consumption on day d	Avian advice 11(2), University of Arkansas. (2009)	g	42.66–31.18
$g(d)$	Broiler's average daily gain in weight on day d	Ross308 performance objectives. (2022)	g	21–100
$I(d)$	Broilers average feed intake on day d	Ross308 performance objectives. (2022)	g	20–189
A	Size of the farm	Assumed	m^2	100
ρ_{farm}	Stocking density	Directive 2007/43/EC — minimum rules for the protection of chickens kept for meat production	kg/m^2	39
w_{tar}	Weight of the broilers at the end of the fattening period (36-th day)	Becker et al. (2022)	kg	2.332
c_{init}	Total CFU of ESBL *E. coli* in colonized one day old chicks	Becker et al. (2022)	CFU	100
p_{init}	Initial prevalence of colonized broilers	Expert opinion, Robé	Fraction	0.01
L	Litter quantity per square meter	Becker et al. (2022)	g/m^2	1000

Module initialization

154   The farm module is initialized using the total number of broilers in a flock computed as,

$$N = \left(\frac{\rho_{\text{farm}}}{w_{\text{tar}}} \times A \right), \tag{1}$$

where ρ_{farm} is the farm stocking density (kg/m^2), w_{tar} is the target weight (in kg) at day D_{max},
and A is the farm area (m^2). Further let $N_{\text{col}}(d)$, for $d = 1, 2, \ldots, D_{\text{max}}$, denote the total number of
colonized broilers in the flock on day d, with an initial proportion p_{init} used to set $N_{\text{col}}(1)$. The
flock is partitioned into two compartments: susceptible (S) and colonized (C), with no recovery
assumed during the production cycle; their respective initial sizes are $(N - N_{\text{col}}(1))$ and $N_{\text{col}}(1)$.

For $i \in C \cup S$ and $d = 1, 2, \ldots, D_{\text{max}}$, let $C_{gut,i}^{init}(d)$ denote the initial total number of CFUs of
ESBL *E. coli* present in the i-th broiler's gut on day d. $C_{gut,i}^{init}(1)$ is set to 0 CFU for every $i \in S$

and $C_{gut,i'}^{init}(1)$ is set to c_{init} CFUs, for every $i' \in C$. Moreover, on day 1, the barn litter is
assumed to be completely free of ESBL *E. coli*, with no CFUs present. Consequently, the value of
both $C_{barn}^{init}(1)$ and $F_{barn}^{cont}(1)$, representing respectively the total initial CFU count and amount of
contaminated feces in the litter on day 1, are set to zero.

Feces excretion

The first barn-level event implemented by the farm module is the production of feces by the
broilers. On day d, for $d = 1, 2, ..., D_{max}$, the i-th broiler's feces output denoted by $F_i(d)$, for $i \in$
$C \cup S$, is calculated as,

$$F_i(d) = w(d) \cdot \alpha + I(d) - g(d). \tag{2}$$

Here $w(d)$ is a uniformly distributed random variable with limits $w_{min}(d)$ and $w_{max}(d)$,
representing respectively the day-specific bounds on water consumption of broilers. α is a
fractional reduction factor for water. $I(d)$ and $g(d)$ respectively denote the average feed intake
(kg/day) and daily weight gain (kg/day) of a broiler Ross 308. After the excretion of day, the total
amount of contaminated feces in the barn environment till the beginning of day $(d + 1)$ is given
by,

$$F_{barn}^{cont}(d + 1) = F_{barn}^{cont}(d) + \sum_{i' \in C,}^{N_{col}(d)} F_{i'}(d). \tag{3}$$

181 **ESBL *E. coli* excretion**

182 The event ESBL *E. coli* excretion models the excretion of ESBL *E. coli* in the barn environment
183 along with the feces by the colonized broilers. The total number of CFUs of ESBL *E. coli*
184 remaining in the i'-th colonized broiler's gut, for $i' \in C$, on day d after excretion, is given by,

$$C_{gut,i'}^{post.exc}(d) = C_{gut,i'}^{init}(d) \times (1\text{-}\epsilon), \tag{4}$$

187 where ϵ is the excretion rate. The total cumulated load of ESBL *E. coli* (in CFUs) in the litter
188 after excretion on day d, for $d = 1, 2, ..., D_{max}$, is obtained as,

$$C_{barn}^{post.exc}(d) = C_{barn}^{init}(d) + \sum_{i' \in C}^{N_{col}(d)} C_{gut,i'}^{init}(d) \times \epsilon, \tag{5}$$

190 **ESBL *E. coli* growth inside broilers' guts**

191 Within colonized broilers, bacterial growth event in the gut is modeled with a logistic growth
192 function (see, e.g., Tsoularis and Wallace 2002). With K being the carrying capacity and the
193 growth rate $r_{i'} \sim \mathcal{U}(r_{\min}, r_{\max})$ distributed uniformly, the number of CFUs of ESBL *E. coli* in the
194 gut of i'-th colonized broiler, for $i' \in C$, on day d after the growth event is given by:

195
$$C_{gut,i'}^{post.growth}(d) = \frac{K}{1 + \left(\dfrac{K - C_{gut,i'}^{post.exc}(d)}{C_{gut,i'}^{post.exc}(d)} \right) \exp(-r_{i'})} \tag{6}$$

This formulation captures saturation effects and within-flock variability through random draws
for $r_{i'}$.

Feces ingestion

The next barn-level event is the ingestion of feces where each broiler is assumed to accidentally
ingest a certain proportion ρ_{ingest} of contaminated fecal material from the barn environment (see,
e.g., Becker et al., 2022), along with their daily feed. This triggers the colonization of susceptible,
non-colonized broilers, while the already colonized ones remain persistently colonized. The total
amount of contaminated feces (in g) accidentally ingested by the i-th broiler, for $i \in C \cup S$, on
day d is given by,

205
$$F_{ingest,i}^{cont}(d) = I(d)\rho_{\text{ingest}}. \tag{7}$$

Let $N_{\text{new}}(d)$ denote the expected number of newly colonized broilers on day d through accidental
ingestion of contaminated feces. At this stage, the farm module converts the same number of
susceptible broilers to the colonized state by simple random sampling without replacement.
Simultaneously at this moment, the flock is re-partitioned into three mutually exclusive and
exhaustive compartments, namely, susceptible (S), newly colonized (C^*) on day d, and
previously colonized (C), with respective sizes $(N - N_{\text{col}}(d) - N_{\text{new}}(d))$, $N_{\text{new}}(d)$ and $N_{\text{col}}(d)$.
The quantity of ESBL *E. coli* ingested by already-colonized broilers is considered negligible
compared with their gut load and is therefore neglected. The gut concentrations for broilers
belonging to all the three compartments are updated as,

215
$$C_{gut,i}^{init}(d+1) = \begin{cases} F_{ingest,i}^{cont}(d) \dfrac{C_{barn}^{post.exc}(d)}{F_{barn}^{cont}(d+1)}, & \text{for } i \in C^*, \\ C_{gut,i}^{post.growth}(d), & \text{for } i \in C, \\ C_{gut,i}^{init}(d), & \text{for } i \in S. \end{cases} \tag{8}$$

216 Simultaneously the ESBL *E. coli* load in the barn environment is also updated as,

217
$$C_{barn}^{post.ingest}(d) = C_{barn}^{post.exc}(d) - \sum_{i'' \in C^*}^{N_{\text{new}}(d)} F_{ingest,i''}^{cont}(d) \dfrac{C_{barn}^{post.exc}(d)}{F_{barn}^{cont}(d+1)}. \tag{9}$$

218 However, the quantity of contaminated feces ingested from the barn environment by all the
219 broilers is considered negligible compared to $F_{barn}^{cont}(d+1)$, and hence it is kept unchanged.

Transmission

The transmission event is used to estimate $N_{\text{new}}(d)$. The framework proposed by Dame-Korevaar (see, e.g., Dame-Korevaar et al. 2020a) is adapted as a conceptual basis but extended to honour flock heterogeneity and mass balance constraints. The module is based on the following assumptions, (1) broilers acquire ESBL *E. coli* only by accidentally ingesting contaminated litter and broiler-to-broiler transmission through direct contact is ignored; (2) the simulation operates on a fixed time-step of 1 day, so each broiler experiences at most one exposure decision per day. The force of infection $\lambda(d)$ relates to the environmental concentration of ESBL *E. coli* to $N_{\text{new}}(d)$, and is derived as,

$$\lambda = \beta \times \log_{10}\left(\frac{C_{barn}^{post.exc}(d)}{F_{barn}^{cont}(d+1)}\right), \tag{10}$$

where β is the transmission coefficient. The expected number of new colonization during the interval on day d is obtained as,

$$N_{\text{new}}(d) = \left(N - N_{\text{col}}(d)\right)\left[1 - \exp(-\lambda(d))\right]. \tag{11}$$

On day $(d+1)$, the two compartments C^* and C are merged as C and the corresponding size is updated as $N_{\text{col}}(d+1) = N_{\text{col}}(d) + N_{\text{new}}(d)$. This ensures that the newly colonized broilers contribute to excretion of contaminated feces and experience growth of ESBL *E.coli* inside their guts.

Environmental decay

The final barn-level event is the environmental decay which models the reduction in ESBL *E. coli* population in the barn environment due to absence of suitable growth conditions and thus preventing indefinite bacterial build-up. For a decay rate ϕ the initial barn load for the next day after environmental decay, is obtained as,

$$C_{barn}^{init}(d+1) = C_{barn}^{post.ingest}(d) \times (1 - \phi). \tag{12}$$

Final concentration calculation

At the end of a single simulation of the farm module on day D_{max}, the final ESBL *E. coli* concentration in litter (CFU/g) denoted by C_{final}, is obtained by dividing the total environmental load by the sum of accumulated feces and the mass of fresh bedding L,

$$C_{\text{final}} = \frac{C_{barn}^{init}(D_{\text{max}}+1)}{F_{\text{total}} + L}, \tag{13}$$

where $F_{\text{total}} = \sum_{d=1}^{D_{\text{max}}} \sum_{i \in C \cup S}^{N} F_i(d)$ is the aggregated feces mass by day D_{max}.

Soil Module

All parameter values and distributional assumptions for the soil and river modules can be found in Table 2.

E. coli decay in soil

Empirical data on *E. coli* decay in litter-amended soil were obtained from Sharma Manan (Sharma Manan et al. 2019). Data consisted of log-transformed *E. coli* concentrations in the soil measured at several time points (days post-inoculation, *dpi*).

To characterize *E. coli* decline, an exponential decay function was assumed to follow:

$$y = C \times exp(-k \times t) \tag{14}$$

where y is the log-transformed *E. coli* concentration at time t (day), C is the initial concentration parameter, and k is the decay rate constant. The parameters C and k were estimated via nonlinear least squares:

$$\operatorname*{argmin}_{C,k} \sum_{i=1}^{n} (y_i - C \times \exp(-k \times t_i))^2 , \tag{15}$$

Litter Application

The temporal evolution of the ESBL *E. coli* load on the soil was modeled for each Monte Carlo simulation i by a first-order decay of the initial litter-derived load. In our model, we therefore describe:

$$L_i(t) = C_{\mathrm{man},i} \times R_{\mathrm{app}} \times 1000 \times \exp(-k_{\mathrm{soil}} t), \tag{16}$$

where $L_i(t)$ (CFU/m²) is the ESBL *E. coli* load at time t (days) after application, $C_{\mathrm{man},i}$ (CFU. g⁻¹) is the Monte Carlo draw of the litter concentration from the farm module output C_{final}, R_{app} (kg/m²) is the application rate, the factor 1000 converts kilograms to grams, and k_{soil} is the decay constant.

River Transport and Decay Model

We route the soil-derived load $L_i(t)$ into a simplified runoff and in-stream decay scheme, for
each Monte Carlo iteration i and day t. Surface wash-off is computed once per day based on a
constant fraction of freely mobile bacteria, then that wash-off load is instantly delivered to the
receiving water body (i.e. travel time is assumed negligible on the daily time-step).

Wash-off load

$$W_i(t) \ = \ L_i(t) \ \times \ A \ \times \ (1 - K_d) \ \times \ f_{\text{wash}} \tag{17}$$

Here, $W_i(t)$ (CFU/day) is the total bacterial load washed off from the field on day t in simulation
i. $L_i(t)$ (CFUm²) is the soil-module output, A (m²) is the field area, K_d is the partition coefficient
(fraction of bacteria sorbed to soil), and f_{wash} is the wash-off fraction of freely mobile bacteria.
We compute wash-off once per day as a fixed fraction of the field load that is free for transport,
representing the expected (event-averaged) daily export due to rainfall/runoff. By routing
$W_i(t)$ directly into the river on the same day, we are effectively assuming that the farm-to-stream
distance is short enough that travel time can be ignored at daily resolution.

In-stream decay rate
Mancini's Equation *(Mancini 1978)* was applied to capture in-river *E. coli* decay as a function of
water temperature (T), salinity, and solar radiation. The daily first-order decay rate k was
described as:

$$k = 0.8 + 0.006 \ \times \ EC \ \times \ 1.07^{(T-20)} + \frac{I \ \times \ (1 - \exp(-e \ \times \ H))}{e \ \times \ H}, \tag{18}$$

where EC is salinity (%), I is the solar radiation (ly/hr), e is the light extinction coefficient, and H
is water column depth (m).

Accumulated river load

$$M_i(t) \ = \ M_i(t-1) \ \exp(-k) \ + \ W_i(t), M_i(0) = W_i(0) \tag{19}$$

Where $M_i(t)$ (CFU) is the cumulative ESBL *E. coli* count in the water site on day t. Each day,
the previous day's count decays by $\exp(-k)$ and the new wash-off input $W_i(t)$ is added.

Dividing the accumulated $M_i(t)$ by the site volume V (L) yields *the* ESBL *E. coli* concentration
$C_i(t)$ In colony-forming units per liter:

$$C_i(t) \ = \ \frac{M_i(t)}{V} (\text{CFU L}^{-1}) \tag{20}$$

Table 2 Key input variables for the soil, river, exposure and lettuce-exposure modules

Variable	Description	Source	Unit	Value
k	First-order die-off rate of ESBL-E. coli in litter-amended top-soil	Sharma Manan (2019)	/Day	0.0362
R_{app}	Broiler litter applied to field surface	Assumed	Kg/m²	2
K_d	Bacteria partition coefficient	Sowah (2020)	Fraction	0.95
W_i	Mobile-phase cells washed off per run-off event	Sowah (2020)	Fraction	0.50
T	River-water temperature	O'Flaherty 2019a	°C	21–28
EC	Salinity	O'Flaherty 2019a	%	0.035–0.075
IA	Global solar irradiance	O'Flaherty 2019a	Ly/h	17.3–25.4
et	Light-extinction coefficient	O'Flaherty 2019a	m	0.26–0.31
H	Mean water-column depth	O'Flaherty 2019a	m	0.5–6
V	Effective pool volume at water site	O'Flaherty 2019a	L	6.75×10^7–8.25×10^7

Time notation (soil and river modules). We define continuous time t as days since litter
application, with $t = 0$ at application. Simulation outputs are stored and reported on
discrete post-application days d $d = 1,2, ...$, where day d represents the interval $t \in (d -$
$1, d]$. The soil module outputs loads $d \geq 1$, the rive**r** module uses the same day index with
zero initial load at $t = 0$ and updates concentrations from the day d export and in-stream
decay.

Lettuce module

To estimate *E. coli* contamination on lettuce, we linked daily irrigation-water concentrations
$C_i(t)$ (CFU/mL) from the river model to leaf-surface adhesion and biphasic die-off on the crop
(Table 3), following a previously proposed framework by O'Flaherty (O'Flaherty et al. 2019b).
We evaluated scenarios with different intervals between land application of litter and lettuce
planting. We define ΔAP as the days between application of litter and planting, with the
environmental time origin at $t = 0$ on the day of application, thus planting occurs at $t = \Delta AP$. In
the lettuce module, irrigation begins on the planting day and proceeds daily through the growth
period (35 days), so ΔAP is operationally equivalent to the interval from application to the first
irrigation event. Leaf die-off is indexed on a leaf clock $\tau = 1 ... 35$.

Table 3. Key input variables for the lettuce module

Variable	Description	Source	Unit	Value
ΔAP	Days between litter application and lettuce planting	User defined	days	1, 20, 50, 100, 150
f	Fast-phase fraction in biphasic decay	O'Flaherty 2019b	Fraction	7.6×10^{-5}
k_1	Fast decay coefficient	O'Flaherty 2019b	day	4.45
k_2	Slow decay coefficient	O'Flaherty 2019b	day	0.0698
$V_{\text{attach,i}}$	Leaf-film water volume	O'Flaherty 2019b	mL/g	$0.006 + \text{LN}(-4.75, 0.50)$

D_{treat}	Producer-level washing reduction factor	O'Flaherty 2019b	fraction	0.826
$D_{\text{wash},i}$	Consumer washing efficiency	O'Flaherty 2019b	fraction	Triangular 0.65, 0.99, 0.99
$S_{portion}$	Portion size for risk estimate	O'Flaherty 2019b	g	100
T	Days from planting to harvest	User defined	day	35
r_{DR}	Exponential dose-response parameter	Tacket 2000	CFU	2.18×10^{-6}
F_{UPEC}	Fraction of *E.coli* that are uropathogenic	Heida 2025	fraction	0.1
$p_{\text{colon|gut}}$	P(urinary colonisation | gut colonisation)	Heida 2025	probability	Uniform 0.35, 0.46
$p_{\text{UTI|colon}}$	P(symptomatic UTI | urinary colonisation)	Heida 2025	probability	0.067
δ	DALY burden per UTI case	Heida 2025	DALY/case	Uniform 3.7, 12.84

339
340 **Leaf-surface adhesion**

$$E_{\text{adh},i}(t) = C_i(t) \times V_{\text{attach,i}} \tag{21}$$

344 where $E_{\text{adh},i}(t)$ (CFU/g) is the bacteria adhering to leaves, and $V_{\text{attach,i}}$ (mL/g) is the water-
345 attachment volume.

346
347 **Biphasic decay on leaves**

348
$$\log_{10}\left[E_{\text{obs},i}(t)\right] = f\,(-k_1\,t) + (1-f)\,(-k_2\,t) \tag{22}$$

350   and

$$E_{\text{decay,i}}(t) = E_{\text{adh,i}}(t) \times 10^{\log_{10}[E_{\text{obs,i}}(t)]} \tag{23}$$

352   where a fraction f of cells decays at the fast rate k_1 (/day) and the remainder $(1 - f)$ at the slower
353   rate k_2 (/day). Here, t is days since first irrigation after planting.

355   **Accumulated contamination at harvest**

$$E_{\text{cum,i}} = \sum_{t=p}^{p+T-1} E_{\text{adh,i}}(t) - \sum_{t=p}^{p+T-1} E_{\text{decay,i}}(t) \tag{24}$$

357   where p is the planting day, T is the growth duration (35 days), and the sums collect adhesion and
358   decay over the growth period to compile the accumulated contamination at harvest $E_{\text{cum,i}}$ **Post-**
359   **harvest washing**

362   A washing reduction factor D_{treat} represents the fraction of bacteria removed by producer washing
363   with water. The concentration at harvest is calculated as:

$$E_{\text{post},i} = E_{\text{cum},i} \times D_{\text{treat}} \tag{25}$$

366   Furthermore, we modelled two scenarios: the consumption of 100g of lettuce $S_{portion}$ by
367   consumer of not further washed lettuce, as:
368
$$E_{\text{unwashed},i} = S_{portion} \times E_{\text{post},i} \tag{26}$$

371   and the consumption of $S_{portion}$ with an additional home washing ($D_{\text{wash},i}$), as:

372
$$E_{\text{wash},i} = E_{\text{post},i} \times (1 - D_{\text{wash},i}), E_{\text{final},i}^{100\text{g}} = S_{portion} \times E_{\text{wash},i} \tag{27}$$

376   **Probability of infection and DALY estimation**
377

We mapped lettuce dose (CFU per 100 g serving) to health outcomes using a pooled exponential
dose-response for enteropathogenic *E. coli* (EPEC) with a fecal-shedding endpoint as a surrogate
for gastrointestinal (GI) colonization. The parameterization ultimately traces to human volunteer
challenge with EPEC O127:H6 (strain E2348/69); colonization in those studies was determined
by recovery/shedding of the challenge strain from stools (primary source and trial details in
Tacket et al., 2000). We adopted this surrogate because ESBL- or UPEC-specific human DR data
are not available, using the approach taken by Heida et al., 2025 for ESBL *E. coli* in recreational
water.
To translate GI colonization to UTI burden, we followed Heida's framework: (i) apply a UPEC-
like fraction to the ingested ESBL population; (ii) use conditional probabilities for urinary tract
colonization given gut colonization and for symptomatic UTI given urinary colonization; and (iii)
compute DALY per serving from outcome probabilities and published disability
weights/durations.
391
392 The probability of GI is calculated as:

393
$$P_{\text{gut}} = 1 - \exp(-r_{\text{DR}} E_k F_{\text{UPEC}}) \tag{28}$$

401
402
where r_{DR} is the exponential dose-response parameter ,E_k is the ingested dose (CFU) in a single
event (lettuce consumption) and F_{UPEC} is the fraction of ESBL *E.coli* that are uropathogenic.
Chicken feces (and thus litter) commonly contain *E. coli* populations in which a minority meet
ExPEC criteria, and some of these are UPEC-like; published prevalences range from ~13%
ExPEC in broiler feces (with many categorized as UPEC) to ~23% ExPEC in retail poultry meat.
We therefore set $F_{\text{UPEC}} = 0.10$ as a realistic, conservative central, in accordance with Heida et al.,
2025 (Stromberg et al. 2017; Mellata 2013; Lyhs et al. 2012; Literák et al. 2012).

The probability of UTI is estimated as:

404
$$P_{\text{UTI}} = P_{\text{gut}} \times p_{\text{colon|gut}} \times p_{\text{UTI|colon}} \tag{29}$$

407 where $p_{\text{colon|gut}}$ is the conditional probability of urinary tract colonization given gut colonization
408 and $p_{\text{UTI|colon}}$ is the conditional probability of symptomatic UTI given urinary tract colonization.

409 Finally, we estimated the DALY as:
410
$$\text{DALY} = P_{\text{UTI}} \times \delta \tag{30}$$

412 where δ is the DALY burden per UTI case (DALY/case).

Sensitivity Analysis

We performed a variance-based sensitivity analysis using Partial Rank Correlation Coefficients (PRCC) to quantify how variability in the model inputs propagates to uncertainty in the litter load and in the final exposure outputs.

First, we identified key parameters spanning from each module, each varied uniformly over ±50 % of its baseline value. We generated 1000 stratified Latin Hypercube Sampling (LHS) draws across the n-dimensional input space, ensuring full coverage of each parameter's marginal distribution without clustering.

For each of the 1000 parameter vectors, we executed the complete simulation chain, extracting the mean human exposure. Runs were parallelized with independent L'Ecuyer- Combined Multiple Recursive Generator substreams (L'Ecuyer, 1999).

Once the vector of output metrics was assembled, we computed PRCCs and derived 95 % confidence intervals through 100 bootstrap replicates. PRCC measures the monotonic relationship between each input and the outcome, controlling for all other parameters; those whose intervals excluded zero were deemed statistically significant.

Software

All analyses and simulations were implemented in R (version 4.3.1) and RStudio (version 2024.09.1 build 394). Key packages including tidyverse (Wickham et al. 2019) for data manipulation and visualization, mc2d (Pouillot et al. 2016) and triangle (Carnell 2022) for distribution sampling, nls2 (Grothendieck 2013) for nonlinear regression, furrr (Bengtsson-Palme et al. 2023) for parallel computation, sensitivity (Iooss et al. 2024), and lhs (Carnell 2016) for sensitivity analysis. The full QMRA model is openly available on ENVIRE GitHub, https://github.com/ENVIRE-JPIAMR.

Results and discussion

Farm module
ESBL *E. coli* concentration in litter rose sharply after day 2 (16×10^{-3} CFU/g, SD 5×10^{-4}, UI 7.6 $\times 10^{-4}$–2.7×10^{-4}), reaching a maximum at day 8 (3.7×10^{4} CFU/g, SD 1.7×10^{3}, UI 3.33–3.90 $\times$

442 10^4), before declining to 1.6×10^4 CFU/g (SD 16.1, UI 1.6–1.60 x10^4) at day 36 (Figure 2).

444 **Figure 2**. ESBL *E. coli* concentration (CFU/g) in litter over the broiler production period. Mean (dark red
445 line) with 95 % uncertainty interval (shaded grey; 2.5th–97.5th percentiles) of 1000 iterations.

447 The proportion of colonized broilers climbed from 65% (UI 11–76%) on day 3 to more than 99.9
448 % by day 7 (UI 99–99%), remaining essentially constant thereafter (Figure 3).

449
Figure 3. Prevalence of colonized broilers over time. Mean (blue line) with 95 % uncertainty interval
(shaded grey; 2.5th–97.5th percentiles) of 1000 iterations.

The farm module simulates a rapid and efficient spread of ESBL *E. coli* within broiler flocks,
consistent with field data and experimental studies (Ceccarelli Daniela et al. 2017;
Kasparaviciene et al. 2025; Huijbers et al. 2016; Robé et al. 2021). In our simulation, flock-level
colonization approached 100% within the first week. This pattern is similar to longitudinal studies
where prevalence in day-old chicks often rises from negligible levels to near saturation within 5
to 7 days (Huijbers et al., 2018; Kasparaviciene et al., 2024; Dierikx et al., 2013). For example,
Kasparaviciene et al. (2024) detected no ESBL *E. coli* on day 0, but found a 57.5% prevalence by
day 5, while Huijbers et al. (2018) reported increases from 0-24% at day 1 to over 96% by day 7.
These dynamics are biologically plausible given the extremely low infectious dose required for
colonization, as few as 10 to 100 CFU can establish persistent colonization in broilers (Robé et al.
2019; Robé et al. 2021).

The concentration of ESBL *E. coli* in the litter estimated by this module aligns with the lower to
mid-range of reported concentrations, though literature values vary widely. Tofani et al. (2022)
measured ESBL/AmpC-*E. coli* loads in broiler cecal contents between 2.85 and 4.17
$\log_{10}$ CFU/g, while Atanasova et al. (2025) reported up to 5.48 $\log_{10}$ CFU/g in fresh broiler litter.
At the lower end, Blaak et al. (2015) found a mean concentration of 2.4×10^4 CFU/kg in feces
from Dutch farms. Also Becker et al. (2022) quantified a lower number of CFU/g in broiler litter.
Such discrepancies are likely influenced by multiple factors including broiler age, diet, specific
bacterial strains, litter conditions, sampling methods, and farm management. The model's
simplified assumption of logistic growth and a constant excretion rate does not fully capture
transient increases in shedding that may occur due to stressors such as thinning, transport, or diet.
As a result, the model may underestimate peak shedding under certain real-world scenarios.
Finally, our model underlines a critical message: once ESBL-producing *E. coli* is introduced in
the barn, it spreads rapidly and is difficult to suppress (Robé et al. 2024; Luyckx et al. 2015).
Because broiler colonization requires only a minimal infectious dose, even small biosecurity
lapses at chick placement can result in widespread contamination.

In our simulation, we applied a Susceptible-Infected (colonized) (SI) transmission model,
assuming that once a broiler becomes colonized it remains so until the end of the production
cycle. Persistent colonization of gut bacteria like *E. coli* in broilers is well documented (Robé et
al. 2019), yet this simplification may overlook more complex within-host dynamics (Galarneau et
al. 2020). Dankittipong et al. (2023) compared both SI and Susceptible-Infectious-Susceptible
(SIS) frameworks to study carbapenemase- and ESBL-producing *E. coli*, highlighting different
transmission rates among resistant strains. Likewise, Becker et al. (2022) and Dame-Korevaar et
al. (2020b) found the SI model adequate for capturing within-flock spread in broilers. By
contrast, Furusawa et al. (2024) developed two compartmental models - one with a time-
dependent decline in susceptibility and one with partial immunity to phylogenetic groups - to
describe ESBL-producing *E. coli* transmission in Dutch broiler chains. Those models also
incorporated environmental contamination between production cycles and within flocks, offering
a more detailed framework for testing interventions and estimating public health risk.

Nevertheless, given the evidence for long-term carriage in broiler chickens, we judged the SI
model the most consistent with our "worst case scenario" approach, since it reflects a permanent
infection state once colonization occurs.
499
The sensitivity analysis of the farm module highlights three overwhelmingly dominant drivers of
within-flock bacterial loads (Figure 4). First, the on-farm decay rate ϕ shows a very strong
negative correlation with mean litter *E. coli* concentration (PRCC ≈ -0.98), indicating that small
increases in bacterial die-off sharply reduce end-cycle loads. Conversely, gut carrying capacity K
(PRCC ≈ +0.93) and the shedding rate ϵ (PRCC ≈ +0.93) both exhibited very strong positive
correlations, underscoring that higher maximal gut loads and greater per-bacterium excretion
rates drive litter contamination. Secondary factors include water metabolic loss α (PRCC ≈ -0.16)
and target broiler weight w_{tar} (PRCC ≈ -0.04), which have moderate negative effects. All other
parameters have PRCCs near zero with confidence intervals spanning zero, indicating negligible
impact under the studied ranges. These results point to the critical importance of both bacterial
die-off dynamics and host-level factors (gut capacity and shedding) in shaping ESBL *E. coli*
loads on broiler farms.

Figure 4. Sensitivity analysis litter ESBL *E. coli* load (CFU/g litter) at day 36; displayed is the mean
Partial Rank Correlation Coefficient including 95% confidence intervals

Soil & River modules

When expressed per m², the mean ESBL *E. coli* loads started at 3.2×10^7 (SD 3.2×10^4, UI 3.2–
3.2×10^7) CFU/m² on day 1 post litter application, fell to 2.2×10^7 (SD 2.2×10^4 ,UI 2.2–2.2 ×
10^7) CFU/m² at day 10, and declined further to 8.6×10^5 (SD 8.8×10^2, UI 8.6–8.6×10^5)
CFU/m² by day 100. This per-area view mirrors the per-mass decay and confirms that fields
receive a substantial bacterial pulse immediately after litter application, but surface-area
concentrations drop substantially within three months. Consequently, while broiler litter is a
pronounced short-term source of soil contamination, its long-term contribution to human
exposure via soil contact is minimal under baseline conditions. A limitation of our model is the
lack of studies to use for validating the soil module. Most of the literature focuses on generic *E.
coli* (Sarnino et al., 2025) or uses CFU/g of soil mixed with manure as quantitative output
(Gekenidis et al. 2020). It is also important to specify that another big limitation of our model is
the simplified litter application, as we assumed the ESBL *E. coli* to spread uniformly following a
fixed application rate.

Simulated ESBL *E. coli* in river water decreased from 8.3×10^{-2} CFU/mL (SD 4.8×10^{-3}, UI 7.6–
9.2×10^{-2}) on day 1 to 6.0×10^{-2} CFU/mL (SD 3.5×10^{-3}, UI 5.5–6.6×10^{-2}) by day 10 and
reached 6.2×10^{-5} CFU/mL (SD 3.6×10^{-6}, UI 5.7–6.9×10^{-5}) by day 200. Early declines were
modest, reflecting dilution and decay; by day 200, concentrations were essentially zero.
These magnitudes fall within reported bathing-water ranges for AR/ESBL *E. coli* (predicted
0.45–345.09 CFU/100 mL as summarized by O'Flaherty et al., 2019a) and are consistent with
ESBL fractions (0.05–1%) applied to recreational criteria (≈0.06–4.1 CFU/100 mL) in Heida et
al., 2025.

Lettuce module

Our model simulated the contamination of lettuce via irrigation with water containing ESBL *E.
coli*, predicting contamination levels at harvest influenced by river contamination, application
practices, and post-irrigation environmental factors, with risk reduction achieved through
standard household washing. In our simulation, we estimated the exposure via ingestion of both
washed by final consumers and unwashed lettuce. Furthermore, we compared harvest-time
exposure across various intervals between litter application and lettuce planting. With a one-day
interval, mean exposure reached 1.7 CFU/100 g (SD 8.5×10^{-2}, UI 1.5–1.8) for unwashed lettuce
and 0.2 CFU/100g (SD 0.1, UI 0.03–0.49) after a consumer wash, indicating that even simple
rinsing may reduce loads by nearly 90 % (Figure 5). As the interval lengthened, exposures fell
markedly: at 20 days, 0.85 (SD 5.0×10^{-2}, UI 0.76–0.95) (unwashed) and 0.1 (SD 7.4×10^{-2}, UI
0.01 – 0.25) (washed); at 150 days, 7.6×10^{-3} (SD 4.5×10^{-4}, 6.7–8.6×10^{-3}) CFU and 8.6×10^{-4}

550   (SD 5.5 × 10⁻⁴, 1.1–2.0 × 10⁻⁴) CFU, respectively.

Figure 5. Predicted ESBL *E. coli* on lettuce at harvest (CFU/100 g) at different discrete intervals between
broiler litter application and planting day. Lines & points show mean exposure; error bars show the 95 %
uncertainty interval (2.5th–97.5th percentiles) across 1000 iterations.

Harvested lettuce doses were converted into infection probabilities and DALYs. Without
washing, mean gut-colonization risks ranged from 3.30×10^{-7} on planting interval 1,
corresponding to UTI risks of 9.02×10^{-9} and DALYs lost per serving of 7.48×10^{-8} (Table 4).
Consumer washing and increasing the interval between litter application and lettuce planting
reduced colonization risks, UTI risks, and DALYs. It appears that the rapid natural decay of *E.
coli* on foliage and the incremental benefit of simple post-harvest washing reduce the risk of UTI
and DALY burden from raw-leaf consumption. Our results are consistent with studies identifying
irrigation water as a plausible route for pre-harvest contamination of fresh produce. However, we
did not model or compare alternative routes and therefore do not rank their relative importance
(Solomon et al.; Jensen et al. 2013; Holvoet et al. 2013; Iwu and Okoh 2019). Several studies
have shown that *E. coli*, including pathogenic strains, can attach to and survive on lettuce surfaces
for extended periods (Islam et al. 2004; Ingham et al. 2004). Environmental factors such as higher
temperatures and light intensity tend to speed inactivation on plant surfaces (Weller et al. 2017;
Ottoson et al. 2011). In our model, increasing the interval between litter application and planting
reduces predicted contamination. This is consistent with studies showing time-dependent decline
of *E. coli* in manure-amended soils and reduced detection on crops with longer intervals after
application (Ingham et al. 2005; Black et al. 2021).

575 **Table 4**. Mean and UI (in brackets) probability of gastrointestinal colonization (GI) and urinary tract
576 infection (UTI), and corresponding DALY per exposure event for each planting interval and exposure
577 scenario (washed vs. unwashed lettuce).

Planting Interval	Risk GI	Risk UTI	DALY	Exposure
1	3.3×10^{-7} (2.9×10^{-7}– 3.6×10^{-7})	9.0×10^{-9} (7.5×10^{-9} – 1.0×10^{-8})	7.5×10^{-8} (3.4×10^{-8} – 1.2×10^{-7})	Unwashed
1	4.3×10^{-8} (6.0×10^{-9} – 9.6×10^{-8})	1.1×10^{-9} (1.6×10^{-10} – 2.6×10^{-9})	1.0×10^{-8} (1.2×10^{-9} – 3.0×10^{-8})	Washed
20	1.6×10^{-7} (1.5×10^{-7} – 1.8×10^{-7})	4.5×10^{-9} (3.6×10^{-9} – 5.2×10^{-9})	3.6×10^{-8} (1.7×10^{-8} – 5.8×10^{-8})	Unwashed
20	2.1×10^{-8} (2.6×10^{-9} – 5.0×10^{-8})	5.8×10^{-10} (7.1×10^{-11} – 1.3×10^{-9})	4.7×10^{-9} (4.9×10^{-10} – 1.3×10^{-8})	Washed
50	5.5×10^{-8} (4.9×10^{-8} – 6.1×10^{-8})	1.5×10^{-9} (1.2×10^{-9} – 1.8×10^{-9})	1.1×10^{-8} (5.6×10^{-9} – 1.8×10^{-8})	Unwashed
50	6.5×10^{-9} (8.2×10^{-10} – 1.5×10^{-8})	1.7×10^{-10} (2.2×10^{-11} – 4.3×10^{-10})	1.5×10^{-9} (1.6×10^{-10} – 3.9×10^{-9})	Washed
100	9.3×10^{-9} (8.3×10^{-9} – 1.0×10^{-8})	2.5×10^{-10} (2.1×10^{-10} – 3.0×10^{-10})	2.2×10^{-9} (9.0×10^{-10} – 3.4×10^{-9})	Unwashed
100	1.1×10^{-9} (1.5×10^{-10} – 2.8×10^{-9})	3.1×10^{-11} (3.6×10^{-12} – 7.8×10^{-11})	2.3×10^{-10} (2.2×10^{-11} – 5.4×10^{-10})	Washed
150	1.5×10^{-9} (1.3×10^{-9} – 1.6×10^{-9})	4.0×10^{-11} (3.4×10^{-11} – 4.9×10^{-11})	3.2×10^{-10} (1.5×10^{-10} – 5.1×10^{-10})	Unwashed
150	1.7×10^{-10} (2.2×10^{-11} – 3.9×10^{-10})	4.6×10^{-12} (5.6×10^{-13} – 1.0×10^{-11})	3.7×10^{-11} (4.2×10^{-12} – 1.0×10^{-10})	Washed

Regarding post-harvest mitigation at the consumer level, the model prediction of limited efficacy
on *E. coli* complete reduction for household washing is strongly supported by experimental data
(Uhlig et al. 2017; Jensen et al. 2015; Lopez et al. 2015). However, properly rinsing green leaves
may reduce the consumer's risk (Ottoson et al. 2011). The limitations of consumer-level washing
practices firmly place the primary responsibility for ensuring the microbial safety of fresh
produce, particularly regarding contaminants introduced via irrigation, on upstream controls
within the agricultural production system.
The sensitivity analysis (Figure 6) showed that lettuce-borne ESBL *E. coli* exposure is
overwhelmingly governed by how readily bacteria sorb to soil particles: the partition coefficient
has a very strong negative association (PRCC ≈ -0.90), meaning that greater sorption reduces
transfer to runoff and, ultimately, to foliage (Sowah et al., 2020). Soil decay kinetics also matter:
the estimated decay rate in soil and the environmental decay rate on the farm both correlate
negatively (PRCC ≈ -0.36 and -0.31, respectively), indicating that faster bacteria die-off sharply
cuts lettuce contamination. On the flip side, parameters that increase bacterial availability such as
wash-off fraction from soil to water (PRCC ≈ +0.17) and the shedding rate in the farm module
(PRCC ≈ +0.17) modestly raise exposure. Litter mass (kg/m^2) and gut carrying capacity also exert
smaller but significant influences. Together, these results suggest that strategies aiming to
enhance soil-borne decay and reduce bacterial release during runoff will be most effective at
reducing ESBL *E. coli* levels on harvested lettuce.

Figure 6. Sensitivity analysis for lettuce consumption (single event). Displayed is the mean Partial Rank
Correlation Coefficient including 95% confidence intervals

601
602

Limitations

Our model incorporates several simplifying assumptions that merit further refinement. In general,
our model represents a worst-case scenario, potentially overestimating human exposure. First, in
the farm module, we treat *E. coli* as uniformly distributed across all fecal matter and assume each
broiler remains colonized from the moment of infection until cycle end, with bacterial growth
governed by a single logistic curve. In reality, fecal deposition is spatially clustered, broiler
immune responses and stress levels vary over time, and diversity among ESBL *E. coli* strains can
differ, factors that can drive transient "hot spots" of shedding and alter transmission dynamics.

Second, the soil module idealizes litter spreading as perfectly homogeneous and represents die-
off with one constant exponential rate. This ignores the influence of temperature fluctuations, soil
moisture, pH gradients, bioturbation by earthworms and other fauna, and incorporation by tillage
all of which can create microsites where bacteria persist or die off more rapidly. Furthermore,
litter is assumed to be spread on fields immediately after flock removal (0 days storage) and
without any treatment (such as composting), ensuring maximal viable ESBL *E. coli* loads at

application. Nevertheless, although manure treatment and storage are common practices (Sarnino
et al., 2025), the Nitrates Directive (91/676/EEC), the main EU legislation regulating manure use
to protect water quality, does not mandate treatments or set a waiting period.

Third, our river transport component applies fixed wash-off fractions and partition coefficients,
yielding a smooth decline in contaminant load. Yet real watersheds undergo storm-driven pulses,
variable flow regimes, sediment resuspension, channel morphology effects, and spatial
heterogeneity in land cover, which together can produce episodic peaks in bacterial
concentrations that our model may miss. As a potential improvement, incorporation of a full
SWAT model could better capture these complex hydrological and land-use dynamics (Douglas-
Mankin et al, 2010).

Fourth, the lettuce module relies on a biphasic decay curve with uniform attachment efficiency to
leaf surfaces and a single removal efficiency for post-harvest washing. It does not account for
differences in leaf microstructure, irrigation method (e.g., overhead versus drip), spray droplet
size, canopy microclimate, or commercial-scale wash protocols.

Furthermore, our framework omits key microbial ecology processes, most notably horizontal
transfer of resistance genes (e.g. via plasmids) among environmental bacteria and interactions
with other microbial populations that may inhibit or promote ESBL *E. coli* survival. We also do
not capture potential regrowth phases under favorable conditions.
635
636 Another dominant uncertainty is the soil-water attachment/partitioning of ESBL *E. coli* under
637 local conditions-i.e., the proportion that is particle-associated versus freely suspended. Ideal data
638 would quantify, after rainfall or irrigation in litter-amended plots, bacteria in the water phase and
639 in the sediment phase, or use simple laboratory tests to estimate the particle-association fraction
640 for local soils.
641
Our dose-response model maps from an EPEC human challenge model (fecal shedding as a proxy
for GI colonization) to UTI via conditional transitions and a UPEC-like fraction. This surrogate is
necessary because no ESBL- or UPEC-specific human DR exists; consequently, risk outputs
should be interpreted as scenario-based indicators with uncertainty concentrated in the UPEC
fraction and conditional transitions rather than as clinical predictions. Heida et al. 2025
emphasized the same caveats for recreational waters; we adopt that caution here. Evidence to date
documents contamination of raw vegetables with ESBL/ExPEC *E. coli* but direct epidemiologic
links between raw-vegetable consumption and UTI are limited (Balbuena-Alonso et al. 2023;
Song et al. 2020) .The biological plausibility rests on GI colonization serving as a reservoir for
UPEC and periurethral transfer, with sex differences in UTI incidence not resolved in our per-
serving DALY (Josephs-Spaulding et al. 2021).

Finally, to our knowledge, this is the first ESBL *E. coli* focused QMRA connecting broiler flock,
litter-amended soil, receiving water and lettuce consumption in a single framework. For this
reason, a head-to-head validation against a prior end-to-end model is not yet possible. We
therefore adopted a modular validation strategy and, where feasible, compared predicted
magnitudes and expected signatures with published models and experimental studies.

Despite these simplifications, the model captures the dominant processes governing ESBL *E. coli*

dissemination to produce and, by intentionally adopting conservative assumptions, provides
exposure estimates that likely bound real-world outcomes. Consequently, we believe the results
remain sufficiently robust to inform risk management and guide intervention strategies.

Future model iterations should introduce spatially explicit litter application, temperature and
moisture driven decay rates, event-based hydrologic inputs, mechanistic leaf-surface colonization
dynamics, and microbial community interactions, including gene transfer dynamics.
Incorporating these elements will enhance realism, reduce uncertainty, and yield more robust
estimates of human exposure risk.

667
668
669

Conclusion
In this study, we developed and applied a four-module QMRA model to trace the journey of
ESBL *E. coli* from broiler flocks through soil, surface water, and fresh produce. The farm module
confirmed rapid amplification in litter, with near-complete flock colonization within one week
and end-cycle litter loads in the low 10^4 CFU/g range-values. Soil levels of ESBL *E. coli* peaked
at about 3.2×10^7 CFU/m² right after spreading, then fell steadily and were substantially lower
three months later. River concentrations showed a similar drop, starting near 8.3×10^{-2} CFU/mL
on day 1, falling to 6.0×10^{-2} CFU/mL by day 10, and reaching under 10^{-4} CFU/mL by day 200.
Lettuce simulations showed harvest loads of 1.7 CFU/100 g at one day interval between
application of litter and planting, dropping to 0.85 CFU/100 g at 20 days, with simple household
washing cutting exposure by ~90 percent. Overall, in the lettuce module, the mean gut-
colonization probabilities per 100 g serving did not exceed 10^{-7}, UTI risks 10^{-9}, and associated
DALYs per serving 10^{-8}, with household washing reducing risk levels. Sensitivity analyses
identified the soil-water partition coefficient and decay rates as the most influential parameters,
pointing to targeted data collection needs. Although natural decay, dilution, and interventions
such as composting can reduce human exposure by orders of magnitude, persistent contamination
calls for robust manure management and strict irrigation-water quality controls. Future work
should introduce spatial heterogeneity in litter spreading, dynamic weather drivers, host immunity
and horizontal gene transfer to refine risk estimates and guide effective on-farm and pre-harvest
interventions.

Funding

This work is a part of the European project ENVIRE funded by the JPIAMR program of the
European Union, and funded by the German Federal Ministry for Research and Education
(Support Code: 01KI2202A).

Acknowledgments

The authors would like to thank all the members of the ENVIRE project consortium for their
support.

Author Contributions
NS: Writing - original draft, Formal analysis, Conceptualization, Visualization, Writing - review

& editing. SB: Formal analysis, Conceptualization, Writing - review & editing. LC:
Conceptualization, Writing - review & editing, Supervision. RM: Conceptualization, Writing -
review & editing, Supervision, Funding acquisition.

Conflict of interest

704 The authors declare that the research was conducted in the absence of any commercial or
705 financial relationships that could be construed as a potential conflict of interest.

Supplementary data

707 The spreadsheet "Sarnino-et-al-2025b-results.xlsx" containing detailed simulation outputs is
708 available online alongside this article.

Publication bibliography

Agersø, Yvonne; Jensen, Jacob Dyring; Hasman, Henrik; Pedersen, Karl (2014): Spread of
Extended Spectrum Cephalosporinase-Producing Escherichia coli Clones and Plasmids from
Parent Animals to Broilers and to Broiler Meat in a Production Without Use of
Cephalosporins. In *Foodborne Pathogens and Disease* 11 (9), pp. 740–746. DOI:
10.1089/fpd.2014.1742.

Alegbeleye, Oluwadara Oluwaseun; Sant'Ana, Anderson S. (2020): Manure-borne pathogens
as an important source of water contamination: An update on the dynamics of pathogen
survival/transport as well as practical risk mitigation strategies. In *International journal of*
hygiene and environmental health 227, p. 113524. DOI: 10.1016/j.ijheh.2020.113524.

Apostolakos, Ilias; Mughini-Gras, Lapo; Fasolato, Luca; Piccirillo, Alessandra (2019):
Assessing the occurrence and transfer dynamics of ESBL/pAmpC-producing Escherichia coli
across the broiler production pyramid. In *PLOS ONE* 14 (5), e0217174. DOI:
10.1371/journal.pone.0217174.

Atanasova, Aleksandra; Amon, Thomas; Friese, Anika; Rösler, Uwe; Merle, Roswitha;
Herrmann, Christiane et al. (2025): Effects of Carbon–to–Nitrogen Ratio and Temperature
on the Survival of Antibiotic-Resistant and Non-Resistant Escherichia coli During Chicken
Manure Anaerobic Digestion. In *Poultry* 4, p. 9. DOI: 10.3390/poultry4010009.

Balbuena-Alonso, Maria G.; Camps, Manel; Cortés-Cortés, Gerardo; Carreón-León, Eder A.;
Lozano-Zarain, Patricia; Del Rocha-Gracia, Rosa Carmen (2023): Strain belonging to an
emerging, virulent sublineage of ST131 Escherichia coli isolated in fresh spinach, suggesting
that ST131 may be transmissible through agricultural products. In *Frontiers in Cellular and*
Infection Microbiology Volume 13 - 2023. Available online at
https://www.frontiersin.org/journals/cellular-and-infection-
microbiology/articles/10.3389/fcimb.2023.1237725.

Becker, Evelyne; Correia-Carreira, Guido; Projahn, Michaela; Käsbohrer, Annemarie (2022):
Modeling the Impact of Management Changes on the Infection Dynamics of Extended-
Spectrum Beta-Lactamase-Producing Escherichia coli in the Broiler Production. In *2076-*
2607 10 (5). DOI: 10.3390/microorganisms10050981.

Bengtsson-Palme, Johan; Abramova, Anna; Berendonk, Thomas U.; Coelho, Luis Pedro;
Forslund, Sofia K.; Gschwind, Rémi et al. (2023): Towards monitoring of antimicrobial
resistance in the environment: For what reasons, how to implement it, and what are the
data needs? In *0160-4120* 178, p. 108089. DOI: 10.1016/j.envint.2023.108089.

Blaak, Hetty; van Hoek, Angela H A M; Hamidjaja, Raditijo A.; van der Plaats, Rozemarijn Q J;
Kerkhof-de Heer, Lianne; de Roda Husman, Ana Maria; Schets, Franciska M. (2015):
Distribution, Numbers, and Diversity of ESBL-Producing E. coli in the Poultry Farm
Environment. In *PLOS ONE* 10 (8), e0135402. DOI: 10.1371/journal.pone.0135402.

Black, Zoe; Balta, Igori; Black, Lisa; Naughton, Patrick J.; Dooley, James S. G.; Corcionivoschi,
Nicolae (2021): The Fate of Foodborne Pathogens in Manure Treated Soil. In *1664-302X* 12.
Available online at
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2021.781357.

Carnell, R. (2022): triangle: Distribution Functions and Parameter Estimates for the
Triangle Distribution. In *R package version* 1.

Carnell, Rob (2016): Package 'lhs'. In *CRAN. https://cran. rproject.*
org/web/packages/lhs/lhs. pdf.

Ceccarelli Daniela; van Essen-Zandbergen Alieda; Smid Bregtje; Veldman Kees T.; Boender
Gert Jan; Fischer Egil A. J. et al. (2017): Competitive Exclusion Reduces Transmission and
Excretion of Extended-Spectrum-β-Lactamase-Producing Escherichia coli in Broilers. In
Applied and Environmental Microbiology 83 (11), e03439-16. DOI: 10.1128/AEM.03439-16.

Çekiç, Samantha K.; De, Jaysankar; Jubair, Mohammad; Schneider, Keith R. (2017):
Persistence of Indigenous Escherichia coli in Raw Bovine Manure-Amended Soil. In *0362-*
028X 80 (9), pp. 1562–1573. DOI: 10.4315/0362-028X.JFP-17-033.

Chantziaras, Ilias; Boyen, Filip; Callens, Bénédicte; Dewulf, Jeroen (2014): Correlation
between veterinary antimicrobial use and antimicrobial resistance in food-producing
animals: a report on seven countries. In *J Antimicrob Chemother* 69 (3), pp. 827–834. DOI:
10.1093/jac/dkt443.

Chapman, B.; Pintar, K.; Smith, B. A. (2018): Multi-Exposure Pathway Model to Compare
Escherichia coli O157 Risks and Interventions. In *0272-4332* 38 (2), pp. 392–409. DOI:
10.1111/risa.12826.

Collineau, Lucie; Chapman, Brennan; Bao, Xu; Sivapathasundaram, Branavan; Carson,
Carolee A.; Fazil, Aamir et al. (2020): A farm-to-fork quantitative risk assessment model for
Salmonella Heidelberg resistant to third-generation cephalosporins in broiler chickens in
Canada. In *International Journal of Food Microbiology* 330, p. 108559. DOI:
10.1016/j.ijfoodmicro.2020.108559.

Dame-Korevaar, Anita; Fischer, Egil A.J.; van der Goot, Jeanet; Velkers, Francisca; Ceccarelli,
Daniela; Mevius, Dik; Stegeman, Arjan (2020a): Early life supply of competitive exclusion
products reduces colonization of extended spectrum beta-lactamase-producing Escherichia
coli in broilers. In *0032-5791* 99 (8), pp. 4052–4064. DOI: 10.1016/j.psj.2020.04.025.

Dame-Korevaar, Anita; Kers, Jannigje G.; Goot, Jeanet; Velkers, Francisca; Ceccarelli, Daniela;
Mevius, Dik et al. (2020b): Competitive Exclusion Prevents Colonization and
Compartmentalization Reduces Transmission of ESBL-Producing Escherichia coli in
Broilers. In *Frontiers in Microbiology* 11. DOI: 10.3389/fmicb.2020.566619.

Dankittipong, Natcha; Alderliesten, Jesse B.; van den Broek, Jan; Dame-Korevaar, M. Anita;
Brouwer, Michael S.M.; Velkers, Francisca C. et al. (2023): Comparing the transmission of
carbapenemase-producing and extended-spectrum beta-lactamase-producing Escherichia
coli between broiler chickens. In *Preventive Veterinary Medicine* 219, p. 105998. DOI:
10.1016/j.prevetmed.2023.105998.

Dierikx, Cindy M.; van der Goot, Jeanet A.; Smith, Hilde E.; Kant, Arie; Mevius, Dik J. (2013):
Presence of ESBL/AmpC -Producing Escherichia coli in the Broiler Production Pyramid: A
Descriptive Study. In *PLOS ONE* 8 (11), e79005. DOI: 10.1371/journal.pone.0079005.

Douglas-Mankin, K. R.; Srinivasan, Raghavan; Arnold, J. G. (2010): Soil and Water
Assessment Tool (SWAT) model: Current developments and applications. In *Transactions of*
the ASABE 53 (5), pp. 1423–1431.

Fastl, Christina; Carvalho Ferreira, Helena C. de; Babo Martins, Sara; Sucena Afonso, João; Di
Bari, Carlotta; Venkateswaran, Narmada et al. (2023): Animal sources of antimicrobial-
resistant bacterial infections in humans: a systematic review. In *0950-2688* 151, e143. DOI:
10.1017/S0950268823001309.

Furusawa, Minori; Widgren, Stefan; Evers, Eric G.; Fischer, Egil A.J. (2024): Quantifying
health risks from ESBL-producing Escherichia coli in Dutch broiler production chains and
potential interventions using compartmental models. In *Preventive Veterinary Medicine* 224,
p. 106121. DOI: 10.1016/j.prevetmed.2024.106121.

Galarneau, Karen D.; Singer, Randall S.; Wills, Robert W. (2020): A system dynamics model
for disease management in poultry production. In *0032-5791* 99 (11), pp. 5547–5559. DOI:
10.1016/j.psj.2020.08.011.

Gekenidis, Maria-Theresia; Rigotti, Serena; Hummerjohann, Jörg; Walsh, Fiona; Drissner,
David (2020): Long-Term Persistence of blaCTX-M-15 in Soil and Lettuce after Introducing
Extended-Spectrum β-Lactamase (ESBL)-Producing Escherichia coli via Manure or Water.
In *2076-2607* 8 (11). DOI: 10.3390/microorganisms8111646.

Grothendieck, Maintainer G. (2013): Package 'nls2'. In *Non-linear regression with brute force*.

Heida, Ashley; Hamilton, Mark T.; Gambino, Julia; Sanderson, Kaylee; Schoen, Mary E.; Jahne,
Michael A. et al. (2025): Population Ecology-Quantitative Microbial Risk Assessment
(QMRA) Model for Antibiotic-Resistant and Susceptible E. coli in Recreational Water. In
0013-936X 59 (9), pp. 4266–4281. DOI: 10.1021/acs.est.4c07248.

Holvoet, Kevin; Sampers, Imca; Callens, Bénédicte; Dewulf, Jeroen; Uyttendaele, Mieke
(2013): Moderate Prevalence of Antimicrobial Resistance in Escherichia coli Isolates from
Lettuce, Irrigation Water, and Soil 79. DOI: 10.1128/AEM.01995-13.

Howard, Keya J.; Martin, Emily; Gentry, Terry; Feagley, Sam; Karthikeyan, Raghupathy
(2016): Effects of Dairy Manure Management Practices on E. coli Concentration and
Diversity. In *Water, Air, & Soil Pollution* 228 (1), p. 4. DOI: 10.1007/s11270-016-3182-7.

Hubbard, L. E.; Givens, C. E.; Griffin, D. W.; Iwanowicz, L. R.; Meyer, M. T.; Kolpin, D. W.
(2020): Poultry litter as potential source of pathogens and other contaminants in
groundwater and surface water proximal to large-scale confined poultry feeding operations.
In *0048-9697* 735, p. 139459. DOI: 10.1016/j.scitotenv.2020.139459.

Huijbers, P. M. C.; Graat, E. A. M.; van Hoek, Aham; Veenman, C.; Jong, M. C. M. de; van
Duijkeren, E. (2016): Transmission dynamics of extended-spectrum β-lactamase and AmpC
β-lactamase-producing Escherichia coli in a broiler flock without antibiotic use. In
Preventive Veterinary Medicine 131, pp. 12–19. DOI: 10.1016/j.prevetmed.2016.07.001.

Ingham, Steven; Losinski, Jill; Andrews, Matthew; Breuer, Jane; Breuer, Jeffry; Wood,
Timothy; Wright, Thomas (2004): Escherichia coli Contamination of Vegetables Grown in
Soils Fertilized with Noncomposted Bovine Manure: Garden-Scale Studies 70, pp. 6420–
6427. DOI: 10.1128/AEM.70.11.6420-6427.2004.

Ingham, Steven C.; Fanslau, Melody A.; Engel, Rebecca A.; Breuer, Jeffry R.; Breuer, Jane E.;
Wright, Thomas H. et al. (2005): Evaluation of Fertilization-to-Planting and Fertilization-to-
Harvest Intervals for Safe Use of Noncomposted Bovine Manure in Wisconsin Vegetable
Production. In *0362-028X* 68 (6), pp. 1134–1142. DOI: 10.4315/0362-028X-68.6.1134.

Iooss, Bertrand; Da Veiga, Sebastien; Janon, Alexandre; Pujol, Gilles; Iooss, Maintainer
Bertrand; Rcpp, LinkingTo et al. (2024): Package 'sensitivity'. In *R Package from*
https://cran. r-project. or/package= sensitivity.

Islam, Mahbub; Doyle, Michael; Phatak, Sharad; Millner, Pat; Jiang, Xiuping (2004):
Persistence of Enterohemorrhagic Escherichia coli O157:H7 in Soil and on Leaf Lettuce and
Parsley Grown in Fields Treated with Contaminated Manure Composts or Irrigation Water
67, pp. 1365–1370. DOI: 10.4315/0362-028X-67.7.1365.

Iwu, Chidozie D.; Okoh, Anthony I. (2019): Preharvest Transmission Routes of Fresh
Produce Associated Bacterial Pathogens with Outbreak Potentials: A Review. In *1660-4601*
16 (22). DOI: 10.3390/ijerph16224407.

Jacobs, Kyle; Wind, Lauren; Krometis, Leigh-Anne; Hession, W. Cully; Pruden, Amy (2019):
Fecal Indicator Bacteria and Antibiotic Resistance Genes in Storm Runoff from Dairy
Manure and Compost-Amended Vegetable Plots. In *0047-2425* 48 (4), pp. 1038–1046. DOI:
10.2134/jeq2018.12.0441.

Jensen, Annette; Storm, C.; Forslund, Anita; Baggesen, Dorte; Dalsgaard, Anders (2013):
Escherichia coli Contamination of Lettuce Grown in Soils Amended with Animal Slurry 76,
pp. 1137–1144. DOI: 10.4315/0362-028X.JFP-13-011.

Jensen, Dane A.; Friedrich, Loretta M.; Harris, Linda J.; Danyluk, Michelle D.; Schaffner,
Donald W. (2015): Cross contamination of Escherichia coli O157:H7 between lettuce and

853 wash water during home-scale washing. In *Food Microbiology* 46, pp. 428–433. DOI:
854 10.1016/j.fm.2014.08.025.

855 Josephs-Spaulding, Jonathan; Krogh, Thøger Jensen; Rettig, Hannah Clara; Lyng, Mark;
856 Chkonia, Mariam; Waschina, Silvio et al. (2021): Recurrent Urinary Tract Infections:
857 Unraveling the Complicated Environment of Uncomplicated rUTIs. In *Frontiers in Cellular*
858 *and Infection Microbiology* Volume 11 - 2021. Available online at
859 https://www.frontiersin.org/journals/cellular-and-infection-
860 microbiology/articles/10.3389/fcimb.2021.562525.

861 Kasparaviciene, Beatrice; Novoslavskij, Aleksandr; Aksomaitiene, Jurgita; Stankeviciene,
862 Jurate; Kasetiene, Neringa; Sinkevicius, Romualdas; Malakauskas, Mindaugas (2025):
863 Prevalence and Antimicrobial Resistance of ESBL E. coli in Early Broiler Production Stage
864 and Farm Environment in Lithuania. In *2076-2607* 13 (2). DOI:
865 10.3390/microorganisms13020425.

866 Leistner, R.; Gürntke, S.; Sakellariou, C.; Denkel, L. A.; Bloch, A.; Gastmeier, P.; Schwab, F.
867 (2014): Bloodstream infection due to extended-spectrum beta-lactamase (ESBL)-positive K.
868 pneumoniae and E. coli: an analysis of the disease burden in a large cohort. In *Infection* 42
869 (6), pp. 991–997. DOI: 10.1007/s15010-014-0670-9.

870 Literák, Ivan; Reitschmied, Tomas; Bujnáková, Dobroslava; Dolejska, Monika; Cizek, Alois;
871 Bardon, Jan et al. (2012): Broilers as a Source of Quinolone-Resistant and Extraintestinal
872 Pathogenic Escherichia coli in the Czech Republic. In *Microbial drug resistance (Larchmont,*
873 *N.Y.)* 19. DOI: 10.1089/mdr.2012.0124.

874 Lopez, K.; Getty, K.J.K.; Vahl, Christopher (2015): Validation of washing treatments to reduce
875 Escherichia coli 0157:H7 and Salmonella spp. on the surface of green leaf lettuce and
876 tomatoes 35, pp. 377–384.

877 Luyckx, Kaat; van Weyenberg, Stephanie; Dewulf, Jeroen; Herman, Lola; Zoons, J.; Vervaet, E.
878 et al. (2015): On-farm comparisons of different cleaning protocols in broiler houses. In
879 *Poultry science* 94. DOI: 10.3382/ps/pev143.

880 Lyhs, Ulrike; Ikonen, Ilona; Pohjanvirta, Tarja; Raninen, Kaisa; Perko-Mäkelä, Päivikki;
881 Pelkonen, Sinikka (2012): Extraintestinal pathogenic Escherichia coli in poultry meat
882 products on the Finnish retail market. In *Acta Veterinaria Scandinavica* 54 (1), p. 64. DOI:
883 10.1186/1751-0147-54-64.

884 MacKinnon, M. C.; Sargeant, J. M.; Pearl, D. L.; Reid-Smith, R. J.; Carson, C. A.; Parmley, E. J.;
885 McEwen, S. A. (2020): Evaluation of the health and healthcare system burden due to
886 antimicrobial-resistant Escherichia coli infections in humans: a systematic review and
887 meta-analysis. In *2047-2994* 9 (1), p. 200. DOI: 10.1186/s13756-020-00863-x.

888 Mallioris, Panagiotis; Teunis, Gijs; Lagerweij, Giske; Joosten, Philip; Dewulf, Jeroen;
889 Wagenaar, Jaap A. et al. (2023): Biosecurity and antimicrobial use in broiler farms across
890 nine European countries: toward identifying farm-specific options for reducing
891 antimicrobial usage. In *0950-2688* 151, e13. DOI: 10.1017/S0950268822001960.

Mancini, John L. (1978): Numerical estimates of coliform mortality rates under various
conditions. In *Journal (Water Pollution Control Federation)*, pp. 2477–2484.

Mellata, Melha (2013): Human and Avian Extraintestinal Pathogenic Escherichia coli:
Infections, Zoonotic Risks, and Antibiotic Resistance Trends. In *Foodborne Pathogens and*
Disease 10. DOI: 10.1089/fpd.2013.1533.

Mootian, Gabriel; Wu, Wen-Hsuan; Matthews, Karl R. (2009): Transfer of Escherichia coli
O157:H7 from soil, water, and manure contaminated with low numbers of the pathogen to
lettuce plants 72 11, pp. 2308–2312. Available online at
https://api.semanticscholar.org/CorpusID:21229382.

Moreno, Miguel A.; García-Soto, Silvia; Hernández, Marta; Bárcena, Carmen; Rodríguez-
Lázaro, David; Ugarte-Ruíz, María; Domínguez, Lucas (2019): Day-old chicks are a source of
antimicrobial resistant bacteria for laying hen farms. In *Veterinary Microbiology* 230,
pp. 221–227. DOI: 10.1016/j.vetmic.2019.02.007.

Nilsson, Oskar; Börjesson, Stefan; Landén, Annica; Bengtsson, Björn (2014): Vertical
transmission of Escherichia coli carrying plasmid-mediated AmpC (pAmpC) through the
broiler production pyramid. In *J Antimicrob Chemother* 69 (6), pp. 1497–1500. DOI:
10.1093/jac/dku030.

O'Flaherty, E.; Borrego, C. M.; Balcázar, J. L.; Cummins, E. (2018): Human exposure
assessment to antibiotic-resistant Escherichia coli through drinking water. In *0048-9697*
616-617, pp. 1356–1364. DOI: 10.1016/j.scitotenv.2017.10.180.

O'Flaherty, E.; Solimini, A.; Pantanella, F.; Cummins, E. (2019a): The potential human
exposure to antibiotic resistant-Escherichia coli through recreational water. In *0048-9697*
650 (Pt 1), pp. 786–795. DOI: 10.1016/j.scitotenv.2018.09.018.

O'Flaherty, E.; Solimini, A. G.; Pantanella, F.; Giusti, M. de; Cummins, E. (2019b): Human
exposure to antibiotic resistant-Escherichia coli through irrigated lettuce. In *0160-4120* 122,
pp. 270–280. DOI: 10.1016/j.envint.2018.11.022.

Oikarainen, Paula E.; Pohjola, Leena K.; Pietola, Eeva S.; Heikinheimo, Annamari (2019):
Direct vertical transmission of ESBL/pAmpC-producing Escherichia coli limited in poultry
production pyramid. In *Veterinary Microbiology* 231, pp. 100–106. DOI:
10.1016/j.vetmic.2019.03.001.

Okada, Elena; Young, Brian Jonathan; Pérez, Débora Jesabel; Pellegrini, María Celeste;
Carciochi, Walter Daniel; Lavallén, Carla Mariela et al. (2024): Effect of on-farm poultry
litter composting processes on physicochemical, biological, and toxicological parameters
and reduction of antibiotics and antibiotic-resistant Escherichia coli. In *Waste Management*
174, pp. 310–319. DOI: 10.1016/j.wasman.2023.12.005.

Ottoson, Jakob; Nyberg, Karin; Lindqvist, Roland; Albihn, Ann (2011): Quantitative
Microbial Risk Assessment for Escherichia coli O157 on Lettuce, Based on Survival Data
from Controlled Studies in a Climate Chamber. In *Journal of food protection* 74, pp. 2000–
2007. DOI: 10.4315/0362-028X.JFP-10-563.

Pouillot, R.; Delignette-Muller, M. L.; Kelly, D. L.; Denis, J. B. (2016): The mc2d package. In *R*
package 30.

Rawson, Thomas; Dawkins, Marian Stamp; Bonsall, Michael B. (2019): A Mathematical
Model of Campylobacter Dynamics Within a Broiler Flock. In *1664-302X* Volume 10 - 2019.
Available online at
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2019.01940.

Robé, Caroline; Blasse, Anja; Merle, Roswitha; Friese, Anika; Roesler, Uwe; Guenther,
Sebastian (2019): Low Dose Colonization of Broiler Chickens With ESBL-/AmpC- Producing
Escherichia coli in a Seeder-Bird Model Independent of Antimicrobial Selection Pressure. In
1664-302X Volume 10 - 2019. Available online at
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2019.02124.

Robé, Caroline; Daehre, Katrin; Merle, Roswitha; Friese, Anika; Guenther, Sebastian; Roesler,
Uwe (2021): Impact of different management measures on the colonization of broiler
chickens with ESBL- and pAmpC- producing Escherichia coli in an experimental seeder-bird
model. In *PLOS ONE* 16 (1), e0245224. DOI: 10.1371/journal.pone.0245224.

Robé, Caroline; Projahn, Michaela; Boll, Katrin; Blasse, Anja; Merle, Roswitha; Roesler, Uwe;
Friese, Anika (2024): Survival of highly related ESBL- and pAmpC- producing Escherichia
coli in broiler farms identified before and after cleaning and disinfection using cgMLST. In
BMC Microbiology 24 (1), p. 143. DOI: 10.1186/s12866-024-03292-7.

Sarnino, Nunzio; Basak, Subhasish; Collineau, Lucie; Merle, Roswitha (2025): Pathways of
Escherichia coli transfer from animal manure: risks and mitigation in agriculture. In
Frontiers in Public Health Volume 13 - 2025. Available online at
https://www.frontiersin.org/journals/public-
health/articles/10.3389/fpubh.2025.1568621.

Sharma Manan; Millner Patricia D.; Hashem Fawzy; Vinyard Bryan T.; East Cheryl L.; Handy
Eric T. et al. (2019): Survival of Escherichia coli in Manure-Amended Soils Is Affected by
Spatiotemporal, Agricultural, and Weather Factors in the Mid-Atlantic United States 85 (5),
e02392‑18. DOI: 10.1128/AEM.02392-18.

Siller, Paul; Daehre, Katrin; Thiel, Nadine; Nübel, Ulrich; Roesler, Uwe (2020): Impact of
short-term storage on the quantity of extended-spectrum beta-lactamase–producing
Escherichia coli in broiler litter under practical conditions. In *0032-5791* 99 (4), pp. 2125–
2135. DOI: 10.1016/j.psj.2019.11.043.

Solomon, Ethan; Yaron, Sima; Matthews, Karl: Transmission of Escherichia coli O157:H7
from Contaminated Manure and Irrigation Water to Lettuce Plant Tissue and Its Subsequent
Internalization 68, pp. 397–400. DOI: 10.1128/AEM.68.1.397-400.2002.

Song, Jihyun; Oh, Sung-Suck; Kim, Junghee; Shin, Jinwook (2020): Extended-spectrum β-
lactamase-producing Escherichia coli isolated from raw vegetables in South Korea. In
Scientific Reports 10 (1), p. 19721. DOI: 10.1038/s41598-020-76890-w.

Sowah, Robert; Bradshaw, Kenneth; Snyder, Blake; Spidle, David; Molina, Marirosa (2020):
Evaluation of the soil and water assessment tool (SWAT) for simulating E. coli
concentrations at the watershed-scale 746, p. 140669. DOI:
10.1016/j.scitotenv.2020.140669.

Stromberg, Zachary R.; Johnson, James R.; Fairbrother, John M.; Kilbourne, Jacquelyn; van
Goor, Angelica; Curtiss, Roy, 3rd; Mellata, Melha (2017): Evaluation of Escherichia coli
isolates from healthy chickens to determine their potential risk to poultry and human
health. In *PLOS ONE* 12 (7), e0180599. DOI: 10.1371/journal.pone.0180599.

Subirats, Jessica; Murray, Roger; Yin, Xiaole; Zhang, Tong; Topp, Edward (2021): Impact of
chicken litter pre-application treatment on the abundance, field persistence, and transfer of
antibiotic resistant bacteria and antibiotic resistance genes to vegetables. In *0048-9697* 801,
p. 149718. DOI: 10.1016/j.scitotenv.2021.149718.

Suzuki, Yoshihiro; Horita, Tomoyuki; Nishimura, Emi; Xie, Hui; Tamai, Soichiro; Kobayashi,
Ikuo et al. (2024): Crop contamination evaluation by antimicrobial-resistant bacteria via
livestock waste compost-fertilized field soil. In *0304-3894* 480, p. 135987. DOI:
10.1016/j.jhazmat.2024.135987.

Thomas, C.; Idler, C.; Ammon, C.; Amon, T. (2024): Applied Research Note: Survival of
Escherichia coli and temperature development during composting of chicken manure with a
typically low carbon/nitrogen ratio and moisture content. In *Journal of Applied Poultry*
Research 33 (2), p. 100402. DOI: 10.1016/j.japr.2024.100402.

Tofani, Silvia; Albini, Elisa; Blasi, Francesca; Cucco, Lucilla; Lovito, Carmela; Maresca,
Carmen et al. (2022): Assessing the Load, Virulence and Antibiotic-Resistant Traits of
ESBL/Ampc E. coli from Broilers Raised on Conventional, Antibiotic-Free, and Organic
Farms. In *Antibiotics* 11 (11). DOI: 10.3390/antibiotics11111484.

Tsoularis, A.; Wallace, J. (2002): Analysis of logistic growth models. In *Mathematical*
Biosciences 179 (1), pp. 21–55. DOI: 10.1016/S0025-5564(02)00096-2.

Uhlig, Elisabeth; Olsson, Crister; He, Jiayi; Stark, Therese; Sadowska, Zuzanna; Molin, Goran
et al. (2017): Effects of household washing on bacterial load and removal of Escherichia coli
from lettuce and "ready‐to‐eat" salads. In *Food Science & Nutrition* 5. DOI:
10.1002/fsn3.514.

van Overbeek, Leo; Duhamel, Marie; Aanstoot, Stefan; van der Plas, Carin Lombaers; Nijhuis,
Els; Poleij, Leo et al. (2021): Transmission of Escherichia coli from Manure to Root Zones of
Field-Grown Lettuce and Leek Plants. In *2076-2607* 9 (11). DOI:
10.3390/microorganisms9112289.

Weller, Daniel L.; Kovac, Jasna; Roof, Sherry; Kent, David J.; Tokman, Jeffrey I.; Kowalcyk,
Barbara et al. (2017): Survival of Escherichia coli on Lettuce under Field Conditions
Encountered in the Northeastern United States. In *0362-028X* 80 (7), pp. 1214–1221. DOI:
10.4315/0362-028X.JFP-16-419.

Wickham, Hadley; Averick, Mara; Bryan, Jennifer; Chang, Winston; McGowan, Lucy
D'Agostino; François, Romain et al. (2019): Welcome to the Tidyverse. In *Journal of open*
source software 4 (43), p. 1686.

Zhang, Yangjunna; Schmidt, John W.; Arthur, Terrance M.; Wheeler, Tommy L.; Zhang, Qi;
Wang, Bing (2022): A Farm-to-Fork Quantitative Microbial Exposure Assessment of β-
Lactam-Resistant Escherichia coli among U.S. Beef Consumers. In *2076-2607* 10 (3). DOI:
10.3390/microorganisms10030661.

1014