Phenotypic Selectivity of Artificial Intelligence-enhanced Electrocardiography

in Cardiovascular Diagnosis and Risk Prediction

Philip M. Croon, MD[1,2,3], Lovedeep S. Dhingra, MBBS, MHS[1,2], Dhruva Biswas, MD,

PhD[1,2], Evangelos K Oikonomou, MD, DPhil[1,2], Rohan Khera, MD, MS[1,2, 4, 5, 6]

[1]Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of

Medicine, New Haven, CT

[2]Cardiovascular Data Science (CarDS) Lab, Yale School of Medicine, New Haven,

CT

[3]Department of Cardiology, Amsterdam Cardiovascular Sciences, Amsterdam

University Medical Centre, University of Amsterdam, Amsterdam, The Netherlands

[4]Section of Health Informatics, Department of Biostatistics, Yale School of Public

Health, New Haven, CT

[5]Section of Biomedical Informatics and Data Science, Yale School of Medicine, New

Haven, CT

[6]Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New

Haven, CT

Correspondence:

Rohan Khera, MD, MS

195 Church St, 6th Floor, New Haven, CT 06510

Word Count: 4792

Figures/Tables: 5

Abstract (348/350)

Introduction: Artificial intelligence (AI)-enhanced electrocardiogram (ECG) models

are designed to detect specific anatomical and functional cardiac abnormalities.

Understanding the selectivity of their phenotypic associations is essential to inform

their clinical use. Here, we sought to assess whether AI-ECG models function as

condition-specific classifiers or broader cardiovascular risk markers.

Methods: We included four distinct study populations, drawn from both electronic

health records (EHR) and prospective cohort studies. We deployed six image-based

AI-ECG models, including five validated models for the detection of left ventricular

systolic dysfunction (LVSD), aortic stenosis (AS), mitral regurgitation (MR), left

ventricular hypertrophy (LVH), a composite model for structural heart disease (SHD),

and a negative control AI-ECG model for biological sex. Additionally, we developed

six experimental models designed to identify non-cardiovascular conditions.

Diagnosis codes from EHR and cohorts were transformed into interpretable

phenotypes using a phenome-wide association study (PheWAS) framework. We

assessed associations of AI-ECG probabilities with cross-sectional phenotypes using

logistic regression, and with new-onset cardiovascular diseases using Cox

regression. Pearson correlation coefficients were calculated to compare phenotypic

signatures.

Results: The study included one random ECG from 233,689 individuals (mean age

59±18 years, 130,084 [56%] women) across sites. Each of the five AI-ECG models

was more likely to be associated with cardiovascular phenotypes compared with

other phenotype groups (odds ratios ranging from 2.16 to 4.41, $p<10^{-\square}$), while the

sex model did not show a similar pattern. All AI-ECG models were significantly

associated with their respective target phenotype, but also showed similar or stronger

associations with a broad range of other cardiovascular phenotypes. Phenotypic

associations were similar across AI-ECG models trained for different conditions,

which was not observed in models for non-cardiovascular conditions. Correlation of

phenotype association patterns between models was high (r = 0.65–0.99). This

pattern was consistent across all models, external datasets, and in both cross-

sectional and prospective analyses.

Conclusions: Despite being developed to detect specific cardiovascular conditions,

AI-ECG models detect the presence and predict the future development of a broad

range of cardiovascular diseases with similar propensity. This challenges their role as

binary diagnostic tools and instead supports their use as broader cardiovascular

biomarkers.

Non-standard Abbreviations and Acronyms

AI – Artificial Intelligence

AI-ECG – Artificial Intelligence-enhanced Electrocardiogram

AS – Aortic Stenosis

AUROC – Area Under the Receiver Operating Characteristic Curve

CNN – Convolutional Neural Network

CVD – Cardiovascular Disease

EHR – Electronic Health Record

ECG – Electrocardiogram

FDA – Food and Drug Administration

ICD – International Classification of Diseases

LVEF – Left Ventricular Ejection Fraction

LVH – Left Ventricular Hypertrophy

LVSD – Left Ventricular Systolic Dysfunction

MR – Mitral Regurgitation

HCM – Hypertrophic cardiomyopathy

HOCM – Hypertrophic obstructive cardiomyopathy

NHS – National Health Service

PheWAS – Phenome-Wide Association Study

SHD – Structural Heart Disease

UKB – UK Biobank

YNHH – Yale New Haven Hospital

YNHHS – Yale New Haven Health System

INTRODUCTION

Artificial intelligence (AI) has expanded the utility of the electrocardiogram (ECG), enabling the detection of subtle disease signatures previously considered unrecognizable even by experts.[1,2] Several AI-ECG models have now been extensively validated for the detection of various specific functional and structural cardiovascular conditions, including left ventricular systolic dysfunction (LVSD), valvular heart disease (VHD), and left ventricular hypertrophy (LVH).[3–7] Moreover, AI-ECG has demonstrated effectiveness as a digital biomarker capable of predicting new-onset cardiovascular diseases (CVD).[8–11] These developments have led to the FDA approval for AI-ECG algorithms designed to detect LVSD and hypertrophic cardiomyopathy, marking its transition from research into clinical application.[12]

While AI-ECG models have been developed and evaluated as condition-specific tools, an assessment of their selectivity in phenotypic associations is required to guide their clinical use. Many CVDs exhibit shared pathophysiological mechanisms and risk factors, which may result in similar conduction and rhythm alterations.[10,13] Visible ECG abnormalities, such as ST-segment changes or prolonged QT intervals, are also frequently associated with multiple cardiac diseases, reflecting overlapping pathological processes.[14–16] As a result, AI-ECG models may not differentiate between distinct disease entities, as suggested by a series of models designed for individual conditions, but instead identify ECG signatures representing broader cardiovascular pathology. With the ongoing transition of AI-ECG into clinical practice, these potential limitations must be systematically evaluated to clarify their clinical implications and ensure safe, effective deployment.[17]

In this study, we evaluated whether image-based AI-ECG models trained on specific cardiovascular disease labels exhibit selective phenotypic association patterns for cross-sectional and new-onset cardiovascular disorders across four diverse cohorts, including a large tertiary hospital, a group of four community hospitals, an outpatient medical network, and the prospective UK Biobank. We also

included a classification model for biological sex and seven experimental models as

negative control outcomes for cardiac phenotypic patterns potentially detectable on

AI-ECG**.**

METHODS

The Yale Institutional Review Board approved the study protocol and waived the

requirement for informed consent, as the study involves secondary analysis of pre-

existing data. The data underlying this study cannot be shared publicly due to patient

privacy concerns and institutional regulations.

Evaluation of AI-ECG Models for Cardiovascular Disease

We leveraged a range of AI-ECG models that were previously developed and

extensively validated in multinational cohorts to detect distinct cardiovascular

abnormalities. These cardiovascular disease models were trained and validated for

specific labels including: (1) LVSD, defined as a left ventricular ejection fraction

(LVEF) < 40% on echocardiography, left-sided valvular heart diseases, namely (2) AS

and (3) MR, graded as mild to severe based on echocardiographic guidelines, (4)

severe LVH, defined by an interventricular septal diameter at end-diastole > 15 mm,

combined with moderate to severe (grade II/III) left ventricular diastolic dysfunction,

and (5) composite SHD including LVSD, valvular heart disease and LVH.[18]

The development and external validation of these models, including cohort

characteristics, has been described in prior studies.[3,19] All models were developed in

the same group of individuals who had undergone an electrocardiogram and

echocardiogram concurrently (within 30 days of each other) in the Yale New Haven

Hospital (YNHH).

The individual disease label models are all convolutional neural networks

(CNN) based on the EfficientNet-B3 architecture, which consists of 384 layers and

over 10 million trainable parameters. The model input consists of 300×300 pixel

image representations of standard 12-lead ECGs. To enhance label efficiency in

model development, CNNs were initialized with weights from a self-supervised,

contrastive learning framework designed to recognize patient-specific ECG patterns

independent of clinical interpretation.[20] Fine-tuning was performed in two stages. In

the initial stage, only the final two layers of the network were unfrozen and optimized,

followed by a second stage in which all layers were unfrozen to allow full model

adaptation to the task-specific objective. This staged training approach was used to

improve optimization stability and efficiency, while preserving the model's capacity to

update all weights based on the disease-specific signal. The AI-ECG model for

detecting composite structural heart disease, called PRESENT-SHD, employs an

XGBoost classifier that integrates predictions from individual CNNs trained to detect

LVSD, valvular heart disease, and LVH, along with an individual's age and sex as

input features.[19,21] All models were externally validated across diverse clinical and

community-based settings and demonstrated state-of-the-art performance.

Development and Evaluation of Experimental AI-ECG Models

To evaluate whether the signature of an AI-ECG model trained on a non-

cardiovascular, biological label differs from that of one trained to detect

cardiovascular diseases, we developed an AI-ECG model to define biological sex

from the ECG, using the same methodology and cohort as the disease-specific

models described in the previous paragraph. Such models have previously been

proposed, and this model demonstrated excellent performance (area under the

receiver operating curve (AUROC)>0.90) across all cohorts, including internal

validation in the YNHH cohort and external validation in community hospitals,

outpatient clinics, and the UK Biobank **(Supplementary Figure 1, Supplementary**

Table 1).[22]

To examine whether the signatures of the AI-ECG models are dependent on

the training label rather than consistently suggesting cardiovascular pathology, we

developed multiple experimental models to serve as negative controls for the

cardiovascular association of AI-ECG models. We used the same methodology and

development population as the validated AI-ECG models for cardiovascular disease

detection. First, we developed a model to classify whether an ECG was recorded

during an odd-numbered month. Next, we derived 6 experimental models designed to

detect the occurrence of non-cardiovascular disease labels, including viral respiratory

infection, transport accident, headache, fracture of the lower leg, being bitten by a

dog, and dermatophytosis. These conditions were defined based on the presence of

diagnosis codes up to one year before or after the ECG **(Supplementary Table 2)**.

None of these experimental models outperformed random chance in the

development cohort (AUROC ~0.50), confirming their role as negative control

outcomes for AI-ECG **(Supplementary Table 3, Supplementary Figure 2)**.

Data Sources

We included individuals from multiple cohorts. The primary dataset was sourced from

YNHH, a large tertiary medical center in Connecticut, USA. For external validation,

we included: (i) four community-based hospitals in the Yale New Haven Health

System (YNHHS): Bridgeport, Greenwich, Lawrence + Memorial and Westerly

Hospitals, (ii) Northeast Medical Group spanning multiple outpatient clinics, and (iii)

participants from the UK Biobank cardiovascular imaging substudy with an ECG

recorded at enrollment. Extensive health and biometric data were available through

linkage with National Health Service (NHS) electronic health records. ECGs included

in the cohorts from the YNHH, community hospitals, and outpatient clinics were

recorded as part of routine clinical care from 2013 to 2024, while the ECGs from UK

Biobank participants were collected between 2014 and 2020. For all cohorts,

diagnostic codes, classified as International Classification of Diseases (ICD)-10

codes, along with the first recorded dates, were extracted.

Study population

In the cohort derived from the YNHH, individuals who were part of the model

development were excluded from our analysis. For the cohorts that originated from

the YNHH, community hospitals, and outpatient clinics, we selected individuals with

at least one ECG as part of routine clinical care. To ensure adequate follow-up for

disease ascertainment, we applied a one-year blanking period starting from each

patient's first hospital encounter. We then randomly selected one ECG recorded after

this period per individual, allowing for consistent follow-up and representation across

the spectrum of disease stages. For the UK Biobank, we included all participants with

an available ECG, using the earliest recorded ECG per individual. To define the

baseline characteristics of the cohorts, we used patient demographics and diagnostic

codes **(Supplementary Table 4).**

Exploring Model Associations with Clinical Phenotypes

We conducted a phenome-wide association study (PheWAS) to explore cross-

sectional associations between AI-ECG model predictions and a wide range of

clinical phenotypes derived from diagnostic codes. PheWAS is an approach used to

systematically evaluate associations between a specific genetic or clinical trait and a

wide range of phenotypes.[23] It has previously been used to identify novel disease

associations, uncover comorbid conditions, and explore the phenotypic spectrum

associated with biomarkers or genetic variants.[24–26]

For cross-sectional analysis, only diagnostic codes recorded before or on the

day the ECG was recorded were considered, while those recorded afterwards were

excluded. Additionally, to maintain statistical robustness, only phenotypes ≥20

occurrences were included in the analysis. These diagnostic codes were transformed

into interpretable clinical phenotypes (e.g. heart failure, aortic valve disease) and

their corresponding phenotype group (e.g. cardiovascular, respiratory) using the

publicly available phecode catalog.[27,28] The use of phenotypes as representations of

clinical diagnosis codes and the reproducibility of phenotype associations have been

validated in several prior reports.[23,29–31] Patients with multiple coexisting conditions

were included in each relevant phenotype category, ensuring our cross-sectional

associations reflect real-world multimorbidity. However, to evaluate how concomitant

diseases influence our findings, we performed a sensitivity analysis in which we

established the cross-sectional associations on cohorts successively restricted to

individuals with no more than one, then no more than two, and finally no more than

three of the four key target phenotypes.

We defined phenotypes as either on-target, reflecting the specific condition

that most closely approximates the disease that each model was trained to detect, or

off-target, representing any other disease the model associated with. The target

phenotypes were defined in consensus with a panel of three clinical cardiologists,

who reviewed the AI-ECG models and available phecodes to identify those most

closely aligned with each phenotype. Target phenotype names for each specific heart

disease model were defined as follows: (1) heart failure for the LVSD model, (2)

aortic valve disease for the AS model, (3) mitral valve disease for the MR model, (4)

cardiomegaly, the phecode-derived phenotype most closely aligned with LVH,

subsequently referred to as LVH and (5) a composite outcome that encompasses all

of the above phenotypes for the SHD model **(Supplementary Table 5)**. Although

these labels are not exhaustive, they provide a reproducible and representative

framework for assessing disease-specific patterns of association and are clinically

meaningful targets that align with prior work by our group and others.[8,19,32]

To assess whether phenotypic associations with cardiovascular phenotypes

were consistent across study cohorts, we compared pairwise associations of effect

estimates between the YNHH and external validation sites. We visualized the

strength and direction of phenotype associations using volcano plots. To compare

model-specific associations with phenotypes, we generated a heatmap of ORs

between each AI-ECG model and the five target cardiovascular phenotypes. Next, we

visualized the correlation of their phenotype association profiles across the same five

target phenotypes, in a second set of heatmaps.

To assess the phenotypic overlap of AI-ECG models trained to detect

conditions with distinct pathophysiology but overlapping clinical manifestations, we

conducted a sensitivity analysis comparing the LVH and AS models from the main

study with a previously published and validated hypertrophic cardiomyopathy (HCM)

model with conditions that represent their target conditions.[33]

Association of AI-ECG with New-Onset Cardiovascular Disease

To evaluate the prognostic value of AI-ECG model predictions for new-onset

cardiovascular disease, we conducted time-to-event analyses using AI-ECG model

predictions, including those for cardiovascular disease and biological sex, as

independent variables and the predefined target phenotypes as dependent variables.

In addition to individual conditions, we included a composite endpoint comprising all

5 target phenotypes. This resulted in five outcomes for which separate Cox

proportional hazards models were fitted: (1) heart failure, (2) aortic valve disease, (3)

mitral valve disease, (4) LVH, and (5) a composite of these conditions. To ensure

analyses were limited to an at-risk sample, individuals with any of these diagnoses

recorded before or on the day of the ECG acquisition were excluded.

Statistical Analysis

The characteristics of the study population, the exposures, and the outcomes were

summarized using means and standard deviations (SD) for continuous variables and

counts and percentages for categorical variables.

Cross-sectional associations between AI-ECG model predictions and clinical

phenotypes were assessed using age and sex adjusted logistic regression, where

each phenotype was the dependent variable and individual AI-ECG probabilities were

the key independent variable. These were reported as odds ratios (ORs) and 95%

confidence intervals (95% CI). These analyses were conducted using the PheWAS R

package (version 0.99.6.1).[34,35]

To evaluate whether the associations within phenotype groups identified by AI-

ECG models exceeded what would be expected by chance, we compared the

observed proportion of significant associations of the included models with the

expected proportions using Fisher's Exact test, with corresponding ORs and p-values

reported. The predicted model probabilities from all models were logit-transformed

and standardized for all analyses. To assess the reproducibility of individual

phenotype associations across healthcare settings, we computed Pearson correlation

coefficients between ORs and compared the results across cohorts.

To evaluate similarity in phenotypic association patterns across AI-ECG

models, we first calculated pairwise Pearson correlation coefficients between ORs

across the five predefined target phenotypes, separately within each cohort. Next, we

repeated this analysis including all cardiovascular phenotypes. The resulting

correlation matrices quantified the pairwise concordance in OR distributions across

models, reflecting overlap in their phenotypic association profiles.

In assessing new-onset diseases, we developed age and sex adjusted Cox

proportional hazards models with the models' predictions as independent variables

and the time to the occurrence of the first event as the dependent variable. Follow-up

time was defined from the date of the ECG to the earliest occurrence of one of the

following events: (i) onset of the tested phenotype, (ii) death, (iii) loss to follow-up, or

(vi) day of data extraction, with a maximum follow-up time of five years. For the Cox

models, hazard ratios (HRs) and 95% confidence intervals (CIs) were reported. To

confirm that proportional‑hazards and linearity assumptions held across all models,

outcomes, and cohorts, we (1) plotted Schoenfeld residuals over time and (2) fitted

five‑knot restricted cubic splines for each predictor. Visual inspection of the disease-

specific models' diagnostics revealed no meaningful violations **(Supplementary**

Figure 3).

To control for multiple comparisons, we applied Bonferroni correction, with an

adjusted significance threshold based on the number of phenotypes analyzed in each

cohort. All statistical analyses were conducted using R (version 4.2.1) and Python

(version 3.12.5).

RESULTS

Study Population

There were 235,685 individuals with at least one ECG of antecedent care within the

YNHH. Of these, 116,540 were eligible for analysis, based on none of their data had

been used in the model development and validation of prior AI-ECG models. These

individuals had a mean age of 55.5 $\pm$ 19.1 years and 65,920 (56.6%) were female,

69,560 (59.7%) self-identified as non-Hispanic White, 22,912 (19.7%) as non-

Hispanic Black, and 18,130 (15.6%) as Hispanic. Among these participants,

hypertension was present in 55,828 (47.9%), diabetes in 20,902 (17.9%), and heart

failure in 8,611 (7.4%) individuals based on the presence of health encounters with

corresponding diagnoses for these conditions in the period preceding the index ECG

(Table 1).

The cohort drawn from the community hospitals included 63,790 individuals

who had a mean age of 61.1 ± 19.5 years, 36,088 (56.6%) were female, 41,401

(64.9%) self-reported as White, 7,760 (12.2%) as Black and 11,687 (18.3%) as

Hispanic. Of these individuals, 32,021 (50.2%) had hypertension, 12,324 (19.3%)

had diabetes, and 5,506 (8.6%) had heart failure. In the 11,005 individuals that were

included from the outpatient clinic cohort, the mean age was 61.4 ± 15.0 years, and

6,241 (56.7%) were women. A total of 8,696 (79.0%) were White, 890 (8.1%) Black,

and 928 (8.4%) Hispanic. Hypertension was present in 6,800 (61.8%), diabetes in

1,793 (16.3%), and heart failure in 408 (3.7%) individuals.

From the UK Biobank, the cohort consisted of 42,354 participants with a mean

age of 64.1 ± 7.8 years, including 21,835 (51.6%) women, and were predominantly

White (n=40,893, 96.6%). Hypertension was present in 6,269 (14.8%) of the

individuals, diabetes in 1,340 (3.2%) and 225 (0.5%) had heart failure **(Table 1)**.

Cross-sectional association of AI-ECG models with cardiovascular phenotypes

Across the 989 clinical phenotypes in the YNHH cohort, CVD-specific AI-ECG models

were significantly more likely to be associated with phenotypes categorized as

cardiovascular, compared with non-cardiovascular phenotypes, with ORs ranging

from 4.41 (p = 2.26 × 10^{-17}) for the AS model to 2.16 (p = 1.39 × $10^{-}6$) for the

composite SHD model **(Figure 1, Supplementary Data 1)**. In contrast, the AI model

trained to predict biological sex was not more likely to be associated with

cardiovascular phenotypes but showed an increase in significant associations with

genitourinary phenotypes (OR = 2.31; p = 1.34 × 10^{-7}).

Similarly, among the available phenotypes from community hospitals (n=868),

outpatient clinics (n=528), and the UK Biobank (n=687) model predictions for CVD

specific AI-ECG models were consistently more likely to be associated with a broad

range of cardiovascular phenotypes. In contrast, the AI-ECG model for biological sex

classification did not show a cardiovascular specific phenotype association pattern

(Supplementary Data 1-4, Supplementary Figures 4-5). Additionally, while some

associations between the cardiovascular disease models and non-cardiovascular

phenotype groups were observed, they lacked consistency across models and

cohorts **(Supplementary Data 1-4, Supplementary Figures 4-5)**.

Off-target Phenotypic Associations of AI-ECG

We evaluated whether the AI-ECG models for detecting specific cardiovascular

diseases were most strongly associated with their respective training labels. In the

YNHH cohort, the LVSD model was strongly associated with its target phenotype,

heart failure (OR per SD 2.91; 95% CI: 2.84-2.98). However, the model also showed

strong associations with several off-target phenotypes, including primary/intrinsic

cardiomyopathies (OR ◻3.58; 95% CI: 3.47-3.70), left bundle branch block (OR◻

3.11; 95% CI 2.96-3.26), and chronic ischemic heart disease (OR ◻2.93; 95% CI:

2.82-3.05) **(Figure 2B, Supplementary Data 5)**. This pattern was consistent across

other AI-ECG models, each showing strong associations with a range of

cardiovascular phenotypes including but not limited to its target phenotype

(Supplementary Figure 6).

The sensitivity analysis restricting this cross-sectional analysis to individuals

with progressively fewer concomitant diseases showed an attenuation of the effect

sizes, but patterns similar to the primary analysis across all models **(Supplementary**

Figure 7). The specific examination of the AS model, LVH model and the HCM model

showed that all 3 models have a high degree of overlap in the identification of each

other's target phenotypes, including the subphenotype of hypertrophic obstructive

cardiomyopathy **(Supplementary Figure 8)**.

Associations between model outputs and phenotypes were consistent across

the external validation cohorts **(Figure 2A; Supplementary Figures 9–11),** and off-

target cardiovascular disease associations frequently exceeded those with the target

phenotype**.** In the community hospitals, the LVSD model showed the strongest

associations with primary/intrinsic cardiomyopathies (OR 3.45; 95% CI: 3.31-3.61),

chronic ischemic heart disease (OR 2.87, 95% CI: 2.72-3.03) followed by its target

heart failure (OR 2.57; 2.49-2.65) **(Supplementary Figure 6; Supplementary Data**

6). Similar patterns were observed in the outpatient clinics and the UK Biobank.

For the AI-ECG model classifying biological sex, the most significant

associations across cohorts were with sex-specific conditions **(Figure 2B;**

Supplementary Figure 6; Supplementary Data 5–8).

Phenotypic Association Patterns of Disease-specific AI-ECG Models

In the YNHH cohort, the phenotype association patterns of the LVSD, MR, LVH, and

SHD models were highly correlated, with Pearson correlation coefficients ranging

from 0.88 to 0.99. In contrast, the AS model showed only moderate correlations with

the other cardiovascular disease models (r = 0.61–0.86). **(Figure 3, Supplementary**

Figure 12). These patterns persisted when extending the analysis to include all

cardiovascular phenotypes **(Supplementary Figure 13).**

On- and Off-Target Cardiovascular Predictions with AI-ECG Models

After the exclusion of individuals with any of the target phenotypes before the ECG,

97659 individuals at risk had a median follow-up of 4.3 years (interquartile range

(IQR), 2.5–5.0) in the YNHH cohort. During this period, 1,9580 individuals (2.0%)

developed heart failure, 1,309 (1.3%) aortic valve disease, 711 (0.7%) mitral valve

disease, and 2,374 (2.4%) LVH, collectively representing 5,177 individuals (5.3%)

with an adverse cardiovascular outcome. The population characteristics, follow-up

times, and incidence of key cardiovascular phenotypes for all the cohorts are detailed

in **Table 1**.

 Across models, heart failure consistently emerged as the top-ranked

phenotype in the YNHH **(Figure 4; Supplementary Figure 14–16; Supplementary**

Data 11–13). The LVSD model was associated 2-fold elevated hazard for new onset

HF (HR 2.09; 95% CI: 2.01–2.17), along with an elevated risk for a range of incident

cardiovascular diseases, including aortic valve disease (HR 1.35; 95% CI: 1.28–

1.42), mitral valve disease (HR 1.67; 95% CI: 1.56–1.79), LVH (HR 1.69; 95% CI:

1.63–1.75) and composite SHD (HR 1.67; 95% CI: 1.63–1.72) for the composite of all

these individual cardiovascular phenotypes.

These findings were consistent across cohorts **(Figure 4; Supplementary**

Figures 14–16; Supplementary Data 11–13). In the community hospitals, all AI-

ECG models were also significantly associated with the target phenotypes. In the

outpatient clinics and the UK Biobank, associations were generally consistent but

less uniformly significant **(Figure 4; Supplementary Figures 14–16)**. The AI-ECG

model trained to predict biological sex was not significantly associated with any new-

onset cardiovascular phenotype in YNHH, a finding that was consistent across all

cohorts.

Experimental Models for Non-Cardiovascular Disease Outcomes

Experimental Models for Non-Cardiovascular Disease Outcomes

Across all cohorts, experimental AI-ECG models trained on non-cardiovascular

outcomes were not significantly associated with clinical phenotypes, consistent with

their role as negative controls. A limited number of associations were observed in the

UK Biobank cohort, predominantly with cardiovascular conditions such as essential

hypertension and coronary atherosclerosis. These findings were not replicated in

other datasets **(Supplementary Figures 17–18)**.

DISCUSSION

In this study of over 230,000 individuals from four distinct clinical and community-

based cohort studies across the US and UK, we demonstrate that a range of AI-ECG

models developed to identify specific structural cardiac disorders have non-specific

associations with a range of cardiovascular conditions. These models used standard

12-lead ECG images as input, with prior validation in multinational cohorts, and

assessments confirming consistency with signal-based models in prior reports.[3,19]

While these models consistently showed stronger associations with cardiovascular

phenotypes, these associations were notable across both the target and non-target

conditions, suggesting that they function better as general cardiovascular screening

tools than as condition-specific diagnostic modalities. This pattern was consistently

observed across cohorts with prevalent and frequently coexisting cardiovascular

conditions, in sensitivity analyses with limited multimorbidity, and in a healthier

population initially free of structural heart disease. Moreover, these tools represent

digital biomarkers of future cardiovascular disease in individuals without overt cardiac

dysfunction. The strongest association was found between these AI-ECG tools and

the risk of new-onset HF, rather than the risk of future development of the target

conditions. Notably, the ensemble model trained to identify a composite of structural

heart diseases exhibited broader and stronger associations than models trained on

individual conditions.

Our study presents essential insights to interpret the evolving field of AI-ECG-

based cardiovascular screening. Currently, there has been a preponderance of AI-

ECG models developed for label specific classification tasks for a variety of

cardiovascular diseases.[1,3,11,33,36–38] This approach has led to numerous AI-ECG

models demonstrating the strong discriminatory performance for the detection of

these specific conditions.[17,39,40] However, our head-to-head evaluation of AI-ECG

models revealed that the limited selectivity of these models hinders their utility for

disease-specific screening.[41] This may be a result of limited understanding of what an

abnormal prediction on AI-ECG models may represent.[8,33] This is particularly relevant

as the exclusion of the target condition in those with abnormal AI-ECG predictions

may still represent predisposition for a range of other cardiovascular conditions.[42]

Before the widespread clinical implementation of AI-ECG, it is essential to inform

clinicians what these positive screens across different AI-ECG models may mean for

patients and to understand whether an algorithm's output reflects a precise signal

linked to its intended target condition or whether it captures broader, non-specific

disease signatures.

In contrast to the prevailing disease-specific paradigm, our findings suggest

that AI-ECG models, despite being trained to detect specific cardiovascular disease

labels, capture a highly correlated, broader physiological signature rather than

distinct disease-specific features, thereby reframing the role of AI-ECG from that of

binary diagnostic classifiers to biomarkers of cardiovascular vulnerability. This fits in

the growing recognition that ECGs encode rich, latent information spanning a wide

range of both cardiovascular and non-cardiovascular conditions.[43,44] While current

approaches fine-tune AI-ECG models to use this information for the abstraction of

disease-specific ECG signatures, our data revealed that this narrow strategy offers

limited clinical utility.[1,45]

To examine these shared signatures, we evaluated a broad number of

phenotypic associations across models. For example, left bundle branch block, a

recognizable conduction abnormality, emerged consistently as a major association

across multiple model outputs and is clinically associated with a range of

cardiovascular diseases.[46,47] Other frequently associated traits included

hypertension, diabetes, and renal dysfunction, conditions known to drive systemic

and cardiac remodeling not limited to a specific disease, further substantiating that

AI-ECG models may be detecting subtle rhythm and conduction disturbances linked

to generalized cardiovascular risk, or confounders of the target disease label used in

model development.[46,47]

Despite the limited utility of disease-specific models, the choice of training

label substantially shaped performance and clinical applicability. In our experiments,

a model trained on a well-defined composite endpoint of structural heart diseases

produced broader, more robust associations than models trained on single disease

labels. Conversely, a biological sex classification model revealed distinct non-

cardiovascular signatures, and models trained on random or ECG-unrelated labels

yielded no meaningful patterns. These results suggest that the future development of

AI-ECG algorithms would be most optimized when focused on detecting well-defined

composite cardiovascular outcomes.

These findings carry important implications for the future development,

regulatory approval, and subsequent implementation and reimbursement of AI-ECG

models. An abnormal AI-ECG-based risk profile should prompt a comprehensive

cardiovascular evaluation, including echocardiography, to uncover structural

pathology beyond the model's target. Moreover, high scores independently predict

future cardiovascular events even in the absence of overt disease, underscoring AI-

ECG's role as a prognostic biomarker. Incorporating these continuous risk outputs

into clinical workflows can guide early preventive strategies, such as lipid-lowering

therapies, intensified blood pressure control, and lifestyle modification.[48] In future

research, developers of AI-ECG tools could specifically target the development of

models that differentiate between key subphenotypes, which may improve their role

in scenarios where this would be clinically meaningful. This strategy has been

effective in other AI domains, including AI-based echocardiography.[49,50]

These findings may also have important implications for the current regulatory

and reimbursement frameworks for AI-ECG. Recently, the FDA has granted 510(k)

clearance to algorithms for models detecting hypertrophic cardiomyopathy and LVSD,

and CMS has introduced a reimbursable code for AI-ECG interpretation. [51] However,

prevailing evaluation frameworks remain anchored in disease-specific, binary

classification, which can lead to separate approval and coverage pathways for

multiple algorithms.[52] Given our observation that models trained on distinct conditions

have a highly correlated phenotypic pattern, it may be important to evaluate whether

such models truly function as disease-specific classifiers, and shift the evaluation of

AI-ECG to using a single AI-ECG model as a broad cardiovascular risk classifier,

rather than relying on multiple disease-specific tools.

Our study should be interpreted considering the following potential limitations.

Phenotype classification relied on ICD diagnostic codes, which are subject to

misclassification and variability across clinical settings, potentially affecting the

accuracy of our outcome definitions. Moreover, the use of diagnostic codes may

result in the potential underrepresentation of certain phenotypes. This is particularly

important in the UK Biobank, where only inpatient codes are available, resulting in

the low phenotype counts for certain phenotypes, such as LVH. However, the

consistent results across distinct external validation populations suggest the

generalizability of the findings despite differences in coding practices. Furthermore,

while there may be indication bias in ECG acquisition within the clinical cohorts

where ECGs are typically performed for specific clinical reasons, potentially leading

to a higher prevalence of CVD with the general population, the UK Biobank cohort

consists of research participants who underwent ECGs as part of a structured

protocol, reducing the likelihood of such bias. Additionally, PheWAS analyses involve

testing a large number of phenotypes, which increases the risk of false positives.

However, we applied strict multiple testing correction and replication in independent

datasets.[3,19,53] Moreover, AI-ECG models were trained on historical datasets with

structured outcome labels, which may not capture the full nuance of real-world

clinical practice or longitudinal disease progression. Since all models utilized in this

study are based on a single architecture and trained in the same population, caution

is warranted when extrapolating our findings to AI-ECG models developed using

different architectures or trained in distinct populations. However, image-based

models, including those used in this study, have been extensively validated and have

demonstrated similar performance to signal-based models in head-to-head

comparison, with high correlation and comparable accuracy across multiple studies

and independent datasets.[3,19,33,54,55] The study, however, suggests the role of such

selectivity assessments across different models before recommending their use as

single disease diagnostic agents. Finally, while our findings suggest predictive utility,

they remain observational. Prospective validation of a composite AI-ECG model in

pragmatic trials will be critical to translate these findings into clinical practice.

Conclusion

Despite being developed for the detection of specific cardiovascular conditions, AI-

ECG models both detect the presence as well as predict the future development of a

broad range of cardiovascular diseases with similar propensity. This challenges their

role as distinct binary diagnostic tools and instead suggests the clinical utility of AI-

ECG as a broader cardiovascular biomarker.

Acknowledgments

Sources of funding

Dr Khera was supported by the National Institutes of Health (R01AG089981,

R01HL167858, and K23HL153775) and the Doris Duke Charitable Foundation

(2022060). Dr Oikonomou was supported by the National Heart, Lung, and Blood

Institute of the National Institutes of Health (F32HL170592).

Disclosures

EKO reported being a cofounder of Evidence2Health, serving as a consultant to

Caristo Diagnostics Ltd and Ensight-AI, having stock options in Caristo Diagnostics

Ltd, receiving a grant from the National Heart, Lung, and Blood Institute of the

National Institutes of Health, and having patents 63/508,315 and 63/177,117 outside

the submitted work. RK reported receiving grants from the National Heart, Lung, and

Blood Institute, National Institutes of Health, Doris Duke Charitable Foundation,

Bristol Myers Squibb, Novo Nordisk, BridgeBio, and Blavatnik Foundation, being an

academic cofounder of Ensight-AI and Evidence2Health, having patents 63/346,610,

WO2023230345A1, US20220336048A1, 63/484,426, 63/508,315, 63/580,137,

63/606,203, 63/619,241, and 63/562,335 pending, and serving as associate editor of

JAMA outside the submitted work. No other disclosures were reported. These

affiliations and potential financial interests have been disclosed and are being

managed in accordance with institutional policies.

Supplemental Material

Supplemental Materials

Supplementary Tables 1–5

Supplementary Figures 1-18

Supplementary Data 1-12

References

1.   Siontis KC, Noseworthy PA, Attia ZI, Friedman PA. Artificial intelligence-
enhanced electrocardiography in cardiovascular disease management. *Nat Rev*
Cardiol. 2021;18:465–478.

2.   Somani S, Russak AJ, Richter F, Zhao S, Vaid A, Chaudhry F, De Freitas JK,
Naik N, Miotto R, Nadkarni GN, Narula J, Argulian E, Glicksberg BS. Deep
learning and the electrocardiogram: review of the current state-of-the-art.
Europace. 2021;23:1179–1191.

3.   Sangha V, Nargesi AA, Dhingra LS, Khunte A, Mortazavi BJ, Ribeiro AH, Banina
E, Adeola O, Garg N, Brandt CA, Miller EJ, Ribeiro ALP, Velazquez EJ, Giatti L,
Barreto SM, Foppa M, Yuan N, Ouyang D, Krumholz HM, Khera R. Detection of
left ventricular systolic dysfunction from electrocardiographic images. *Circulation*.
2023;148:765–777.

4.   Tsai D-J, Lou Y-S, Lin C-S, Fang W-H, Lee C-C, Ho C-L, Wang C-H, Lin C.
Mortality risk prediction of the electrocardiogram as an informative indicator of
cardiovascular diseases. *Digit Health*. 2023;9:20552076231187250.

5.   Attia ZI, Kapa S, Lopez-Jimenez F, McKie PM, Ladewig DJ, Satam G, Pellikka
PA, Enriquez-Sarano M, Noseworthy PA, Munger TM, Asirvatham SJ, Scott CG,
Carter RE, Friedman PA. Screening for cardiac contractile dysfunction using an
artificial intelligence-enabled electrocardiogram. *Nat Med*. 2019;25:70–74.

6.   Kwon J-M, Kim K-H, Akkus Z, Jeon K-H, Park J, Oh B-H. Artificial intelligence for
detecting mitral regurgitation using electrocardiography. *J Electrocardiol*.
2020;59:151–157.

7.   Kwon J-M, Jeon K-H, Kim HM, Kim MJ, Lim SM, Kim K-H, Song PS, Park J,
Choi RK, Oh B-H. Comparing the performance of artificial intelligence and
conventional diagnosis criteria for detecting left ventricular hypertrophy using
electrocardiography. *Europace*. 2020;22:412–419.

8.   Dhingra LS, Aminorroaya A, Sangha V, Pedroso AF, Asselbergs FW, Brant LCC,
Barreto SM, Ribeiro ALP, Krumholz HM, Oikonomou EK, Khera R. Heart failure
risk stratification using artificial intelligence applied to electrocardiogram images:
a multinational study. *Eur Heart J*. 2025;46:1044–1053.

9.   Noseworthy PA, Attia ZI, Behnken EM, Giblon RE, Bews KA, Liu S, Gosse TA,
Linn ZD, Deng Y, Yin J, Gersh BJ, Graff-Radford J, Rabinstein AA, Siontis KC,
Friedman PA, Yao X. Artificial intelligence-guided screening for atrial fibrillation
using electrocardiogram during sinus rhythm: a prospective non-randomised
interventional trial. *Lancet*. 2022;400:1206–1212.

10. Sau A, Pastika L, Sieliwonczyk E, Patlatzoglou K, Ribeiro AH, McGurk KA,
Zeidaabadi B, Zhang H, Macierzanka K, Mandic D, Sabino E, Giatti L, Barreto
SM, Camelo L do V, Tzoulaki I, O'Regan DP, Peters NS, Ware JS, Ribeiro ALP,
Kramer DB, Waks JW, Ng FS. Artificial intelligence-enabled electrocardiogram

for mortality and cardiovascular risk estimation: a model development and
validation study. *Lancet Digit Health*. 2024;6:e791–e802.

11. Sau A, Barker J, Pastika L, Sieliwonczyk E, Patlatzoglou K, McGurk KA, Peters
NS, O'Regan DP, Ware JS, Kramer DB, Waks JW, Ng FS. Artificial intelligence-
enhanced electrocardiography for prediction of incident hypertension. *JAMA*
Cardiol. 2025;10:214–223.

12. Anumana ECG-AI LEF [Internet]. [cited 2025 Feb 11];Available from:
https://anumana.ai/ecg-ai-lef

13. Kumar V, Venkataraman R, Aljaroudi W, Osorio J, Heo J, Iskandrian AE, Hage
FG. Implications of left bundle branch block in patient treatment. *Am J Cardiol*.
2013;111:291–300.

14. Sahranavard T, Alimi R, Arabkhazaei J, Nasrabadi M, Alavi Dana SMM, Gholami
Y, Izadi-Moud A, Esmaily H, Ebrahimi M, Ferns GA, Moohebati M, Saffar Soflaei
S, Ghayour Mobarhan M. Association of major and minor ECG abnormalities
with traditional cardiovascular risk factors in the general population: a large scale
study. *Sci Rep*. 2024;14:11289.

15. Jørgensen PG, Jensen JS, Marott JL, Jensen GB, Appleyard M, Mogelvang R.
Electrocardiographic changes improve risk prediction in asymptomatic persons
age 65 years or above without cardiovascular disease. *J Am Coll Cardiol*.
2014;64:898–906.

16. Hari KJ, Singleton MJ, Ahmad MI, Soliman EZ. Relation of minor
electrocardiographic abnormalities to cardiovascular mortality. *Am J Cardiol*.
2019;123:1443–1447.

17. Croon PM, Pedroso AF, Khera R. The emerging role of AI in transforming
cardiovascular care. *Future Cardiol*. 2025;1–4.

18. Zoghbi WA, Adams D, Bonow RO, Enriquez-Sarano M, Foster E, Grayburn PA,
Hahn RT, Han Y, Hung J, Lang RM, Little SH, Shah DJ, Shernan S,
Thavendiranathan P, Thomas JD, Weissman NJ. Recommendations for
noninvasive evaluation of native valvular regurgitation: A report from the
American society of echocardiography developed in collaboration with the
society for cardiovascular magnetic resonance. *J Am Soc Echocardiogr*.
2017;30:303–371.

19. Dhingra LS, Aminorroaya A, Sangha V, Pedroso AF, Shankar SV, Coppi A,
Foppa M, Brant LCC, Barreto SM, Ribeiro ALP, Krumholz HM, Oikonomou EK,
Khera R. Ensemble deep learning algorithm for structural heart disease
screening using electrocardiographic images. *J Am Coll Cardiol*. 2025;85:1302–
1313.

20. Sangha V, Khunte A, Holste G, Mortazavi BJ, Wang Z, Oikonomou EK, Khera R.
Biometric contrastive learning for data-efficient deep learning from
electrocardiographic images. *J Am Med Inform Assoc*. 2024;31:855–865.

21. Kossaify A, Nasr M. Diastolic dysfunction and the new recommendations for
echocardiographic assessment of left ventricular diastolic function: Summary of
guidelines and novelties in diagnosis and grading. *J Diagn Med Sonogr*.
2019;35:317–325.

22. Attia ZI, Friedman PA, Noseworthy PA, Lopez-Jimenez F, Ladewig DJ, Satam G,
Pellikka PA, Munger TM, Asirvatham SJ, Scott CG, Carter RE, Kapa S. Age and
sex estimation using artificial intelligence from standard 12-lead ECGs. *Circ*
Arrhythm Electrophysiol. 2019;12:e007284.

23. Denny JC, Ritchie MD, Basford MA, Pulley JM, Bastarache L, Brown-Gentry K,
Wang D, Masys DR, Roden DM, Crawford DC. PheWAS: demonstrating the
feasibility of a phenome-wide scan to discover gene-disease associations.
Bioinformatics. 2010;26:1205–1210.

24. Boland MR, Alur-Gupta S, Levine L, Gabriel P, Gonzalez-Hernandez G. Disease
associations depend on visit type: results from a visit-wide association study.
BioData Min. 2019;12:15.

25. Fan M, Li N, Huang L, Chen C, Dong X, Gao W. Exploring potential drug targets
in multiple cardiovascular diseases: A study based on proteome-wide Mendelian
randomization and colocalization analysis. *Cardiovasc Ther*.
2025;2025:5711316.

26. Wan NC, Grabowska ME, Kerchberger VE, Wei W-Q. Exploring beyond
diagnoses in electronic health records to improve discovery: a review of the
phenome-wide association study. *JAMIA Open*. 2025;8:ooaf006.

27. Denny JC, Bastarache L, Ritchie MD, Carroll RJ, Zink R, Mosley JD, Field JR,
Pulley JM, Ramirez AH, Bowton E, Basford MA, Carrell DS, Peissig PL, Kho AN,
Pacheco JA, Rasmussen LV, Crosslin DR, Crane PK, Pathak J, Bielinski SJ,
Pendergrass SA, Xu H, Hindorff LA, Li R, Manolio TA, Chute CG, Chisholm RL,
Larson EB, Jarvik GP, Brilliant MH, McCarty CA, Kullo IJ, Haines JL, Crawford
DC, Masys DR, Roden DM. Systematic comparison of phenome-wide
association study of electronic medical record data and genome-wide
association study data. *Nat Biotechnol*. 2013;31:1102–1110.

28. PheWAS Resources [Internet]. [cited 2025 Mar 21];Available from:
https://phewascatalog.org/phewas/#phe12

29. Ritchie MD, Denny JC, Crawford DC, Ramirez AH, Weiner JB, Pulley JM,
Basford MA, Brown-Gentry K, Balser JR, Masys DR, Haines JL, Roden DM.
Robust replication of genotype-phenotype associations across multiple diseases
in an electronic medical record. *Am J Hum Genet*. 2010;86:560–572.

30. Wu P, Gifford A, Meng X, Li X, Campbell H, Varley T, Zhao J, Carroll R,
Bastarache L, Denny JC, Theodoratou E, Wei W-Q. Mapping ICD-10 and ICD-
10-CM codes to phecodes: Workflow development and initial evaluation. *JMIR*
Med Inform. 2019;7:e14325.

31. Wei W-Q, Bastarache LA, Carroll RJ, Marlo JE, Osterman TJ, Gamazon ER,
Cox NJ, Roden DM, Denny JC. Evaluating phecodes, clinical classification

software, and ICD-9-CM codes for phenome-wide association studies in the electronic health record. *PLoS One*. 2017;12:e0175508.

32. Akbilgic O, Butler L, Karabayir I, Chang PP, Kitzman DW, Alonso A, Chen LY, Soliman EZ. ECG-AI: electrocardiographic artificial intelligence model for prediction of heart failure. *Eur Heart J Digit Health*. 2021;2:626–634.

33. Sangha V, Dhingra LS, Aminorroaya A, Croon PM, Sikand NV, Sen S, Martinez MW, Maron MS, Krumholz HM, Asselbergs FW, Oikonomou EK, Khera R. Identification of hypertrophic cardiomyopathy on electrocardiographic images with deep learning. *Nat Cardiovasc Res*. 2025;1–10.

34. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for phenome-wide association studies in the R environment. *Bioinformatics*. 2014;30:2375–2376.

35. PheWAS: The PheWAS R package [Internet]. Github; [cited 2025 Feb 12]. Available from: https://github.com/PheWAS/PheWAS

36. Yao X, Rushlow DR, Inselman JW, McCoy RG, Thacher TD, Behnken EM, Bernard ME, Rosas SL, Akfaly A, Misra A, Molling PE, Krien JS, Foss RM, Barry BA, Siontis KC, Kapa S, Pellikka PA, Lopez-Jimenez F, Attia ZI, Shah ND, Friedman PA, Noseworthy PA. Artificial intelligence-enabled electrocardiograms for identification of patients with low ejection fraction: a pragmatic, randomized clinical trial. *Nat Med*. 2021;27:815–819.

37. Oikonomou EK, Sangha V, Shankar SV, Coppi A, Krumholz HM, Nasir K, Miller EJ, Gallegos-Kattan C, Al-Kindi S, Khera R. Tracking the pre-clinical progression of transthyretin amyloid cardiomyopathy using artificial intelligence-enabled electrocardiography and echocardiography. *medRxiv* [Internet]. 2024 [cited 2024 Oct 9];Available from: https://pubmed.ncbi.nlm.nih.gov/39252891/

38. Kwon J-M, Kim K-H, Jo Y-Y, Jung M-S, Cho Y-H, Shin J-H, Lee Y-J, Ban J-H, Lee SY, Park J, Oh B-H. Artificial intelligence assessment for early detection and prediction of renal impairment using electrocardiography. *Int Urol Nephrol*. 2022;54:2733–2744.

39. Palermi S, Vecchiato M, Ng FS, Attia Z, Cho Y, Anselmino M, De Ferrari GM, Saglietto A, International AI-ECG Working Group. Artificial intelligence and the electrocardiogram: A modern renaissance. *Eur J Intern Med* [Internet]. 2025 [cited 2025 Jun 17];Available from: http://dx.doi.org/10.1016/j.ejim.2025.04.036

40. Mossavarali S, Vaezi A, Gholami Z, Molaei A, Yekaninejad MS, Asselbergs FW, Shafiee A. Determinants of artificial intelligence electrocardiogram-derived age and its association with cardiovascular events and mortality: a systematic review and meta-analysis. *NPJ Digit Med*. 2025;8:322.

41. Lampert J, Bhatt DL, Vaid A, Kon K, Feinman J, Jou S, Kauffman J, Nadkarni G, Reddy VY. Calibration of ECG-based deep-learning algorithm scores for patients flagged as high risk for hypertrophic cardiomyopathy. *NEJM AI* [Internet]. 2025 [cited 2025 May 5];2. Available from: http://dx.doi.org/10.1056/aioa2400421

42. Yoon M, You SC. Implementing ECG-AI in clinical care: From pixel to practice. *J*
Am Coll Cardiol. 2025;85:1314–1316.

43. Friedman SF, Khurshid S, Venn RA, Wang X, Diamant N, Di Achille P, Weng L-C,
Choi SH, Reeder C, Pirruccello JP, Singh P, Lau ES, Philippakis A, Anderson
CD, Maddah M, Batra P, Ellinor PT, Ho JE, Lubitz SA. Unsupervised deep
learning of electrocardiograms enables scalable human disease profiling. *NPJ*
Digit Med. 2025;8:23.

44. Hughes JW, Theurer J, Vukadinovic M, Rogers AJ, Somani S, Kang G,
Ghazizadeh Z, O'Sullivan JW, Jain SS, Gomes B, Salerno M, Ashley E, Zou JY,
Perez MV, Ouyang D. A deep learning phenome wide association study of the
electrocardiogram. *Eur Heart J Digit Health.* 2025;ztaf047.

45. Muzammil MA, Javid S, Afridi AK, Siddineni R, Shahabi M, Haseeb M, Fariha
FNU, Kumar S, Zaveri S, Nashwan AJ. Artificial intelligence-enhanced
electrocardiography for accurate diagnosis and management of cardiovascular
diseases. *J Electrocardiol.* 2024;83:30–40.

46. Fuchs FD, Whelton PK. High blood pressure and cardiovascular disease.
Hypertension. 2020;75:285–292.

47. Deferrari G, Cipriani A, La Porta E. Renal dysfunction in cardiovascular diseases
and its consequences. *J Nephrol.* 2021;34:137–153.

48. Antoniades C, Chan K. Using artificial intelligence to spot heart failure from
ECGs: is it prime time? *Eur Heart J.* 2025;46:1054–1056.

49. Slivnick JA, Hawkes W, Oliveira J, Woodward G, Akerman A, Gomez A, Hamza
I, Desai VK, Barrett-O'Keefe Z, Grogan M, Dispenzieri A, Scott CG, Davison HN,
Cotella J, Maurer M, Helmke S, Scherrer-Crosbie M, Soltani M, Goyal A, Zareba
KM, Cheng RK, Kirkpatrick JN, Kitano T, Takeuchi M, Tiemi Hotta V, Campos
Vieira ML, Elissamburu P, Ronderos RE, Prado A, Koutroumpakis E, Deswal A,
Pursnani A, Sarswat N, Patel AR, Addetia K, Ruberg FL, Randazzo M, Asch FM,
O'Driscoll J, Al-Roub N, Strom JB, Kidd L, Cuddy S, Upton R, Lang RM, Pellikka
PA. Cardiac amyloidosis detection from a single echocardiographic video clip: a
novel artificial intelligence-based screening tool. *Eur Heart J* [Internet].
2025;Available from: http://dx.doi.org/10.1093/eurheartj/ehaf387

50. Vrudhula A, Stern L, Cheng PC, Ricchiuto P, Daluwatte C, Witteles R, Patel J,
Ouyang D. Impact of case and control selection on training artificial intelligence
screening of cardiac amyloidosis. *JACC Adv.* 2024;3:100998.

51. Babic B, Glenn Cohen I, Stern AD, Li Y, Ouellet M. A general framework for
governing marketed AI/ML medical devices. *NPJ Digit Med.* 2025;8:328.

52. Windecker D, Baj G, Shiri I, Kazaj PM, Kaesmacher J, Gräni C, Siontis GCM.
Generalizability of FDA-approved AI-enabled medical devices for clinical use.
JAMA Netw Open. 2025;8:e258052.

53. Bastarache L, Denny JC, Roden DM. Phenome-wide association studies. *JAMA.*
2022;327:75–76.

54. Croon PM, Boonstra MJ, Allaart CP, Arends BKO, Dhingra LS, Huang Y-C, Mast
T, Khera R, Kuo C-F, Kwon J-M, Lee H-S, Lee MS, van de Leur RR, Liu Z-Y,
Oikonomou EK, Selder JL, Winter MM, Asselbergs FW. AI-ECG for LVSD
detection: a systematic review and first-in-kind multinational head-to-head
comparison [Internet]. medRxiv. 2025;Available from:
http://dx.doi.org/10.1101/2025.07.08.25331129

55. Sau A, Zeidaabadi B, Patlatzoglou K, Pastika L, Ribeiro AH, Sabino E, Peters
NS, Ribeiro ALP, Kramer DB, Waks JW, Ng FS. A comparison of artificial
intelligence-enhanced electrocardiography approaches for prediction of time-to-
mortality using electrocardiogram images. *Eur Heart J Digit Health*.
2024;ztae090.

Table 1: Baseline characteristics of the included cohorts

	Yale New Haven Hospital	Community Hospitals	Outpatient clinics	UK Biobank
Total N	116540	63790	11005	42354
Age	55.5 ± 19.1	61.1 ± 19.5	61.4 ± 15.0	64.1 ± 7.8
Sex Female	65920 (56.6%)	36088 (56.6%)	6241 (56.7%)	21835 (51.6%)
Race/Ethnicity				
White	69560 (59.7%)	41401 (64.9%)	8696 (79.0%)	40893 (96.6%)
Black	22912 (19.7%)	7760 (12.2%)	890 (8.1%)	305 (0.7%)
Hispanic	18130 (15.6%)	11687 (18.3%)	928 (8.4%)	
Missing	2624 (2.3%)	1325 (2.1%)	268 (2.4%)	116 (0.3%)
Asian	2054 (1.8%)	958 (1.5%)	146 (1.3%)	603 (1.4%)
Others	1260 (1.1%)	659 (1.0%)	77 (0.7%)	437 (1.0%)
Heart Failure n (%)	8611 (7.4%)	5506 (8.6%)	408 (3.7%)	225 (0.5%)
Acute Myocardial Infarction n (%)	1617 (1.4%)	1351 (2.1%)	161 (1.5%)	120 (0.3%)
Ischemic Heart Disease n (%)	9703 (8.3%)	6434 (10.1%)	1013 (9.2%)	2233 (5.3%)
Stroke n (%)	7710 (6.6%)	5636 (8.8%)	593 (5.4%)	488 (1.2%)
Hypertension n (%)	55828 (47.9%)	32021 (50.2%)	6800 (61.8%)	6269 (14.8%)
Diabetes n (%)	20902 (17.9%)	12324 (19.3%)	1793 (16.3%)	1340 (3.2%)
Phenotypes				
Heart failure n (%)	8357 (7.2%)	5336 (8.7%)	377 (3.4%)	183 (0.4%)
Aortic valve disease n (%)	5526 (4.8%)	3108 (5.0%)	899 (8.2%)	69 (0.2%)
Mitral valve disease n (%)	9137 (7.9%)	6564 (10.6%)	2798 (25.5%)	212 (0.5%)
Left ventricular hypertrophy n (%)	9909 (8.6%)	3310 (5.4%)	734 (6.7%)	153 (0.4%)
New-onset phenotypes				
N at risk	97659	53046	8953	42014
Median FU (yrs (iqr))	4.3 (2.5;5.0)	3.7 (2.2;5.0)	4.1 (3.0;5.0)	5.0 (4.7;5.0)
Heart failure n (%)	1980 (2.0%)	1817 (3.4%)	113 (1.3%)	181 (0.4%)
Aortic valve disease n (%)	1309 (1.3%)	916 (1.7%)	349 (3.9%)	65 (0.2%)

Mitral valve disease n (%)	711 (0.7%)	687 (1.3%)	292 (3.3%)	65 (0.2%)
Left ventricular hypertrophy n (%)	2374 (2.4%)	1346 (2.5%)	387 (4.3%)	163 (0.4%)
Composite n (%)	5177 (5.3%)	4052 (7.6%)	968 (10.8%)	395 (0.9%)

This table presents the baseline characteristics of all cohorts and prevalence of both

cross-sectional and new-onset key phenotypes. Baseline characteristics are based

on demographics and ICD codes **(Supplementary Table 4)**. Phenotypes are based

on the publicly available PheWAS catalog which uses ICD codes and phenotype

mapping **(Supplementary Table 5)**. Binary variables are expressed as n (%), while

continuous variables are reported as mean (Standard deviation). Abbreviations: FU

follow-up; iqr, inter-quartile range.

Figure 1: Overrepresentation of Associations with Cardiovascular Phenotypes by AI-ECG

Panel A: Distribution of significant associations per phenotype groups from the age sex adjusted serial logistic regression for each AI-ECG model in the Yale New Haven Health System and the four community hospitals. Each pie chart represents the proportion of significant phenotype associations across different organ systems or disease categories, as indicated by the color legend. The "Expected" distribution reflects the proportion of all tested phenotypes before applying significance thresholds. Across models, most associations relate to the circulatory system, except for the sex model. **Panel B:** The odds ratios (OR) for each phenotype category being significantly associated with the AI-ECG models compared with the expected distribution in the Yale New Haven Health System and the four community hospitals. Each marker represents the OR for a significant phenotype within a disease category. The y-axis represents the OR, while the x-axis corresponds to different AI-ECG models. Colors correspond to disease categories, diamonds indicate significant enrichment, and circles represent non-significant enrichment as shown as displayed in the legend. Abbreviations: AS, aortic stenosis; LVH, left ventricular hypertrophy; LVSD, left ventricular systolic dysfunction; MR, mitral regurgitation; SHD, structural heart disease; OR, odds ratio.

Figure 2: Associations between phenotypes and AI-ECG

Panel A: Scatter plots comparing odds ratios (ORs) between Yale New Haven Hospital (YNHH) and Community Hospitals for Sex and Mitral Regurgitation (MR). Data points corresponding to the circulatory system are highlighted in red. OR values are clipped at the 99th percentile to reduce the influence of outliers. The diagonal dashed line represents perfect agreement between the two cohorts. **Panel B:** Volcano plots displaying the associations between various phenotypes and AI-ECG model predictions across four cohorts: YNHH (blue), Community Hospitals (green), Outpatient Clinics (red), and UK Biobank (purple). The x-axis represents the odds ratios and are presented on the logarithmic scale, reflecting the effectsize and directionality of the association. The y-axis shows the $-\log_{10}$(p-value) representing the significance of the association. Labeled points indicate the top 6 statistically significant associations. Due to maximal computational precision limits, p values are capped at 1×10^{-300}. Abbreviations: AS, aortic stenosis; LVH, left ventricular hypertrophy; LVSD, left ventricular systolic dysfunction; MR, mitral regurgitation; SHD, structural heart disease; OR, odds ratio.

Figure 3: Correlation of on-target phenotype association profiles across AI-ECG models in four independent cohorts

These heatmaps show Pearson correlation coefficients between odds ratio (OR) profiles and AI-ECG models across all phenotypes that we defined as being the target for any of the models for each pair of AI-ECG models. Higher correlation coefficients indicate that models exhibit similar phenotypic association patterns across target cardiovascular conditions. Correlations are shown for four cohorts: Yale New Haven Hospitals, community hospitals, outpatient clinics, and the UK Biobank. Strong correlations were observed among cardiovascular disease (CVD) models, with the SHD model consistently showing high similarity with other disease-specific models. The model trained to predict biological sex showed consistently weaker correlation with cardiovascular models, particularly in the UK Biobank where inverse correlations were observed. AS: Aortic stenosis, LVH: Left ventricular hypertrophy, LVSD: Left ventricular systolic dysfunction, MR: Mitral regurgitation, SHD: Structural heart disease composite model, Sex: Biological sex prediction model.

Figure 4: Forest plot displaying the results of the Cox Proportional Hazards models

Forest plot displaying the hazard ratios (HR) with 95% confidence intervals (CI) for various cardiovascular conditions across four cohorts: UK Biobank (UKB, red), Yale New Haven Hospital (YNHH, blue), Community Hospitals (green), and NEMG outpatient clinics (yellow). Each panel represents a different AI-ECG model prediction, including Sex, Mitral Regurgitation (MR), Left Ventricular Systolic Dysfunction (LVSD), Left Ventricular Hypertrophy (LVH), Aortic Stenosis (AS), and Structural Heart Disease (SHD). The x-axis represents the hazard ratio, with the vertical dashed line at HR = 1.0 indicating no association. The analysis was conducted using age- and sex-adjusted Cox regression models. Abbreviations: LVSD, Left Ventricular Systolic Dysfunction; LVH, Left Ventricular Hypertrophy; MR, Mitral Regurgitation; AS, Aortic Stenosis; SHD, Structural Heart Disease.

Yale New Haven Hospitals

	AS	LVH	LVSD	MR	SHD	Sex
AS	1	0.87	0.68	0.65	0.76	0.85
LVH	0.87	1	0.95	0.9	0.97	0.68
LVSD	0.68	0.95	1	0.97	0.99	0.46
MR	0.65	0.9	0.97	1	0.96	0.31
SHD	0.76	0.97	0.99	0.96	1	0.52
Sex	0.85	0.68	0.46	0.31	0.52	1

Community hospitals

	AS	LVH	LVSD	MR	SHD	Sex
AS	1	0.97	0.85	0.79	0.85	0.77
LVH	0.97	1	0.96	0.92	0.95	0.58
LVSD	0.85	0.96	1	0.99	0.99	0.32
MR	0.79	0.92	0.99	1	1	0.23
SHD	0.85	0.95	0.99	1	1	0.32
Sex	0.77	0.58	0.32	0.23	0.32	1

Outpatient clinics

	AS	LVH	LVSD	MR	SHD	Sex
AS	1	0.99	0.97	0.93	0.96	0.41
LVH	0.99	1	0.99	0.97	0.99	0.33
LVSD	0.97	0.99	1	0.98	0.99	0.25
MR	0.93	0.97	0.98	1	1	0.11
SHD	0.96	0.99	0.99	1	1	0.19
Sex	0.41	0.33	0.25	0.11	0.19	1

UK Biobank

	AS	LVH	LVSD	MR	SHD	Sex
AS	1	0.78	0.62	0.48	0.58	0.71
LVH	0.78	1	0.8	0.41	0.73	0.64
LVSD	0.62	0.8	1	0.84	0.99	0.12
MR	0.48	0.41	0.84	1	0.89	-0.17
SHD	0.58	0.73	0.99	0.89	1	0.042
Sex	0.71	0.64	0.12	-0.17	0.042	1