Fathers' and Mothers' Support Needs and Support Experiences

after Rapid Genome Sequencing

Authors

Helen Dolling[1 2 3], Sophie Rowitch[1], Malachy Bromham[1], Stephanie Archer[2 4], Sara O'Curry[5]

[6], David H. Rowitch[3 5], F. Lucy Raymond [5 8], Claire Hughes[1 2], Kate Baker[5 7 8 9]

Affiliations

1. Centre for Child, Adolescent, and Family Research, University of Cambridge, UK.

2. Department of Psychology, University of Cambridge, UK.

3. Department of Paediatrics, University of Cambridge, Cambridge, UK.

4. Department of Public Health & Primary Care, University of Cambridge, UK.

5. Cambridge University Hospitals NHS Foundation Trust, Cambridge Biomedical

Campus, Hills Road, Cambridge CB2 0QQ, UK

6. Paediatric Psychological Medicine Services, Cambridgeshire and Peterborough NHS

Foundation Trust, Cambridge, UK.

7. MRC Cognition & Brain Sciences Unit, University of Cambridge, UK.

8. Department of Genomic Medicine, University of Cambridge, UK.

9. Department of Pathology, University of Cambridge, UK.

Corresponding author

Helen Dolling

Centre for Child, Adolescent, and Family Research,

Department of Psychology, University of Cambridge,

Free School Lane,

Cambridge, CB2 3RQ, UK

E: hd300@cam.ac.uk

M: +44 (0)7779747516

ABSTRACT

As early rapid genomic sequencing (rGS) is adopted in paediatric medicine, there is an urgency to understand and address family support needs. This mixed methods study (*Peregrin**) examined the experiences of 96 parents, 1–5 years after receiving trio rGS results for their child with a severe early-onset condition. Quantitative outcome measures assessed parental well-being, life satisfaction, and family impact, comparing results to non-clinical population data, between mothers and fathers, and according to child's diagnostic outcome. Qualitative semi-structured interviews explored parents' satisfaction with support, engagement with support networks, and unmet needs. Quantitatively, mothers exhibited elevated anxiety and depression relative to population norms, and there was a lack of strong correlation in well-being metrics within couples. Parents of children with a genomic diagnosis reported poorer well-being, explained by greater medical complexity. Qualitatively, insufficient support was more frequently reported by those whose child had received a genomic diagnosis (36%) compared to those without (6%). Families drew on a range of formal and informal support sources, including condition-specific groups, though these were accessed by a minority of fathers. These findings highlight persistent and evolving support needs in families affected by complex childhood health conditions, which persist after rGS. Parents' support needs are highly individual, vary over time and across children's illness trajectory. There remain important gaps between parental needs and support, impacting on family well-being.

Keywords: NICU, PICU, rapid genome sequencing; support; parental psychosocial outcomes

INTRODUCTION

Genomic testing is becoming available in healthcare settings worldwide; rapid genome sequencing (rGS) is now offered in neonatal and paediatric intensive care units (NICU/PICU) in England. More families are receiving diagnoses of rare or ultra-rare conditions, many being life-threatening or life-limiting, early in their child's life (1). Early genomic testing and diagnosis can have clinical and personal benefits (2,3), but can also cause anxiety, distress, uncertainty and upheaval (4,5). Studies of parental support needs in the context of paediatric rare diseases and genomic diagnostics remain scarce (6,7).

Having a child admitted to a NICU/PICU with an unexpected severe illness can be traumatic, with long-term implications for families (8,9). A range of risk and protective factors explain why some families adjust and adapt, while others struggle to maintain family functioning. Factors include problem-solving and coping styles, family relationships, socioeconomic status (SES) and resource availability (10,11). Caregiver well-being is also influenced by a child's medical complexity, adaptive functioning and behaviour (12–14). Support is consistently identified as a protective factor for family adjustment and, as a modifiable factor, provides a natural focus for interventions (15).

As genomic testing becomes more common, it is important to understand the perspectives of caregivers regarding the need for and availability of support. The transition to parenthood, irrespective of a child's health, is often accompanied by reductions in parental relationship quality and mutual support (18,19). Parents differ in their responses to childhood illness, with mothers reporting higher levels of anxiety, depression and stress symptoms than fathers (20,21). Previous research has highlighted possible gender and cultural differences in values and norms around stress, coping, and the role of support (22,23). Disease rarity may constrain support availability and amplify psychosocial impacts of childhood-onset rare disease on parents (24).

To address these issues, the Peregrin* (Parental Experiences of Rapid Early Genomic

Results in Infancy) study followed-up parents whose children had received rGS because of

early-onset illness. The first objective was to evaluate parents' psychosocial needs, using

validated questionnaires. We compared parental well-being, life satisfaction and family

impact with non-clinical population data, between mothers and fathers, and according to

presence / absence of a genomic diagnosis. The second objective was to explore parents'

support experiences through qualitative interviews, focusing on perceptions and sources of

support, and attitudes to support groups.

Support is a multidimensional construct that includes emotional, informational and

instrumental elements, and encompasses both actual and perceived support (16). Adopting

Cohen's definition of support as "*a social network's provision of psychological and material*

resources intended to benefit an individual's ability to cope with stress" (17), we applied a

multi-layered approach incorporating functional (i.e. perceived availability of support) and

structural (i.e. extent, diversity and frequency of interactions) components. By integrating

these objectives and approaches, unmet support needs can be identified, and

recommendations drawn.

MATERIALS AND METHODS

Participants

This article reports a subset of data from Peregrin*, a follow-up study of 96 parents of infants

and children who had received rGS within the Next Generation Children project (NGC)

(25,26). NGC was a translational research study (November 2016 – August 2020) with broad

inclusion criteria to explore feasibility, acceptability, clinical utility, and logistics of

implementing rGS in NICU/PICU within tertiary level hospitals in the UK. Families (n =

521, 90% trios) were recruited from three NICU, two PICU, and by rapid referrals from

Paediatric Neurology and Clinical Genetics in the East and Southeast of England. The cohort

reflected the distribution of SES within the populations served by the recruiting hospitals.

Consent was provided for long-term follow-up via NIHR Rare Diseases BioResource. The

Peregrin* cohort (data collected December 2021 – January 2023) thus represents the first

large group of UK-based infants and children with any severe, complex, and unexplained

symptoms to receive early rGS in hospital settings, before NHS implementation.

Parents were eligible for Peregrin* if they had participated in NGC and they were a parent

of (a) an infant (≤2 years at rGS), whether or not a genomic diagnosis had been received; or

(b) a child (>2 years at rGS), who had received a genomic diagnosis. Parents whose child met

the above criteria and had died > 12 months previously were eligible. Parents of children >2

years at rGS without a genomic diagnosis were excluded because their experiences are

represented within existing literature on diagnostic odyssey (27,28).

Recruitment followed a cluster stratification approach (29), with child's age (≤2 years or

>2 years at rGS) as the stratification variable and Genomic Diagnosis/No Genomic Diagnosis

as cluster boundaries. We aimed to recruit 30 families per cluster to enable subgroup analyses

and capture diverse experiences. Parents were contacted by the NIHR Bioresource via a

personalised email and letter, followed by a phone call within two weeks. Invitations were

sent in batches, prioritising older children. Each participant received a £15 gift card upon

study completion.

Study Design

This study used a convergent parallel mixed methods design to generate a multi-faceted

understanding of parental perspectives, and to allow comparisons with clinical and non-

clinical cohorts (30). Data integration occurred at multiple phases, with qualitative data given

primacy alongside demographic data for group analyses. Quantitative measures were

collected one day to a week before interviews; qualitative analyses were completed first,

followed by statistical analyses. This method was chosen to inform data interpretation and

triangulation.

Measures

Parents completed questionnaires and provided demographic information online via

Qualtrics™. Supplementary Table 1 lists questionnaire measures, internal validity metrics

and scoring information. Questionnaire measures were selected after consultation with

clinicians and researchers, and with a hospital parents' group. Individual in-depth interviews

(average length 95 minutes, range 46 min. – 2h 46 min.) were conducted via Zoom Video

Communications (https://zoom.us/) with 91 parents by the primary researcher (H.D.) and

recorded for transcription and analysis. A semi-structured interview guide, based on Aldridge

et al. (31) was applied to elicit information about psychosocial impact of trio rGS, from

which the questions on support are described here (Supplementary Table 2). Interview

questions were piloted with parents of children with complex health conditions from two

families (one mother, one father).

Questionnaire analyses

Analyses were conducted using IBM SPSS Statistics (version 28.0). Where permitted by the

scoring manual, we used mean imputation to estimate missing values; otherwise pairwise

deletion was applied, indicated by the number of individuals in reported analyses.

Distributions were explored for each variable, and parametric tests were appropriate.

Peregrin* group scores on outcome measures were compared with UK population

norms (32,33). Where UK cohort data was unavailable, other non-clinical samples were

utilised (34,35). Effect sizes were calculated as the difference between Peregrin* mean and

comparison sample mean, divided by pooled standard deviation (36). Diagnosis and No

Diagnosis sub-groups were compared using independent samples t-tests. To compare

mothers' and fathers' well-being, within-couple analyses were carried out. Pearson

correlations were used to compute bivariate associations and paired-samples t-tests or chi-

squared (McNemar's) were used to compare within-couple differences in means. Bonferroni

corrections were applied for multiple testing.

Interview Analyses

Interviews were transcribed verbatim, and all transcripts were checked for accuracy, with

68% reviewed by two researchers. Analyses were conducted in NVivo (Version 12) and in

Microsoft Excel, using inductive and deductive content analysis (CA) (37,38), and

framework analysis (FA) (39). Participant responses were coded line-by-line across all

transcripts. The codebook for support sources, and framework for support group use, were

developed from existing literature and from themes derived from close reading of 20

transcripts, using inductive and deductive approaches, and then applied to the entire data set.

Transcripts were independently double coded by research assistants (S.R., M.B.), using the

codebook and framework, and discrepancies resolved by consensus. All quotes have been

anonymised and lightly edited for clarity, with an ellipsis in square brackets […] indicating

that text has been shortened.

Support levels were categorised with four response options: "no support," "minimal,"

"sufficient," or "high level." Responses were dichotomised (none/minimal cf. sufficient/high)

for comparison between Diagnosis/No Diagnosis (Dx / NDx) and Fathers/Mothers (F / M).

Broader responses were analysed thematically. Parental engagement with support groups was

categorised based on overall use, current use, group type (e.g., condition-specific social

media group, peer messaging group, rare disease charity), and perceived benefits and

drawbacks. Single episodes of support group contact, independent online information-

seeking, and unspecified support sources were excluded (40).

RESULTS

Cohort characteristics

755 parents were contacted (22 returned to sender). Of 192 respondents (25.5%), 139 expressed interest, 52 declined, 1 was ineligible, and 96 consented to participate. Reasons given for non-participation were time constraints and family commitments. Participants (32 couples; 32 parents participating alone) were parents of 67 affected children, representing 63 families (including 6 parents of 4 deceased children, and 4 sibling pairs). 91 parents participated in an interview; 5 parents completed questionnaires only. Parental self-reported demographic and health characteristics are described in Supplementary Table 3. Fathers' (n = 37) mean age at interview was 41.0 years (SD = 5.6, range 31 - 53); mothers' (n = 56) mean age at interview was 38.7 years (SD = 5.4, range 29 - 55).

Time since rGS at interview ranged from 1 to 5 years (mean 3.5 years, SD = 1.0); 57% of families had received a genomic diagnosis by the time of participation in Peregrin*. Children (30 male, 33 female) were aged 2 - 15 years (mean 5.6 years, SD = 2.7, excluding 4 deceased children) at the time of interviews. All children lived at home, with mean 2.14 (range 1 - 5) children per household.

Children's clinical conditions and genetic diagnoses were heterogeneous, and often rare or ultra-rare. The most frequently affected body systems were nervous system (n = 41), digestive and respiratory systems (n = 35 each), cardiovascular system (n = 34) and growth (n = 27), i.e. many children had multi-system disorders. Children's Health-Related Quality of Life (HRQL, PedsQL™ V4.0 Generic Core Scales) was significantly impaired relative to population norms for overall impact, physical functioning and psychosocial functioning (Supplementary Table 4).

Questionnaire results – Family well-being and parents' mental health

Comparison to general population norms

Table 1 compares Peregrin* parents' well-being indices with UK population norms (or

equivalent non-clinical samples). Significant differences were observed in PedsQL™ Family

Impact Module (PedsQL™ 2.0 FIM) total scale and several parent HRQL subscales (36). No

differences emerged for parents' physical functioning, family functioning total score, or

family relationship scores. Life satisfaction scores remained within the average to high range

and did not differ from norms (35,41). Table 2 reports Hospital Anxiety and Depression Scale

(HADS) (42) comparisons - mothers reported significantly higher levels of anxiety (d = -

0.34) and depression (d = -0.28) compared to UK norms, while fathers did not.

Associations with diagnostic group

Parents in the Diagnosis group reported significantly higher total family impact, and

expressed more worry, compared with the No Diagnosis group (Supplementary Tables 5 and

6). These two groups also differed significantly on Total Family Function, impacting on

Daily Activities to a greater extent than Family Relationships. There were no group

differences in anxiety, depression, or life satisfaction.

Children in the Diagnosis group had significantly lower HRQL compared to the No

Diagnosis group on the Paediatric Quality of Life Inventory (PedsQL™ V4.0) (Total $t(84)$ =

5.12, $p < .001$, one-sided), prompting us to re-examine well-being differences. *Post hoc* one-

way analysis of covariance showed no significant difference in FIM total scores between

Diagnostic groups ($F_{1, 83}$ = 3.74, p = .057) but a significant covariate effect for child's HRQL

($F_{1, 83}$ = 80.72, p <.001). This indicates that parents' psychosocial functioning may be

explained by the complexity of children's conditions, rather than the presence or receipt of a

genomic diagnosis.

Comparisons between fathers and mothers and associations within couples

Within-couple comparisons (Supplementary Table 7) confirmed that mothers reported

significantly higher anxiety symptoms than fathers, while parents' depression levels did not

differ as measured by HADS. Mothers did, however, report more emotional difficulties on

the FIM Emotional Functioning subscale. Mothers also reported greater worry regarding their

child's medical treatments, side effects, others' reaction to child's condition, its impact on

other family members, and on child's future. No significant differences between mothers and

fathers emerged in Social Functioning or Communication about their child's health and

family situation.

Bivariate analysis of maternal and paternal variables showed a moderate within-

couple association for Worry (Supplementary Table 7), but not for parents' HRQL scores,

mental health symptoms or total family impact scores. Instead, each parent's FIM scores

correlated strongly with their own HRQL ($r_{(30)}$ = .949, .960, $p <$ .001 for mothers and fathers,

respectively), highlighting the individual nature of physical and mental health, and appraisal

of family impact.

Interview findings – understanding parents' support

Sources and Perceptions of Support

Participants defined support in their own terms and named varied sources of primary support.

Figure 1 presents the five most prevalent sources by parent gender (A) and by child's

diagnostic status (B), inspired by Gise et al. (43). In total, 19 categories of support were

identified (Supplementary Figure 1). Table 3 summarises and illustrates the key qualitative

findings on parental perceptions of support. Parents with adequate support cited emotional,

informational, instrumental, esteem, and companionate support from both formal sources

(e.g., hospital staff, counsellors, GPs, schools, hospices) and informal networks (e.g., spouse,

parents, friends, colleagues). Some parents relied on a few key sources or were self-

sufficient, knowing where to seek help when needed:

"Whenever I've had any concerns, and I've gone to the consultant, they've come straight back

to us." INT13, F Dx

"We have a very close strong family base who we can rely on to keep us out of any trouble."

INT22, F, NDx

"If I have concerns, I feel that I have people that I can talk to. [...] I've sought some solace in

being able to reach out and get a hold of doctor [expert in child's condition] and his team."

INT28, M, Dx

Partners and family were similarly important to most parents, while healthcare

professionals were a more frequent support source for mothers than for fathers. Many parents

reported a lack of emotional support from their extended family. In particular, grandparents

were often perceived as either unable to understand their situation, emotionally distant, or in

need of protection from distress. Some parents also lacked emotional support from their

spouse/partner due to differing coping styles or fear of burdening an already overwhelmed

partner:

"It's mostly me and my husband that talk about things and I'll speak to my sisters, and I

speak to my mum [...]. But [...] sometimes I think they don't really understand how she is."

INT2, M, Dx

"I'm the man of the house sort of thing and, you know, you gotta let it wash over you sort of,

and I don't wanna bleat. [...] There are times I'm thinking - I can't talk to her anymore about

it, because she does so much all the time!" INT5, F, Dx

For fathers, friends and the workplace were key sources of emotional and social

support, providing opportunities for venting, distraction, and self-mastery. Some employers

offered flexible working arrangements, which helped to reduce stress related to childcare,

medical appointments and financial pressures, while a few provided counselling services. In

contrast, only 5.4% of mothers cited work as a source of support, possibly reflecting lower

workforce participation (69.5% vs. 100%) and higher part-time employment rates (65.9% vs.

5.5%).

"Through the whole thing, we continue to both work, and we had very flexible and

sympathetic employer." INT6, M, Dx

Mothers valued friends and peer parents for emotional, informational, and practical

support, though many lacked time or energy for socialising, and some lost friendships due to

differing parenting experiences. For some, peers (parents of children with the same condition

or other health conditions/disabilities) became new closest friends:

"So, we've been - me and the mum - become friendly and we're helping each other out,

because there's things that she knew, and I didn't know." INT65, M, NDx

Satisfaction with Support

74.7% parents reported "sufficient" or "high" levels of support, while 25.3% reported "no" or

"minimal" support, with 6 parents giving mixed responses, which were averaged for the

analyses (illustrative quotes, Table 4). Parents of children with a genomic diagnosis were

much more likely to report insufficient support (36%) compared to those without a genomic

diagnosis (6%). The differences between fathers' (31%) and mothers' (21%) satisfaction with

support were small.

Parents reported receiving "no" or "minimal" support when either multiple support

needs (e.g., medical, practical, and emotional) were unmet, their needs were acute (e.g.,

mental health deterioration, change in child's health), available provision failed to meet

expectations, the support was granted as the result of parental "*battling*" or "*fighting*", or

parent felt emotionally and socially isolated:

"No one's spoken to us about it, no one has given us any information on it, and the new

doctor - the Geneticist, he didn't seem to know anything." INT30, F, Dx

"Actually, I recently went to a private therapist because I thought I had hit my ceiling. I

thought I was at breaking point." INT14, M, Dx

"You could go and talk to any sort of counsellor or whatever, but unless they've actually

experienced what it is like to live with a child with these really challenging points of view, it

doesn't matter how many - you know - how well qualified this counsellor is. If you just don't

get it, there's no point." INT20, F, NDx

"With the genetic? Hell, "no" - not really. It's just having to kind of fight for the speech and

language and the physio. I kind of feel like I'm just getting fobbed off all the time." INT60, M,

VUS

"We're kind of isolated really, you know. And I think probably if I had, if I could do more

things with friends, then maybe I wouldn't feel so… maybe isolated." INT2, M, Dx

Key elements of insufficient support included: (i) lack of practical home-based

support, often due to distant or elderly family members, or lack of information about social

care options; (ii) feeling emotionally unsupported by healthcare providers, and experiencing

long wait times for mental health assessments or family counselling due to lack of

psychological service provision; (iii) limited organisational support and service co-ordination,

requiring significant effort to obtain necessary health and educational services for their child;

(iv) unmet informational or emotional support needs, with uncertainty about where to seek

help, reluctance to burden others, or a preference for self-reliance. Some reported adequate

medical support for the child but unmet parental support needs, or changes in support over

time. Despite individual and gender differences in parents' willingness, need, or capacity to

seek or accept support, parents expected support to be offered to *all* parents.

Engagement with Support Groups

Parents accessed diverse support groups including online condition-specific, general

(inter)national rare disease organisations, disability-related networks, and in-person specialist

baby/toddler groups. Online platforms, particularly Facebook groups, were the most

frequently used, offering flexibility and accessibility, although availability varied by

condition, prompting some parents to create new groups, tailored to their child's diagnosis.

Mothers engaged in support groups more than twice as often as fathers (78.2% vs. 34.3%).

However, both parents acknowledged the value of such groups, often supporting other

individuals/families themselves. In most couples, at least one parent participated, allowing

the other to benefit indirectly.

"My partner is member of a group - the [syndrome] group on Facebook. [...] So, we… well,

[wife] more so… uses that just to keep abreast of the whole, you know, the condition and, [...]

with research and all that sort of stuff really." INT24, F, Dx

Parents identified both benefits and challenges to support group involvement.

Membership facilitated shared experiences and offered guidance on managing the child's

condition. Emotional and informational support helped to reduce parental isolation and

fostered a sense of belonging, especially when emotional support from family and friends

was lacking. Practical advice focused on therapies, diet, applying for disability benefits,

schooling, equipment, disability access in social spaces, and travel.

"Like she's nonverbal, so I never really know… Any epilepsy sites are helpful because for

people who've got it, they can tell you how they feel like before they have one and after. [...] A

lot of times I'll go on there to see what people think about various medications." INT18, M,

Dx

"People have said before: "Oh, do you not want to speak to other parents?" I really haven't

wanted to speak to other parents because I really wanted us to manage our situation as it is

for us as a family and for [daughter] as it is for her. I didn't want to take on other people

tales of woe and listen to how awful everything is. Anyway, having since joined this Facebook

group, it's been great in terms of knowledge and information sharing and knowing what else

is going on." INT7, M, Dx

However, support groups were described as a "double-edged sword," providing

valuable resources but also heightening anxiety about the child's future. They were

unsuitable for parents who described themselves "unsociable". Additionally, not all parents

identified as "support group users", preferring to use terminology such as "parent forums" or

"networks" instead. Some parents disengaged due to emotional distress, lack of shared

experiences, or changing needs over time.

"[...] I did find a page on Facebook of all places - a [syndrome] Facebook. And there are

people all over the world who've got kids with the same diseases [...] And people put videos

of their kids on there who are the same age as [child], and [...] it'll just give you a bit of a

boost. And then you see the other side of the flip story, where they've got a little lens on there

who are a lot worse than [child], and it hits home sort of thing." INT5, F, Dx

"Now on this group - I remember signing up to it, and I couldn't believe what I was seeing. It

was quite scary. And yeah, I'm still on the group, but they don't like me on there." INT10, M,

Dx

"Who's got the time to go and attend support group? My day is full on. I started 7:00 o'clock

and I ended 11:00 o'clock." INT12, M, Dx

Parents' engagement with groups fluctuated / evolved: some parents withdrew after

initial participation (13/55, 23.6%), while others joined after an adjustment period of months

to years, as their needs changed. Reasons for disengagement included returning to work and

time constraints, variability within condition-specific groups causing distress, and lack of

shared experiences. Some parents were excluded due to eligibility criteria (e.g., bereavement

groups for foetal vs. infant loss). Others found peer support relationships unfulfilling,

outgrew the need for support, or encountered group discontinuation.

"There's Facebook groups to join: dads' groups, bereavement groups […] - and I have joined

a couple of them. [Wife] joined one, and she said it just feels like it's just little women

349   *bitching. […] I don't go on it anymore." INT91, F, Dx*

"I tried to join a group actually through SAND's and I wasn't allowed." INT90, M, Dx

After hospital discharge, many parents found it difficult to access peer support and

missed out on early socialising and intervention opportunities. Barriers included lack of

information or absence of suitable groups, time constraints, work commitments, or a

preference for alternative support sources such as health professionals, family or friends.

Noticeable differences in one's child's development within non-specialist (mainstream)

baby/toddler groups, and lack of father-specific support groups further hindered participation

(Supplementary Table 8).

DISCUSSION

The Peregrin* study identified vulnerability and variability in well-being and quality of life

among parents of children who had received early rGS - the well-being of the cohort was

compromised, and support needs were not fully met. Parents faced barriers to support,

reinforcing the need for timely, inclusive, accessible and varied resources for parents of

children with complex medical conditions (44). Contrasts in needs emerged between

diagnostic groups, with parents of children with rare genetic, medically-complex conditions

especially at risk of emotional distress. A third of these parents reported minimal or no

support, contrasting with the expectation that obtaining a genetic diagnosis will enhance

access to support and improve well-being (24,45). This is consistent with previous findings

that parents in receipt of (rapid) genomic sequencing results, and those caring for children

with (rare) complex conditions, frequently experience caregiver stress due to emotional,

social and financial pressures, but often lack the right support at the time of highest need

(6,7,46). Across follow up studies, a subset of parents reports ongoing high anxiety and/or

depression (14,47,48). This mismatch between the needs of most-impacted families and low

support availability necessitates a holistic and dynamic approach, ensuring sustained family

support that addresses both child-focused medical care and parental well-being.

An important element of this study was the comparison between mothers' and fathers' needs

and support experiences. Consistent with previous findings (49), mothers reported above

general population levels of anxiety, worry and emotional difficulties, whereas this was not

seen at group level for fathers. Within-couple analysis indicated independent mental health

symptom scores between parents, emphasising the need for individualised psychological

interventions for parents. Fathers, similar to those of typically developing children, described

stress rather than mental health difficulties and often questioned their entitlement for

professional support, prioritising the needs of their partner (50). The differences between

parents' adjustment and adaptation deserve careful consideration in future research (20).

Parents defined support based on their specific needs and assessed its adequacy in relation

to their child, themselves, and their families. Both formal and informal support networks

were described as crucial in fostering a sense of belonging and facilitating shared experiences

among families. As observed previously in families managing medical complexity and

genetic conditions (43,51,52) perceived (functional) and actual (structural) parental support

changed over time. Informational and emotional support was most critical during the

diagnostic phase and at key transitions, while instrumental (e.g., respite care) and social

support became essential for long-term family well-being. Despite observable gender

differences in support utilisation, there was considerable overlap, with some parents engaging

in social networks typically associated with the opposite gender.

Family and partners were primary support sources for most parents, indicating the need to

foster strong family relationships, particularly when under strain from a child's diagnosis.

However, partnership status alone is an unreliable proxy for social support (53). Parents who

are heavily dependent on partners may require professional intervention when relationship

strain or mental health concerns arise. While many parents reported satisfactory social

functioning, a considerable minority experienced isolation and difficulties securing support.

Parents with high self-reliance often viewed seeking support as a weakness, despite

experiencing isolation and insufficient emotional connection. Primary caregivers would

benefit from tailored interventions to maintain friendships and mitigate social isolation,

especially during diagnosis and transitional periods (46,54).

It has been reported that rare disease diagnosis can facilitate access to condition-specific

support groups (6), yet many parents in Peregrin* did not access support groups, emphasising

the need for alternative sources of support. Some families preferred guidance from trusted

healthcare providers rather than charitable organisations or peer networks. Additionally, not

all social support was perceived as beneficial. Support groups could be distressing for parents

confronted with potential deterioration or loss, were considered unsuitable by parents who

described themselves as "*unsociable*" or "*self-reliant"*, or who preferred their child's

condition not to dominate their spare time and life experience. Thus, while support groups

benefit many, they should be supplementary rather than the primary mode of support.

Bolstering partner and family relationships should be a priority in support provision, while

professionals, employers and friends also contribute to meeting families' long-term needs

(52).

Limitations

The complexity and heterogeneity of conditions diagnosed within our study group may limit

generalisability to less severe or more common genetic conditions. The study's cross-

sectional design precluded assessment of temporal changes, with potential biases of

retrospective reporting, in the context of evolving adjustment. Despite a diverse cohort and

data from both mothers and fathers, sample size limits quantitative analyses. There are also

limitations of demographic representation - a high proportion of parents had degree-level

education and were either married or cohabiting, and participants were recruited from a

tertiary university hospital, potentially differing in access to healthcare and social services.

While we followed the methodology adopted in non-clinical samples, maximising

participation and comparisons by including both mothers and fathers, non-independence

within the dataset introduces a potential bias. Longitudinal research is needed to examine

changes in parental support needs following a child's diagnosis of a rare genomic condition.

Studies should also investigate resilience and burnout in parents who exhibit high

independence but face caregiving stress (55).

Conclusion and Practice Implications

Based on Peregrin*, healthcare professionals can improve their support of parental well-being

in several ways. First, practitioners can recognise the differential impacts of a child's

diagnosis on individual parents and validate each parent's unique experience and role within

their family. Second, self-reliant and highly independent parents may benefit from

normalising support-seeking behaviours and sharing of research evidence linking support

with improved well-being. Alternative terminology to "support" (e.g., networks, resources, or

forums) may increase engagement (of fathers in particular), without affecting parental self-

esteem. Third, support assessments and practical referrals should be integrated and their

importance reinforced, benefiting families most in need of and least able to access support.

Lastly, psychological support should be a core component of initial diagnostic and long-term

care for families who are managing complex medical and genetic conditions. Our findings

highlight the diversity of parental support preferences and the importance of offering flexible,

timely, and tailored support options at the time of genomic testing and across the child's

illness trajectory, to accommodate families' evolving needs and circumstances.

Data Availability

Additional data supporting the findings of this study are available in the Apollo repository

[https://doi.org/10.17863/CAM.121146], as part of the author's PhD thesis. Deidentified data

are available upon request and should be addressed to the corresponding author.

Acknowledgements

We thank NIHR BioResource volunteers for their participation, and gratefully acknowledge

NIHR BioResource centres, NHS Trusts and staff for their contribution. We thank the

National Institute for Health Research, NHS Blood and Transplant, and Health Data Research

UK as part of the Digital Innovation Hub Programme. The views expressed are those of the

author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and

Social Care.

We thank Madeline Connell (M.C.), Charlotte Burdge (C.B.), and Kate Poulter (K.P.) for data

cleaning and research support.

Funding Statement

This research was supported by the NIHR Cambridge Biomedical Research Centre

(NIHR203312). The views expressed are those of the authors and not necessarily those of the

Author Contributions

Conceptualisation: H.D., K.B., C.H.; Data Curation and Project Administration: H.D.;

Funding Acquisition: H.D., L.R., D.H.R., C.H., K.B.; Formal Analysis: H.D., S.R., M.B.;

Resources: H.D., S.R., M.B., L.R.; Supervision: C.H., K.B., S.O.C., S.A.; Writing-original

draft: H.D., K.B., C.H., S.O.C.; Writing-review and editing: H.D., K.B., C.H., L.R., S.O.C.,

D.H.R., S.A., S.R., M.B.

Ethics Declaration

This Peregrin* follow-up study with the parents from the Next Generation Children

project (NGC, REC reference 13/EE/0325) was approved by the local research ethics

committee (REC reference: 21/EE/0154, HVS/2020/3264, IRAS: 300380), by the NIHR

Bioresource (NBR133), and by the Cambridge University Hospitals NHS Foundation Trust

institutional review board (A095947). Written electronic informed consent was obtained from

all participants prior to study commencement and archived. All data was de-identified. The

study adhered to the principles set out in the Declaration of Helsinki.

Competing Interests

The authors declare no competing interests.

References

1. Kingsmore SF, Nofsinger R, Ellsworth K. Rapid genomic sequencing for genetic
disease diagnosis and therapy in intensive care units: a review. NPJ Genom Med .
2024 Feb 27;9(1):17.

2. Stark Z, Ellard S. Rapid genomic testing for critically ill children: time to become
standard of care? European Journal of Human Genetics. 2022 Feb 8;30(2):142–9.

3. Hayeems RZ, Luca S, Assamad D, Bhatt A, Ungar WJ. Utility of Genetic Testing
from the Perspective of Parents/Caregivers: A Scoping Review. Children. 2021 Mar
27;8(4):259.

4. Berrios C, Koertje C, Noel-MacDonnell J, Soden S, Lantos J. Parents of newborns in
the NICU enrolled in genome sequencing research: hopeful, but not naïve. Genetics in
Medicine. 2020 Feb 1;22(2):416–22.

5. S Pereira, HS Smith, LA Frankel, KD Christensen, R Islam JR. Psychosocial effect
of newborn genomic sequencing on families in the babyseq project: a randomized
clinical trial. JAMA Pediatr. 2021;175:1132–41.

6. Pelentsov LJ, Laws TA, Esterman AJ. The supportive care needs of parents caring
for a child with a rare disease: A scoping review. Disabil Health J. 2015 Oct
1;8(4):475–91.

7. Lynch F, Nisselle A, Stark Z, Gaff CL, McClaren B. Genetics follow up after rapid
genomic sequencing in intensive care: current practices and recommendations for
service delivery. European Journal of Human Genetics. 2022 Nov 11;30(11):1276–82.

8. Wyatt T, Shreffler KM, Ciciolla L. Neonatal intensive care unit admission and
maternal postpartum depression. J Reprod Infant Psychol. 2019 May 27;37(3):267–76.

9. Quadir A, Festa M, Gilchrist M, Thompson K, Pride N, Basu S. Long-term follow-
up in pediatric intensive care-a narrative review. Front Pediatr. 2024;12:1430581.

10. Hilliard ME, McQuaid EL, Nabors L, Hood KK. Resilience in Youth and Families
Living With Pediatric Health and Developmental Conditions: Introduction to the
Special Issue on Resilience. J Pediatr Psychol. 2015 Oct 27;40(9):835–9.

11. McConnell D, Savage A, Breitkreuz R. Resilience in families raising children with
disabilities and behavior problems. Res Dev Disabil. 2014 Apr;35(4):833–48.

12. Raina P, O'Donnell M, Schwellnus H, Rosenbaum P, King G, Brehaut J, et al.
Caregiving process and caregiver burden: Conceptual models to guide research and
practice. BMC Pediatr. 2004 Jan 14

13. Grunberg VA, Geller PA, Bonacquisti A, Patterson CA. NICU infant health
severity and family outcomes: a systematic review of assessments and findings in
psychosocial research. J Perinatol. 2019;39(2):156–72

14. Mann K, Alvey JC, Marty C, Murphy NA. Health-Related Quality of Life and
Family Functioning of Parents of Children with Medical Complexity. Curr Phys Med
Rehabil Rep. 2019 Mar 25

15. Fang Y, Luo J, Boele M, Windhorst D, van Grieken A, Raat H. Parent, child, and
situational factors associated with parenting stress: a systematic review. Eur Child
Adolesc Psychiatry. 2022 Jul 25

16. Wethington E, Kessler RC. Perceived Support , Received Support , and
Adjustment to Stressful Life Events. 1986;27(1):78–89.

17. Cohen S. Social relationships and health. Am Psychol. 2004 Nov;59(8):676–84.

18. Belsky J, Rovine M. Patterns of Marital Change across the Transition to
Parenthood: Pregnancy to Three Years Postpartum. J Marriage Fam. 1990 Feb;52(1):5.

19. Volling BL, Yu T, Gonzalez R, Tengelitsch E, Stevenson MM. Maternal and
paternal trajectories of depressive symptoms predict family risk and children's
emotional and behavioral problems after the birth of a sibling. Dev Psychopathol. 2019
Oct 5;31(04):1307–24.

20. Wade CM, Matthews J, Forbes F, Vertkas L, Burn M, Cann WG. Focus on
Fathers: Exploring the parenting experiences of fathers using a large population-level
sample. Child Youth Care Forum. 2023 Aug 1;52(4):801–28.

21. Van Oers HA, Haverman L, Limperg PF, van Dijk-Lokkart EM, Maurice-Stam H,
Grootenhuis MA. Anxiety and Depression in Mothers and Fathers of a Chronically Ill
Child. Matern Child Health J. 2014 Oct 4;18(8):1993–2002.

22. Uhing A, Williams JS, Garacci E, Egede LE. Gender differences in the relationship
between social support and strain and mortality among a national sample of adults. J
Behav Med. 2021;44(5):673–81.

23. Burleson BR. The experience and effects of emotional support: What the study of
cultural and gender differences can tell us about close relationships, emotion, and
interpersonal communication. Pers Relatsh. 2003 Mar 12;10(1):1–23.

24. Atkins JC, Padgett CR. Living with a Rare Disease: Psychosocial Impacts for
Parents and Family Members – a Systematic Review. J Child Fam Stud. 2024 Feb
1;33(2):617–36.

25. French CE, Delon I, Dolling H, Sanchis-Juan A, Shamardina O, Mégy K, et al.
Whole genome sequencing reveals that genetic conditions are frequent in intensively
ill children. Intensive Care Med. 2019 May 1;45(5):627–36.

26. French CE, Dolling H, Mégy K, Sanchis-Juan A, Kumar A, Delon I, et al.
Refinements and considerations for trio whole-genome sequence analysis when
investigating Mendelian diseases presenting in early childhood. Human Genetics and
Genomics Advances. 2022 Jul 25;3(3):100113.

27. Carmichael N, Tsipis J, Windmueller G, Mandel L, Estrella E. "Is it Going to
Hurt?": The Impact of the Diagnostic Odyssey on Children and Their Families. J
Genet Couns. 2015 Apr 4;24(2):325–35.

28. Bauskis A, Strange C, Molster C, Fisher C. The diagnostic odyssey: insights from
parents of children living with an undiagnosed condition. Orphanet J Rare Dis . 2022
Dec 18;17(1):233.

29. Thompson SK. Stratified adaptive cluster sampling. Vol. 78, Biometrika. 1991
[cited 2021 Apr 21].

30. Creswell JW, Inoue M. A process for conducting mixed methods data analysis. J
Gen Fam Med. 2025 Jan 1;26(1):4–11.

31. Aldridge CE, Osiovich H, (Hal) Siden H, Elliott AM. Rapid genome-wide
sequencing in a neonatal intensive care unit: A retrospective qualitative exploration of
parental experiences. J Genet Couns. 2021 Apr;30(2):616–29.

32. Upton P, Eiser C, Cheung I, Hutchings HA, Jenney M, Maddocks A, et al.
Measurement properties of the UK-English version of the Pediatric Quality of Life
InventoryTM 4.0 (PedsQLTM) generic core scales. Health Qual Life Outcomes. 2005
Dec 1;3(1):22.

33. Breeman S, Cotton S, Fielding S, Jones GT. Normative data for the Hospital
Anxiety and Depression Scale. Quality of Life Research. 2015 Feb 27;24(2):391–8.

34. Medrano GR, Berlin KS, Davies WH. Utility of the PedsQLTM family impact
module: Assessing the psychometric properties in a community sample. Quality of
Life Research. 2013 Dec;22(10):2899–907.

35. Siedlecki KL, Tucker-Drob EM, Oishi S, Salthouse TA. Life satisfaction across
adulthood: different determinants at different ages? J Posit Psychol. 2008 Jul
;3(3):153–64.

36. Varni JW, Sherman SA, Burwinkle TM, Dickinson PE, Dixon P. The PedsQLTM
Family Impact Module: Preliminary reliability and validity. Health Qual Life
Outcomes. 2004 Sep 27;2(1):55.

37. Forman J, Damschroder L. Qualitative Content Analysis. In: Advances in
Bioethics. 2007. p. 39–62.

38. Graneheim UH, Lundman B. Qualitative content analysis in nursing research:
Concepts, procedures and measures to achieve trustworthiness. Nurse Educ Today .
2004 Feb;24(2):105–12.

39. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework
method for the analysis of qualitative data in multi-disciplinary health research. BMC
Med Res Methodol. 2013 Sep 18;13(1):117.

40. Sapiets SJ, Hastings RP, Stanford C, Totsika V. Families' Access to Early
Intervention and Supports for Children With Developmental Disabilities. J Early Interv
. 2023 Jun 8;45(2):103–21.

41. Diener E, Inglehart R, Tay L. Theory and Validity of Life Satisfaction Scales. Soc
Indic Res. 2013 Jul 13;112(3):497–527.

42. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. Acta
Psychiatr Scand. 1983;67(6):361–70.

43. Gise J, Cohen LL. Social Support in Parents of Children with Cancer: A
Systematic Review. J Pediatr Psychol. 2022 Oct 13;47(3):292–305.

44. Sapiets SJ, Hastings RP, Totsika V. Predictors of Access to Early Support in
Families of Children with Suspected or Diagnosed Developmental Disabilities in the
United Kingdom. J Autism Dev Disord. 2024 Apr 4;54(4):1628–41.

45. Witt S, Schuett K, Wiegand-Grefe S, Boettcher J, Quitmann J. Living with a rare
disease - experiences and needs in pediatric patients and their parents. Orphanet J Rare
Dis. 2023 Dec 1;18(1):1–16.

46. Edelstein H, Schippke J, Sheffe S, Kingsnorth S. Children with medical
complexity: a scoping review of interventions to support caregiver stress. Child Care
Health Dev. 2017 May 29;43(3):323–33.

47. Patel R, Friedrich B, Sanderson SC, Ellard H, Lewis C. Anxiety and quality-of-life
for parents of children with undiagnosed rare conditions: A multi-site quantitative
survey study. J Genet Couns. 2025 Aug 1;34(4):e70085.

48. Robinson JO, Wynn J, Biesecker B, Biesecker LG, Bernhardt B, Brothers KB, et
al. Psychological outcomes related to exome and genome sequencing result disclosure:
a meta-analysis of seven Clinical Sequencing Exploratory Research (CSER)
Consortium studies. Genet Med. 2019;21:2781–90.

49. Cohn LN, Pechlivanoglou P, Lee Y, Mahant S, Orkin J, Marson A, et al. Health
Outcomes of Parents of Children with Chronic Illness: A Systematic Review and
Meta-Analysis. J Pediatr. 2020 Mar 1;218:166-177.e2.

50. Darwin Z, Galdas P, Hinchliff S, Littlewood E, McMillan D, McGowan L, et al.
Fathers' views and experiences of their own mental health during pregnancy and the
first postnatal year: a qualitative interview study of men participating in the UK Born
and Bred in Yorkshire (BaBY) cohort. BMC Pregnancy Childbirth. 2017 Jan
26;17(1):45.

51. Bradshaw S, Bem D, Shaw K, Taylor B, Chiswell C, Salama M, et al. Improving
health, wellbeing and parenting skills in parents of children with special health care
needs and medical complexity - a scoping review. BMC Pediatr. 2019 Aug
30;19(1):301.

52. Plumridge G, Metcalfe A, Coad J, Gill P. The role of support groups in facilitating
families in coping with a genetic condition and in discussion of genetic risk
information. Health Expectations. 2012 Sep 17;15(3):255–66.

53. Perelli-Harris B, Hoherz S, Lappegård T, Evans A. Mind the "Happiness" Gap:
The Relationship Between Cohabitation, Marriage, and Subjective Well-being in the
United Kingdom, Australia, Germany, and Norway. Demography. 2019 Aug
15;56(4):1219–46.

54. Mead N, Lester H, Chew-Graham C, Gask L, Bower P. Effects of befriending on
depressive symptoms and distress: systematic review and meta-analysis. Br J
Psychiatry. 2010 Feb 2;196(2):96–101.

55. Faw MH. Supporting the supporter. J Soc Pers Relat. 2018 Feb 1;35(2):202–23.

Figure Legends

Figure 1 Five Most Prevalent Sources of Support by **(A)** Parent Gender, and by **(B)** Child's

Diagnostic Status

Note: Abbreviations HCP = healthcare professionals, C = Counselling / Mental Health

Support, Self = Self-Reliant.

Supplementary Figure 1. All Primary Sources of Support for Fathers **(A)** and Mothers **(B)**

Note: 19 categories of support, listed in order of frequency.

Table 1. Family Well-Being: Comparison of Peregrin* Cohort to Population Norms

Scale	Population Norms *M (SD)* N variable	Total Cohort *M (SD)* N variable	Difference	Effect Size
Family Impact[1]	Non-clinical *N in italics* [a]	**FIM** N = 86	FIM	FIM
Total Impact	70.8 (14.5) *726*	61.7 (20.8)	9.1	0.60 ***
Parent HRQL	69.4 (15.5) *809*	64.5 (20.7)	4.9	0.30 *
Physical	64.9 (17.4) *901*	64.7 (22.3)	0.2	0.01
Emotional	67.6 (17.9) *897*	62.2 (23.9)	5.4	0.29 *
Social	74.4 (19.1) *896*	62.7 (26.2)	11.7	0.59 ***
Cognitive	73.5 (18.6) *871*	67.9 (22.1)	5.6	0.30 *
Communication	81.9 (17.7) *894*	57.4 (26.9)	24.5	1.31 ***
Worry	78.1 (20.1) *876*	50.6 (27.9)	27.5	1.32 ***
Family Functioning	65.5 (18.5) *870*	63.6 (25.3)	1.9	0.10
Daily Activities	63.2 (22.5) *901*	51.0 (27.9)	12.2	0.53 ***
Family Relationships	67.0 (19.4) *881*	71.2 (23.2)	-4.2	-0.21
Subjective Well-Being[2]	23.46 (6.86) [4]	SWLS N = 94 24.5 (6.4)	SWLS -1.04	SWLS -0.15

*** $p < 0.0001$; ** $p < .001$; * $p < .05$. Effect sizes are designated as small (d ~ 0.2), medium (d ~ 0.5), or large (d ~0.8 or more).

Note. Higher values equal better health-related quality of life, family functioning, and higher subjective wellbeing. FIM = Family Impact Module; HRQL = health-related quality of life; SWLS = Satisfaction with Life Scale. [1]Data from Medrano et al., 2013. [2]Data from Siedlecki et al., 2008.

[a] Ages 18 - 39 (*N* = 200) are reported to match the mean ages in the Peregrin* sample of 32.9 for mothers (range 29 - 55) and 35.7 for fathers (31 - 53); the mean and standard deviation for ages 40 - 59 are 22.37 (*7.29*) (*N* = 310).

Table 2. Parents' Mental Health: Comparison of Peregrin* Cohort to Population Norms

HADS	UK Normative Data[b]		Peregrin* $N = 94$		Difference		Effect Size	
	M (SD)		M (SD)					
Anxiety	Female	Male	Female	Male	Female	Male	Female	Male
	6.78	5.51	8.21	5.78	-1.43	-0.27	-0.34 *	-0.07
	(4.23)	(4.04)	(4.33)	(4.26)				
Depression	4.12	3.83	5.19	4.28	-1.07	-0.45	-0.28 *	-0.12
	(3.78)	(3.74)	(4.17)	(3.23)				

* $p < .05$

Note. Higher values equal increased anxiety and depression on HADS. HADS = Hospital Anxiety and Depression Scale. Anxiety and Depression subscales (7 questions each) are scored from 0 - 21. Cronbach's α for Anxiety = 0.85, α for Depression = 0.82. Data are from Breeman et al., 2015.

[b] The UK Population Norms for age range 25 - 65: Anxiety, Female $N = 3,491$; Male $N = 2,698$. Depression, Female $N = 3,503$; Male $N = 2,695$.

Table 3. Key Qualitative Findings on Parental Perceptions of Support

Main findings	Illustrative quotes
Parents evaluated support adequacy through the lens of personal, child, and family needs	*Well, we're all functioning and surviving and fine; so, I guess we have sufficient support.* INT33, M, Dx *I suppose when we think about support where our kids are concerned, we think about them being able – well, for [affected daughter] - her being able to access the things that she needs in order to have a good life and how well we can provision that. And we rely very heavily on support.* INT4, M, Dx
Parents sometimes lacked emotional support due to differing coping styles or fear of burdening others	*You can talk to your family and your mum as much as you like, but they don't know, they're not experts. [...] And yeah, it's like dealing with grief, isn't it? Yeah, everybody deals with it differently. And I yeah, I tried to say things to him [husband], but it was too much; it's too overwhelming for him to hear the worst end of the spectrum or what could potentially be the future.* INT25, M, Dx
Peer parents became an important source of support for many mothers	*[I turn] to another Mum who's got a child with the same diagnosis, for the most part, yeah, and she's somebody I've met through social groups, and we've met up several times, actually. But she will be my first point - for the most part. If it was something obviously very medical and urgent, then my son's liver team and the clinical nurse specialist there and yeah. So, I guess her and then the clinical nurse specialists are my saving graces every day.* INT84, M, Dx
Parents' and child's support needs and support provision changed over time	*Uh, sufficient now, definitely. Like now I've got the 12 hours and now he's at school - the special school - yeah, I'll get a break.* INT32, M, Dx *Yeah, I'd say it's sufficient support, only because at the moment it's all in medical terms.* INT27, M, Dx *Well, at the minute there's no support, no. Well, in a sense we don't have any support from the system as such, or from managers, or from the doctors. There's no support in terms of the mental [health] support.* INT15, F, Dx *I suppose we're just perhaps [at] a bit of a crossroads of what the… how the support that we've got around us… is it going to be adequate enough to get [affected son] to the next phase of his life? which is sort of going into*

	secondary school and things like that. Yeah, I think there's always room to add more support. INT25, M, Dx
Parents utilised a variety of sources of support	*We know that we can contact the hospital if we need anything, and we'll get a response. We've got school that we can call on. We've got parents.* INT8, M, Dx
Parents' willingness, need, or capacity to seek or accept support differed – but parents expected support to be offered to all individuals regardless	*I've got a high level of support. It's whether I ask for it is the big question.* INT87, M, Dx *I find it very difficult having to be reliant on so many people. I really, really dislike it, like <u>really</u> dislike. I hate being dependent on other people, and I've never had to be, and I really hate it. I mean, I'm at the mercy of so many people!* INT62, M, NDx *I feel that I personally I'm a person I like to find out as much information as I can. So, I do a lot of reading [...] At times it feels isolating and it's overwhelming [...] And so, I think I've sought some solace in being able to reach out and get a hold of doctor [specialist in son's rare condition] and his team, they're very responsive via e-mail. [...]and I feel going forward that that's going to be a really useful avenue to keep on with because I feel like those guys are [in] a better place to answer some of these difficult questions.* INT28, M, Dx *What could make it better is to guide us as parents on the various steps, things that you should do: like the DLA, the EHCP, which is a jungle in itself. And in the beginning, even simple things like: how do you request prescription medication? We had no idea about anything.* INT52, M, Dx
Parenting a child with CMC was experienced as a lonely journey without "right" support	*I think it was just about connecting with other families who have gone through similar things. I think at the time I was feeling a bit alone, like all my friends' children were healthy and yet hadn't... didn't know anyone who had a similar experience.* INT83, M, NDx
Getting or finding support was a "battle" for several families	*It is down to us to do - to fight the fight, if you like. [...] in terms of physiotherapy and medical research, there is minimal support.* INT7, M, Dx

Note. Abbreviations: CMC = child(ren) with medical complexity; INT followed by a number = interviewee number; M = mother, F = father; Dx = genetic diagnosis, VUS = variant of uncertain significance, NDx = no genetic diagnosis identified.

Table 4. Key Qualitative Findings on Parents' Satisfaction with Support

Main Findings	Illustrative Quotes
Satisfaction with level of support:	
"None" or "Minimal"	*At the time [of rGS] I would have said "sufficient"; now I would say "minimal", because it's very much: "deal with it on your own now". [...] it's very hard for them to keep track of everyone and to do everything and there's a lot more problems nowadays with different diagnoses and stuff that weren't available back in the day. But yeah, I do still feel there's very much a case of, yeah, the NHS is very stretched and they're not able to offer the services that they perhaps would like?* INT70, F, VUS
"Sufficient" to "High"	*Again, specialist wise [affected daughter] is pretty lucky. She's in good hands and she's got the right people in place, and then, like I say, support-wise for me and [husband], you know, it's not necessarily the quantity we have, but it's the quality that we have, and we know that there is always someone there. [...] I think we sometimes feel like we're imposing or, you know, it should be something we should be dealing with ourselves anyway, but you know, we get told off quite a lot [...] for obviously not asking for enough help. But you know, that's just how I think me and [husband] are.* INT52, M, Dx
"Mixed"	*Medically I have high support. Personally, I don't have any support, no. Medically, I have, I've got high support in terms of her die[t], like you know, the sleep studies that she gets, the OT, the physiotherapy - all that is pretty high - the nutritionist; and they're very responsive. So, I really appreciate that, but personally, I haven't got any support in terms of the social net or it's like, you know, the backup. I haven't got that.* INT12, M, Dx
Satisfaction with types of support:	
Lack of instrumental support	*Support in a way we don't have any support for him. Me and my husband. We both are managing him because he is dependent for all the basic things. [...] We have emotional support from our families, but all are in [another country].* INT16, M, Dx
Lack of emotional support and access to mental health resources	*I mean the GP is fairly useless, so it's just a case of relying on your support network. I rely on my work colleagues quite a bit to vent and I can always turn to my mum if*

	anything really gets to me, 'cause I don't like to burden the wife with it, she's got her own issues. INT51, F, Dx
	So, I think with a diagnosis like that, counselling would have been incredibly useful. Part of me thinks that having some supports in how to deal with the diagnosis, someone that knows the condition quite well, someone to say: "There will still be good days," and, "this is the support that you will have along the way," [...] Because when you receive a diagnosis like that, you don't look at the near future, you look at the whole life. INT14, M, Dx
Lack of formal support and service provision, at time of diagnosis and across illness trajectory (including death)	*Ultimately, you think when you got - when I got the test results even - okay, we're gonna get people help us now and, you know, they're gonna help us on our journey and do this, do that, and do the other, but we always seem to be fighting every single minute and every single appointment for [affected child].* INT5, F, Dx
	I never was given information about the support groups, so I wouldn't even know where to start looking. INT77, M, Dx
	So, I actually found her support myself by going to her school and getting it. But I do think that there needs to be something done with siblings because actually there is a - that is a gap. It's really hard just for young children to - they don't understand. INT90, M, Dx, Deceased
Unmet support needs, coupled with reluctance to burden others, and a preference for self-reliance	*I mean there is an old saying that you can find it easier to talk to strangers. I found it easier to talk to you right now about my feelings than I would do my own Mum and Dad or my sister or anyone that I'm close to. Because the people you're close to, you know, they've got their own issues and they've got their own problems sort of thing you don't want it, like I said with [wife], you don't want to burden them with what's going on.* INT5, F, Dx
	I'm very conscious of the fact that my mum suffers with the same [mental health] issues that I suffer with, and so I tend to pick and choose if I talk to my mum about things. INT87, M, Dx