Goal-Directed Control in Schizophrenia: Loss-Biased Engagement of the Anterior

Cingulate Relates to Negative-Symptom Outcomes

Running title: Anterior Cingulate Loss Bias in Schizophrenia

Wolfgang Omlor [1]*, Giacomo Cecere [1]*, Akhil Ratan Misra [1]*, Gao-Yang Huang [2], Xiaoyan

Wu [1], Victoria Edkins [1], Giulio Pergola [4], Boris Quednow [3,5], Philippe Tobler [3,6,7], Philipp

Homan [1,3]

[1]Psychiatric Hospital, University of Zurich, Lenggstrasse 31, Zurich, 8032, Switzerland

[2]Swiss Federal Institute of Technology Zurich, Zurich, Switzerland

[3]Neuroscience Center Zurich, University of Zurich & Swiss Federal Institute of Technology Zurich,

Zurich, Switzerland

[4]Lieber Institute for Brain Development, Johns Hopkins Medical Campus, Baltimore, Maryland;

Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine,

Baltimore, Maryland.

[5]Experimental Pharmacopsychology and Psychological Addiction Research, Department of Adult

Psychiatry and Psychotherapy, University Hospital of Psychiatry Zurich, University of Zurich, Zurich,

Switzerland

[6]Zurich Center for Neuroeconomics, Department of Economics, University of Zurich, Zurich 8006,

Switzerland

[7]University Research Priority Program "Adaptive Brain Circuits in Development and Learning" (URPP

AdaBD), University of Zurich, Zurich 8057, Switzerland.

*These authors contributed equally

Correspondence concerning this article should be addressed to:

Wolfgang Omlor, MD, PhD, Psychiatric Hospital, University of Zurich, Lenggstrasse 31, 8032, Zurich,

Switzerland, E-Mail: omlor@bli.uzh.ch; Tel.: +41 58 384 73 17.

Philipp Homan, MD, PhD, Psychiatric Hospital, University of Zurich, Lenggstrasse 31, 8032 Zurich,

Switzerland. E-mail: philipp.homan@bli.uzh.ch; Tel.: +41 58 384 33 65

Word count abstract: 248.

Word count text body: 5988.

Abstract:

Background and Hypothesis: Individuals with schizophrenia spectrum disorders (SSD)

often show diminished reward pursuit, whereas loss avoidance is relatively preserved. The

neural mechanisms of this dissociation and its relation to negative symptoms remain

unclear. We hypothesized that in SSD, cognitive resources are preferentially directed toward

avoiding losses rather than pursuing rewards, potentially limiting reward processing and

contributing to negative symptoms.

Study Design: Using computational modeling of behavior during a two-stage decision task

which distinguished between goal-directed (model-based) and habitual (model-free)

strategies under reward and loss conditions, we studied 42 stable individuals with SSD and

48 healthy controls (HC) during functional magnetic resonance imaging.

Study Results: In individuals with SSD, model-based control was shifted toward loss

avoidance relative to HC, with corresponding changes in prefrontal circuitry. In anterior

cingulate, orbitofrontal, and dorsolateral prefrontal regions, individuals with SSD showed

increased activation during model-based control in the loss condition. Within this group, loss-

biased activation in the right anterior cingulate region was associated with anhedonia. In 25

patients with available follow-up data, loss-biased activation in the right anterior cingulate

region at baseline was prospectively related to worsening of motivation and social

engagement over the subsequent year.

Conclusions: Our findings suggest that, compared to HC, those with SSD allocate their

limited cognitive resources more toward loss avoidance relative to reward pursuit. The

association between loss-biased anterior cingulate engagement and anhedonia supports a

neurocomputational account of diminished pleasure in psychotic disorders, with potential

implications for developing motivation-targeted treatments and early prediction of negative-

symptom worsening.

Keywords: Schizophrenia spectrum disorders; reinforcement learning; decision making; model-based control; functional magnetic resonance imaging; anhedonia.

Introduction:

How does the mind decide when it is worth investing cognitive effort to chase a reward versus avoid a loss? Everyday choices—double-checking a tax return to avoid a penalty or staying late at work to secure a bonus—force the brain to weigh potential gains against possible costs[1]. In schizophrenia spectrum disorders (SSD), which are marked by profound disturbances in cognition and goal-directed behavior[2-4], this balance between reward pursuit and loss avoidance appears to be altered and related to negative symptoms[5]. Strikingly, the drive to pursue rewards is blunted, whereas the capacity to avoid losses remains intact or is even heightened[5, 6]. This asymmetry raises the possibility that cognitive resources for goal-directed behavior are preferentially used to avoid losses at the expense of reward pursuit[7, 8]. Yet this cognitive resource-reallocation hypothesis has not been tested within a computational framework that separates goal-directed from habitual control and manipulates motivational context in individuals with SSD.

The two-stage reinforcement-learning task, coupled with computational modeling, allows researchers to quantify the balance between two principal decision processes that place different demands on cognitive resources: model-based control is goal-directed, forward-looking, cognitively demanding, slow, and flexible, whereas model-free control is habitual, backward-looking, resource-efficient, fast and rigid[9-30]. The two-stage decision task has been used to demonstrate that compulsive conditions, including obsessive–compulsive disorder and substance dependence, are associated with reduced model-based control in reward contexts, but not in loss contexts[31, 32]. Because model-based control hinges on cognitive resources[14, 15] and dopaminergic signaling[33-36], both often compromised in psychotic disorders[3], individuals on the schizophrenia spectrum are expected to exhibit impaired

model-based learning[4]. Indeed, in reward variants of the two-stage decision task, individuals

with SSD have shown impairments in model-based control[37].

It remains unclear, however, how reward and loss contexts modulate the balance between

model-based and model-free control in individuals with SSD. Resolving this knowledge gap

is essential to determine whether psychosis shifts goal-directed cognitive control toward loss

avoidance at the expense of reward pursuit, and whether this distortion drives reward

devaluation and negative symptoms. We therefore aimed to test whether model-based

control and its neural implementation are redirected from reward pursuit to loss avoidance in

individuals with SSD, and how this reallocation relates to distinct symptom domains. Guided

by evidence that SSD entail restricted cognitive capacity[2, 3, 38], value representation is

dependent on cognition[7, 8, 39], and reward pursuit is attenuated relative to loss avoidance[5],

we derived a set of specific questions for behavior, neural engagement and their clinical

correlates. We hypothesized that limited cognitive resources in psychotic disorders are

preferentially allocated to goal-directed avoidance of negative outcomes rather than reward

pursuit, and therefore asked whether:

(i) individuals with SSD would show relatively preserved model-based control when choices

avert losses alongside attenuated model-based behavior when they secure gains; (ii) any

such goal-directed control asymmetry would be accompanied by a corresponding neural

signature—specifically, stronger loss- than reward-related activation in three prefrontal hubs

implicated in model-based computations[4]: the anterior cingulate cortex (ACC) for allocating

cognitive resources according to the expected value of control[40-43], the orbitofrontal cortex

(OFC) for prospective value representation[5, 44, 45], and the dorsolateral prefrontal cortex

(DLPFC) for planning and working memory[46, 47]; and (iii) the magnitude of loss-biased

recruitment of model-based control regions would be associated with negative-symptom

severity. Given the link between blunted reward learning with preserved loss avoidance and

negative symptoms in SSD[5], we considered that this motivational asymmetry could reflect

aberrant allocation of goal-directed control that is linked to negative symptoms.

To address these questions, we combined functional magnetic resonance imaging (fMRI)

with the canonical two-step sequential decision task[9] (**Fig. 1a**) in medicated, clinically stable

individuals with SSD and healthy controls (HC). Separate reward and loss versions of the

task were used during fMRI to map prefrontal activation in the ACC, DLPFC, and OFC (**Fig.**

1b) during model-based control. This design allowed us to quantify context-dependent shifts

between model-based and model-free control and to characterize their neural signatures

within these regions (**Fig. 1c**). In a subset of participants with SSD who had longitudinal

clinical data, we investigated in an additional explorative analysis whether context biased

activation during model-based control predicted one-year follow-up levels of negative-

symptom–related domains. This mechanistic–prognostic approach links circuit-level

computations to both cross-sectional symptom variation and the longitudinal course of

negative-symptom–related outcomes.

Methods

Participants

50 individuals with SSD were recruited from the University Hospital of Psychiatry in Zurich,

and 48 HC were recruited through local advertisements. Diagnoses were established with a

structured clinical interview and according to ICD-10 criteria. Eligible patients met criteria for

SSD and were within five years of their first psychotic episode. In addition, eligible patients

showed no evidence of substance-induced psychosis. All patients were clinically stable, in

remission, and receiving antipsychotic medication at the time of testing. HC had no lifetime

history of psychiatric or neurological disorders. The study was approved by the Cantonal

Ethics Committee of Zurich, and all procedures adhered to its guidelines and complied with

the Declaration of Helsinki. All participants provided written informed consent prior to

participation, after receiving a complete description of the study. The final analysis sample

comprised 48 HC and 42 participants with SSD. Eight participants with SSD were excluded a

priori for quality reasons: four for insufficient task engagement and four for excessive head

motion or incomplete fMRI acquisition (details below).

Clinical Assessments

Symptoms were assessed using several rating scales. The Brief Negative Symptom Scale

(BNSS) was used to assess the different negative symptom domains of the apathy (avolition,

asociality, anhedonia) and diminished expression (blunted affect, alogia) dimensions[48]. To

complement this domain-specific measure, overall psychopathology was evaluated with the

Brief Psychiatric Rating Scale (BPRS). Detailed cognitive functioning was characterized with

the MATRICS Consensus Cognitive Battery (MCCB). All participants were required to pass a

urine drug screen and a breathalyzer test. In addition, AMDP clinical ratings (0–3 scale; 0 =

absent, 1 = mild, 2 = moderate, 3 = severe) were obtained at baseline and at follow-up

assessments at 3, 6, 9, and 12 months if individuals with SSD were available. Although

AMDP ratings provide a standardized, time-efficient clinician-rated assessment across a

broad range of psychopathological symptoms, ratings were available only for a subset of

patients (n=25) who attended outpatient follow-up visits or were clinically stable during

subsequent hospitalizations, and within this subset they were not consistently obtained at all

prespecified time points (**Supplementary Tables 1–4**). As apathy-related proxies (avolition,

asociality, anhedonia), we used the AMDP items amotivation and social withdrawal (AMDP

does not include a direct anhedonia item). As diminished-expression proxies (blunted affect,

alogia), we used constricted affect and affective rigidity (AMDP does not include a direct

alogia item).

Task Design

We used the two-stage task as described in previous research[9]. In total, participants

completed 400 trials (200 trials per context) with 3 seconds allocated for each choice. The

interval between first and second-stage stimuli was 1 second, followed by 1 second of

feedback presentation after the second-stage choice. The sequence of events in each trial is

illustrated in **Fig. 1a**. We employed the simplified version of the two-stage decision task

described by Decker et al.[49], which replaces abstract fractal cues with easily discriminable

"spaceship" and "alien" stimuli. This adaptation with an embedded story line lowers

perceptual and working-memory demands, accommodating the cognitive impairments often

observed in SSD, and provides a more intuitive, engaging environment for participants.

Each trial included two stages: In the first stage, participants used an MRI-compatible button

box to select one of two spaceships. Choices had to be made within 2 seconds; otherwise,

the trial was aborted. Once a choice was made, the selected option moved to the top of the

screen, while the unselected option disappeared. In the second stage, participants were

presented with one of two new states (two red or two purple aliens) each containing two

additional options. The participants selected one option, and the outcome was displayed. In

the reward setting, the outcome was either a monetary reward (represented by a coin) or no

reward (represented by a red cross). In the loss setting, the outcome was either a monetary

loss (represented by a broken coin) or no loss (represented by a green cross). Trials were

separated by an inter-trial interval of randomized duration, averaging approximately one

repetition time (TR). The second-stage state presented was probabilistically determined by

the first-stage choice, based on the transition scheme shown in **Fig. 1a**. The position (left or

right) of the options at each state was randomly varied between trials. Reward probabilities

for each bottom-stage option were dynamic, fluctuating with independent Gaussian noise[9].

This dynamic structure encouraged ongoing learning. The order of reward and loss☐trial

blocks (a block of 200 trials in each context) was randomized for each participant.

Before the fMRI task, participants completed a computerized training session to understand

the task structure. In this training session, participants completed 50 practice trials. They

were informed that reward probabilities would vary throughout the task, but the transition

probabilities between the first and second stages would remain fixed. Participants were also

told that each first-stage choice was predominantly associated with one of the two second-

stage states, and the specific mapping was disclosed. Participants were informed that they

would receive 50 CHF for task completion. Four participants with SSD did not engage with

the task (always selected the same response option/pressed the same button during one

stage)—and were therefore excluded from further analysis.

Behavior analysis

To examine learning effects without presupposing a particular reinforcement-learning model,

we first asked how events on the previous trial shaped choice on the subsequent trial. We

computed a stay-probability (StP) index based on the combinations of transition type

(common/rare) and outcome (reward/loss). From this, we derived straightforward and

intuitive "raw" metrics of model-based (MB_{raw}) and model-free (MF_{raw}) control

(**Supplementary Fig. 1**). Model-based and model-free raw scores were calculated

separately for reward and loss contexts as follows:

Reward context:

$$MB_{raw} = StP_{reward, common} - StP_{reward, rare} - StP_{no\ reward, common} + StP_{no\ reward, rare}$$

$\qquad$ $MF_{raw} = StP_{reward, common} + StP_{reward, rare} - StP_{no\ reward, common} - StP_{no\ reward, rare}$

Loss context:

$\qquad$ $MB_{raw} = StP_{no\ loss, common} - StP_{no\ loss, rare} - StP_{loss, common} + StP_{loss, rare}$

$\qquad$ $MF_{raw} = StP_{no\ loss, common} + StP_{no\ loss, rare} - StP_{loss, common} - StP_{loss, rare}$

Separately for each context, we defined the balance between model-based and model-free

control (B_{raw}) as the difference between the respective raw scores, providing a

straightforward and intuitive metric to quantify their relative contributions:

$\qquad$ $B_{raw} = MB_{raw} - MF_{raw}$

To quantify the context-dependent difference in the deployment of model-based versus

model-free control, we defined the raw context bias (CB_{raw}) as the difference between the

B_{raw} scores obtained in the reward and loss contexts, computed as follows:

$\qquad$ $CB_{raw} = B_{raw, reward} - B_{raw, loss}$

Computational modeling provides a powerful framework for systematically quantifying latent

cognitive processes underlying learning tasks, allowing the precise characterization of

mechanisms that cannot be directly inferred from behavioral data alone[9, 50, 51]. Therefore, we

additionally deployed a hierarchical Bayesian reinforcement-learning model that incorporates

the full choice history and individual learning rates, providing a generative, mechanistically

explicit estimate of model-based versus model-free control. We fitted a hybrid reinforcement-

learning model that integrates two algorithms: (i) model-based control, which computes

action values through prospective (recursive) planning over the task's transition structure;

and (ii) model-free control, which updates action values via reward-prediction errors,

strengthening actions that are rewarded and weakening those followed by loss[9, 52]. A key

parameter, denoted as ω, determines the balance between these systems in shaping first-

stage choices, where their predicted values diverge most. This weighting parameter is fixed

across trials, with $\omega = 1$ reflecting fully model-based control, and $\omega = 0$ indicating purely

model-free influence. Further model specifications and fitting procedures are detailed in the

supplementary materials. Similar to ω, B_{raw} quantifies the relative contribution of model-

based versus model-free learning, with higher values indicating greater reliance on model-

based control. To assess the correspondence between the two metrics, we computed their

Spearman correlation. To quantify the context-dependent difference in the deployment of

model-based relative to model-free control, we defined the context bias (CB) as the

difference between the ω scores obtained in the reward and loss contexts (**Fig. 1c**):

$$CB = \omega_{reward} - \omega_{loss}$$

Hierarchical Bayesian parameter estimation was implemented through Stan (v. 2.18.2). Prior

research has demonstrated that hierarchical parameter estimation provides more stable and

reliable estimates compared to individual-level estimation[51]. Posterior distributions were

approximated using the Markov Chain Monte Carlo (MCMC) algorithm. Each model was run

with four chains, comprising 1000 burn-in samples and 5000 post-burn-in samples. Mixing

and convergence of the chains to stationary values were assessed by computing the

convergence index ($\square$) and through visual inspection[51].

For the two-stage decision task, we utilized the *ts_par7* model that is implemented in

hBayesDM package in R and incorporated both reward and loss contexts[51]. This model

quantifies trial-by-trial decision-making dynamics, using seven parameters: α_1 (learning rate

in stage 1), β_1 (inverse temperature in stage 1), α_2 (learning rate in stage 2), β_2 (inverse

temperature in stage 2), pi (perseverance), ω (balance between model-based and model-

free learning) and lambda (eligibility trace). To determine the optimal specification, we also

fitted four- and six-parameter variants of the two-stage reinforcement-learning model and

evaluated their performance using the leave-one-out information criterion (LOOIC)[51]. LOOIC

favored the seven-parameter model over both the four- and six-parameter models

(**Supplementary Table 5**). Following the approach of previous work[53], we assessed the

identifiability of the seven-parameter two-stage reinforcement-learning model by simulating

datasets from 100 synthetic agents, separately for the reward and loss settings. Each agent

completed 200 trials (matching the empirical reward and loss blocks), and the same model

was re-fit at the individual (non-hierarchical) level. Recovery was quantified as the Spearman

rank correlation between true and recovered parameters across agents (**Supplementary**

Figs 2 and 3).

MRI processing.

Imaging was acquired on a Philips Achieva 3□T scanner with a 32-channel head coil during

resting state and during the reward and loss versions of the two-stage decision task.

Preprocessing followed an fMRIPrep-based workflow. Full acquisition parameters,

preprocessing steps, quality-control criteria, and exclusions are detailed in the

Supplementary Material. Four SSD participants were excluded from fMRI analyses: two for

excessive head motion and two for scanner malfunction resulting in incomplete data.

Analysis of fMRI Data.

To investigate neural activity associated with task conditions, an event-related analysis was

implemented using generalized linear modeling in FSL-FEAT. The analysis was based on

the timeseries of Reward Prediction Errors (RPEs) as generated from the simulation of the

model over each subject's task choices. This approach followed a framework inspired by

Daw, Gershman, Seymour, Dayan, Dolan [9], concentrating on prediction errors occurring at

two critical moments within each trial: the onset of second-stage stimuli and the onset of

reward delivery.

For the first-level analysis, two onset regressors were included in the model (i) first-stage

onset (ii) second stage and feedback onset time incorporated into the model as a single

regressor. The first onset regressor was parametrically modulated by the value of the

chosen first-stage action and its partial derivative with respect to the ω-parameter from

computational modeling (see also Supplementary Information). On the other hand, the

second onset regressor was modulated by model-free prediction error and by the difference

of model-free and model-based reward prediction error and its partial derivative with respect

to the ω-parameter. Note that this difference regressor is zero at reward delivery as both

algorithms converge at this point. To avoid confounding of the neural results due to activity

differences between these two time points per se, within subject mean centering of the

difference regressor was carried out. The analysis also included nuisance onset at feedback.

Additionally, to correct for artefacts caused by physiological noise we included 6 motion

parameters as estimated by fMRIPrep: a regressor for framewise displacement and an

additional five noise regressors calculated using anatomical CompCor. To quantify context-

dependent differences in neural activation associated with model-based versus model-free

control, we defined the neural context bias (nCB) as the difference between β coefficients for

model-based reward prediction errors estimated in reward versus loss contexts (**Fig. 1c**):

$$nCB = \beta_{reward} - \beta_{loss}$$

We defined activation as 'loss-biased' in a given group when the significant group difference

in nCB indicated a numerical reward bias in the comparison group shifting toward a

numerical loss bias in the target group. Conversely, we defined activation as 'reward-biased'

when a numerical loss bias in the comparison group shifted toward a numerical reward bias

in the target group.

Region of interest and whole brain analysis

Guided by convergent evidence across mammalian studies, we selected the three prefrontal

regions most consistently implicated in model-based control—the OFC, ACC and DLPFC—

as primary regions of interest (ROIs) to examine their context-specific activation patterns[4, 40,

43, 44, 46]. Although some studies implicate additional regions such as the hippocampus in

model-based control[54], a quantitative whole-brain meta-analysis in human participants found

no consistent hippocampal involvement in standard two-stage decision tasks and concluded

that the hippocampus plays a substantial role primarily when tasks include explicit spatial

relations between decision stages[44]. We excluded striatal regions as primary ROIs because

they encode a mixture of model-based and model-free signals[9, 44], whereas our focus was on

regions selective for model-based control. Our results also confirm and corroborate earlier

evidence[9] that the ventral striatum encodes a mixture of model-based and model-free value

signals (**Supplementary Fig. 4**). ROI masks were taken from the Harvard-Oxford cortical

atlas to ensure anatomical precision (see also Supplemental Data). Due to the absence of

explicit labels for certain prefrontal subregions in the Harvard–Oxford atlas, the middle

frontal gyrus was used as a proxy for the DLPFC[55]. Each ROI was further divided into left

and right hemispheres to allow for lateralized analysis. These resulting binary masks were

applied to extract the mean BOLD activity within each region for subsequent analyses. This

anatomically guided approach ensured consistency across subjects and enhanced

interpretability of ROI-based neural activation patterns during task performance. We

additionally examined the spatial distribution of activation within each ROI. This analysis

provided a finer-grained view of how each region supports model-based control in the

reward relative to loss contexts. Because our ROI analysis was hypothesis-driven and

therefore might overlook regions that contribute to model-based control, we also conducted

an exploratory whole-brain analysis.

Statistical analyses

Potential differences between individuals with SSD and healthy participants were assessed

using two-tailed Wilcoxon rank-sum tests. Within-group differences across contexts (e.g.,

omega parameters in reward versus loss contexts within the schizophrenia group) were

evaluated using two-tailed Wilcoxon signed-rank tests. Associations between context-

dependent ROI activation bias/behavior bias and different variables (negative symptom

domains, age, sex, chlorpromazine equivalents, illness duration, occupancy of $5\text{-}HT_{2A}$ and

D_2-receptors[56] were initially explored with Spearman's rank correlations to obtain robust

estimates. Significant associations were further examined using general linear models

(GLMs), adjusting for age, sex, antipsychotic dose, and illness duration. ROIs were further

examined in a longitudinal analysis in which we tested whether baseline context-biased

activation predicted the follow-up course of negative-symptom–related features derived from

AMDP ratings. To aid interpretation, we recoded the AMDP items so that higher scores

indicate better functioning and related them to context-biased activation (positive = reward

bias, negative = loss bias): motivation (inverse of reduced drive), social engagement

(inverse of social withdrawal), affective range (inverse of constricted affect), and affective

flexibility (inverse of affective rigidity). In the longitudinal AMDP analysis (see above), we

computed per domain (e.g. motivation) the mean follow-up severity (FU; arithmetic mean

across all available assessments between 3 and 12 months) for each participant with at least

1 follow-up. We also recorded K, the per-participant number of available follow-up

assessments (range 1–4), to account for unequal follow-up density. To investigate the

prospective association of context-biased ROI activation with follow-up features that are

related to negative symptoms, an ANCOVA was used in the SSD group:

$$FU_i = \beta_0 + \beta_1\,ROI_i + \beta_2\,BASE_i + \beta_3\,K_i + \varepsilon_i$$

where FU_i is the mean 3-12 months domain severity for participant i, ROI_i is the context bias

of participant i in the given ROI at baseline, $BASE_i$ and K_i are mean-centered baseline level

and follow-up count, and $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$. We report unstandardized coefficients (β) with 95%

confidence intervals, model R^2, and partial R^2 for the ROI term (unique variance beyond

baseline and K). Added-variable plots (**Fig. 4c-f**) display follow-up scores adjusted for

baseline and K versus rACC context bias. Two-sided tests were used. The association

between true and recovered model parameters was investigated with Spearman's rank

correlations. For all analyses involving multiple comparisons, p-values were corrected using

FDR correction according to Benjamini-Hochberg. Statistical significance was defined as a

FDR-corrected $P < 0.05$. Unless noted otherwise, reported p-values are FDR-adjusted.

Results

The SSD cohort and HC showed no significant differences in age, sex, or ethnicity (**Table 1**).

However, participants with SSD had lower educational attainment scores compared to

controls (SSD: 1.66 ± 1.17; controls: 2.23 ± 1.29, $p = 0.02$). All participants with SSD

received antipsychotic medication. Symptom severity was mild to moderate (BPRS total

score: 39.04 ± 10.43). Negative symptom ratings on the BNSS showed both apathy-related

symptoms (anhedonia: 10.68 ± 4.63; avolition: 6.5 ± 2.76; asociality: 6.06 ± 2.65) and

diminished-expression components (blunted affect: 9.5 ± 4.55; alogia: 4.34 ± 2.48).

Decision behavior in reward and loss contexts:

In the raw behavioral analysis (B_{raw}, see Methods), individuals with SSD showed reduced

model-based control compared to HC in both reward ($p = 0.005$) and loss contexts ($p = $

0.014; **Supplementary Fig. 1**). Within-group comparisons revealed that HCs exhibited a

numerically higher contribution of model-based control in the reward relative to the loss

context, whereas individuals with SSD showed the opposite pattern—a numerically higher

contribution in the loss relative to the reward context (**Fig. 2a, Supplementary Fig. 5a**).

However, these context-dependent differences were not statistically significant ($p = 0.28$).

To obtain a more nuanced estimate of model-based versus model-free contributions, we

applied a hybrid reinforcement learning model. Consistent with the raw behavioral findings,

the hybrid model confirmed reduced model-based control in individuals with SSD in the

reward context ($p = 7.9 \times 10^{-13}$), but not in the loss context, when comparing individual-level

ω parameters across groups (**Supplementary Fig. 5b**). The model further revealed a

significant reward bias in HCs ($p = 3.6 \times 10^{-\square}$) and a significant loss bias in the SSD group

($p = 2.9 \times 10^{-\square}$). The difference in context-dependent biases between groups was

significant ($p = 2.8 \times 10^{-1\square}$; **Fig. 2b**). We also examined whether ω-context bias was related

to age, sex, illness duration, antipsychotic dose (chlorpromazine equivalents), or 5-HT_{2A} and

D_2 receptor occupancy (see Methods), but none of these associations reached significance.

Across all contexts and groups, ω and B_{raw} were strongly correlated ($r_s = 0.60$–0.70, $p \leq 1 \times$

$10^{-\square}$ in every case), indicating that B_{raw} reliably approximates the model-derived estimate

and similarly captures the balance between model-based and model-free control. In the

reward context, the posterior distribution of group mean differences confirmed that

individuals with SSD had a lower model-based weight parameter (ω), as well as lower

learning rates at both the first (α_1) and second stage (α_2), and reduced exploitation at the first

stage (β_1) compared to HC (HC minus SSD, 95% HDIs; Δω: $0.16 - 0.68$; Δα1: $0.04 - 0.86$;

Δα2: $0.16 - 0.65$; Δβ1: $0.23 - 3.29$; **Supplementary Fig. 6**). In the loss context, the posterior

distribution indicated significantly lower second-stage learning rates and reduced exploitation

at both stages in the SSD group, while the model-based weight parameter (ω) did not differ

between groups (HC minus SSD, 95% HDIs: Δα2: $0.31 - 0.9$; Δβ1: $0.61 - 3.57$; Δβ2: $0.06 -$

1.37; **Supplementary Fig. 7**). Both in the reward and loss setting, MCMC samples of all

model parameters were well-mixed and converged to stationary distributions

(**Supplementary Figs 8 and 9**). Convergence to stationary distributions was additionally

indicated by the index of convergence ($\square$ across parameters: 1.00 to 1.04 in the reward and

loss setting). Parameter-recovery analysis confirmed that all seven hybrid-model parameters

were identifiable: across contexts, all Spearman correlations between true and recovered

values of the seven parameters were positive and significant ($p \square \leq \square 0.001$ in every case; ω:

$r_s = 0.72$ in the reward context; $r_s = 0.76$ in the loss context; for more details, see

Supplementary Figs $\square$2 - 3).

Context-dependent neural activations:

In the ACC, HC showed a significant reward bias in both hemispheres (*signrank test*; left and

right: p = 0.001), while individuals with SSD exhibited a numerical loss bias. Group

comparisons revealed a significant difference in context bias for both the left and right ACC

(*ranksum test*; left and right: $p = 6*10^{-4}$, **Fig. 3a, b**). In the DLPFC, a significant reward bias

was found in HC in both hemispheres (*signrank test*; left: p = 0.002; right: p = 0.01). In the

left hemisphere, this reward bias differed significantly from the numerical loss bias observed

in individuals with SSD (ranksum test: p = 0.02, **Fig. 3a**). In the OFC, HC showed a

significant reward bias in both hemispheres (*signrank test*; left: $p = 0.001$; right: $p = 0.01$),

while individuals with SSD demonstrated a significant shift toward a numerical loss bias

(*ranksum test*; left: $p = 0.001$; right: $p = 0.002$, **Fig. 3a, b**). Context-biased activation within

ROIs was predominantly observed in central portions of the DLPFC and OFC in both groups.

Within the ACC, individuals with SSD exhibited loss-biased activation predominantly in

dorsal regions, whereas reward-biased activation in controls was stronger in ventral regions

(**Fig. 3c, d**).

We next examined whether context bias in any of these regions was associated with

negative symptom domains. Only the context bias in the right ACC showed a significant

negative correlation with the symptom domain of anhedonia (r_s = -0.41; p = 0.048; **Fig. 4a**).

Thus, greater loss-biased activation in rACC was associated with greater anhedonia. As an

additional control, we regressed anhedonia on rACC context bias while adjusting for age,

sex, illness duration and antipsychotic dose (see Methods); the bias remained a significant

negative predictor ($\beta = -4.26 \pm 1.51 SE, p = 0.008$). We also tested whether regional

context bias correlated with age, sex, illness duration, antipsychotic dose (chlorpromazine

equivalents), HT_{2A}- and D_2-occupancy according to their medications (see Methods); none of

these associations reached significance. Apart from the a priori regions of interest (ACC,

OFC, and DLPFC), our exploratory fMRI analysis identified additional areas showing reward-

or loss-biased activation during model-based control (**Supplementary Fig. 10**). During

model-based choices, individuals with SSD showed loss-biased activity centered in anterior

and posterior cingulate as well as temporal regions, whereas reward-biased responses were

observed mainly in occipito-temporal regions (**Supplementary Table 6**). By contrast, HC

expressed a broad reward-biased network spanning for instance cingulate and paracingulate

regions, parietal, orbitofrontal and supramarginal regions, thalamus, angular gyrus as well as

further medial and dorsolateral prefrontal regions, with only a small occipital-pole cluster

showing the opposite loss-dominant pattern (**Supplementary Table 7**).

Baseline rACC context bias predicts follow-up negative-symptom–related outcomes:

In the SSD subset with longitudinal clinical data (n=25), particularly baseline anterior

cingulate context bias during goal-directed control was prospectively associated with

negative-symptom-related outcomes over follow-up (**Fig. 4b-f**). In ANCOVA, greater loss

bias in rACC predicted worse baseline-adjusted follow-up levels (mean 3–12 months) in

several negative-symptom–related domains. The association was strongest for motivation (β

= 1.83, 95% CI 1.20–2.46, p = 3.18×10^{-5}; partial R^2 = 0.64; model R^2 = 0.79), and significant

effects were also observed for social engagement (β = 1.11, 95% CI 0.31–1.92, p = 0.03;

partial R^2 = 0.28; model R^2 = 0.54, **Fig. 4c, d**). A similar but smaller effect on motivation (β =

1.55, 95% CI 0.86–2.25, p = 4x10^{-3}; partial R^2 = 0.51; model R^2 = 0.718) and a similar effect

on social engagement (β = 1.15, 95% CI 0.4–1.91, p = 0.03; partial R^2 = 0.33, model R^2 =

0.57) was present in the lACC. In addition, the lOFC showed a weak yet significant

association between context-biased activation and social engagement (β = 0.91, 95% CI

0.171–1.65, p = 0.04; partial R^2 = 0.238; model R^2 = 0.51). No other ROI showed a

significant association with the course of negative-symptom–related domains, and no ROI

was significantly related to the course of affective range or affective flexibility. The largest

associations with affective range and affective flexibility were also observed in the rACC,

positive in direction but did not survive multiple-comparison correction (**Fig. 4e, f**). In

summary, the longitudinal analysis indicates that greater baseline loss bias, particularly in

anterior cingulate regions, forecasts poorer subsequent apathy-related outcomes, most

prominently in motivation.

Discussion

The present study provides converging behavioral and neural evidence that SSD are characterized by a systematic re-allocation of goal-directed cognitive resources away from reward pursuit and toward loss avoidance. Relative to the reward-biased deployment of model-based strategies in HC, individuals with SSD showed model-based control primarily when potential losses were at stake. This loss-biased strategy was mirrored at the neural level by preferential engagement of the bilateral ACC, OFC and left DLPFC during model-based learning in the loss condition. Only the degree of loss-biased ACC activation tracked the severity of a negative symptom domain at baseline, given the association with anhedonia. Critically, baseline loss-biased engagement in the right anterior cingulate cortex also carried prognostic information: A more loss-biased ACC context-bias signal predicted poorer follow-up levels in several negative-symptom–related domains, with the strongest effect for motivation. Together, these results offer a mechanistic explanation for the clinical observation that individuals with schizophrenia show impaired pursuit of positive outcomes yet intact loss avoidance[5, 6], while linking this asymmetry to the negative symptom domain anhedonia and to the subsequent course of negative symptom-related outcomes. It is worth mentioning that the raw behavioral analysis showed only a non-significant numerical context effect, while the hierarchical reinforcement learning model revealed a robust difference. In contrast to the raw behavioral analysis, the hierarchical reinforcement-learning model fits the full choice history and jointly estimates learning rate, perseveration, and lapse parameters, and is therefore better suited to distill the goal-directed context effect. Hierarchical pooling further reduces within-group noise through shrinkage, increasing power to detect between-group differences[51]. Given these analytic differences, an underlying context effect that is subtle in the coarse, previous-trial-based summary analysis of raw behavior is likely to emerge more robustly when estimated with the full hierarchical hybrid model.

Canonical reinforcement-learning theory assumes that model-based control is deployed when the expected benefits of flexible planning outweigh its cognitive costs[4, 9]. In healthy

adults, appetitive contexts typically enhance the perceived value of planning, leading to

robust model-based control in reward contexts, as observed in our control sample and in

previous studies[44, 46]. The loss-biased pattern in participants with SSD suggests that for

these individuals the subjective cost–benefit calculus is distorted: avoiding losses is deemed

more valuable than obtaining gains. One parsimonious interpretation is that patients discount

prospective rewards because they anticipate diminished pleasure from realizing potential

gains[57], whereas losses remain fully salient. In line with this idea, low "decision acuity"—a

trait-like reduction in overall planning capacity marked by noisier, short-sighted choices[58]—

could further restrict cognitive resources and channel them chiefly into preventing losses,

thereby producing the loss-biased goal-directed control we observe. A complementary line of

evidence shows that people with SSD are less willing to expend cognitive effort even for

large rewards, pointing to a domain-general undervaluation of appetitive incentives[59, 60].

These findings align with evidence for impaired value representation during reward

processing relative to preserved loss avoidance in SSD[5], and our results frame this as the

consequence of a strategic shift toward loss-oriented cognitive control.

Our fMRI results reveal how the cognitive bias towards loss avoidance in individuals with

SSD is implemented in the prefrontal cognitive network. In controls, reward-related increases

in model-based control were accompanied by greater activation of bilateral ACC, OFC and

left DLPFC—regions known to orchestrate cost-dependent allocation of cognitive resources,

value representation, and prospective planning according to goal-representation[4, 47]. In

individuals with SSD, the same circuit was preferentially engaged for model-based control

when potential losses, rather than rewards, were at stake. These findings extend prior

observations of blunted OFC and DLPFC responses to positive outcomes[8] by demonstrating

that computations underpinning flexible decision-making are context-dependent and retuned

toward loss avoidance in psychotic disorders. While loss-dominated activation in the OFC of

individuals with SSD may signal a heightened valuation of avoiding negative outcomes, a

similar activation pattern in the left DLPFC might reflect increased recruitment of cognitive

resources during the planning of loss avoidance compared to reward seeking[4, 42, 43, 46]. The

ACC emerged as a particularly pivotal hub: its loss-biased engagement was the only neural

measure that covaried with a negative symptom domain. The ACC is ideally positioned to

integrate motivational signals with task-control demands and to evaluate whether engaging

the additional cognitive effort required for model-based learning is worthwhile[40, 42, 43]. A loss-

dominated encoding of action value within the ACC could therefore funnel limited cognitive

resources into preventing negative outcomes, leaving insufficient capacity for goal-directed

reward processing and manifesting clinically as anhedonia. Our longitudinal analysis

suggests that this anterior cingulate control signature is not merely epiphenomenal but

captures variance in later negative-symptom-related outcomes beyond baseline levels,

consistent with a candidate neurocomputational marker for risk stratification. Interestingly,

the association between rACC loss bias and current as well as future negative-symptom–

related outcomes was evident for apathy-related domains but not for diminished-expression-

related components. This pattern is consistent with accounts positing a link between

aberrant valence processing and negative symptoms specifically within the apathy

dimension[48, 61], and it further specifies an rACC-centered goal-directed control mechanism

with prognostic relevance.

Our findings converge with mounting evidence that the ACC is central to psychotic disorders,

with ACC dysfunction contributing to impairments in proactive control, and uniform structural

features characterizing this region in schizophrenia[4, 41, 62]. Identifying a loss-biased planning

mode in schizophrenia has potential translational relevance. First, it provides a

neurocomputational phenotype that could be quantified and used to stratify patients or

monitor treatment responses. Second, the association between anhedonia and ACC loss

bias positions the anterior cingulate cortex as a promising target for circuit-based

interventions. Non-invasive brain stimulation targeting the ACC or novel pharmacological

interventions may re-balance resource allocation and thereby ameliorate motivational

deficits. Our findings also extend earlier active-inference results showing elevated forgetting

and reduced policy precision during action selection in early psychosis[63], and place the

concomitant computational profile in a motivational framework that distinguishes loss from

reward contexts.

This study has several limitations. First, our neurocomputational findings are cross-sectional

and cannot establish whether loss-biased planning predates illness onset, emerges

progressively or reflects a secondary adaptation to chronic disability. Longitudinal studies in

high-risk and first-episode cohorts are needed to pinpoint the emergence of loss-biased

cognitive processing in schizophrenia. Second, participants were medicated, and systematic

modulation of cognitive processing and behavioral strategies by the complex receptor

modulation of antipsychotic agents[56, 64, 65] cannot be excluded. However, we found no

associations between context-dependent brain activation or behavior and chlorpromazine

equivalents, 5-HT2A and D2 receptor occupancy. These findings argue against the

interpretation that loss-biased behavior and brain activation observed in individuals with SSD

merely reflect medication effects. Third, a recent study showed that modeling parameters in

reinforcement tasks can vary over time, potentially limiting their capacity for precision

psychiatry[66]. In our case, a within-subject contrast of reward and loss context may attenuate

common-mode fluctuations such as global response scaling and lapse-related noise, thereby

yielding greater longitudinal stability than single-context parameters. Finally, the monetary

outcomes used here do not capture the rich spectrum of rewards and losses encountered in

daily life, and virtual-reality tasks embedding social, vocational and other incentives could

enhance ecological validity.

In summary, by integrating computational modeling with neuroimaging, we demonstrate that

SSD are not marked by a uniform deficit in goal-directed cognition but by a selective

redirection of cognitive resources toward loss avoidance. This behavioral shift is instantiated

in a prefrontal network comprising ACC, OFC and DLPFC, and loss-biased engagement of

the ACC is coupled to the negative symptom domain anhedonia. Our findings provide a

mechanistic explanation for the preservation of avoidance learning alongside impaired

reward pursuit in schizophrenia and outline circuit-level targets for interventions aimed at

restoring adaptive motivation.

Acknowledgements

The authors thank Dr. Franziska Knolle for her critical reading of the manuscript and her

valuable scientific advice.

Funding/Support:

This work was supported by a NARSAD grant from the Brain & Behavior Research

Foundation (28445, awarded to Philipp Homan) and by a grant from the Baugarten Stiftung

via Stiftung für wissenschaftliche Forschung an der Universität Zürich (STWF-22-007,

awarded to Wolfgang Omlor).

Competing interests: Dr. P. Homan has received grants and honoraria from Boehringer

Ingelheim, Janssen, Lundbeck, Mepha, Neurolite, and Novartis. The other authors declared

no competing interests.

Author Contributions: W.O. and P.H. designed the experiments. G.C. and W.O. performed

the experiments. W.O., A.M., G.H. and G.C. analyzed the data. X.W., V.E., G.P., B.Q., P.T.

contributed to the concept of the data analysis and manuscript. P.H. initiated and supervised

the study. All authors approved, refined and edited the final version of the manuscript.

Data and Code availability

Upon publication, all de-identified data underlying the results and the analysis code will be

deposited in a public repository with persistent identifiers (DOIs) and an open license;

repository links will be added to the final article.

References

1. Reitich-Stolero T, Aberg KC, Halperin D, et al. Rate and noise in human amygdala drive increased exploration in aversive learning. *Nature* Aug 27 2025.

2. Marder SR, Cannon TD. Schizophrenia. *N Engl J Med* Oct 31 2019;381(18):1753-1761.

3. McCutcheon RA, Reis Marques T, Howes OD. Schizophrenia-An Overview. *JAMA Psychiatry* Feb 1 2020;77(2):201-210.

4. Barch DM, Culbreth AJ, Sheffield JM. Cognitive Control in Schizophrenia: Advances in Computational Approaches. *Curr Dir Psychol Sci* Feb 2024;33(1):35-42.

5. Gold JM, Waltz JA, Matveeva TM, Kasanova Z, Strauss GP, Herbener ES, Collins AG, Frank MJ. Negative symptoms and the failure to represent the expected reward value of actions: behavioral and computational modeling evidence. *Arch Gen Psychiatry* Feb 2012;69(2):129-138.

6. Marder SR, Umbricht D. Negative symptoms in schizophrenia: Newly emerging measurements, pathways, and treatments. *Schizophr Res* Aug 2023;258:71-77.

7. Kring AM, Elis O. Emotion deficits in people with schizophrenia. *Annu Rev Clin Psychol* 2013;9:409-433.

8. Strauss GP, Waltz JA, Gold JM. A review of reward processing and motivational impairment in schizophrenia. *Schizophr Bull* Mar 2014;40 Suppl 2(Suppl 2):S107-116.

9. Daw ND, Gershman SJ, Seymour B, Dayan P, Dolan RJ. Model-based influences on humans' choices and striatal prediction errors. *Neuron* Mar 24 2011;69(6):1204-1215.

10. Lee SW, Shimojo S, O'Doherty JP. Neural computations underlying arbitration between model-based and model-free learning. *Neuron* Feb 5 2014;81(3):687-699.

11. Shahar N, Moran R, Hauser TU, Kievit RA, McNamee D, Moutoussis M, Consortium N, Dolan RJ. Credit assignment to state-independent task representations and its relationship with model-based decision making. *Proc Natl Acad Sci U S A* Aug 6 2019;116(32):15871-15876.

12. Deserno L, Moran R, Michely J, Lee Y, Dayan P, Dolan RJ. Dopamine enhances model-free credit assignment through boosting of retrospective model-based inference. *Elife* Dec 9 2021;10.

13. Feher da Silva C, Lombardi G, Edelson M, Hare TA. Rethinking model-based and model-free influences on mental effort and striatal prediction errors. *Nat Hum Behav* Jun 2023;7(6):956-969.

14. Otto AR, Gershman SJ, Markman AB, Daw ND. The curse of planning: dissecting multiple reinforcement-learning systems by taxing the central executive. *Psychol Sci* May 2013;24(5):751-761.

15. Otto AR, Raio CM, Chiang A, Phelps EA, Daw ND. Working-memory capacity protects model-based learning from stress. *Proc Natl Acad Sci U S A* Dec 24 2013;110(52):20941-20946.

16. Karthik SK, Pereira KMK, Culbreth AJ, Barch DM, Moran EK. Higher
motivation and pleasure scores predict more reliance on model-free decision
making. *Cogn Affect Behav Neurosci* May 22 2025.
17. Daw ND. Are we of two minds? *Nat Neurosci* Nov 2018;21(11):1497-1499.
18. Kool W, Cushman FA, Gershman SJ. When Does Model-Based Control Pay
Off? *PLoS Comput Biol* Aug 2016;12(8):e1005090.
19. Gillan CM, Kosinski M, Whelan R, Phelps EA, Daw ND. Characterizing a
psychiatric symptom dimension related to deficits in goal-directed control.
Elife Mar 1 2016;5.
20. Seow TXF, Benoit E, Dempsey C, Jennings M, Maxwell A, O'Connell R, Gillan
CM. Model-Based Planning Deficits in Compulsivity Are Linked to Faulty
Neural Representations of Task Structure. *J Neurosci* Jul 28
2021;41(30):6539-6550.
21. Vaghi MM, Moutoussis M, Vasa F, et al. Compulsivity is linked to reduced
adolescent development of goal-directed control and frontostriatal functional
connectivity. *Proc Natl Acad Sci U S A* Oct 13 2020;117(41):25911-25922.
22. Grosskurth ED, Bach DR, Economides M, Huys QJM, Holper L. No
substantial change in the balance between model-free and model-based
control via training on the two-step task. *PLoS Comput Biol* Nov
2019;15(11):e1007443.
23. Sebold M, Nebe S, Garbusow M, et al. When Habits Are Dangerous: Alcohol
Expectancies and Habitual Decision Making Predict Relapse in Alcohol
Dependence. *Biol Psychiatry* Dec 1 2017;82(11):847-856.
24. Drummond N, Niv Y. Model-based decision making and model-free learning.
Curr Biol Aug 3 2020;30(15):R860-R865.
25. Doll BB, Simon DA, Daw ND. The ubiquity of model-based reinforcement
learning. *Curr Opin Neurobiol* Dec 2012;22(6):1075-1081.
26. Collins AGE, Cockburn J. Beyond dichotomies in reinforcement learning. *Nat
Rev Neurosci* Oct 2020;21(10):576-586.
27. Karagoz AB, Moran EK, Barch DM, Kool W, Reagh ZM. Evidence for shallow
cognitive maps in Schizophrenia. *Cogn Affect Behav Neurosci* Mar 20 2025.
28. Konovalov A, Krajbich I. Gaze data reveal distinct choice processes
underlying model-based and model-free reinforcement learning. *Nat Commun*
Aug 11 2016;7:12438.
29. Economides M, Kurth-Nelson Z, Lubbert A, Guitart-Masip M, Dolan RJ.
Model-Based Reasoning in Humans Becomes Automatic with Training. *PLoS
Comput Biol* Sep 2015;11(9):e1004463.
30. Knolle F, Sen P, Culbreth AJ, et al. Investigating disorder-specific and
transdiagnostic alterations in model-based and model-free decision making. *J
Psychiatry Neurosci* 2024;49(6):E389-E401.
31. Voon V, Derbyshire K, Ruck C, et al. Disorders of compulsivity: a common
bias towards learning habits. *Mol Psychiatry* Mar 2015;20(3):345-352.
32. Voon V, Baek K, Enander J, et al. Motivation and value influences in the
relative balance of goal-directed and habitual behaviours in obsessive-
compulsive disorder. *Transl Psychiatry* Nov 3 2015;5(11):e670.
33. Wunderlich K, Smittenaar P, Dolan RJ. Dopamine enhances model-based
over model-free choice behavior. *Neuron* Aug 9 2012;75(3):418-424.
34. Deserno L, Huys QJ, Boehme R, et al. Ventral striatal dopamine reflects
behavioral and neural signatures of model-based control during sequential
decision making. *Proc Natl Acad Sci U S A* Feb 3 2015;112(5):1595-1600.

35. Doll BB, Bath KG, Daw ND, Frank MJ. Variability in Dopamine Genes Dissociates Model-Based and Model-Free Reinforcement Learning. *J Neurosci* Jan 27 2016;36(4):1211-1222.

36. Sharp ME, Foerde K, Daw ND, Shohamy D. Dopamine selectively remediates 'model-based' reward learning: a computational approach. *Brain* Feb 2016;139(Pt 2):355-364.

37. Culbreth AJ, Westbrook A, Daw ND, Botvinick M, Barch DM. Reduced model-based decision-making in schizophrenia. *J Abnorm Psychol* Aug 2016;125(6):777-787.

38. Collins AGE, Albrecht MA, Waltz JA, Gold JM, Frank MJ. Interactions Among Working Memory, Reinforcement Learning, and Effort in Value-Based Choice: A New Paradigm and Selective Deficits in Schizophrenia. *Biol Psychiatry* Sep 15 2017;82(6):431-439.

39. Collins AG, Brown JK, Gold JM, Waltz JA, Frank MJ. Working memory contributions to reinforcement learning impairments in schizophrenia. *J Neurosci* Oct 8 2014;34(41):13747-13756.

40. Akam T, Rodrigues-Vaz I, Marcelo I, Zhang X, Pereira M, Oliveira RF, Dayan P, Costa RM. The Anterior Cingulate Cortex Predicts Future States to Mediate Model-Based Action Selection. *Neuron* Jan 6 2021;109(1):149-163 e147.

41. Omlor W, Rabe F, Fuchs S, et al. Estimating Multimodal Structural Brain Variability in Schizophrenia Spectrum Disorders: A Worldwide ENIGMA Study. *Am J Psychiatry* Apr 1 2025;182(4):373-388.

42. Shenhav A, Botvinick MM, Cohen JD. The expected value of control: an integrative theory of anterior cingulate cortex function. *Neuron* Jul 24 2013;79(2):217-240.

43. Shenhav A, Cohen JD, Botvinick MM. Dorsal anterior cingulate cortex and the value of control. *Nat Neurosci* Sep 27 2016;19(10):1286-1291.

44. Huang Y, Yaple ZA, Yu R. Goal-oriented and habitual decisions: Neural signatures of model-based and model-free learning. *Neuroimage* Jul 15 2020;215:116834.

45. Wilson RC, Takahashi YK, Schoenbaum G, Niv Y. Orbitofrontal cortex as a cognitive map of task space. *Neuron* Jan 22 2014;81(2):267-279.

46. Smittenaar P, FitzGerald TH, Romei V, Wright ND, Dolan RJ. Disruption of dorsolateral prefrontal cortex decreases model-based in favor of model-free control in humans. *Neuron* Nov 20 2013;80(4):914-919.

47. Stein DJ, Costa DLC, Lochner C, Miguel EC, Reddy YCJ, Shavitt RG, van den Heuvel OA, Simpson HB. Obsessive-compulsive disorder. *Nat Rev Dis Primers* Aug 1 2019;5(1):52.

48. Begue I, Kaiser S, Kirschner M. Pathophysiology of negative symptom dimensions of schizophrenia - Current developments and implications for treatment. *Neurosci Biobehav Rev* Sep 2020;116:74-88.

49. Decker JH, Otto AR, Daw ND, Hartley CA. From Creatures of Habit to Goal-Directed Learners: Tracking the Developmental Emergence of Model-Based Reinforcement Learning. *Psychol Sci* Jun 2016;27(6):848-858.

50. Homan P, Levy I, Feltham E, et al. Neural computations of threat in the aftermath of combat trauma. *Nat Neurosci* Mar 2019;22(3):470-476.

51. Ahn WY, Haines N, Zhang L. Revealing Neurocomputational Mechanisms of Reinforcement Learning and Decision-Making With the hBayesDM Package. *Comput Psychiatr* Oct 2017;1:24-57.

52. Worbe Y, Palminteri S, Savulich G, Daw ND, Fernandez-Egea E, Robbins
TW, Voon V. Valence-dependent influence of serotonin depletion on model-
based choice strategy. *Mol Psychiatry* May 2016;21(5):624-629.
53. Shahar N, Hauser TU, Moutoussis M, Moran R, Keramati M, consortium N,
Dolan RJ. Improving the reliability of model-based decision-making estimates
in the two-stage decision task with reaction-times and drift-diffusion modeling.
PLoS Comput Biol Feb 2019;15(2):e1006803.
54. Miller KJ, Botvinick MM, Brody CD. Dorsal hippocampus contributes to model-
based planning. *Nat Neurosci* Sep 2017;20(9):1269-1276.
55. Schienle A, Kogler W, Wabnegger A. A randomized trial that compared brain
activity, efficacy and plausibility of open-label placebo treatment and cognitive
reappraisal for reducing emotional distress. *Sci Rep* Aug 26
2023;13(1):13998.
56. Nyberg S, Farde L, Eriksson L, Halldin C, Eriksson B. 5-HT2 and D2
dopamine receptor occupancy in the living human brain. A PET study with
risperidone. *Psychopharmacology (Berl)* 1993;110(3):265-272.
57. Kring AM, Barch DM. The motivation and pleasure dimension of negative
symptoms: neural substrates and behavioral outputs. *Eur
Neuropsychopharmacol* May 2014;24(5):725-736.
58. Moutoussis M, Garzon B, Neufeld S, et al. Decision-making ability,
psychopathology, and brain connectivity. *Neuron* Jun 16 2021;109(12):2025-
2040 e2027.
59. Barch DM, Culbreth AJ, Ben Zeev D, Campbell A, Nepal S, Moran EK.
Dissociation of Cognitive Effort-Based Decision Making and Its Associations
With Symptoms, Cognition, and Everyday Life Function Across
Schizophrenia, Bipolar Disorder, and Depression. *Biol Psychiatry* Sep 15
2023;94(6):501-510.
60. Treadway MT, Peterman JS, Zald DH, Park S. Impaired effort allocation in
patients with schizophrenia. *Schizophr Res* Feb 2015;161(2-3):382-385.
61. Waltz JA, Xu Z, Brown EC, Ruiz RR, Frank MJ, Gold JM. Motivational Deficits
in Schizophrenia Are Associated With Reduced Differentiation Between Gain
and Loss-Avoidance Feedback in the Striatum. *Biol Psychiatry Cogn Neurosci
Neuroimaging* Mar 2018;3(3):239-247.
62. Brugger SP, Howes OD. Heterogeneity and Homogeneity of Regional Brain
Structure in Schizophrenia: A Meta-analysis. *JAMA Psychiatry* Nov 1
2017;74(11):1104-1111.
63. Knolle F, Sterner E, Moutoussis M, et al. Action selection in early stages of
psychosis: an active inference approach. *J Psychiatry Neurosci* Jan-Feb
2023;48(1):E78-E89.
64. Omlor W, Cecere G, Huang GY, et al. Exploratory analysis of the relationship
between striatal connectivity and apathy during phosphodiesterase 10
inhibition in schizophrenia: findings from a randomized crossover trial. *BMC
Med* Mar 28 2025;23(1):187.
65. Winkelbeiner S, Leucht S, Kane JM, Homan P. Evaluation of Differences in
Individual Treatment Response in Schizophrenia Spectrum Disorders: A
Meta-analysis. *JAMA Psychiatry* Oct 1 2019;76(10):1063-1073.
66. Vrizzi S, Najar A, Lemogne C, Palminteri S, Lebreton M. Behavioral,
computational and self-reported measures of reward and punishment
sensitivity as predictors of mental health characteristics. *Nature Mental Health*
2025.

840   **Figures and Tables**

Figure 1: Experimental setting and modeling hypothesis. (a) *Two-stage decision-making task.* Adapted from Decker, Otto, Daw, Hartley [49]. Participants made two consecutive choices to either earn play coins (*reward context*) or avoid losing play coins (*loss context*). Probabilistic transitions between the first and second stages (common vs. rare) allowed for the quantification of model-based and model-free control. In stage 1, each spaceship was associated predominantly (70% probability, 'common' transitions) with one of the two alien pairs in stage 2, and less frequently (30% probability, 'rare' transitions) with the other pair. **(b)** Schematic of the human prefrontal network supporting model-based reinforcement learning. Dorsolateral prefrontal cortex (DLPFC) supports planning and working-memory processes; orbitofrontal cortex (OFC) encodes prospective option values; anterior cingulate cortex (ACC) integrates these signals to allocate cognitive control. **(c)** Schematic showing the computation of context biases in goal-directed control. For each participant, the model-based weight ω and the task-evoked BOLD parameter β were estimated separately in *reward* and *loss* contexts (grey bars). Context bias was defined as the difference $\Delta\omega = \omega_{reward} - \omega_{loss}$ for behavior and $\Delta\beta = \beta_{reward} - \beta_{loss}$ for neural activation. Positive Δ indicates a *reward bias* (black bar with arrow pointing up), whereas negative Δ indicates a *loss bias* (black bar with arrow pointing down).

887
Figure 2: Analysis of decision behavior: (a) *Simplified raw behavior scores:* Regarding the relative
contribution of model-based learning, healthy controls (HC blue) exhibited a numerically higher mean
contribution of model-based control in the reward relative to the loss context, whereas individuals with
schizophrenia spectrum disorders (SSD) showed a numerically higher mean contribution of model-
based control in the loss relative to the reward context. When the context bias of raw behavior scores
was compared between groups the difference was not significant. **(b)** *ω-scores from the modeling
analysis:* Regarding the relative contribution of model-based learning, the ω-scores indicated a
significant reward bias in HC and a significant loss bias in individuals with SSD. In addition, the
context bias of ω-values differed significantly between groups. Mean and 95% confidence intervals
are shown. Asterisks indicate *p* < 0.05.
898
899

915
916
Figure 3: Context-dependent BOLD activation during model-based control: (**a-b**). The context
bias of activation in the anterior cingulate cortex (ACC), dorsolateral prefrontal cortex (DLPFC) and
orbitofrontal cortex (OFC) for healthy controls (HC, blue) and individuals with schizophrenia spectrum
disorders (SSD, orange) is shown, plotted separately for the left and right hemispheres. Symbols
indicate group means and 95 % confidence intervals. Asterisks indicate p < 0.05. (**c-d**) Brain maps
displaying context-biased activation within ACC, DLPFC, and OFC masks, separately for the left (c)
and right hemisphere (d). Maps are presented unthresholded for visualization purposes.

927
928
Figure 4: Right ACC context bias and negative-symptom outcomes. (a) Baseline associations:
Correlations between the goal-directed context bias in the right anterior cingulate cortex (rACC) and
BNSS domains. Black marks indicate the apathy cluster (anhedonia, asociality, avolition); gray marks
indicate the diminished-expression cluster (blunted affect, alogia). Lines show 95% confidence
intervals. **(b)** Schematic of the longitudinal analysis: rACC context bias measured at baseline (fMRI)
was related to negative-symptom features derived from AMDP chart ratings at 3, 6, 9, and 12 months.
For each feature, we summarized the average follow-up severity across the available assessments; K
denotes the number of follow-up assessments for each participant. **(c–f)** Relationship between
baseline goal-directed context bias in the rACC and follow-up motivation, social engagement,
affective range, and affective flexibility. Points show follow-up scores adjusted for baseline and K, and
lines show linear fits. In apathy-related domains, a more loss-biased rACC signal (more negative
values on the x-axis) is associated with worse adjusted outcomes over the next year, with the
strongest association for motivation. Abbreviations: Anhed.: Anhedonia; Asoc: Asociality; Avol.:
Avolition; B.A.: Blunted affect; Alog.: Alogia. Asterisks indicate $P < 0.05$.

Characteristic	HC	SSD	p-value
Demographics			
Age (years), mean (SD)	28.00 (5.57)	27.14 (6.42)	0.481
Sex (% female)	20 (41.7)	11 (22.0)	0.061
Ethnicity, n (% Caucasian)	38 (79.2)	40 (80.0)	
% Asian	6 (12.5)	6 (12.0)	0.995
% African	4 (8.3)	4 (8.0)	
Educational attainment, mean (SD)	2.23 (1.29)	1.66 (1.17)	0.024
Illness Characteristics			
Disease Duration (months), mean (SD)	NA	18.26 (29.63)	
Hospitalizations, mean (SD)	NA	6.96 (4.98)	
Medication Status			
Dosage, antipsychotic (CPZ eq.)	NA	277.20 (160.93)	
Clinical Ratings			
BPRS Total Score, mean (SD)	NA	39.04 (10.43)	
Suspicion	NA	1.94 (1.10)	
Hallucination	NA	1.84 (1.13)	
BNSS Total Score, mean (SD)	NA	40.44 (15.37)	
Asociality	NA	6.06 (2.65)	
Avolition	NA	6.50 (2.76)	
Anhedonia	NA	10.68 (4.63)	
Distress	NA	3.32 (1.49)	
Blunted Affect	NA	9.50 (4.55)	
Alogia	NA	4.34 (2.48)	
Cognitive Performance			
MATRICS - Attention	NA	0.97 (1.06)	
MATRICS - Working Memory	NA	1.47 (0.86)	
MATRICS - Verbal Learning	NA	1.59 (1.13)	
MATRICS - Reasoning	NA	1.44 (1.05)	
MATRICS - Visual Learning	NA	1.50 (0.90)	

Table 1. Participant characteristics. Clinical symptoms were assessed using the Brief Psychiatric
Rating Scale (BPRS) for general psychopathology and the Brief Negative Symptom Scale (BNSS) for
negative symptoms. Cognitive performance was evaluated with the MATRICS Consensus Cognitive
Battery (MCCB), reported as standardized T-scores. Antipsychotic medication dosage is reported in
chlorpromazine equivalents (CPZ eq.). Data are presented as mean (standard deviation) or count
(percentage). Educational attainment is coded as follows: 0 = mandatory schooling, 1 = secondary
education (high school or equivalent), 2 = vocational training, 3 = higher professional education, 4 =
university degree. HC: Healthy controls. SSD: Schizophrenia spectrum disorders.

957